

## Lower bounds for index of Wente tori

*Dedicated to Katsuhiko Shiohama on the occasion of his sixtieth birthday*

Levi Lopes de LIMA, Vicente Francisco de SOUSA NETO and  
Wayne ROSSMAN

(Received February 29, 2000)

(Revised October 2, 2000)

**ABSTRACT.** We show that any of the constant mean curvature tori first found by Wente must have index at least eight. The proof uses a numerical computation by Mathematica.

### 1 Introduction

The Hopf conjecture asked if all closed surfaces immersed in  $\mathbf{R}^3$  with constant mean curvature  $H$  must be round spheres. It was proven true when either the surface has genus zero by Hopf himself [H], or the immersion is actually an embedding by Alexandrov [H]. However, it does not hold in general, and the first counterexamples, of genus 1, were found by Wente [We]. Abresch [A] and Walter [Wa] made more explicit descriptions for these surfaces of Wente, which all have one family of planar curvature lines [Sp]. We call these surfaces the *original Wente tori*.

Constant mean curvature surfaces are critical for area, but not necessarily area minimizing, for all compactly supported volume-preserving variations. Hence the index—loosely speaking, the *dimension* of area-reducing volume-preserving variations, to be defined in Section 3—can be positive. If it is zero, the surface is stable. Do Carmo and Peng [CP] showed that the only complete stable minimal surface is a plane. Fischer-Colbrie [FC] showed that a complete minimal surface in  $\mathbf{R}^3$  has finite index if and only if it has finite total curvature, and that the catenoid and Enneper's surface have index 1. Likewise, for surfaces with constant mean curvature  $H \neq 0$ , Barbosa and Do Carmo [BC] showed that only round spheres are stable, and Lopez and Ros [LR] and Silveira [Si] independently showed that they have finite index if and only if they are compact. This leaves open the question of whether there

---

2000 *Mathematical Subject Classification.* 53A10, 53A35

*Key words and phrases.* constant mean curvature surfaces, Wente tori, Morse index.

exist surfaces with constant mean curvature  $H \neq 0$  and low positive index, for example with index 1.

The third author [R1], [R2] showed numerically that the most natural candidates for unstable surfaces of constant mean curvature  $H \neq 0$  with low index—the original Wente tori—all have index at least 7, and with a numerical experiment suggested that the sharpest lower bound is either 8, 9, or 10, and is most likely 9. This leads one to conjecture that all closed surface with constant mean curvature  $H \neq 0$  have index at least 9.

The purpose of this article is to show that the original Wente tori all have index at least 8, improving the lower bound of [R1], [R2]. The final part of our argument relies on numerics.

## 2 The original Wente tori

In this section we shall give a brief description of the original Wente tori, based on [Wa]. Later, we shall assume that the mean curvature  $H$  is  $1/2$ , but in this and the next section we shall only assume that  $H$  is a nonzero constant.

Let  $\mathfrak{X} : \mathbf{C}/\Gamma \rightarrow \mathbf{R}^3$  be a conformal immersion of class  $C^\infty$  where  $\mathbf{C}/\Gamma$  is a compact 2-dimensional torus determined by the 2-dimensional lattice  $\Gamma$ . Note that  $(x, y)$  are then isothermal coordinates on  $\mathbf{C}/\Gamma$ . The fundamental forms and the Gauss and mean curvature functions are

$$I = E(dx^2 + dy^2), \quad II = L dx^2 + 2M dx dy + N dy^2, \quad K = \frac{LN - M^2}{E^2}, \quad H = \frac{L + N}{2E}.$$

Since  $H$  is constant, the Hopf differential  $\Phi dz^2$  is holomorphic, where  $\Phi = \frac{1}{2}(L - N) - iM$  and  $z = x + iy$ . Thus  $\Phi$  is constant and  $\mathfrak{X}$  has no umbilics points. Moreover, by a change of the coordinates  $(x, y)$ , we may assume  $\Phi = 1$  and so  $M = 0$ ,  $L = e^F + 1$ ,  $N = e^F - 1$ , and  $(x, y)$  become curvature line parameters, where  $F : \mathbf{C}/\Gamma \rightarrow \mathbf{R}$  is defined by  $HE = e^F$ . We have the equations of Gauss and Weingarten:

$$\begin{aligned} \mathfrak{X}_{xx} &= \frac{1}{2}F_x \mathfrak{X}_x - \frac{1}{2}F_y \mathfrak{X}_y - (e^F + 1)\mathcal{N}, \\ \mathfrak{X}_{yy} &= -\frac{1}{2}F_x \mathfrak{X}_x + \frac{1}{2}F_y \mathfrak{X}_y - (e^F - 1)\mathcal{N}, \\ \mathfrak{X}_{xy} &= \frac{1}{2}F_y \mathfrak{X}_x + \frac{1}{2}F_x \mathfrak{X}_y, \\ \mathcal{N}_x &= H(1 + e^{-F})\mathfrak{X}_x, \quad \mathcal{N}_y = H(1 - e^{-F})\mathfrak{X}_y \end{aligned} \tag{1}$$

$$\Delta F + 4H \sinh F = 0, \tag{2}$$

where  $\mathcal{A} = \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2}$  and  $\mathcal{N} : \mathbf{C}/\Gamma \rightarrow R^3$  is the unit normal vector field, i.e. the Gauss map. Therefore the problem of finding constant mean curvature immersed tori in  $\mathbf{R}^3$  reduces to solving the PDE system (1) and (2) by real analytic functions  $F, \mathcal{N}, \mathfrak{X}$  defined on  $\mathbf{R}^2$  and doubly periodic with respect to some fundamental lattice  $\Gamma \subset \mathbf{R}^2$ .

In the case of the original Wente tori, in Walter's notation, the solution  $F$  of (2) is:

$$(3) \quad \tanh\left(\frac{F}{4}\right) = \gamma \cdot \bar{\gamma} \cdot cn_k(\alpha x) cn_{\bar{k}}(\bar{\alpha} y),$$

where  $cn_k$  denotes the Jacobi amplitudinus cosinus function with modulus  $k$ , and  $k = \sin \theta$ ,  $\bar{k} = \sin \bar{\theta}$ , for  $\theta, \bar{\theta} \in (0, \pi/2)$  and  $\theta + \bar{\theta} < \pi/2$ , and

$$\gamma = \sqrt{\tan \theta}, \quad \bar{\gamma} = \sqrt{\tan \bar{\theta}}, \quad \alpha = \sqrt{4H \frac{\sin 2\bar{\theta}}{\sin 2(\theta + \bar{\theta})}}, \quad \bar{\alpha} = \sqrt{4H \frac{\sin 2\theta}{\sin 2(\theta + \bar{\theta})}}.$$

LEMMA 1 ([A], [Wa]). *The set of all original Wente tori are in a one-to-one correspondence with the set of reduced fractions  $\ell/n \in (1, 2)$ .*

For each  $\ell/n$ , we call the corresponding Wente torus  $\mathcal{W}_{\ell/n}$ . Following Walter's notation, each  $\mathcal{W}_{\ell/n}$  has either one or two planar geodesic loops in the central symmetry plane: two loops if  $\ell$  is odd, and one loop if  $\ell$  is even. Each loop can be partitioned into  $2n$  congruent curve segments, and  $\ell$  is the total winding order of the Gauss map along each loop.

The conditions for double periodicity of the position vector function  $\mathfrak{X}$  are expressed in terms of  $\theta$  and  $\bar{\theta}$ . Walter determined that there is exactly one

$$\bar{\theta} \cong 65.354955^\circ$$

that solves one period problem. The other period problem is solved with the correct choice of  $\theta \in (0, (\pi/2) - \bar{\theta})$ , and, for any  $\ell/n \in (1, 2)$ , this correct choice is the unique solution  $\theta$  of

$$(4) \quad \int_0^{\pi/2} \frac{1 + \tan \theta \tan \bar{\theta} \cos^2 \varphi}{1 - \tan \theta \tan \bar{\theta} \cos^2 \varphi} \frac{d\varphi}{\sqrt{1 - \sin^2 \theta \sin^2 \varphi}} = \frac{\ell}{n} \frac{\pi}{2} \sqrt{\frac{\sin 2\bar{\theta}}{\sin 2(\theta + \bar{\theta})}}.$$

For any  $\ell/n \notin (1, 2)$ , there is no solution  $\theta \in (0, (\pi/2) - \bar{\theta})$  of (4). In Table 2 we give some values of  $\theta$  with respect to  $\ell/n$ .

Now, if  $x_{\ell/n}$  (resp.  $y_{\ell/n}$ ) denotes the length of the period of  $cn_k(\alpha x)$  (resp.  $cn_{\bar{k}}(\bar{\alpha} y)$ ), then we have the following lemma:

LEMMA 2 ([Wa]).  $\mathfrak{X} : \mathbf{C}/\Gamma \rightarrow W_{\ell/n} \subset \mathbf{R}^3$  is a conformal immersion ( $W_{\ell/n}$  denotes the image of  $\mathfrak{X}$ ), where

$$\Gamma = \text{span}_{\mathbf{Z}}\{(nx_{\ell n}, 0), (0, y_{\ell n})\} \quad \text{when } \ell \text{ is odd, and}$$

$$\Gamma = \text{span}_{\mathbf{Z}}\left\{\left(\frac{nx_{\ell n}}{2}, \frac{y_{\ell n}}{2}\right), (0, y_{\ell n})\right\} \quad \text{when } \ell \text{ is even.}$$

The curves  $\{[x_0, y] \mid x_0 = \text{constant}\}$  are mapped by  $\mathfrak{X}$  to planar curvature lines of  $W_{\ell/n}$ .

The lengths  $x_{\ell n}$  and  $y_{\ell n}$  can be computed as follows:

$$(5) \quad x_{\ell n} = \frac{4}{\alpha} \int_0^{\pi/2} \frac{d\varphi}{\sqrt{1 - k^2 \sin^2 \varphi}}, \quad y_{\ell n} = \frac{4}{\bar{\alpha}} \int_0^{\pi/2} \frac{d\varphi}{\sqrt{1 - \bar{k}^2 \sin^2 \varphi}}.$$

### 3 The definition of index and preliminary results

The Jacobi operator associated to  $\mathcal{W}_{\ell/n}$  is  $-\Delta_I - |II|^2$  on  $\mathbf{C}/\Gamma$ , with  $|II|^2 = E^{-2}(L^2 + 2M^2 + N^2) = 2H^2(1 + e^{-2F})$  and  $\Delta_I$  the Laplace-Beltrami operator associated to the metric  $I$ . The corresponding quadratic form is

$$(6) \quad Q(u, u) = \int_{\mathbf{C}/\Gamma} u \mathcal{L}(u) dx dy,$$

where

$$\mathcal{L}u = -\Delta u - Vu \quad \text{with} \quad V = 4H \cosh(F)$$

and  $\Delta$  the Euclidean Laplacian. Note that in equation (6), we are integrating with respect to the flat metric on  $\mathbf{C}/\Gamma$ .

Consider a smooth volume-preserving variation  $\mathfrak{X}_t$  of the immersion  $\mathfrak{X}$  with parameter  $t$  so that  $\mathfrak{X}_0$  is the surface  $\mathcal{W}_{\ell/n}$ . By reparametrizing the surfaces of the variation, we may assume that the variation vector field at  $t = 0$  is  $u\mathcal{N}$  for some  $u \in C^\infty(\mathbf{C}/\Gamma)$ . Then

$$\frac{\partial}{\partial t} \text{area}(\mathfrak{X}_t)|_{t=0} = 0 \quad \text{and} \quad \frac{\partial^2}{\partial t^2} \text{area}(\mathfrak{X}_t)|_{t=0} = Q(u, u).$$

Furthermore, the volume-preserving condition implies  $\int_{\mathbf{C}/\Gamma} u dA = 0$ . Thus, if

$$\mathcal{V} = \left\{ u \in C^\infty(\mathbf{C}/\Gamma); \int_{\mathbf{C}/\Gamma} u dA = 0 \right\},$$

then we can give the following definition (see [BC]):

DEFINITION 1. We define  $\text{Ind}(\mathfrak{X}(\mathbf{C}/\Gamma))$ , the *index* of the immersion  $\mathfrak{X}$  of  $\mathbf{C}/\Gamma$ , to be the maximum of the dimensions of the subspaces of  $\mathcal{V}$  restricted to which  $Q$  is negative definite.

Since the first derivative of area is zero, and the second derivative is  $Q(u, u)$ , the index in a sense measures the amount of area-reducing volume-preserving variations.

Let  $L^2 = L^2(\mathbf{C}/\Gamma) = \{u \in C^\infty(\mathbf{C}/\Gamma) \mid \int_{\mathbf{C}/\Gamma} u^2 dx dy < \infty\}$  provided with the inner product  $\langle u, v \rangle_{L^2} = \int_{\mathbf{C}/\Gamma} uv dx dy$ . It follows from the standard spectral theorem that the operator  $\mathcal{L} = -\Delta - V$  on  $\mathbf{C}/\Gamma$  has a discrete spectrum of eigenvalues

$$\beta_1 < \beta_2 \leq \dots \nearrow \infty$$

and corresponding eigenfunctions

$$v_1, v_2 \dots \in C^\infty(\mathbf{C}/\Gamma),$$

which form an orthonormal basis for  $L^2$ . Moreover, we have the following variational characterization for the eigenvalues:

$$\beta_j = \inf_{V_j} \left( \sup_{\phi \in V_j, \|\phi\|_{L^2}=1} \int_{\mathbf{C}/\Gamma} \phi \mathcal{L} \phi dx dy \right),$$

where  $V_j$  runs through all  $j$  dimensional subspaces of  $C^\infty(\mathbf{C}/\Gamma)$ .

LEMMA 3 ([R1], [R2]). *If  $\mathcal{L}$  has  $k$  negative eigenvalues, then  $\text{Ind}(W_{\ell/n})$  is either  $k$  or  $k - 1$ . Furthermore, if there exists a subspace  $\mathcal{S} \subset L^2$  such that  $\mathcal{S} \subset C^\infty(\mathbf{C}/\Gamma)$  and  $\dim(\mathcal{S}) = k$  and  $Q$  restricted to  $\mathcal{S}$  is negative definite, then  $\text{Ind}(W_{\ell/n}) \geq k - 1$ .*

By Lemma 3, our goal becomes to compute the number of negative eigenvalues of  $\mathcal{L}$ .

Now, we use a convenient fact: For the flat torus  $\mathbf{C}/\Gamma$ , with  $\Gamma = \text{span}_{\mathbf{Z}}\{(a_1, a_2), (b_1, b_2)\}$ , the complete set of eigenvalues of  $-\Delta u_i = \alpha_i u_i$  are

$$\frac{4\pi^2}{(a_1 b_2 - a_2 b_1)^2} ((m_2 b_2 - m_1 a_2)^2 + (m_1 a_1 - m_2 b_1)^2),$$

with corresponding orthonormal eigenfunctions

$$c_{m_1, m_2} \cdot (\sin \text{ or } \cos) \left( \frac{2\pi}{a_1 b_2 - a_2 b_1} ((m_2 b_2 - m_1 a_2)x + (m_1 a_1 - m_2 b_1)y) \right),$$

where  $m_1, m_2 \in \mathbf{Z}$ ,  $c_{m_1, m_2} = \sqrt{2/(a_1 b_2 - a_2 b_1)}$  if  $|m_1| + |m_2| > 0$ ,  $c_{0,0} = \sqrt{1/(a_1 b_2 - a_2 b_1)}$ . With the aid of Lemma 2 we list 17 of the  $\alpha_i$  and  $u_i$  in Table 1.

**Table 1:** The first 17 eigenvalues and eigenfunctions of  $-\Delta u_i = \alpha_i u_i$ .

| eigenvalues $\alpha_i$<br>for $\ell$ odd                                      | eigenfunctions $u_i$<br>for $\ell$ odd   | eigenvalues $\alpha_i$<br>for $\ell$ even                                      | eigenfunctions $u_i$<br>for $\ell$ even  |
|---|--|--|--|
| $\alpha_1 = 0$  | $u_1 = \frac{1}{\sqrt{nx_{\ell n} y_{\ell n}}}$  | $\alpha_1 = 0$   | $u_1 = \frac{\sqrt{2}}{\sqrt{nx_{\ell n} y_{\ell n}}}$   |
| $\alpha_2 = \frac{4\pi^2}{n^2 x_{\ell n}^2}$                                  | $u_2 = \frac{\sin(2\pi x / (nx_{\ell n}))}{\sqrt{nx_{\ell n} y_{\ell n} / 2}}$   | $\alpha_2 = \frac{16\pi^2}{n^2 x_{\ell n}^2}$                                  | $u_2 = \frac{\sin(4\pi x / (nx_{\ell n}))}{\sqrt{nx_{\ell n} y_{\ell n} / 4}}$   |
| $\alpha_3 = \frac{4\pi^2}{n^2 x_{\ell n}^2}$                                  | $u_3 = \frac{\cos(2\pi x / (nx_{\ell n}))}{\sqrt{nx_{\ell n} y_{\ell n} / 2}}$   | $\alpha_3 = \frac{16\pi^2}{n^2 x_{\ell n}^2}$                                  | $u_3 = \frac{\cos(4\pi x / (nx_{\ell n}))}{\sqrt{nx_{\ell n} y_{\ell n} / 4}}$   |
| $\alpha_4 = \frac{4\pi^2}{y_{\ell n}^2}$                                      | $u_4 = \frac{\sin(2\pi y / y_{\ell n})}{\sqrt{nx_{\ell n} y_{\ell n} / 2}}$  | $\alpha_4 = \frac{4\pi^2}{n^2 x_{\ell n}^2} + \frac{4\pi^2}{y_{\ell n}^2}$     | $u_4 = \frac{\sin\left(\frac{2\pi x}{nx_{\ell n}} + \frac{2\pi y}{y_{\ell n}}\right)}{\sqrt{nx_{\ell n} y_{\ell n} / 4}}$    |
| $\alpha_5 = \frac{4\pi^2}{y_{\ell n}^2}$                                      | $u_5 = \frac{\cos(2\pi y / y_{\ell n})}{\sqrt{nx_{\ell n} y_{\ell n} / 2}}$  | $\alpha_5 = \frac{4\pi^2}{n^2 x_{\ell n}^2} + \frac{4\pi^2}{y_{\ell n}^2}$     | $u_5 = \frac{\cos\left(\frac{2\pi x}{nx_{\ell n}} + \frac{2\pi y}{y_{\ell n}}\right)}{\sqrt{nx_{\ell n} y_{\ell n} / 4}}$    |
| $\alpha_6 = \frac{16\pi^2}{n^2 x_{\ell n}^2}$                                 | $u_6 = \frac{\sin(4\pi x / (nx_{\ell n}))}{\sqrt{nx_{\ell n} y_{\ell n} / 2}}$   | $\alpha_6 = \frac{4\pi^2}{n^2 x_{\ell n}^2} + \frac{4\pi^2}{y_{\ell n}^2}$     | $u_6 = \frac{\sin\left(\frac{2\pi x}{nx_{\ell n}} - \frac{2\pi y}{y_{\ell n}}\right)}{\sqrt{nx_{\ell n} y_{\ell n} / 4}}$    |
| $\alpha_7 = \frac{16\pi^2}{n^2 x_{\ell n}^2}$                                 | $u_7 = \frac{\cos(4\pi x / (nx_{\ell n}))}{\sqrt{nx_{\ell n} y_{\ell n} / 2}}$   | $\alpha_7 = \frac{4\pi^2}{n^2 x_{\ell n}^2} + \frac{4\pi^2}{y_{\ell n}^2}$     | $u_7 = \frac{\cos\left(\frac{2\pi x}{nx_{\ell n}} - \frac{2\pi y}{y_{\ell n}}\right)}{\sqrt{nx_{\ell n} y_{\ell n} / 4}}$    |
| $\alpha_8 = \frac{4\pi^2}{n^2 x_{\ell n}^2} + \frac{4\pi^2}{y_{\ell n}^2}$    | $u_8 = \frac{\sin\left(\frac{2\pi x}{nx_{\ell n}} + \frac{2\pi y}{y_{\ell n}}\right)}{\sqrt{nx_{\ell n} y_{\ell n} / 2}}$    | $\alpha_8 = \frac{16\pi^2}{y_{\ell n}^2}$                                      | $u_8 = \frac{\sin(4\pi y / y_{\ell n})}{\sqrt{nx_{\ell n} y_{\ell n} / 4}}$  |
| $\alpha_9 = \frac{4\pi^2}{n^2 x_{\ell n}^2} + \frac{4\pi^2}{y_{\ell n}^2}$    | $u_9 = \frac{\cos\left(\frac{2\pi x}{nx_{\ell n}} + \frac{2\pi y}{y_{\ell n}}\right)}{\sqrt{nx_{\ell n} y_{\ell n} / 2}}$    | $\alpha_9 = \frac{16\pi^2}{y_{\ell n}^2}$                                      | $u_9 = \frac{\cos(4\pi y / y_{\ell n})}{\sqrt{nx_{\ell n} y_{\ell n} / 4}}$  |
| $\alpha_{10} = \frac{4\pi^2}{n^2 x_{\ell n}^2} + \frac{4\pi^2}{y_{\ell n}^2}$ | $u_{10} = \frac{\sin\left(\frac{2\pi x}{nx_{\ell n}} - \frac{2\pi y}{y_{\ell n}}\right)}{\sqrt{nx_{\ell n} y_{\ell n} / 2}}$ | $\alpha_{10} = \frac{64\pi^2}{n^2 x_{\ell n}^2}$                               | $u_{10} = \frac{\sin(8\pi x / (nx_{\ell n}))}{\sqrt{nx_{\ell n} y_{\ell n} / 4}}$  |
| $\alpha_{11} = \frac{4\pi^2}{n^2 x_{\ell n}^2} + \frac{4\pi^2}{y_{\ell n}^2}$ | $u_{11} = \frac{\cos\left(\frac{2\pi x}{nx_{\ell n}} - \frac{2\pi y}{y_{\ell n}}\right)}{\sqrt{nx_{\ell n} y_{\ell n} / 2}}$ | $\alpha_{11} = \frac{64\pi^2}{n^2 x_{\ell n}^2}$                               | $u_{11} = \frac{\cos(8\pi x / (nx_{\ell n}))}{\sqrt{nx_{\ell n} y_{\ell n} / 4}}$  |
| $\alpha_{12} = \frac{16\pi^2}{y_{\ell n}^2}$                                  | $u_{12} = \frac{\sin(4\pi y / y_{\ell n})}{\sqrt{nx_{\ell n} y_{\ell n} / 2}}$   | $\alpha_{12} = \frac{36\pi^2}{n^2 x_{\ell n}^2} + \frac{4\pi^2}{y_{\ell n}^2}$ | $u_{12} = \frac{\sin\left(\frac{6\pi x}{nx_{\ell n}} + \frac{2\pi y}{y_{\ell n}}\right)}{\sqrt{nx_{\ell n} y_{\ell n} / 4}}$ |
| $\alpha_{13} = \frac{16\pi^2}{y_{\ell n}^2}$                                  | $u_{13} = \frac{\cos(4\pi y / y_{\ell n})}{\sqrt{nx_{\ell n} y_{\ell n} / 2}}$   | $\alpha_{13} = \frac{36\pi^2}{n^2 x_{\ell n}^2} + \frac{4\pi^2}{y_{\ell n}^2}$ | $u_{13} = \frac{\cos\left(\frac{6\pi x}{nx_{\ell n}} + \frac{2\pi y}{y_{\ell n}}\right)}{\sqrt{nx_{\ell n} y_{\ell n} / 4}}$ |

**Table 1:** (Continued)

| eigenvalues $\alpha_i$<br>for $\ell$ odd                                      | eigenfunctions $u_i$<br>for $\ell$ odd  | eigenvalues $\alpha_i$<br>for $\ell$ even                                      | eigenfunctions $u_i$<br>for $\ell$ even   |
|---|---|--|---|
| $\alpha_{14} = \frac{36\pi^2}{n^2x_{\ell n}^2}$                               | $u_{14} = \frac{\sin(6\pi x/nx_{\ell n})}{\sqrt{nx_{\ell n}y_{\ell n}/2}}$  | $\alpha_{14} = \frac{36\pi^2}{n^2x_{\ell n}^2} + \frac{4\pi^2}{y_{\ell n}^2}$  | $u_{14} = \frac{\sin\left(\frac{6\pi x}{nx_{\ell n}} + \frac{2\pi y}{y_{\ell n}}\right)}{\sqrt{nx_{\ell n}y_{\ell n}/4}}$ |
| $\alpha_{15} = \frac{36\pi^2}{n^2x_{\ell n}^2}$                               | $u_{15} = \frac{\cos(6\pi x/nx_{\ell n})}{\sqrt{nx_{\ell n}y_{\ell n}/2}}$  | $\alpha_{15} = \frac{36\pi^2}{n^2x_{\ell n}^2} + \frac{4\pi^2}{y_{\ell n}^2}$  | $u_{15} = \frac{\cos\left(\frac{6\pi x}{nx_{\ell n}} + \frac{2\pi y}{y_{\ell n}}\right)}{\sqrt{nx_{\ell n}y_{\ell n}/4}}$ |
| $\alpha_{16} = \frac{16\pi^2}{n^2x_{\ell n}^2} + \frac{4\pi^2}{y_{\ell n}^2}$ | $u_{16} = \frac{\sin\left(\frac{4\pi x}{nx_{\ell n}} + \frac{2\pi y}{y_{\ell n}}\right)}{\sqrt{nx_{\ell n}y_{\ell n}/2}}$ | $\alpha_{16} = \frac{16\pi^2}{n^2x_{\ell n}^2} + \frac{16\pi^2}{y_{\ell n}^2}$ | $u_{16} = \frac{\sin\left(\frac{4\pi x}{nx_{\ell n}} + \frac{4\pi y}{y_{\ell n}}\right)}{\sqrt{nx_{\ell n}y_{\ell n}/4}}$ |
| $\alpha_{17} = \frac{16\pi^2}{n^2x_{\ell n}^2} + \frac{4\pi^2}{y_{\ell n}^2}$ | $u_{17} = \frac{\cos\left(\frac{4\pi x}{nx_{\ell n}} + \frac{2\pi y}{y_{\ell n}}\right)}{\sqrt{nx_{\ell n}y_{\ell n}/2}}$ | $\alpha_{17} = \frac{16\pi^2}{n^2x_{\ell n}^2} + \frac{16\pi^2}{y_{\ell n}^2}$ | $u_{17} = \frac{\cos\left(\frac{4\pi x}{nx_{\ell n}} + \frac{4\pi y}{y_{\ell n}}\right)}{\sqrt{nx_{\ell n}y_{\ell n}/4}}$ |

With the orderings for the eigenvalues as chosen in Table 1, we do not necessarily have  $\alpha_i \leq \alpha_j$  for  $i \leq j$ . However, we still have  $\alpha_i \nearrow \infty$  as  $i \nearrow \infty$ . Choose  $\alpha_{\rho_{\ell/n}(1)}, \alpha_{\rho_{\ell/n}(2)}, \dots$  the complete set of eigenvalues with multiplicity 1 of the operator  $-\Delta$  on the flat torus  $\mathbf{C}/\Gamma$  reordered by the permutation  $\rho_{\ell/n}$  of  $\mathbf{N}$  so that  $\alpha_{\rho_{\ell/n}(1)} < \alpha_{\rho_{\ell/n}(2)} \leq \dots \nearrow \infty$ .

The first of the following two lemmas follows from the variational characterization for eigenvalues, and the second follows from Lemma 3, the Courant nodal domain theorem, and geometric properties of the surfaces  $W_{\ell/n}$ :

LEMMA 4 ([R1], [R2]). *Choose  $\mu \in \mathbf{Z}^+$  so that  $\alpha_{\rho_{\ell/n}(\mu)} < 4H$ . Then  $\text{Ind}(W_{\ell/n}) \geq \mu - 1$ .*

LEMMA 5 ([R1], [R2]). *For all  $n \in \mathbf{Z}^+$ ,  $n \geq 2$  we have that  $\text{Ind}(W_{\ell/n}) \geq 2n - 2$  if  $\ell$  is odd, and  $\text{Ind}(W_{\ell/n}) \geq n - 2$  if  $\ell$  is even.*

#### 4 The lower bound 8 for $\text{Ind}(W_{\ell/n})$

We now show the following:

**Numerical Result:**  $\text{Ind}(W_{\ell/n}) \geq 8$  for all  $\ell/n$ .

Observe that, although the eigenvalues of  $\mathcal{L}$  depend on the choice of  $H$ , the number of negative eigenvalues is independent of  $H$ . So without loss of generality we fix  $H = 1/2$ .

**Table 2:**  $x_{\ell/n}, y_{\ell/n}$  are computed using the value  $H = 1/2$ .

| $W_{\ell/n}$ | $\theta$ | $x_{\ell/n}$ | $y_{\ell/n}$ | Lemma 4<br>lower bound<br>for $\text{Ind}(W_{\ell/n})$ | Lemma 5<br>lower bound<br>for $\text{Ind}(W_{\ell/n})$ |
|--------------|----------|--------------|--------------|--|--|
| $W_{3/2}$    | 17.7324° | 2.5556       | 4.2131       | 2  | 2  |
| $W_{4/3}$    | 12.7898° | 3.2767       | 6.3355       | 6  | 1  |
| $W_{5/3}$    | 21.4807° | 1.7557       | 2.6402       | 2  | 4  |
| $W_{7/4}$    | 22.8449° | 1.3315       | 1.9447       | 2  | 6  |
| $W_{8/5}$    | 20.1374° | 2.0842       | 3.2321       | 2  | 3  |
| $W_{12/7}$   | 22.3044° | 1.5150       | 2.2380       | 2  | 5  |
| $W_{14/9}$   | 19.1243° | 2.2970       | 3.6514       | 4  | 7  |
| $W_{16/9}$   | 23.2182° | 1.1872       | 1.7208       | 2  | 7  |

By Lemma 5,  $\text{Ind}(W_{\ell/n})$  can be less than 8 only if  $\ell/n$  is one of  $3/2, 4/3, 5/3, 5/4, 7/4, 6/5, 8/5, 8/7, 10/7, 12/7, 10/9, 14/9, 16/9$ . Lemma 4 also gives explicit lower bounds for the index, since we know the values of  $x_{\ell/n}$  and  $y_{\ell/n}$  numerically by formula (5), and hence we know the  $\alpha_{\rho_{\ell/n}(i)}$  (see Table 1). Lemma 4 implies that the index is at least 8 when  $\ell/n$  is  $5/4, 6/5, 8/7, 10/7, 10/9$ . Thus we only need to consider the following eight surfaces:

$$W_{3/2}, W_{4/3}, W_{5/3}, W_{7/4}, W_{8/5}, W_{12/7}, W_{14/9}, \text{ and } W_{16/9}.$$

For these surfaces we list, in Table 2, the corresponding  $\theta, x_{\ell/n}, y_{\ell/n}$  and lower bounds for index. These approximate values for  $\theta, x_{\ell/n}$ , and  $y_{\ell/n}$  were computed numerically using formulas (4) and (5) and the software Mathematica. Recall that always  $\bar{\theta} \cong 65.354^\circ$ .

We will find specific spaces on which  $\mathcal{L}$  is negative definite, for these eight surfaces.

Let  $N$  be an arbitrary positive integer. Consider a finite subset  $\{\tilde{u}_1 = u_{i_1}, \dots, \tilde{u}_N = u_{i_N}\}$  of the eigenfunctions  $u_i$  of  $-\mathcal{A}$  on  $\mathbf{C}/\Gamma$ , defined in Section 3, with corresponding eigenvalues  $\tilde{\alpha}_j = \alpha_{i_j}, j = 1, \dots, N$ . If we consider any  $u = \sum_{i=1}^N a_i \tilde{u}_i \in \text{span}\{\tilde{u}_1, \dots, \tilde{u}_N\}, a_1, \dots, a_N \in \mathbf{R}$ , then  $\int_{\mathbf{C}/\Gamma} u \mathcal{L}(u) dx dy = \sum_{i,j=1}^N a_i \cdot a_j (\tilde{\alpha}_j \delta_{ij} - \tilde{b}_{ij}) a_j$ , where  $\tilde{b}_{ij} := \int_{\mathbf{C}/\Gamma} V \tilde{u}_i \tilde{u}_j dx dy$ . So we have  $\int_{\mathbf{C}/\Gamma} u \mathcal{L}(u) dx dy < 0$  for all nonzero  $u \in \text{span}\{\tilde{u}_1, \dots, \tilde{u}_N\}$  if and only if the matrix  $(\tilde{\alpha}_j \delta_{ij} - \tilde{b}_{ij})_{i,j=1,\dots,N}$  is negative definite. Lemma 3 then implies:

**THEOREM 1** ([R1], [R2]). *If the  $N \times N$  matrix  $(\tilde{\alpha}_j \delta_{ij} - \tilde{b}_{ij})_{i,j=1,\dots,N}$  is negative definite, then  $\text{Ind}(W_{\ell/n}) \geq N - 1$ .*

DEFINITION 2. Given  $A, B$  even integers and  $\ell, n \in \mathbf{Z}^+$ , we now define the following basic integrals:

$$\begin{aligned}
I_0(\ell, n, A, B) &= \frac{1}{nx_{\ell n}y_{\ell n}} \int_0^{x_{\ell n}/4} \int_0^{y_{\ell n}/4} V \left( \cos \frac{2\pi x}{x_{\ell n}} \right)^A \left( \cos \frac{2\pi y}{y_{\ell n}} \right)^B dydx, \\
I_1(\ell, n) &= \frac{8}{nx_{\ell n}y_{\ell n}} \int_0^{nx_{\ell n}/4} \int_0^{y_{\ell n}/4} V \cos \left( \frac{4\pi x}{nx_{\ell n}} \right) dydx, \\
I_2(\ell, n) &= \frac{8}{nx_{\ell n}y_{\ell n}} \int_0^{nx_{\ell n}/4} \int_0^{y_{\ell n}/4} V \cos \left( \frac{8\pi x}{nx_{\ell n}} \right) dydx, \\
I_3(\ell, n) &= \frac{8}{nx_{\ell n}y_{\ell n}} \int_0^{nx_{\ell n}/4} \int_0^{y_{\ell n}/4} V \cos \left( \frac{16\pi x}{nx_{\ell n}y_{\ell n}} \right) dydx, \\
I_4(\ell, n) &= \frac{8}{nx_{\ell n}y_{\ell n}} \int_0^{nx_{\ell n}/4} \int_0^{y_{\ell n}/4} V \cos \left( \frac{4\pi x}{nx_{\ell n}} \right) \cos \left( \frac{4\pi y}{y_{\ell n}} \right) dydx, \\
I_5(\ell, n) &= \frac{8}{nx_{\ell n}y_{\ell n}} \int_0^{nx_{\ell n}/4} \int_0^{y_{\ell n}/4} V \cos \left( \frac{4\pi x}{nx_{\ell n}} \right) \cos \left( \frac{8\pi x}{nx_{\ell n}} \right) dydx, \\
I_6(\ell, n) &= \frac{8}{nx_{\ell n}y_{\ell n}} \int_0^{nx_{\ell n}/4} \int_0^{y_{\ell n}/4} V \cos \left( \frac{8\pi x}{nx_{\ell n}} \right) \cos \left( \frac{4\pi y}{y_{\ell n}} \right) dydx, \\
I_7(\ell, n) &= \frac{8}{nx_{\ell n}y_{\ell n}} \int_0^{nx_{\ell n}/4} \int_0^{y_{\ell n}/4} V \cos \left( \frac{12\pi x}{nx_{\ell n}} \right) \cos \left( \frac{4\pi y}{y_{\ell n}} \right) dydx.
\end{aligned}$$

Now, for each surface  $W_{\ell/n}$  given in Table 2, we will fix  $N = 9$  and choose the subset  $\{\tilde{u}_1, \dots, \tilde{u}_9\}$  such that the matrix  $(\tilde{\alpha}_j \delta_{ij} - \tilde{b}_{ij})_{i,j=1,\dots,9}$  is negative definite. These choices are given in Table 3. With these choices for  $\tilde{u}_i$ , we have the following lemma:

LEMMA 6. *With the choices given in Table 3, all elements of the eight matrices  $\mathcal{M}(\ell, n) := (\tilde{\alpha}_j \delta_{ij} - \tilde{b}_{ij})_{i,j=1,\dots,9}$  can be expressed in terms of the basic integrals  $I_0(\ell, n, A, B)$  and  $I_j(\ell, n)$  for  $A, B$  even and  $j = 1, 2, \dots, 7$ .*

PROOF. The symmetries  $V(x, y) = V(-x, y) = V(x, -y) = V\left(\frac{x_{\ell n}}{2} - x, y\right) = V\left(x, \frac{y_{\ell n}}{2} - y\right)$  of  $V$  and the identities  $\cos(a \pm b) = \cos(a) \cos(b) \mp \sin(a) \sin(b)$ ,  $\sin(a \pm b) = \sin(a) \cos(b) \pm \sin(b) \cos(a)$  give the relations shown in Table 4, proving the lemma.

By numerical methods, we can estimate that all of the relevant  $I_j(\ell, n)$  for  $j \geq 1$  are approximately zero, and that



**Table 4:** Elements  $M_{i,j}$  of the symmetric matrices  $\mathcal{M}(\ell, n)$  expressed in terms of the basic integrals. We have chosen here to index the  $M_{i,j}$  using the counters associated to  $\alpha_j$  and  $u_j$ , rather than  $\tilde{\alpha}_j$  and  $\tilde{u}_j$ .

|   |   |
|---|---|
| For $W_{3/2}$   | $M_{1,1} = -32I_0(3, 2, 0, 0)$ , $M_{4,4} = \alpha_4 - 64(I_0(3, 2, 0, 0) - I_0(3, 2, 0, 2))$<br>$M_{5,5} = \alpha_5 - 64I_0(3, 2, 0, 2)$ , $M_{7,7} = \alpha_7 - 64I_0(3, 2, 2, 0)$<br>$M_{17,17} = \alpha_{17} - 64(I_0(3, 2, 0, 0) - I_0(3, 2, 0, 2) - I_0(3, 2, 2, 0) + 2I_0(3, 2, 2, 2))$<br>$M_{i,i} = \alpha_i - 32I_0(3, 2, 0, 0)$ , for $i = 2, 3, 8, 9$<br>$M_{i,j} = 0$ for $i < j$ with $i, j \in \{1, 2, 3, 4, 5, 7, 8, 9, 17\}$ .   |
| For $W_{4/3}$   | $M_{1,1} = -48I_0(4, 3, 0, 0)$ , $M_{2,2} = \alpha_2 - 48I_0(4, 3, 0, 0) + 2I_2(4, 3)$<br>$M_{3,3} = \alpha_3 - 48I_0(4, 3, 0, 0) - 2I_2(4, 3)$ , $M_{4,4} = \alpha_4 - 48I_0(4, 3, 0, 0) + 2I_4(4, 3)$<br>$M_{5,5} = \alpha_5 - 48I_0(4, 3, 0, 0) - 2I_4(4, 3)$ , $M_{6,6} = \alpha_6 - 48I_0(4, 3, 0, 0) + 2I_4(4, 3)$<br>$M_{7,7} = \alpha_7 - 48I_0(4, 3, 0, 0) - 2I_4(4, 3)$<br>$M_{8,8} = \alpha_8 - 384(I_0(4, 3, 0, 2) - I_0(4, 3, 0, 4))$<br>$M_{9,9} = \alpha_9 - 96(4I_0(4, 3, 0, 4) - 4I_0(4, 3, 0, 2) + I_0(4, 3, 0, 0))$<br>$M_{1,3} = -2\sqrt{2}I_1(4, 3)$ , $M_{1,9} = -48\sqrt{2}(-I_0(4, 3, 0, 0) + 2I_0(4, 3, 0, 2))$<br>$M_{4,6} = -48(-I_0(4, 3, 0, 0) + 2I_0(4, 3, 0, 2)) + 2I_1(4, 3)$<br>$M_{5,7} = -48(-I_0(4, 3, 0, 0) + 2I_0(4, 3, 0, 2)) - 2I_1(4, 3)$<br>$M_{3,9} = -4I_4(4, 3)$ , all other $M_{i,j}$ with $i < j \leq 9$ are zero.                                 |
| For $W_{5/3}$   | $M_{1,1} = -48I_0(5, 3, 0, 0)$ , $M_{2,2} = \alpha_2 - 48I_0(5, 3, 0, 0) + 2I_1(5, 3)$<br>$M_{3,3} = \alpha_3 - 48I_0(5, 3, 0, 0) - 2I_1(5, 3)$ , $M_{5,5} = \alpha_5 - 96I_0(5, 3, 0, 2)$<br>$M_{6,6} = \alpha_6 - 48I_0(5, 3, 0, 0) + 2I_2(5, 3)$ , $M_{7,7} = \alpha_7 - 48I_0(5, 3, 0, 0) - 2I_2(5, 3)$<br>$M_{8,8} = \alpha_8 - 48I_0(5, 3, 0, 0) + 2I_4(5, 3)$ , $M_{9,9} = \alpha_9 - 48I_0(5, 3, 0, 0) - 2I_4(5, 3)$<br>$M_{15,15} = \alpha_{15} - 96I_0(5, 3, 2, 0)$ , $M_{1,7} = -2\sqrt{2}I_1(5, 3)$<br>$M_{3,15} = -2(I_1(5, 3) + I_2(5, 3))$ , all other pertinent $M_{i,j}$ with $i < j$ are zero.  |
| For $W_{7/4}$   | $M_{i,i} = \alpha_i - 64I_0(7, 4, 0, 0)$ for $i = 1, 2, 3, 6, 7, 8, 9, 14, 15$<br>all other pertinent $M_{i,j}$ with $i < j$ are zero.  |
| For $W_{8/5}$ ,<br>$W_{12/7}$ ,<br>$W_{14/9}$ ,<br>$W_{16/9}$ | $M_{1,1} = -16nI_0(\ell, n, 0, 0)$ , $M_{2,2} = \alpha_2 - 16nI_0(\ell, n, 0, 0) + 2I_1(\ell, n)$<br>$M_{3,3} = \alpha_3 - 16nI_0(\ell, n, 0, 0) - 2I_1(\ell, n)$ , $M_{4,4} = \alpha_4 - 16nI_0(\ell, n, 0, 0) + 2I_4(\ell, n)$<br>$M_{5,5} = \alpha_5 - 16nI_0(\ell, n, 0, 0) - 2I_4(\ell, n)$<br>$M_{10,10} = \alpha_{10} - 16nI_0(\ell, n, 0, 0) + 2I_3(\ell, n)$<br>$M_{11,11} = \alpha_{11} - 16nI_0(\ell, n, 0, 0) - 2I_3(\ell, n)$<br>$M_{12,12} = \alpha_{12} - 16nI_0(\ell, n, 0, 0) + 2I_7(\ell, n)$<br>$M_{13,13} = \alpha_{13} - 16nI_0(\ell, n, 0, 0) - 2I_7(\ell, n)$ , $M_{1,3} = -2\sqrt{2}I_1(\ell, n)$<br>$M_{1,11} = -2\sqrt{2}I_2(\ell, n)$ , $M_{2,10} = -4I_1(\ell, n) + 4I_5(\ell, n)$<br>$M_{3,11} = -4I_5(\ell, n)$ , $M_{4,12} = -2I_1(\ell, n) + 2I_6(\ell, n)$<br>$M_{5,13} = -2I_1(\ell, n) - 2I_6(\ell, n)$ , all other pertinent $M_{i,j}$ with $i < j$ are zero. |



$\mathcal{M}(8,5)$  is approximately the diagonal matrix

$$\mathcal{M}(8,5) \approx \begin{pmatrix} -15.0 & 0 & \mathcal{O} & 0 & 0 & 0 & \mathcal{O} & 0 & 0 \\ 0 & -13.6 & 0 & 0 & 0 & \mathcal{O} & 0 & 0 & 0 \\ \mathcal{O} & 0 & -13.6 & 0 & 0 & 0 & \mathcal{O} & 0 & 0 \\ 0 & 0 & 0 & -10.9 & 0 & 0 & 0 & \mathcal{O} & 0 \\ 0 & 0 & 0 & 0 & -10.9 & 0 & 0 & 0 & \mathcal{O} \\ 0 & \mathcal{O} & 0 & 0 & 0 & -9.2 & 0 & 0 & 0 \\ \mathcal{O} & 0 & \mathcal{O} & 0 & 0 & 0 & -9.2 & 0 & 0 \\ 0 & 0 & 0 & \mathcal{O} & 0 & 0 & 0 & -8.0 & 0 \\ 0 & 0 & 0 & 0 & \mathcal{O} & 0 & 0 & 0 & -8.0 \end{pmatrix}.$$

$\mathcal{M}(12,7)$  is approximately the diagonal matrix

$$\mathcal{M}(12,7) \approx \begin{pmatrix} -29.7 & 0 & \mathcal{O} & 0 & 0 & 0 & \mathcal{O} & 0 & 0 \\ 0 & -28.3 & 0 & 0 & 0 & \mathcal{O} & 0 & 0 & 0 \\ \mathcal{O} & 0 & -28.3 & 0 & 0 & 0 & \mathcal{O} & 0 & 0 \\ 0 & 0 & 0 & -21.5 & 0 & 0 & 0 & \mathcal{O} & 0 \\ 0 & 0 & 0 & 0 & -21.5 & 0 & 0 & 0 & \mathcal{O} \\ 0 & \mathcal{O} & 0 & 0 & 0 & -24.1 & 0 & 0 & 0 \\ \mathcal{O} & 0 & \mathcal{O} & 0 & 0 & 0 & -24.1 & 0 & 0 \\ 0 & 0 & 0 & \mathcal{O} & 0 & 0 & 0 & -18.7 & 0 \\ 0 & 0 & 0 & 0 & \mathcal{O} & 0 & 0 & 0 & -18.7 \end{pmatrix}.$$

$\mathcal{M}(14,9)$  is approximately the diagonal matrix

$$\mathcal{M}(14,9) \approx \begin{pmatrix} -12.1 & 0 & \mathcal{O} & 0 & 0 & 0 & \mathcal{O} & 0 & 0 \\ 0 & -11.7 & 0 & 0 & 0 & \mathcal{O} & 0 & 0 & 0 \\ \mathcal{O} & 0 & -11.7 & 0 & 0 & 0 & \mathcal{O} & 0 & 0 \\ 0 & 0 & 0 & -9.1 & 0 & 0 & 0 & \mathcal{O} & 0 \\ 0 & 0 & 0 & 0 & -9.1 & 0 & 0 & 0 & \mathcal{O} \\ 0 & \mathcal{O} & 0 & 0 & 0 & -10.6 & 0 & 0 & 0 \\ \mathcal{O} & 0 & \mathcal{O} & 0 & 0 & 0 & -10.6 & 0 & 0 \\ 0 & 0 & 0 & \mathcal{O} & 0 & 0 & 0 & -8.3 & 0 \\ 0 & 0 & 0 & 0 & \mathcal{O} & 0 & 0 & 0 & -8.3 \end{pmatrix}.$$

$\mathcal{M}(16,9)$  is approximately the diagonal matrix

$$\mathcal{M}(16,9) \approx \begin{pmatrix} -49.2 & 0 & \emptyset & 0 & 0 & 0 & \emptyset & 0 & 0 \\ 0 & -47.9 & 0 & 0 & 0 & \emptyset & 0 & 0 & 0 \\ \emptyset & 0 & -47.9 & 0 & 0 & 0 & \emptyset & 0 & 0 \\ 0 & 0 & 0 & -35.6 & 0 & 0 & 0 & \emptyset & 0 \\ 0 & 0 & 0 & 0 & -35.6 & 0 & 0 & 0 & \emptyset \\ 0 & \emptyset & 0 & 0 & 0 & -43.7 & 0 & 0 & 0 \\ \emptyset & 0 & \emptyset & 0 & 0 & 0 & -43.7 & 0 & 0 \\ 0 & 0 & 0 & \emptyset & 0 & 0 & 0 & -32.8 & 0 \\ 0 & 0 & 0 & 0 & \emptyset & 0 & 0 & 0 & -32.8 \end{pmatrix}.$$

All eight of these matrices are  $9 \times 9$  and negative definite. Hence Theorem 1 implies the numerical result.

## 5 Appendix: the Mathematica code

The following is a Mathematica code for computing the values  $I_0(4, 3, 0, 0)$ ,  $I_0(4, 3, 0, 2)$ ,  $I_0(4, 3, 0, 4)$ ,  $I_1(4, 3)$ ,  $I_2(4, 3)$ ,  $I_4(4, 3)$ , and the elements of the matrix  $\mathcal{M}(4, 3)$ . The seven other needed codes for different  $\ell$  and  $n$  were written similarly.

```
H = 1/2; k1 = Sin[theta1]; k2 = Sin[theta2];
gamma1 = Sqrt[Tan[theta1]]; gamma2 = Sqrt[Tan[theta2]];
alpha1 = Sqrt[4 H Sin[2theta2]/Sin[2(theta1 + theta2)]];
alpha2 = Sqrt[4 H Sin[2theta1]/Sin[2(theta1 + theta2)]];
F = 4ArcTanh[gamma1 gamma2 JacobiCN[alpha1 x, k1^2]
JacobiCN[alpha2 y, k2^2]];
V = 4 H Cosh[F];

ell = 4; n = 3;
theta1 = 2 Pi (12.7898/360);
theta2 = 2 Pi (65.354955354/360);
x0 = 3.2767; y0 = 6.3355;

Print["I_0(4,3,0,0) is ", I0x4c3c0c0x = (1/(n x0 y0))
NIntegrate[V, {x, 0, x0/4}, {y, 0, y0/4}]];

Print["I_0(4,3,0,2) is ", I0x4c3c0c2x = (1/(n x0 y0))
NIntegrate[V (Cos[2 Pi x/x0])^0 (Cos[2 Pi y/y0])^2,
{x, 0, x0/4}, {y, 0, y0/4}]];

Print["I_0(4,3,0,4) is ", I0x4c3c0c4x = (1/(n x0 y0))
```

```

NIntegrate[V (Cos[2 Pi x/x0])^0(Cos[2 Pi y/y0])^4
{x, 0, x0/4}, {y, 0, y0/4}]];

Print["I_1(4,3) is ", I1x4c3x = (8/(n x0 y0))
(NIntegrate[V (Cos[4 Pi x/(n x0)]), {x, 0, x0/4},
{y, 0, y0/4}] + NIntegrate[V (Cos[4 Pi x/(n x0)]),
{x, x0/4, 2 x0/4}, {y, 0, y0/4}] +
NIntegrate[V (Cos[4 Pi x/(n x0)]), {x, 2 x0/4, n x0/4},
{y, 0, y0/4}])]];

Print["I_2(4,3) is ", I2x4c3x = (8/(n x0 y0))
(NIntegrate[V (Cos[8 Pi x/(n x0)]), {x, 0, x0/4},
{y, 0, y0/4}] + NIntegrate[V (Cos[8 Pi x/(n x0)]),
{x, x0/4, 2 x0/4}, {y, 0, y0/4}] + NIntegrate[V (Cos[8
Pi x/(n x0)]), {x, 2 x0/4, n x0/4}, {y, 0, y0/4}])]];

Print["I_4(4,3) is ", I4x4c3x = (8/(n x0 y0))
(NIntegrate[V (Cos[4 Pi x/(n x0)]) (Cos[4 Pi y/y0]),
{x,0,x0/4}, {y,0,y0/4}]+NIntegrate[V (Cos[4 Pi x/(n x0)])
(Cos[4 Pi y/y0]), {x,x0/4,2 x0/4}, {y,0,y0/4}]+
NIntegrate[V (Cos[4 Pi x/(n x0)]) (Cos[4 Pi y/y0]),
{x, 2 x0/4, n x0/4}, {y, 0, y0/4}])]];

aa = 0; bb = 0; alpha1 = aa(4 N[Pi^2]/(n^2 x0^2)) +
bb (4 N[Pi^2]/(y0^2));

aa = 4; bb = 0; alpha2 = aa(4 N[Pi^2]/(n^2 x0^2)) +
bb (4 N[Pi^2]/(y0^2)); alpha3 = alpha2;

aa = 1; bb = 1; alpha4 = aa(4 N[Pi^2]/(n^2 x0^2)) +
bb (4 N[Pi^2]/(y0^2)); alpha5 = alpha4; alpha6 = alpha4;
alpha7 = alpha4;

aa = 0; bb = 4; alpha8 = aa(4 N[Pi^2]/(n^2 x0^2)) +
bb (4 N[Pi^2]/(y0^2)); alpha9 = alpha8;

Print["M(1,1) is ", -48 I0x4c3c0c0x];
Print["M(2,2) is ", alpha2 - 48 I0x4c3c0c0x];
Print["M(3,3) is ", alpha3 - 48 I0x4c3c0c0x];
Print["M(4,4) is ", alpha4 - 48 I0x4c3c0c0x];
Print["M(5,5) is ", alpha5 - 48 I0x4c3c0c0x];
Print["M(6,6) is ", alpha6 - 48 I0x4c3c0c0x];

```

```

Print["M(7,7) is ", alpha7 - 48 I0x4c3c0c0x];
Print["M(8,8) is ", alpha8 - 384 (I0x4c3c0c2x -
I0x4c3c0c4x)];
Print["M(9,9) is ", alpha9 - 96 (I0x4c3c0c0x +
4 I0x4c3c0c4x - 4 I0x4c3c0c2x)];
Print["M(1,9) is ", -48 N[Sqrt[2]] (-I0x4c3c0c0x +
2 I0x4c3c0c2x)];
Print["M(4,6) is ", -48 (-I0x4c3c0c0x + 2 I0x4c3c0c2x)];
Print["M(5,7) is ", -48 (-I0x4c3c0c0x + 2 I0x4c3c0c2x)];

```

### References

- [A] U. Abresch, Constant mean curvature tori in terms of elliptic functions, *J. Reine Angew. Math.* **374** (1987), 169–192.
- [BC] L. Barbosa, M. do Carmo, Stability of hypersurfaces with constant mean curvature, *Math. Z.* **185** (1984), 339–353.
- [CP] M. do Carmo, C. K. Peng, Stable minimal surfaces in  $\mathbf{R}^3$  are planes, *Bull. Amer. Math. Soc.* **1** (1979), 903–906.
- [FC] D. Fischer-Colbrie, On complete minimal surfaces with finite Morse index in three manifolds, *Invent. Math.* **82** (1985), 121–132.
- [H] H. Hopf, Differential geometry in the large, *Lecture Notes in Math.* **1000** Springer-Verlag (1983).
- [LR] F. J. Lopez, A. Ros, Completenimal surfaces with index one and stable constant mean curvature surfaces, *Comment. Math. Helv.* **64** (1989), 34–43.
- [R1] W. Rossman, The Morse index of Wente tori, to appear in *Geom. Dedicata*.
- [R2] W. Rossman, Wente Tori and Morse Index, *An. Acad. Bras. Ci.* **71(4)** (1999), 607–613.
- [Si] A. M. Silveira, Stability of complete noncompact surfaces with constant mean curvature, *Math. Ann.* **277** (1987), 629–638.
- [Sp] J. Spruck, The elliptic sinh-Gordon equation and the construction of toroidal soap bubbles, *Lecture Notes in Math.* **1340** Springer-Verlag (1988), 275–301.
- [Wa] R. Walter, Explicit examples to the  $H$ -problem of Heinz Hopf, *Geom. Dedicata* **23** (1987), 187–213.
- [We] H. Wente, Counterexample of a conjecture of H. Hopf, *Pacific J. Math.* **121** (1986), 193–243.

*Levi Lopes de Lima*  
*Departamento de Matemática*  
*Universidade Federal do Ceará*  
*Campus do Pici, 60455-760 Fortaleza, Brazil*  
*levi@mat.ufc.br*

*Vicente Francisco de Sousa Neto*  
*Departamento de Matemática*  
*Universidade Católica de Pernambuco*  
*Recife, Brazil*  
*vicente@unicap.br*

*Wayne Rossman*  
*Department of Mathematics*  
*Faculty of Science*  
*Kobe University*  
*Rokko, Kobe 657-8501, Japan*  
*wayne@math.kobe-u.ac.jp*  
*<http://www.math.kobe-u.ac.jp/HOME/wayne/wayne.html>*