

# Grid–symmetric generalized quadrangles\*

William M. Kantor

## Abstract

A generalized quadrangle is classical if it has a grid of axes of symmetry.

In a finite generalized quadrangle  $\mathbf{Q}$  of order  $(s, t)$  with  $s, t > 1$ , a line  $L$  is called an *axis of symmetry* if the group  $T(L)$  of all automorphisms (“symmetries”) that fix every line meeting  $L$  has the maximal possible order  $s$ . Moreover,  $\mathbf{Q}$  is called *span–symmetric* if there are two disjoint axes of symmetry; we will call  $\mathbf{Q}$  *grid–symmetric* if there are two further disjoint axes of symmetry, each of which meets  $L$  and  $M$ .

Span–symmetric generalized quadrangles were first studied in [Pa] (cf. [PT1]), in view of the known examples  $Q(4, q)$  and  $Q(5, q)$ , arising respectively from quadrics in 4– and 5–dimensional projective spaces. More than 20 years ago it was shown that the generalized quadrangles  $Q(4, q)$  are the only span–symmetric ones with  $t \neq s^2$  (cf. [Ka, Th1]). While nonclassical examples exist if  $t = s^2$ , this is not so in the grid–symmetric case:

**Theorem.** *Any grid–symmetric generalized quadrangle of order  $(s, t)$  is isomorphic to  $Q(4, s)$  or  $Q(5, s)$ .*

*Proof.* By the result just noted, we may assume that  $t = s^2$ . There are sets  $\Lambda$  and  $\Lambda^\perp$ , each consisting of  $s + 1$  lines of symmetry, where each line in  $\Lambda$  meets each line in  $\Lambda^\perp$ . Let  $A$  and  $B$  be the groups generated by the symmetries corresponding to  $\Lambda$  and  $\Lambda^\perp$ , respectively. By [Th2, 12.5.5],  $A \cong B \cong \text{SL}(2, s)$ . If  $L \in \Lambda$  and  $M \in \Lambda^\perp$  then  $T(L)$  fixes  $M$  and hence normalizes  $T(M)$ . Also  $T(M)$  normalizes  $T(L)$ , so that these two groups commute since  $T(L) \cap T(M) = 1$ . Thus,  $A$  and  $B$  are commuting groups each of which is isomorphic to  $\text{SL}(2, s)$ .

---

\*This research was supported in part by the National Science Foundation.

Received by the editors August 2002.

Communicated by J. Thas.

2000 *Mathematics Subject Classification* : Primary 51E12.

*Key words and phrases* : generalized quadrangle, quadric, ovoid.

Let  $\Omega$  denote the set of points on all lines of  $\Lambda$ , and hence of  $\Lambda^\perp$ . If  $x$  is any point not in  $\Omega$  then  $\Omega \cup x^A$  is the set of points of a  $Q(4, s)$ -subquadrangle  $\mathbf{Q}_x$  [Th2, 12.5.5]. If  $M \in \Lambda^\perp$  then  $T(M)$  fixes each line of  $\mathbf{Q}_x$  meeting  $M$  and hence acts on the union  $\mathbf{Q}_x$  of these lines. Thus,  $AB$  acts on  $\mathbf{Q}_x$ , and hence acts in the natural manner as  $\Omega^+(4, s)$  on the space  $\mathbf{P}_x = \text{PG}(4, s)$  underlying  $\mathbf{Q}_x$ , fixing the point  $m$  of  $\mathbf{P}_x \setminus \mathbf{Q}_x$  perpendicular to  $\langle \Omega \rangle$ . Note that  $AB \cong \Omega^+(4, s)$ : if  $s$  is odd and  $z_A$  and  $z_B$  are the involutions in  $A$  and  $B$ , respectively, then  $z_A z_B = 1$  on  $\mathbf{Q}_x$  for each point  $x \notin \Omega$ , and hence is 1 on  $\mathbf{Q}$ .

Note that, if  $x \notin \Omega$  as above, then  $(AB)_x \cong \text{PSL}(2, s)$ . For,  $x$  lies on the line of  $\mathbf{P}_x$  joining  $m$  and some point  $n$  of  $\langle \Omega \rangle \setminus \Omega$ , so that the stabilizer  $(AB)_x$  fixes  $n$ . However,  $(AB)_n \cong \Omega(3, s) \cong \text{PSL}(2, s)$  has no proper subgroup of index  $(2, s - 1)$ . Since  $(AB)_n$  permutes the  $(2, s - 1)$  points of  $\mathbf{Q}_x$  on the line  $\langle m, n \rangle$ , it follows that  $(AB)_x = (AB)_n \cong \text{PSL}(2, s)$ .

Now consider any point  $y$  of  $\mathbf{Q}$  not in  $\Omega \cup x^A$  and the resulting point-orbit  $y^A$  and subquadrangle. As in the preceding paragraph,  $G := (AB)_y \cong \text{PSL}(2, s)$ . Here  $G$  acts on  $\mathcal{O} := y^\perp \cap \mathbf{Q}_x$ , which is an ovoid of  $\mathbf{Q}_x$  [PT2, p. 26]: each of the  $s^2 + 1$  lines through  $y$  meets  $\mathbf{Q}_x$ , and no two of the resulting  $s^2 + 1$  points are perpendicular.

Under the Klein correspondence for a suitable quadric of  $\mathbf{P} = \text{PG}(5, q)$  containing  $\mathbf{Q}_x$ , the ovoid  $\mathcal{O}$  produces a spread of lines in  $\text{PG}(3, s)$  and hence also a translation plane  $\pi$  of order  $s^2$ , with kernel containing  $\text{GF}(s)$ . Moreover, under this correspondence, the group  $AB \cong \Omega^+(4, s)$  produces a subgroup of  $\text{GL}(4, s)$ , isomorphic to  $A \times B$ , that has a subgroup  $\hat{G} \cong \text{PSL}(2, s)$  or  $\text{SL}(2, s)$  produced by  $G$ ; moreover  $\hat{G}$  preserves the spread. If  $q$  is odd then  $\hat{G} \not\cong \text{PSL}(2, s)$  since all involutions in  $A \times B$  lie in its center. For all  $q$  it follows that  $G$  produces a collineation group  $\hat{G} \cong \text{SL}(2, s)$  of  $\pi$ .

All translation planes having the preceding properties are known [Sch, Wa]: the nondesarguesian ones are Hall, Hering, Walker and Ott-Schaeffer planes. It is easy to check that, for each of these nondesarguesian planes, the corresponding ovoid spans  $\mathbf{P}$ , whereas our ovoid  $\mathcal{O}$  lies in  $\mathbf{Q}_x$  and hence in the hyperplane  $\mathbf{P}_x$  of  $\mathbf{P}$ . Hence  $\pi$  is desarguesian and  $\mathcal{O}$  is an elliptic quadric.

Thus,  $y^\perp \cap \mathbf{Q}_x$  is an elliptic quadric of  $\mathbf{Q}_x$  for each point  $y$  of  $\mathbf{Q} \setminus \mathbf{Q}_x$ . Consequently, our original generalized quadrangle is classical [TP, Br]. ■

## References

- [Br] M. R. Brown, A characterisation of the generalized quadrangle  $Q(5, q)$  using cohomology. *J. Algebraic Combin.* 15 (2002) 107–125.
- [Ka] W. M. Kantor, Note on span-symmetric generalized quadrangles. *Adv. Geom.* 2 (2002) 197–200.
- [Pa] S. E. Payne, Span-symmetric generalized quadrangles, pp. 231–242 in: *The geometric vein (The Coxeter Festschrift; eds. C. Davis et al.)*, Springer, New York-Heidelberg-Berlin 1981.
- [PT1] S. E. Payne and J. A. Thas, Generalized quadrangles and the Higman-Sims technique. *Eur. J. Comb.* 2 (1982) 79–89.

- [PT2] S. E. Payne and J. A. Thas, Finite generalized quadrangles. Pitman, Boston-London-Melbourne 1984.
- [Sch] H.-J. Schaeffer, Translationsebenen auf denen die Gruppe  $SL(2, p^n)$  operiert, Diplomarbeit, Univ. Tübingen 1975/76.
- [TP] J. A. Thas and S. E. Payne, Spreads and ovoids in finite generalized quadrangles. *Geom. Dedicata* 52 (1994) 227–253.
- [Th1] K. Thas, Classification of span-symmetric generalized quadrangles of order  $s$ . *Adv. Geom.* 2 (2002) 189–196.
- [Th2] K. Thas, Automorphisms and combinatorics of finite generalized quadrangles. Ph.D. thesis, Univ. Ghent 2002.
- [Wa] M. Walker, A characterization of some translation planes. *Abh. Math. Sem. Univ. Hamburg* 49 (1979) 216–233.

William M. Kantor  
Department of Mathematics  
University of Oregon  
Eugene, OR 97403, USA  
email: [kantor@math.uoregon.edu](mailto:kantor@math.uoregon.edu)