

Two-intersection sets with respect to lines on the Klein quadric

F. De Clerck N. De Feyter* N. Durante†

Abstract

We construct new examples of sets of points on the Klein quadric $Q^+(5, q)$, q even, having exactly two intersection sizes 0 and α with lines on $Q^+(5, q)$. By the well-known Plücker correspondence, these examples yield new $(0, \alpha)$ -geometries embedded in $\text{PG}(3, q)$, q even.

1 Preliminaries

A $(0, \alpha)$ -geometry $\mathcal{S} = (\mathcal{P}, \mathcal{L}, I)$ is a connected partial linear space of order (s, t) (i.e., every line is incident with $s + 1$ points, while every point is incident with $t + 1$ lines) such that for every anti-flag $\{p, L\}$ the number of lines through p and intersecting L is 0 or α . The concept of a $(0, \alpha)$ -geometry, introduced by Debroey, De Clerck and Thas [5, 20], generalizes a lot of well-studied classes of geometries such as semipartial geometries [8], partial geometries [2] and generalized quadrangles [16].

A $(0, \alpha)$ -geometry $\mathcal{S} = (\mathcal{P}, \mathcal{L}, I)$ is fully embedded in $\text{PG}(n, q)$ if \mathcal{L} is a set of lines of $\text{PG}(n, q)$ not contained in a proper subspace and \mathcal{P} is the set of all points of $\text{PG}(n, q)$ on the lines of \mathcal{S} . In [20] the $(0, \alpha)$ -geometries ($\alpha > 1$) fully embedded in $\text{PG}(n, q)$, $n > 3, q > 2$, are classified. For $\alpha = 1$ as well as for the $(0, \alpha)$ -geometries with $q = 2$ a classification of the embeddings is out of reach as explained for instance in [6, 20]. As for $\text{PG}(3, q)$, in [5] it is proven that if \mathcal{S} is a $(0, \alpha)$ -geometry ($\alpha > 1$) fully embedded in $\text{PG}(3, q)$, $q > 2$, then every planar pencil of $\text{PG}(3, q)$ (i.e., the

*This research was supported by a BOF ("Bijzonder Onderzoeksfonds") grant at Ghent University.

†This author acknowledges support from the project "Strutture geometriche, Combinatoria e loro applicazioni" of the Italian M.U.R.S.T.

$q + 1$ lines through a point in a plane) contains 0 or α lines of \mathcal{S} . Conversely one easily verifies that a set of lines of $\text{PG}(3, q)$ which shares 0 or α ($\alpha > 1$) lines with every pencil of $\text{PG}(3, q)$ yields a $(0, \alpha)$ -geometry fully embedded in $\text{PG}(3, q)$.

We can use the well-known Plücker correspondence, in order to see the set of lines of the $(0, \alpha)$ -geometry as a set of points on the Klein quadric $Q^+(5, q)$.

For the remainder of the paper we will always assume that $\alpha > 1$ and $q > 2$, and we may conclude that the following objects are equivalent.

- A $(0, \alpha)$ -geometry fully embedded in $\text{PG}(3, q)$.
- A set of lines of $\text{PG}(3, q)$ sharing 0 or α lines with every pencil of $\text{PG}(3, q)$.
- A set of points on the Klein quadric $Q^+(5, q)$ sharing 0 or α points with every line on $Q^+(5, q)$. We call such a set a $(0, \alpha)$ -set on $Q^+(5, q)$.

A *maximal arc of degree α* in $\text{PG}(2, q)$ is a set of points such that every line of $\text{PG}(2, q)$ intersects it in 0 or α points. Examples of maximal arcs in $\text{PG}(2, 2^h)$ were first constructed by Denniston [10]. Examples of maximal arcs in $\text{PG}(2, 2^h)$ which are not of Denniston type were constructed by Thas [18, 19] and by Mathon [15]. Ball, Blokhuis and Mazzocca [1] proved that maximal arcs of degree $1 < \alpha < q$ in $\text{PG}(2, q)$ do not exist if q is odd.

Let \mathcal{K} be a $(0, \alpha)$ -set on $Q^+(5, q)$. Clearly every plane on $Q^+(5, q)$ is either disjoint from \mathcal{K} or intersects \mathcal{K} in a maximal arc of degree α . Consider the $(0, \alpha)$ -geometry $\mathcal{S} = (\mathcal{P}, \mathcal{L}, \mathcal{I})$, fully embedded in $\text{PG}(3, q)$, which corresponds to \mathcal{K} . Then every plane of $\text{PG}(3, q)$ contains either no line of \mathcal{S} or $q\alpha - q + \alpha$ lines of \mathcal{S} which constitute a dual maximal arc of degree α . Similarly through every point p of $\text{PG}(3, q)$ there are either 0 lines of \mathcal{S} or $q\alpha - q + \alpha$ lines of \mathcal{S} which intersect a plane not containing p in a maximal arc of degree α . Let π be a plane of $\text{PG}(3, q)$ containing $q\alpha - q + \alpha$ lines of \mathcal{S} , and let d be such that π contains $q^2 + q + 1 - d$ points of \mathcal{S} . Then counting the lines of \mathcal{S} by their intersection with π we get that $|\mathcal{L}| = |\mathcal{K}| = (q\alpha - q + \alpha)(q^2 + 1 - d)$. We call d the *deficiency* of the $(0, \alpha)$ -geometry \mathcal{S} and of the $(0, \alpha)$ -set \mathcal{K} .

In this paper we will give an overview of the known examples so far and we will give new examples, α being any proper divisor of q , q even.

2 The known examples

It is clear that the design of all points and lines of $\text{PG}(3, q)$ is the only $(0, q + 1)$ -geometry fully embedded in $\text{PG}(3, q)$.

On the other hand let \mathcal{S} be a $(0, q)$ -geometry fully embedded in $\text{PG}(3, q)$ and let Π be the set of planes containing at least two lines of \mathcal{S} . Then for every plane $\pi \in \Pi$, the incidence structure of points and lines of \mathcal{S} in π is a dual affine plane, while the incidence structure with point set the set of lines of \mathcal{S} through a fixed point p of \mathcal{S} and with line set the set of planes of Π through p is an affine plane. These geometries are classified, there are two non-isomorphic examples, see for instance [9, 13, 14]. Here we summarize the result in the terminology of a $(0, q)$ -set on the Klein quadric $Q^+(5, q)$.

Theorem 2.1. *The points of $Q^+(5, q)$ not on a hyperplane U of $\text{PG}(5, q)$, $q > 2$, are the only $(0, q)$ -sets on $Q^+(5, q)$. If U is a tangent hyperplane, then the deficiency is 1. If U is a secant hyperplane then the deficiency is 0.*

Remark

For the $(0, q)$ -set of deficiency 1 the corresponding $(0, q)$ -geometry in $\text{PG}(3, q)$ is the well known dual net denoted by H_q^3 . For the $(0, q)$ -set of deficiency 0 the corresponding $(0, q)$ -geometry in $\text{PG}(3, q)$ is the semipartial geometry denoted by $\overline{W(3, q)}$. For a detailed description of both examples as $(0, q)$ -geometries embedded in $\text{PG}(3, q)$ we refer for instance to [6].

In [1] it is proved that in desarguesian planes of order q , q odd, maximal arcs of degree α , $1 < \alpha < q$, do not exist. Hence we can conclude that if q is odd, no other $(0, \alpha)$ -set, $\alpha > 1$, on the Klein quadric $Q^+(5, q)$ exists. Hence, for other examples we may restrict ourselves to the case q even, $1 < \alpha < q$.

Here is an other example. The points of $Q^+(5, q)$, q even, corresponding to the external lines of a nonsingular hyperbolic quadric in $\text{PG}(3, q)$ form a $(0, \alpha)$ -set on $Q^+(5, q)$ with $\alpha = q/2$ and deficiency $q + 1$. The corresponding $(0, q/2)$ -geometry is denoted by $\text{NQ}^+(3, q)$.

It was conjectured in [5] that H_q^3 , $\overline{W(3, q)}$ and $\text{NQ}^+(3, q)$ are the only $(0, \alpha)$ -geometries, with $\alpha > 1$, fully embedded in $\text{PG}(3, q)$, $q > 2$. This conjecture is false as will be clear from the remainder of the paper.

A first counterexample has been given by Ebert, Metsch and Szőnyi [11]. A k -cap in $\text{PG}(n, q)$ is a set of k points, no three on a line. It is called *maximal* if it is not contained in a larger cap. Quite some research has been done on caps in $\text{PG}(5, q)$ that are contained in the Klein quadric $Q^+(5, q)$. Since the maximum size of a cap in $\text{PG}(2, q)$ is $q + 1$ if q is odd and $q + 2$ if q is even, a cap in $Q^+(5, q)$ has size at most $(q + 1)(q^2 + 1)$ if q is odd and at most $(q + 2)(q^2 + 1)$ if q is even. Glynn [12] constructs a cap of size $(q + 1)(q^2 + 1)$ in $Q^+(5, q)$ for any prime power q (see also [17]). Ebert, Metsch and Szőnyi construct caps of size $q^3 + 2q^2 + 1 = (q + 2)(q^2 + 1) - q - 1$ in $Q^+(5, q)$ for q even. They show that a cap in $Q^+(5, q)$, q even, of size $q^3 + 2q^2 + 1$ is either maximal in $Q^+(5, q)$ and is then a $(0, 2)$ -set of deficiency 1 together with one extra point, or it is contained in a cap of size $(q + 2)(q^2 + 1)$. One easily verifies that caps of size $(q + 2)(q^2 + 1)$ in $Q^+(5, q)$, q even, and $(0, 2)$ -sets of deficiency 0 are equivalent. A cap of size $(q + 2)(q^2 + 1)$ is only known to exist for $q = 2$.

The construction of Ebert, Metsch and Szőnyi is as follows. Let Σ be a 3-space intersecting $Q^+(5, q)$ in a nonsingular elliptic quadric E . Let $L = \Sigma^\beta$ where β is the symplectic polarity associated with $Q^+(5, q)$. Then the line L is external to $Q^+(5, q)$. Consider an ovoid O in Σ which has the same set of tangent lines as E . Let \mathcal{K} be the intersection of $Q^+(5, q)$ with the cone with vertex L and base $E \cup O$. Then \mathcal{K} is a cap of size $(q + 1)|O \setminus E| + q^2 + 1$ which is maximal in $Q^+(5, q)$ [11], and $\mathcal{K} \setminus (E \cap O)$ is a $(0, 2)$ -set in $Q^+(5, q)$ of deficiency $|E \cap O|$.

We have the following possibilities for O . The ovoid O can be an elliptic quadric. Then E and O intersect in either one point or $q + 1$ points which form a conic in a plane of Σ (Types 1(i) and 3(g)(ii) in Table 2 of [3]). We will denote the corresponding $(0, 2)$ -set by \mathcal{E}_1 if $|E \cap O| = 1$ and by \mathcal{E}_{q+1} if $|E \cap O| = q + 1$. On

the other hand when q is an odd power of 2 the ovoid O can be a Suzuki-Tits ovoid. Then E and O intersect in $q \pm \sqrt{2q} + 1$ points and both intersection sizes do occur [7]. We will denote the corresponding $(0, 2)$ -set by $\mathcal{T}_{q-\sqrt{2q}+1}$ if $|E \cap O| = q - \sqrt{2q} + 1$ and by $\mathcal{T}_{q+\sqrt{2q}+1}$ if $|E \cap O| = q + \sqrt{2q} + 1$.

3 Unions of elliptic quadrics

Consider a $(0, 2)$ -set $\mathcal{K} \in \{\mathcal{E}_1, \mathcal{E}_{q+1}\}$ in $Q^+(5, q)$, $q = 2^h$. Let Π be a hyperplane containing Σ and let $p = \Pi \cap L$, where $L = \Sigma^\beta$. Then Π intersects $Q^+(5, q)$ in a nonsingular parabolic quadric $Q(4, q)$ with nucleus p . Since \mathcal{K} is the intersection of $Q^+(5, q)$ with the cone with vertex L and base the symmetric difference $E \Delta O$ we find that $\mathcal{K} \cap \Pi$ is the intersection of $Q(4, q)$ with the cone with vertex p and base $E \Delta O$.

The projection of $Q(4, q)$ from p on Σ yields an isomorphism from the classical generalized quadrangle $Q(4, q)$ to the classical generalized quadrangle $W(q)$ consisting of the points of Σ and the lines of Σ that are tangent to E . This isomorphism induces a bijection from the set of ovoids of $Q(4, q)$ to the set of ovoids of $W(q)$. Since the ovoid O has the same set of tangent lines as E , it is an ovoid of the generalized quadrangle $W(q)$. Hence O is the projection from p on Σ of an ovoid \bar{O} of $Q(4, q)$. So $\mathcal{K} \cap \Pi$ is the symmetric difference $E \Delta \bar{O}$. Since O is a nonsingular elliptic quadric in Σ , \bar{O} is a nonsingular elliptic quadric in a 3-space $\bar{\Sigma} \subseteq \Pi$. Now Σ and $\bar{\Sigma}$ intersect in a plane $\bar{\pi}$ and we may also write $\mathcal{K} \cap \Pi = Q(4, q) \cap (\Sigma \cup \bar{\Sigma}) \setminus \bar{\pi}$.

From the definition of \mathcal{E}_1 and \mathcal{E}_{q+1} it follows that there is exactly one plane $\pi \subseteq \Sigma$ such that $\pi \cap Q(4, q) = E \cap O$. Indeed, if $\mathcal{K} = \mathcal{E}_1$ then E and O intersect in exactly one point and π is the unique tangent plane in Σ to E at this point. If $\mathcal{K} = \mathcal{E}_{q+1}$ then E and O intersect in a nondegenerate conic and π is the ambient plane of this conic. We prove that $\bar{\pi} = \pi$. Since O is the projection of \bar{O} from p on Σ , $E \cap \bar{O} = E \cap O$. Since $\bar{O} = \bar{\Sigma} \cap Q(4, q)$, $\bar{\pi} \cap Q(4, q) = \Sigma \cap \bar{\Sigma} \cap Q(4, q) = \Sigma \cap \bar{O} = E \cap \bar{O} = E \cap O$. So $\bar{\pi}$ is a plane in Σ such that $\bar{\pi} \cap Q(4, q) = E \cap O$. This means that $\bar{\pi} = \pi$.

So $\mathcal{K} \cap \Pi$ is the symmetric difference of elliptic quadrics E and \bar{O} on $Q(4, q)$ with ambient 3-spaces Σ and $\bar{\Sigma}$ intersecting in the plane π . Since this holds for all hyperplanes Π containing Σ we conclude that there exist 3-spaces $\Sigma_0 = \Sigma, \Sigma_1, \dots, \Sigma_{q+1}$ mutually intersecting in the plane π , such that each intersects $Q^+(5, q)$ in an elliptic quadric and such that

$$\mathcal{K} = Q^+(5, q) \cap (\Sigma_0 \cup \Sigma_1 \cup \dots \cup \Sigma_{q+1}) \setminus \pi.$$

What remains to be verified is the position of the 3-spaces Σ_i . Consider a plane π' spanned by L and a point $r \in O \setminus E$. One verifies in the respective cases $\mathcal{K} = \mathcal{E}_1$ and $\mathcal{K} = \mathcal{E}_{q+1}$ that $\pi \cap O = \pi \cap E = E \cap O$, so $r \notin \pi$. Hence π' is skew to π . We determine the points of intersection of Σ_i , $i = 0, \dots, q+1$, with π' . Clearly $\Sigma_0 \cap \pi' = \Sigma \cap \pi' = r$. Let $i \in \{1, \dots, q+1\}$ and let $p_i \in L$ be such that $\Sigma_i \subseteq \langle p_i, \Sigma \rangle$. Let r_i be the unique point of $Q^+(5, q)$ on the line $\langle p_i, r \rangle$. Since $r \in O \setminus E$, r_i is a point of \mathcal{K} and hence of Σ_i . But also $r_i \in \pi'$, so $\Sigma_i \cap \pi' = r_i$. Repeating this reasoning for all points p_i on L we see that the 3-spaces Σ_i , $i = 1, \dots, q+1$, intersect π' in the points of the nondegenerate conic $C' = \pi' \cap Q^+(5, q)$ and that Σ intersects π' in the point r which is the nucleus of the conic C' . We have now proven the following theorem which completely determines the structure of the $(0, 2)$ -sets \mathcal{E}_1 and \mathcal{E}_{q+1} .

Theorem 3.1. *Let $\mathcal{K} \in \{\mathcal{E}_1, \mathcal{E}_{q+1}\}$ and let π be the unique plane in Σ such that $\pi \cap Q^+(5, q) = E \cap O$. Then*

$$\mathcal{K} = (E \cup O_1 \cup \dots \cup O_{q+1}) \setminus \pi,$$

where O_i , $1 \leq i \leq q+1$, is a nonsingular 3-dimensional elliptic quadric on $Q^+(5, q)$ such that its ambient space Σ_i intersects Σ in the plane π . In particular the 3-spaces $\Sigma_1, \dots, \Sigma_{q+1}$ intersect each plane $\pi' = \langle r, L \rangle$ with $L = \Sigma^\beta$ and $r \in O \setminus E$ in the points of the nondegenerate conic $C' = \pi' \cap Q^+(5, q)$, while Σ intersects π' in the nucleus r of the conic C' .

Remark

We can apply the same reasoning to the $(0, 2)$ -sets $\mathcal{T}_{q \pm \sqrt{2q+1}}$. We find then that $\mathcal{T}_{q \pm \sqrt{2q+1}}$ can be written as

$$(E \cup O_1 \cup \dots \cup O_{q+1}) \setminus (E \cap O),$$

where O_1, \dots, O_{q+1} are Suzuki-Tits ovoids in the hyperplanes containing Σ , such that for every $p_i \in L = \Sigma^\beta$, there is exactly one $O_i \subseteq \langle p_i, \Sigma \rangle$, and then O is the projection of O_i from p_i on Σ . However this was already known [4].

4 A new construction

The following construction is inspired by Theorem 3.1. Let π be a plane of $\text{PG}(5, q)$, $q = 2^h$, which does not contain any line of $Q^+(5, q)$ and let π' be a plane skew to π . Let \mathcal{D} denote the set of points $p \in \pi'$ such that $\langle p, \pi \rangle$ intersects $Q^+(5, q)$ in a nonsingular elliptic quadric, and suppose that A is a maximal arc of degree α in π' such that $A \subseteq \mathcal{D}$. Then we define the set $\mathcal{M}^\alpha(A)$ to be the intersection of $Q^+(5, q)$ with the cone with vertex π and base A , minus the points of $Q^+(5, q)$ in π .

Theorem 4.1. *The set $\mathcal{M}^\alpha(A)$ is a $(0, \alpha)$ -set on $Q^+(5, q)$.*

Proof. Let L be a line on $Q^+(5, q)$ which intersects the plane π . Then the subspace $\Sigma = \langle L, \pi \rangle$ has dimension 3 and it contains a line of $Q^+(5, q)$. Hence $\Sigma \cap \pi' \notin A$. So there are no points of $\mathcal{M}^\alpha(A)$ in Σ and hence also none on L .

Let L be a line on $Q^+(5, q)$ which is skew to π . A point p on L is in $\mathcal{M}^\alpha(A)$ if and only if $\langle p, \pi \rangle \cap \pi' \in A$ if and only if the projection of p from π on π' is a point of A . So if L' is the projection of L from π on π' then $|L \cap \mathcal{M}^\alpha(A)| = |L' \cap A| \in \{0, \alpha\}$. So every line on $Q^+(5, q)$ intersects $\mathcal{M}^\alpha(A)$ in 0 or α points. \square

Since the plane π does not contain any line of $Q^+(5, q)$, there are two possibilities: either $\pi \cap Q^+(5, q)$ is a single point or it is a nondegenerate conic. In the former case the $(0, \alpha)$ -set has deficiency 1 and it is denoted by $\mathcal{M}_1^\alpha(A)$. In the latter case the $(0, \alpha)$ -set has deficiency $q+1$ and it is denoted by $\mathcal{M}_{q+1}^\alpha(A)$.

In order to prove that there do exist $(0, \alpha)$ -sets of deficiency 1 and $q+1$ for every $\alpha \in \{2, 2^2, \dots, 2^{h-1} = q/2\}$ we must show that the set \mathcal{D} in the plane π' contains a maximal arc of degree α for every $\alpha \in \{2, 2^2, \dots, 2^{h-1}\}$, and this for both

the case where $\pi \cap Q^+(5, q)$ is a single point and the case where $\pi \cap Q^+(5, q)$ is a nondegenerate conic.

If $\pi \cap Q^+(5, q)$ is a single point p then \mathcal{D} is the set of points of π' which are not on the line $\pi' \cap T_p$, where T_p is the tangent hyperplane to $Q^+(5, q)$ at p . Clearly in this case the set \mathcal{D} contains a maximal arc of degree α for every $\alpha \in \{2, 2^2, \dots, 2^{h-1}\}$.

If $\pi \cap Q^+(5, q)$ is a nondegenerate conic then the plane π^β also intersects $Q^+(5, q)$ in a nondegenerate conic C . Furthermore β induces an anti-automorphism between the projective plane π^β and the projective plane having as points the 3-spaces through π and as lines the hyperplanes through π . This anti-automorphism is such that a 3-space containing π intersects $Q^+(5, q)$ in a nonsingular elliptic quadric if and only if the corresponding line of π^β is external to the conic C . Hence the set \mathcal{D} in the plane π' is the dual of the set of external lines to a nondegenerate conic. It follows that \mathcal{D} is a Denniston type maximal arc [10] of degree $q/2$, and hence that \mathcal{D} contains a maximal arc of degree α for every $\alpha \in \{2, 2^2, \dots, 2^{h-1}\}$. We have proven the following theorem.

Theorem 4.2. *There exist $(0, \alpha)$ -sets on $Q^+(5, q)$, $q = 2^h$, of deficiency 1 and $q + 1$ for all $\alpha \in \{2, 2^2, \dots, 2^{h-1}\}$.*

Corollary 4.3. *There exist $(0, \alpha)$ -geometries fully embedded in $\text{PG}(3, q)$, $q = 2^h$, of deficiency 1 and $q + 1$ for all $\alpha \in \{2, 2^2, \dots, 2^{h-1}\}$.*

By Theorem 3.1 the $(0, 2)$ -set \mathcal{E}_d , $d = 1, q + 1$, is of the form $\mathcal{M}_d^2(H)$ with H a regular hyperoval. Let \mathcal{K} be the $(0, q/2)$ -set corresponding to the $(0, q/2)$ -geometry $\text{NQ}^+(3, q)$, q even. Then \mathcal{K} corresponds to the set of external lines to a nonsingular hyperbolic quadric $Q^+(3, q)$ in $\text{PG}(3, q)$. Let C be the set of points of $Q^+(5, q)$ corresponding to one of the two reguli of lines contained in $Q^+(3, q)$. Then C is a nondegenerate conic in a plane π , and \mathcal{K} is the set of all points of $Q^+(5, q)$ which are not collinear in $Q^+(5, q)$ with any of the points of C . So a point p of $Q^+(5, q)$ is in \mathcal{K} if and only if $p \notin \pi$ and $\langle p, \pi \rangle$ intersects $Q^+(5, q)$ in a nondegenerate elliptic quadric. Hence $\text{NQ}^+(3, q)$ corresponds to the $(0, q/2)$ -set $\mathcal{M}_{q+1}^{q/2}(\mathcal{D})$.

We conclude this paper with a list of all the known distinct examples of $(0, \alpha)$ -sets \mathcal{K} in $Q^+(5, q)$, $\alpha > 1$, $q > 2$. In this list d is the deficiency of the $(0, \alpha)$ -set \mathcal{K} .

- $\alpha = q + 1$, $d = 0$, and \mathcal{K} is the set of all points of $Q^+(5, q)$.
- $\alpha = q$, $d = 0$, and \mathcal{K} corresponds to $\overline{W(3, q)}$.
- $\alpha = q$, $d = 1$, and \mathcal{K} corresponds to H_q^3 .
- $q = 2^h$, $\alpha \in \{2, 2^2, \dots, 2^{h-1}\}$, $d \in \{1, q + 1\}$ and $\mathcal{K} = \mathcal{M}_d^\alpha(A)$.
- $q = 2^{2h+1}$, $\alpha = 2$, $d = q \pm \sqrt{2q} + 1$, and $\mathcal{K} = \mathcal{T}_{q \pm \sqrt{2q} + 1}$.

References

- [1] S. Ball, A. Blokhuis, and F. Mazzocca. Maximal arcs in Desarguesian planes of odd order do not exist. *Combinatorica*, 17(1):31–41, 1997.
- [2] R. C. Bose. Strongly regular graphs, partial geometries and partially balanced designs. *Pacific J. Math.*, 13:389–419, 1963.
- [3] A. A. Bruen and J. W. P. Hirschfeld. Intersections in projective space. II. Pencils of quadrics. *European J. Combin.*, 9(3):255–270, 1988.
- [4] A. Cossidente. Caps embedded in the Klein quadric. *Bull. Belg. Math. Soc. Simon Stevin*, 7(1):13–19, 2000.
- [5] F. De Clerck and J. A. Thas. The embedding of $(0, \alpha)$ -geometries in $\text{PG}(n, q)$. I. In *Combinatorics '81 (Rome, 1981)*, volume 78 of *North-Holland Math. Stud.*, pages 229–240. North-Holland, Amsterdam, 1983.
- [6] F. De Clerck and H. Van Maldeghem. Some classes of rank 2 geometries. In *Handbook of Incidence Geometry*, pages 433–475. North-Holland, Amsterdam, 1995.
- [7] V. De Smet and H. Van Maldeghem. Intersections of Hermitian and Ree ovoids in the generalized hexagon $H(q)$. *J. Combin. Des.*, 4(1):71–81, 1996.
- [8] I. Debroey and J. A. Thas. On semipartial geometries. *J. Combin. Theory Ser. A*, 25(3):242–250, 1978.
- [9] A. Del Fra and D. Ghinelli. $Af^*.Af$ geometries, the Klein quadric and \mathcal{H}_q^n . *Discrete Math.*, 129(1-3):53–74, 1994. Linear spaces (Capri, 1991).
- [10] R. H. F. Denniston. Some maximal arcs in finite projective planes. *J. Combinatorial Theory*, 6:317–319, 1969.
- [11] G. L. Ebert, K. Metsch, and T. Szőnyi. Caps embedded in Grassmannians. *Geom. Dedicata*, 70(2):181–196, 1998.
- [12] D. G. Glynn. On a set of lines of $\text{PG}(3, q)$ corresponding to a maximal cap contained in the Klein quadric of $\text{PG}(5, q)$. *Geom. Dedicata*, 26(3):273–280, 1988.
- [13] M. P. Hale, Jr. Finite geometries which contain dual affine planes. *J. Combin. Theory Ser. A*, 22(1):83–91, 1977.
- [14] J. I. Hall. Classifying copolar spaces and graphs. *Quart. J. Math. Oxford Ser. (2)*, 33(132):421–449, 1982.
- [15] R. Mathon. New maximal arcs in Desarguesian planes. *J. Combin. Theory Ser. A*, 97(2):353–368, 2002.
- [16] S. E. Payne and J. A. Thas. *Finite Generalized Quadrangles*, volume 110 of *Research Notes in Mathematics*. Pitman (Advanced Publishing Program), Boston, MA, 1984.

- [17] L. Storme. On the largest caps contained in the Klein quadric of $\text{PG}(5, q)$, q odd. *J. Combin. Theory Ser. A*, 87(2):357–378, 1999.
- [18] J. A. Thas. Construction of maximal arcs and partial geometries. *Geometriae Dedicata*, 3:61–64, 1974.
- [19] J. A. Thas. Construction of maximal arcs and dual ovals in translation planes. *European J. Combin.*, 1(2):189–192, 1980.
- [20] J. A. Thas, I. Debroey, and F. De Clerck. The embedding of $(0, \alpha)$ -geometries in $\text{PG}(n, q)$. II. *Discrete Math.*, 51(3):283–292, 1984.

Frank De Clerck; Nikias De Feyter
Department of Pure Mathematics and Computer Algebra
Ghent University
Krijgslaan 281 - S22
B-9000 Gent
Belgium
E-mail: fdc@cage.UGent.be; ndfeyter@cage.UGent.be

Nicola Durante
Dipartimento di Matematica e Applicazioni “R. Caccioppoli”
Università di Napoli “Federico II”
Complesso M. S. Angelo, Ed.T
Via Cintia
I-80126 Napoli
Italy
E-mail: ndurante@unina.it