

## The $m$ -Derivations of Analytic Vector Fields Lie Algebras

Randriambolondrantomalala P\*

Department of Mathematics and Computer Science, Faculty of Science, University of Antananarivo, BP 906, 101- Ambohitsaina Antananarivo, Madagascar

### Abstract

We consider a (real or complex) analytic manifold  $M$ . Assuming that  $F$  is a ring of all analytic functions, full or truncated with respect to the local coordinates on  $M$ ; we study the ( $m \geq 2$ )-derivations of all involutive analytic distributions over  $F$  and their respective normalizers.

**Keywords:**  $m$ -derivations; Analytic vector fields Lie algebras; Distributions; Generalized foliations; Stein manifolds; Compact holomorphic manifolds; Chevalley-Eilenberg's cohomology; Compactly supported vector fields

### Introduction and Preliminary

We know several embedding theorems in differential geometry, some of them are of John F. Nash in Riemannian manifolds [1,2], of Whitney [3] in differentiable manifolds and of Grauert in analytic manifolds cf. [4]. They make easy certain study on a differentiable manifold. Here, we are interested to a real or complex analytic  $n$ -dimensional manifold  $M$  and let  $F(M)$  be the ring of all analytic functions on  $M$ . We know that these manifolds can be considered as smooth manifolds. But certain property on a smooth manifold cannot be true on  $M$ , for example the global representation of a smooth function germ theorem. Grabowski had this problem when he studied derivations of the real or complex analytic vector fields Lie algebra cf. [5] and he used Stein manifolds to avoid technical difficulties in them. Here, we examine not only the derivations but the ( $m \geq 2$ )-derivations (generalization of derivation's notion) of a Lie subalgebra of the real or complex analytic vector fields Lie algebra on  $M$ , using Lie algebra tools. In advance, we state that the considered Lie algebras have enough sections more than constant ones in the Lie algebra of all analytic vector fields. Then, we consider only Stein spaces unless expressly stated in a complex analytic case. In the real analytic one, we don't need more hypothesis because of the imbedding theorem of Grauert and Cartan theorems [6]. More precisely, any real analytic manifold can be considered as a closed submanifold of a certain  $\mathbb{R}^l$  (a "real Stein manifold"). Now, an  $m$ -derivation of a Lie algebra  $A$  is a linear map from  $A$  to itself which is distributive on the brackets  $[X_1, [X_2, \dots [X_{m-1}, X_m] \dots]]$ , where all  $X_i$  are in  $A$ . On the one hand, we have studied the  $m$ -derivations of polynomial vector fields Lie algebras on  $\mathbb{R}^n$  in studies of 7. Randriambolondrantomalala [7], an important Lie subalgebra of analytic vector fields, we found that Lie algebras of derivations are different to those of ( $m > 2$ )-derivations. One can see the following example, on  $\mathbb{R}^3$ , the Lie  $\mathbb{R}$ -algebra is spanned by  $\frac{\partial}{\partial x}, \frac{\partial}{\partial y}, \frac{\partial}{\partial z}, x \frac{\partial}{\partial x} + y \frac{\partial}{\partial y} + z \frac{\partial}{\partial z}, y \frac{\partial}{\partial x}, (y)^2 \frac{\partial}{\partial x}$  and let's define the  $\mathbb{R}$ -linear map  $D$  by  $D\left((y)^2 \frac{\partial}{\partial x}\right) = \frac{\partial}{\partial x}$  which is zero otherwise. It's clear that  $D$  is not a derivation, but a 3-derivation. On the other hand, all  $m$ -derivations of a distribution over the full or truncated rings of smooth functions on a differentiable manifold in literature of Randriambolondrantomalala [8], are derivations. These facts lead us to ask if a distribution Lie algebra on an analytic manifold has results as the one or the other above results. So, we will divide our paper into three parts. First, we take a real or complex analytic involutive distribution  $\Omega$  over  $M$ . That is to say,  $\Omega$  is a  $F(M)$ -submodule of the analytic vector fields Lie algebra

$\chi(M)$  on  $M$ . We can find some examples of these distributions and the interests for studying their derivations in literature of Grabowski and Cartan [5,6]. Here, we find the  $\Omega$ 's centralizer and the derivative ideal of  $\Omega$ . We can say also that the normalizer of  $\Omega$  is a Lie subalgebra of analytic vector fields. In addition, we find out that all  $m$ -derivations of  $\Omega$  (resp. of the normalizer of  $\Omega$ ) are inner with respect to a normalizer's vector field (resp. are inner). Second, assuming that  $\Omega$  is an involutive distribution on  $M$  over a subring  $F$  of  $F(M)$ , namely an  $F$ -submodule of  $\chi(M)$  stable by the vector fields bracket, where  $F \neq F(M)$ . One can consider a system of commuting vector fields on  $M$  as in studies of Randriambolondrantomalala [8] and all distribution Lie subalgebras of the Lie algebra of analytic vector fields which commute with this system. The normalizer of  $\Omega$  is an analytic vector fields Lie algebra and contains locally all constant vector fields and Euler's vector field. But in general, we can't use the reasoning by Randriambolondrantomalala [7] to characterize  $m$ -derivations of  $\Omega$ . We make more explicit all  $m$ -derivations of  $\Omega$  and of some of its normalizer. Whereas, in the end, we discuss the Lie algebras of holomorphic vector fields, especially when the holomorphic manifold is not a Stein one, and Lie algebras of locally polynomial vector fields on an analytic manifold  $M$ . Their  $m$ -derivations as well as their normalizers can be characterized by using some results of Randriambolondrantomalala [7].

Therefore, we have found the  $m$ -derivations of all distributions over a set of full or truncated analytic functions with respect to the local coordinates on  $M$ . When  $m = 2$ , we deduce from our results some [5]'s theorems. Third, we can apply our theorems on Lie algebras of real or complex analytic vector fields on  $M$ , of generalized foliation on  $M$  cf. [9], a Lie subalgebra of analytic vector fields on  $S^2$  and on  $\mathbb{T}^2$ , Riemann surfaces, etc. Relations between the Lie algebra of compactly supported vector fields and the compactness of  $M$  are discussed. Moreover, we emphasize the extensions of our theorems when the studied distributions are singular, by using the complexification of a real analytic manifold, Hartogs and Riemann extension theorems. Of course, in these circumstances, we can use theory of coherent sheaves

**\*Corresponding author:** Randriambolondrantomalala P, Department of Mathematics and Computer Science, Faculty of Science, University of Antananarivo, BP 906, 101- Ambohitsaina Antananarivo, Madagascar, Tel: +261331246249; E-mail: [princypcpc@yahoo.fr](mailto:princypcpc@yahoo.fr)

**Received** November 23, 2015; **Accepted** December 17, 2015; **Published** December 19, 2015

**Citation:** Randriambolondrantomalala P (2015) The  $m$ -Derivations of Analytic Vector Fields Lie Algebras. J Generalized Lie Theory Appl S2: 002. doi:10.4172/1736-4337.S2-002

**Copyright:** © 2015 Randriambolondrantomalala P. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

made by Cartan [6] in a Stein case or pass into Grabowski's conjecture cf. [9]. We interpret our results in Chevalley-Eilenberg cohomology sense when  $m = 2$ .

Following the above notations, let  $M$  be a real or complex analytic  $n$ -dimensional manifold. In complex case, we regard a Stein manifold unless special mention. We denote by  $\chi(M)$  the Lie algebra of analytic vector fields on  $M$  and  $F(M)$  the ring of analytic functions on  $M$ . Throughout this paper, we take an atlas in which every chart are connected. Then, the open subset of a chart  $U$  where a non-trivial subset of  $\chi(M)$  doesn't vanish, is dense on  $U$  (non-trivial means different to  $\{0\}$ ). We can use certain results of Randriambolondrantomalala [7,8] because in the proofs of theorem of these papers we consider only analytic functions (polynomials, exponentials). In the same way, we don't need partition of the unity to make global some local results cf. [10]. In all sections of this article, we set an integer  $m \geq 2$ , recall that  $D$  is an  $m$ -derivation of a Lie algebra  $A$  if for  $(X_i)_{1 \leq i \leq m} \subset A$ , we get  $D[X_1, [X_2, \dots, [X_{m-1}, X_m] \dots]] = [D(X_1), [X_2, \dots, [X_{m-1}, X_m] \dots]] + \dots + [X_1, [X_2, \dots, [X_{m-1}, D(X_m)] \dots]]$ . This  $D$  is said inner on a Lie algebra  $\mathbb{B}$  containing  $A$ , if  $D$  is a Lie derivative with respect to an element of  $\mathbb{B}$ . Recall us another basic definition cf. [11].

**Definition 1.1.** A complex manifold  $M$  is a Stein manifold, if we have simultaneously the three following conditions: For every  $x \neq y$ , both in  $M$ ; there is a holomorphic function  $f$  over  $M$  such that  $f(x) \neq f(y)$ . For all  $x \in M$ , it exists  $n$  holomorphic functions  $(f_j)$  over  $M$  such that  $df_j$  are linearly independent over  $\mathbb{C}$  on  $x$ . If  $K$  is a compact set of  $M$ , the following set is compact (holomorphic convexity of  $M$ )

$$\left\{ x \in M \mid |f(x)| \leq \sup_{y \in K} |f(y)|, \text{ for all holomorphic functions } f \text{ over } M \right\}.$$

From these assertions, every local ring of holomorphic functions around  $x \in M$  is spanned by holomorphic functions on  $M$  cf. [12].

Some results of the Lie algebra of compactly supported vector fields  $C_c$  relative to a Stein manifold are the following,

**Proposition 1.2.** A complex analytic manifold  $M$  is compact iff  $C_c$  is non trivial, particularly if  $M$  is a Stein holomorphic manifold,  $C_c$  is trivial.

*Proof.* It's obvious that if  $M$  is compact, then  $C_c = \chi(M)$  and  $C_c$  is not trivial. Conversely, suppose that  $M$  isn't a compact set and there is  $X \in C_c$  such that  $K = \text{Supp}(X) \neq \emptyset$ . We can consider  $K \neq M$  because  $M$  is not compact. Then, we have the nullity of  $X$  in the open set  $C \setminus K \neq \emptyset$ . By analyticity,  $X$  vanishes in whole  $M$ . Hence, we have a contradiction about  $K \neq \emptyset$  and we obtain  $M$  is a compact set. It's clear that a Stein space is never a compact set by definition, then its Lie algebra of compactly supported vector fields is trivial.

### The $m$ -derivations defined by distributions on $F(M)$

Let  $\Omega$  be a non-trivial involutive analytic distribution over the analytic functions ring on  $M$ . Let  $N$  be the normalizer of  $\Omega$  in  $\chi(M)$ , that is to say that the set of all  $X \in \chi(M)$  such that  $[X, \Omega] \subset \Omega$ , and  $\mathfrak{B} = \{x \in M \mid \Omega(x) \neq \{0\}\}$ . We can choose a connected domain  $U_i$  of a chart. If we suppose that it exists an open set  $O_i$  in  $U_i$  where  $\Omega$  vanishes, then  $\Omega_{U_i} = \{0\}$  by analyticity. Otherwise, every open set in  $U_i$  contains an element of  $\mathfrak{B}$ . So,  $\mathfrak{B} \cap U_i$  is dense over  $U_i$ . Moreover, the collection of  $U_i$  forms an atlas of  $M$ , then  $\mathfrak{B}$  is dense over  $M$ . The set  $\mathfrak{B}$  is an open analytic submanifold of  $M$ . Particularly,  $\mathfrak{B}$  is a Stein cf. [13]. Thus, every vector field defined over  $\mathfrak{B}$  admits a continuous extension on  $M$ , and if this last one is analytic, then it's necessarily an element of the normalizer of  $\Omega$ . We use this last fact when we deal with extension theorems.

We know by literature of Nagano's [14] result that  $\Omega$  is integrable, then it yields a generalized foliation  $F$  on  $M$  cf. [10]. So,  $\Omega$  is the Lie algebra of tangent vector fields to the foliation and  $L_F$  the one of all foliation preserving vector fields. It is known that the normalizer  $N$  in  $\chi(M)$  of  $\Omega$  contains  $L_F$  cf. [10]. Hence, the restriction of the foliation in  $\mathfrak{B}$  is non singular.

**Proposition 2.1.** The centralizer of  $\Omega$  vanishes and the derivative ideal of  $\Omega$  coincides with  $\Omega$  itself.

*Proof.* We say that  $X \in \chi(M)$  is in the centralizer of  $\Omega$  if  $[X, \Omega] = \{0\}$ ; and the derivative ideal of denoted by  $[\Omega, \Omega]$  is the Lie algebra spanned by all brackets of two elements of  $\Omega$ . Suppose there is a non zero element  $X$  of the centralizer, we have  $[X, f\Omega] = (X(f))\Omega = \{0\}$ , for all  $f \in F(M)$ . It's not possible in a Stein manifold or in a real analytic manifold if  $X$  doesn't vanish identically over  $M$  and if  $\Omega \neq \{0\}$ . Along with this result, we can adapt the proof of Proposition 2.28 of studies of Randriambolondrantomalala [15] and assert that  $[\Omega, \Omega] = \Omega$ .

Let's recall an Hartogs's extension theorem and Riemann extension theorem.

**Theorem 2.2.** (Hartogs [16]) Let be  $t \geq 2$  and  $D$  be a bounded domain in  $\mathbb{C}^t$ . In addition,  $K$  be a compact subset of  $D$  such that  $D - K$  is a connected set. Then all holomorphic functions  $f$  over  $D - K$  can be extended holomorphically to  $D$ .

**Theorem 2.3.** (Riemann extension theorem) Let  $U$  be an open set in  $\mathbb{C}$  and  $z_0 \in U$ . If  $f : U - \{z_0\} \rightarrow \mathbb{C}$  is holomorphic function such that  $z_0$  is a removable singularity of  $f$ , then  $f$  can be extended into a unique holomorphic function  $\bar{f}$  in  $U$  where  $\bar{f}(z_0) = \lim_{z \rightarrow z_0} f(z)$ .

**Theorem 2.4.** In holomorphic case, all  $m$ -derivations of  $\Omega$ ,  $L_{\mathfrak{B}}$  and of  $N$  are Lie derivatives with respect to elements belonging to  $N$ . In real analytic one, we have the same results if  $\mathfrak{B} = M$ .

*Proof.* We can prove this assertion over  $\mathfrak{B}$  by Theorem 2.1 of studies of Randriambolondrantomalala [8] using Proposition 2.1 and partially Theorem 3.7 of literature of Randriambolondrantomalala [10]. For the corresponding extension theorem over  $M$ , we adopt the following arguments. We know that  $\mathfrak{B}$  is dense over  $M$ , then the restriction of  $\mathfrak{B}$  in each domain of a chart  $U$  is dense over  $U$  ( $U$  is a bounded set). The complement of this  $\mathfrak{B} \cap U$  in  $U$  can be considered as a compact set of the chart such that  $\mathfrak{B} \cap U$  is connected. In holomorphic case, when  $n \geq 2$ , we use Hartogs's theorem in a domain of the chart, so the extension theorem over  $M$  holds. If  $n = 1$ , we know by the isolated zeros principle that the domain of chart contains only a finite number of zeros in the corresponding restriction of  $\mathfrak{B}$ . By continuity at these zeros, which are removable singularities, the Riemann extension theorem can be used. Of course, if  $\mathfrak{B} = M$  in real analytic situation, the extension theorem is applicable.

**Definition 2.5.** The complexification of a real analytic manifold  $M$  is a holomorphic manifold  $\mathbb{H}$  such that there is a real analytic embedding  $f: M \rightarrow \mathbb{H}$  where  $\mathbb{H}$  has a holomorphic atlas  $(U_i, \varphi_i)$ , and  $\varphi_i(f(M) \cap U_i) = \varphi_i(U_i) \cap \mathbb{R}^n$ . We have a Stein complexification if  $\mathbb{H}$  is Stein.

The next theorem is due by Grauert cf. [4,12].

**Theorem 2.6.** Every real analytic manifold has a Stein complexification and can be analytically properly embedded into an Euclidean space  $\mathbb{R}^n$ .

The following complexification of a Lie subalgebra  $\mathbb{G}$  of the real analytic vector fields Lie algebra of  $M$  is in the following sense: if  $M$  can

be embedded in a holomorphic manifold  $\mathbb{H}$ , the complexification  $\overline{\mathbb{G}}$  of  $\mathbb{G}$  in  $\mathbb{H}$  is such that  $\overline{\mathbb{G}}_M = \mathbb{G}$ .

**Theorem 2.7.** *If the complexification of  $\Omega$  in a Stein space  $T$  is still an involutive distribution, then the first result of the Theorem 2.4 holds in real analytic case.*

*Proof.* We use the complexification of  $M$  on a Stein space  $T$ . Consequently, let be  $\overline{\Omega}$  the complexification over  $T$  of  $\Omega$ . Recall that  $\overline{\Omega}$  is an involutive distribution over  $T$  where its normalizer on  $\chi(T)$  is denoted by  $N_{\overline{\Omega}}$ . So, all  $m$ -derivations of  $\Omega$ , of  $L_{\overline{\Omega}}$ , and of  $N$  have their complexified  $m$ -derivations over  $T$  on respectively  $\overline{\Omega}$ , the Lie algebra of all foliation preserving vector fields  $\overline{L}_{\overline{\Omega}}$  on  $T$  and  $N_{\overline{\Omega}}$ . By the results of Theorem 2.4, these last  $m$ -derivations are Lie derivatives with respect to elements belonging to  $N_{\overline{\Omega}}$ . We can affirm that  $\mathfrak{N}_{\overline{\Omega}M} = \mathfrak{N}$  and  $\overline{L}_{\overline{\Omega}M} = L_{\overline{\Omega}}$  by  $\overline{\Omega}_M = \Omega$ . Thus, we have the same result as in the first part of Theorem 2.4.

By definition, the first space of Chevalley-Eilenberg's cohomology of a Lie algebra  $\mathcal{A}$  denoted by  $H^1(\mathcal{A})$  is  $\text{Der}(\mathcal{A})/\text{ad}_{\mathcal{A}}$  with  $\text{Der}(\mathcal{A})$  the Lie algebra of all derivations of  $\mathcal{A}$  and  $\text{ad}_{\mathcal{A}}$  the set of inner derivations of  $\mathcal{A}$ .

Throughout this paper, we suppose that all hypothesis of Theorem 2.7 are satisfied or  $\mathfrak{B} = M$ , in real analytic case.

Following ideas of Theorem 3.7, Corollary 2.14 and Remark 2.15 of literature of Randriambololondrantomalala [10], we state

**Corollary 2.8.** *The first space of Chevalley-Eilenberg's cohomology of  $\Omega$ ,  $L_{\overline{\Omega}}$  and of  $\mathfrak{N}$  is respectively isomorphic to the following respective Lie algebras,  $\mathfrak{N}/\Omega$ ,  $\mathfrak{N}/L_{\overline{\Omega}}$ ,  $\{0\}$ .*

**Remark 2.9.** *By Theorem 2.4, we deduce Theorem 3.2 and 4.1 of studies of Grabowski [5].*

### The $m$ -derivations associated to a distribution over a subring of $F(M)$

Let be an atlas of  $M$  such that  $\Omega$  is locally spanned by  $\left(\frac{\partial}{\partial x^i}\right)_{1 \leq i \leq n}$  over the ring  $F_0$  of real or complex functions depending only on  $(x^j)_{1 \leq j \leq k}$  with respect to the atlas (where  $1 \leq k < n$ ). We can consider  $\Omega$  to be a Lie algebra which commutes with a system  $S$  of commuting vector fields by the usual bracket. That is to say,  $S = \{X_1, \dots, X_q\}$  such that  $[X_i, X_j] = 0$  and  $S$  is locally of rank  $n - k$  ( $0 < q \leq n$ ). It is easy to check that  $[\Omega, \Omega] = \Omega$  because of Randriambololondrantomalala's [8] result. So with the same reason, every  $m$ -derivation of  $\Omega$  is local. Moreover, the normalizer  $N$  of  $\Omega$  is locally isomorphic to  $\Omega \oplus gl(n - k, \mathbb{R}$  or  $\mathbb{C})$  as a vector space. We consider the closed 1-differential forms  $\alpha^i$  and  $\omega^j$  over a  $(n - k)$ -dimensional distinguished connected chart of the generalized foliation generated by  $S$ , where  $i = k + 1, \dots, n$  and an  $m$ -derivation of

$$\Omega, \quad D^{(\alpha, \omega)} = (\alpha^j + \omega^j) \otimes \frac{\partial}{\partial x^j} \quad \text{such that} \quad \ker(\omega^j) \supset F_0(U) \left\langle \frac{\partial}{\partial x^i} \right\rangle_{1 \leq i \leq k} \quad \text{and}$$

$$\ker(\alpha^i) \supset F_0(U) \left\langle \frac{\partial}{\partial x^i} \right\rangle_{k+1 \leq i \leq n} \quad (S \text{ in this chart is } \left\{ \frac{\partial}{\partial x^i} \right\}_{k+1 \leq i \leq n}, \quad F \langle A \rangle$$

denotes a module spanned by  $A$  over a ring  $F$  cf. [8]. We have omitted all singular charts of the foliation because the open set  $R$  of all regular points is dense over  $M$  cf. [10], we have no problem for the extension of our results from  $R$  towards  $M$  as in the previous section. By adapting Theorem 3.12 of literature of Randriambololondrantomalala [8], we obtain easily

**Theorem 3.1.** *All  $m$ -derivations of  $\Omega$  (resp. of  $\mathfrak{N}$ ) are a sum of a Lie derivative with respect to one element of  $\mathfrak{N}$  and a derivation  $D^{(\alpha, \omega)}$  as denoted before (resp. are similar to  $m$ -derivations of  $\Omega$ ).*

Hence, adopting the arguments of Theorem 3.19 of studies of Ravelonirina [17], we hold the following

**Corollary 3.2.** *When the rank of  $S$  is a positive constant  $n - k$ , the first spaces of Chevalley-Eilenberg's cohomology of  $\Omega$  and of  $N$  are both isomorphic to  $(H_R(B))^{n-k} \times (\mathbb{R}^{(n-k)^2+n-k}$  or  $\mathbb{C}^{(n-k)^2+n-k})$  with  $H_R(B)$  is the de Rham cohomology of foliation basic forms of  $M$ .*

As we know, we can split the above  $\Omega$  into a semi-direct sum of Lie algebras  $\Omega_S^1$  and  $\Omega_S^2$  as in studies of Randriambololondrantomalala [8], where they are modules on the ring  $F_0(M)$  of constant functions over the leaves relative to the above generalized foliation. We notice that  $\Omega_S^2$  is spanned by  $S$  on  $F_0(M)$ . We can reason on a distinguished chart  $U$  with the coordinates  $(x^i)_{1 \leq i \leq n}$ . The  $F_0(U)$  is the set of all analytic functions depending only on  $(x^i)_{1 \leq i \leq k}$ ,  $\Omega_{S|U}^1$  is spanned by  $\left(\frac{\partial}{\partial x^i}\right)_{1 \leq i \leq k}$  and  $\Omega_{S|U}^2$  by  $\left(\frac{\partial}{\partial x^i}\right)_{k+1 \leq i \leq n}$  on  $F_0(U)$ .

Now, we discuss the  $m$ -derivations of  $\Omega_S^1$ . The normalizer  $N^1$  of this Lie algebra can be written as a direct sum of Lie algebras  $\mathfrak{N}^1 = [\Omega_S^1, \Omega_S^1] \oplus [Q, Q]$ , where  $Q$  is the centralizer of  $\Omega_S^1$  in  $\chi(M)$  and the center of  $\Omega_S^1$  is zero ( $Q$  is locally spanned by  $\left(\frac{\partial}{\partial x^i}\right)_{k+1 \leq i \leq n}$  on the ring of all analytic functions depending only on  $(x^i)_{k+1 \leq i \leq n}$ ). By a similar argument of Nakanishi [18], we deduce that all  $m$ -derivations of  $N^1$  are a direct sum of those of  $\Omega_S^1$  and of  $Q$ . By Theorem 2.4, it's clear that

**Theorem 3.3.** *Each  $m$ -derivation of  $\Omega_S^1$  (resp. of  $\mathfrak{N}^1$ ) is a Lie derivative with respect to an element of  $N^1$ .*

The normalizer of  $\Omega_S^2$  is locally the sum of the  $F_0(U)$ -module spanned by  $\frac{\partial}{\partial x^i}$  and a vector space generated by  $x^j \frac{\partial}{\partial x^i}$ ,  $k+1 \leq i, l \leq n$ . That is to say, its normalizer is  $\mathfrak{N}$ . In addition, its centralizer is  $\Omega_S^2$  itself. Because of  $[\Omega_S^2, \Omega_S^2] = \{0\}$  or  $\Omega_S^2$  is nilpotent of order 1, we obtain easily

**Theorem 3.4.** *Every endomorphism of  $\Omega_S^2$  is an  $m$ -derivation of  $\Omega_S^2$ .*

So, it's immediate that

**Corollary 3.5.** *The first space of Chevalley-Eilenberg's cohomology of  $\Omega_S^1$ ,  $\Omega_S^2$  and of  $N^1$  are respectively isomorphic to the following respective Lie algebras,  $\mathfrak{N}^1 / \Omega_S^1$ ,  $\text{End}(\Omega_S^2) / \Omega_S^2$ ,  $\{0\}$ .*

Let's consider  $\Omega_S^3$  the Lie subalgebra of  $\Omega$ , spanned by  $X_i$  over a ring  $F \subseteq F(M)$ . That is to say,  $F$  is locally the set of all analytic functions depending only on  $(x^i)_{1 \leq i \leq n}$  where  $0 < l < k + 1$  (resp.  $k + 1 < l < n + 1$ ). When  $F = F(M)$  (resp.  $F = F_0(M)$ ), it is a special case of Lie algebras defined in Theorem 2.4 (resp. in Theorem 3.1) when the submodule is generated by  $X_i$ . In the distinguished local coordinates,  $\Omega_S^3$  is spanned by  $\left(\frac{\partial}{\partial x^i}\right)_{k+1 \leq i \leq n}$  over  $F$ . The normalizer  $\mathfrak{N}^3$  of  $\Omega_S^3$  coincides with the sum of  $\Omega_S^3$  and  $\Omega_S^4$  where the element of this last one is locally the following analytic vector fields  $f^i(x^j, 1 \leq t \leq k) \frac{\partial}{\partial x^i} + g^j(x^t, 1 \leq t \leq k) \frac{\partial}{\partial x^j}$ ,  $1 \leq i \leq k$ ,  $1 \leq j \leq l - 1$ .

$\left( \text{resp. } f^l(x^j, 1 \leq t \leq k) \frac{\partial}{\partial x^t} + h^l(x^j, l \leq t \leq n) x^j \frac{\partial}{\partial x^l} \right)$ . In the first case, we can

adapt Theorem 2.4 because all analytic functions depending on  $(x^j)$  where  $k+1 \leq j \leq n$ , are in the base ring of  $\Omega_S^3$ . In the following case, it is easy to see that Theorem 3.1 can be adapted to  $\Omega_S^3$ . Thus

**Theorem 3.6.** *In the first case, every  $m$ -derivation  $D$  of  $\Omega_S^3$  is a Lie derivative with respect to a  $\mathfrak{N}^3$ 's element; in the second, it is a sum of a Lie derivative of an element of  $\mathfrak{N}^3$  and a  $D^{(a,w)}$  analogous to that of Theorem 3.1. In addition, the corresponding extension theorems hold.*

**Corollary 3.7.** *The first space of Chevalley-Eilenberg's cohomology of  $\Omega_S^3$  is respectively isomorphic to  $\Omega_S^4$  in the first case;*

$$\Omega_S^4 \oplus \left( (H_R(B'))^{l-k-1} \times (\mathbb{R}^{(l-k)(l-k-1)} \text{ ou } \mathbb{C}^{(l-k)(l-k-1)}) \right)$$

in the other one if  $S$  has a constant rank ( $\oplus$  is a module direct sum and  $B'$  is the set of the corresponding foliation basic forms of  $M$ ).

When we regard all the above normalizers on a distinguished chart, they contain locally all constant fields and Euler vector field. So, we ask one question: could we adapt Theorem 3.6 and Theorem 3.9 in [7] to these normalizers? The following remark shows us that this argument is false.

**Remark 3.8.** On  $\mathbb{R}^3$ , we consider the Lie  $\mathbb{R}$ -algebra  $A$  spanned by  $\frac{\partial}{\partial x}, \frac{\partial}{\partial y}, \frac{\partial}{\partial z}, x \frac{\partial}{\partial x} + y \frac{\partial}{\partial y} + z \frac{\partial}{\partial z}, y \frac{\partial}{\partial x}, (y^2) \frac{\partial}{\partial x}, e^y \frac{\partial}{\partial x}, y e^y \frac{\partial}{\partial x}, y^2 e^y \frac{\partial}{\partial x}, \dots$

Lemma 2.3 of literature of Randriambolondrantomalala [10] is not true for  $A$ , so the arguments proposed in the proof of Theorem 3.6 of Princy [7] don't hold in this situation.

Whereas, let  $P$  be a Lie subalgebra consisting locally of polynomial vector fields in  $\chi(M)$ , where the Euler vector field and all constant vector fields are locally in  $P$ . Especially,  $M$  is not supposed to be a Stein in the holomorphic case. Let's recall a well known theorem,

**Theorem 3.9.** *(The maximum principle) [12] Let be  $M$  a connected holomorphic manifold and  $f$  a holomorphic function on  $M$  such that  $|f(z)| \leq |f(z_0)|$ , where  $z_0 \in M$  for all  $z \in M$ ; then  $f$  is a constant function.*

One consequence of the maximum principle is the following, if the holomorphic manifold  $M$  is compact, every holomorphic function on  $M$  is constant in every connected component of  $M$ . We know that  $M$  is locally connected, then each function over  $M$  is locally constant. Therefore, it's clear that if  $M$  is a compact and connected holomorphic manifold, the ring of all holomorphic functions on  $M$  is the complex constant functions ring. It's confirm that results of the following theorem complete our study about an involutive analytic distribution when  $F(M)$  is reduced to  $\mathbb{C}$ .

By adapting Randriambolondrantomalala's [7] theorems and taking account that the vector field found in the proof of Theorem 3.6 of Princy [7] is analytic, it follows that

**Theorem 3.10.** *When  $m$  is even, all  $m$ -derivations of  $\mathfrak{P}$  (resp. of the normalizer  $\mathcal{N}$  of  $\mathfrak{P}$  in  $\chi(M)$ ) are a Lie derivative with respect to one and only one vector field belonging to  $\mathcal{N}$  (resp. to the normalizer of  $\mathcal{N}$  in  $\chi(M)$ ). If  $m$  is an odd number, they are sum of a Lie derivative with respect to one element of  $\mathcal{N}$  (resp. of the normalizer of  $\mathcal{N}$ ) and an  $m$ -derivation of local homogeneous degree  $-2$  of  $\mathfrak{P}$ .*

So, taking into account: the vanishing of the centralizer of  $\mathfrak{P}$  cf. [19] p.91; both the proofs of Theorem 2.12 of Ravelonirina [19],

Corollary 3.12 of Randriambolondrantomalala [7] and Theorem 3.7 in literature of Randriambolondrantomalala [10], we obtain

**Corollary 3.11.** *The first space of Chevalley-Eilenberg's cohomology of  $\mathfrak{P}$ , of  $\mathcal{N}$  and of  $\mathfrak{N}$  is respectively isomorphic to the following respective Lie algebras  $\mathcal{N} / \mathfrak{P}, \mathfrak{N} / \mathcal{N}, \{0\}$ , where  $\mathfrak{N}$  is the normalizer of  $\mathcal{N}$ .*

### Illustrations

Some illustrations of our theorems are given in this section.

**Example 4.1.** It's clear that Theorem 2.4 works well on the Lie algebra of all analytic vector fields  $\chi(M)$ , that is to say, all  $m$ -derivations of  $\chi(M)$  are Lie derivatives by elements of  $\chi(M)$ . We can define the Lie algebra of compactly supported real analytic vector fields  $\mathbb{C}$  and this theorem holds for this last one. In particular,  $H^1(\mathbb{C}) = \chi(M) / \mathbb{C}$  for a non-trivial  $\mathbb{C}$ . More,  $H^1(\mathbb{C}) = \{0\}$  if and only if  $M$  is compact. Obviously, we can use the above cited theorem on the Lie algebras of vector fields relative to a generalized foliation over  $M$ . We can cite some well known Stein spaces,  $\mathbb{C}^n$ , an open poly-disc in  $\mathbb{C}^n$ , non-compact Riemann surfaces, ... and build our results in these.

**Example 4.2.** Let be  $\mathbb{S}^2$  a holomorphic compact connected manifold. It's not a Stein manifold nor a submanifold of  $\mathbb{C}^n$  for any  $n$ , it's a compact Riemann surface. We choose the modified stereographic coordinates over this manifold. The  $\mathbb{S}^2$  has an atlas composed by two charts  $(U, z_1)$  and  $(V, z_2)$  with the overlap map  $\psi(z) = z^{-1}$  in  $U \cap V$ . We remark that the Lie algebra  $\mathbb{L}$  of vector fields on  $M$  spanned over  $\mathbb{C}$  by  $Y_1, Y_2$  and  $Y_3$  is the one of all polynomial vector fields in  $\mathbb{S}^2$ , where

$$Y_1 : \begin{cases} \frac{\partial}{\partial z_1} & \text{in } U \\ -(z_2)^2 \frac{\partial}{\partial z_2} & \text{in } V \end{cases}, \quad Y_2 : \begin{cases} -(z_1)^2 \frac{\partial}{\partial z_1} & \text{in } U \\ \frac{\partial}{\partial z_2} & \text{in } V \end{cases}, \quad Y_3 : \begin{cases} -z_1 \frac{\partial}{\partial z_1} & \text{in } U \\ z_2 \frac{\partial}{\partial z_2} & \text{in } V \end{cases}.$$

By Theorem 3.10, all  $m$ -derivations of  $\mathbb{L}$  are Lie derivatives with respect to a vector field in  $\mathbb{L}$  itself. That is to say, if  $D$  is an  $m$ -derivation of  $\mathbb{L}$  defined by  $D(Y_1) = \alpha^1 Y_1, D(Y_2) = \beta^1 Y_1, D(Y_2) = \gamma^1 Y_1$ ; we have  $D = L_{-\gamma^1 Y_1 + \beta^1 Y_2 + \alpha^1 Y_3}$ .

When we look at  $\mathbb{S}^2$  as a real analytic manifold, we set the charts  $(U, (x^1, x^2))$  and  $(V, (y^1, y^2))$  with the overlap map

$$\phi(x^1, x^2) = \left( y^1 = \frac{x^1}{(x^1)^2 + (x^2)^2}, y^2 = \frac{x^2}{(x^1)^2 + (x^2)^2} \right).$$

We set the real analytic vector field  $Y_3 : \begin{cases} -x^1 \frac{\partial}{\partial x^1} - x^2 \frac{\partial}{\partial x^2} & \text{in } U \\ y^1 \frac{\partial}{\partial y^1} + y^2 \frac{\partial}{\partial y^2} & \text{in } V \end{cases}$  on  $\mathbb{S}^2$

and the Lie algebra  $\mathfrak{A}$  of real analytic vector fields which commute with  $Y_3$ . This  $\mathfrak{A}$  consists of real analytic vector fields  $Y$  such that

$$Y : \begin{cases} F_1 \left( \frac{x^2}{x^1} \right) x^1 \frac{\partial}{\partial x^1} + F_2 \left( \frac{x^1}{x^2} \right) x^2 \frac{\partial}{\partial x^2} & \text{in } U \\ F_3 \left( \frac{y^2}{y^1} \right) y^1 \frac{\partial}{\partial y^1} + F_4 \left( \frac{y^1}{y^2} \right) y^2 \frac{\partial}{\partial y^2} & \text{in } V \end{cases}$$

where in the  $U \cap V$ ,

$$F_3 \left( \frac{y^2}{y^1} \right) = \frac{1}{(y^1)^2 + (y^2)^2} \left( \left( -2F_2 \left( \frac{y^1}{y^2} \right) + F_1 \left( \frac{y^2}{y^1} \right) \right) (y^2)^2 - F_1 \left( \frac{y^2}{y^1} \right) (y^1)^2 \right) \text{ and}$$

$$F_4\left(\frac{y^1}{y^2}\right) = \frac{1}{(y^1)^2 + (y^2)^2} \left( \left( -2F_1\left(\frac{y^2}{y^1}\right) + F_2\left(\frac{y^1}{y^2}\right) \right) (y^1)^2 - F_2\left(\frac{y^1}{y^2}\right) (y^2)^2 \right)$$

with  $F_k$  are arbitrary convenient functions of one variable. So, we can apply all theorems in Section 3 to  $\mathfrak{A}$ . Particularly  $H^1(\mathbb{L}) = \{0\}$  and  $H^1(\mathfrak{A}) = H^1_R(B) \times \mathbb{R}^2$ .

**Example 4.3.** Indeed, Theorem 3.10 can be applied to a polynomial vector fields Lie algebra on the real analytic manifold  $\mathbb{R}^n$  or the Stein manifold  $\mathbb{C}^n$  having the corresponding hypothesis.

**Example 4.4.** We set the Lie algebra  $\mathbb{A}$  over the Stein manifold  $\mathbb{C}^3$  spanned over  $\mathbb{C}$  by  $\frac{\partial}{\partial z^1}, \frac{\partial}{\partial z^2}, \frac{\partial}{\partial z^3}, z^1 \frac{\partial}{\partial z^1} + z^2 \frac{\partial}{\partial z^2} + z^3 \frac{\partial}{\partial z^3}, z^2 \frac{\partial}{\partial z^1}, (z^2)^2 \frac{\partial}{\partial z^1}$ . The normalizer of  $\mathbb{A}$  is  $\mathcal{N}_0 = \mathbb{A} + R$ , where  $R$  is the space over  $\mathbb{C}$  generated by  $z^1 \frac{\partial}{\partial z^1}, z^2 \frac{\partial}{\partial z^2}, z^3 \frac{\partial}{\partial z^3}, z^2 \frac{\partial}{\partial z^3}, z^3 \frac{\partial}{\partial z^1}$ . It is permit to use Theorem 3.10 and when  $m$  is even, every  $m$ -derivation of  $\mathbb{A}$  is inner on  $\mathcal{N}_0$ . If  $m$  is odd, the  $m$ -derivation is a sum of an inner derivation on  $\mathcal{N}_0$  with a  $\mathbb{C}$ -linear map  $D$  defined by  $D\left((z^2)^2 \frac{\partial}{\partial z^1}\right) = \alpha \frac{\partial}{\partial z^1}$  which is zero otherwise ( $\alpha \in \mathbb{C}$ ). Moreover, all  $m$ -derivations of  $\mathcal{N}_0$  are inner for all  $m \geq 2$ . So,  $H^1(\mathbb{A}) \cong R \ominus \left\langle z^1 \frac{\partial}{\partial z^1} + z^2 \frac{\partial}{\partial z^2} + z^3 \frac{\partial}{\partial z^3} \right\rangle$  and  $H^1(\mathcal{N}_0) \cong \{0\}$ .

**Remark 4.5.** In the following example, Theorem 3.10 cannot be applied. We take the 2-torus  $\mathbb{T}^2 = \mathbb{C}/(\mathbb{Z} + i\mathbb{Z})$ , which is a holomorphic connected compact manifold cf. [20], it's not a Stein. All overlap maps are translations, that is to say, they are holomorphic. We can define globally the Lie algebra of all constant vector fields  $\mathfrak{Q}$  on  $M$  and find that  $\mathfrak{Q}$  is the Lie algebra of all holomorphic vector fields over  $M$ . All endomorphisms of each Lie subalgebra of  $\mathfrak{Q}$ , which is inevitably nilpotent of order 1, are  $m$ -derivations of this subalgebra. The normalizer of this subalgebra or its centralizer is the Lie algebra of all vector fields over  $\mathbb{T}^2$ . But  $H^1(\mathfrak{Q}) \cong \text{End}(\mathfrak{Q})/\text{ad}_{\mathfrak{Q}}$  and  $H^1(\mathcal{X}(M)) \cong \text{End}(\mathcal{X}(M))/\text{ad}_{\mathcal{X}(M)}$  since  $H^1(\mathcal{X}(M)) = \{0\}$  in smooth cases.

## References

- Nash J (1954)  $C^1$ -isometric imbeddings. Annals of Mathematics 60: 383-396.
- Nash J (1956) The imbedding problem for Riemannian manifolds. Annals of Mathematics 63: 20-63.

- Whitney H (1936) Differentiable manifolds. The Annals of Math 37: 645-680.
- Grauert H (1958) On Levi's problem and the imbedding of real analytic manifolds. Ann of Math 68: 460-472.
- Grabowski J (1981) Derivations of the Lie algebras of analytic vector fields. Compositio Mathematica 43: 239-252.
- Cartan H (1957) Real analytic varieties and complex analytic varieties. Bulletin de la S.M.F 85: 77-99.
- Randriambolondrantomalala P (2014) Les  $m$ -dérivations des algèbres de Lie de champs de vecteurs polynomiaux. In final revision at Afrika Matematika.
- Randriambolondrantomalala P (2015) The  $m$ -derivations of distribution Lie algebras. Journal of Generalized Lie Theory and Applications 9: 1.
- Grabowski J (1993) Lie algebras of vector fields and generalized foliations. Publicacions Matemàtiques 37: 359-367.
- Randriambolondrantomalala P, Ravelonirina HSG, Anona M (2010) Of the Sur les algèbres de Lie d'une distribution et d'un feuilletage généralisé. African Diaspora Journal of Mathematics 10: 135-144.
- Stein K (1951) Analytic functions of several complex variables at predetermined periodicity and the second Cousin problem. Math Ann 123: 201-222.
- Fosternic F (2011) Stein manifolds and holomorphic mappings. Results of mathematics and its border areas. Folge/A Series of Modern Surveys in Mathematics, Springer-Verlag Berlin Heidelberg 56.
- Fritzsche K, Grauert H (2002) From holomorphic functions to complex manifolds. Graduate texts in Mathematics, Springer-Verlag, New York.
- Nagano T (1966) Linear differential systems with singularities and an application to transitive Lie algebras. J Math Soc Japan 18: 398-404.
- Randriambolondrantomalala P (2010) Sur les algèbres de Lie d'une distribution et d'un feuilletage généralisé. Phd thesis, Université d'Antananarivo.
- Hartogs F (1906) Some consequences of the Cauchy integral formula for functions of several variables. Munch Ber 36: 223-242.
- Ravelonirina HSG, Randriambolondrantomalala P, Anona M (2012) Of the Sur les algèbres de Lie d'un système de champs de vecteurs permutables. Italian Journal of Pure and Applied Mathematics 29: 163-174.
- Nakanishi N (1982) The first cohomology groups of infinite dimensional Lie algebras. Pub RIMS. Kyoto Univ 18: 1-16.
- Ravelonirina HSG, Randriambolondrantomalala P, Anona M (2010) Sur les algèbres de Lie des champs de vecteurs polynomiaux. African Diaspora Journal of Mathematics 10: 87-95.
- Bouche T (1996) Introduction to differential geometry of complex analytic manifolds. Ecole d'été de Mathématiques, Université Joseph Fourier.

Citation: Randriambolondrantomalala P (2015) The  $m$ -Derivations of Analytic Vector Fields Lie Algebras. J Generalized Lie Theory Appl S2: 002. doi:10.4172/1736-4337.S2-002

This article was originally published in a special issue, **Recent Advances of Lie Theory in differential Geometry, in memory of John Nash** handled by Editor. Dr. Princy Randriambolondrantomalala, University of Antananarivo, Madagascar

## OMICS International: Publication Benefits & Features

### Unique features:

- Increased global visibility of articles through worldwide distribution and indexing
- Showcasing recent research output in a timely and updated manner
- Special issues on the current trends of scientific research

### Special features:

- 700 Open Access Journals
- 50,000 editorial team
- Rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at PubMed (partial), Scopus, EBSCO, Index Copernicus and Google Scholar etc.
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.omicsonline.org/submit>