

Matrix Representation for Seven-Dimensional Nilpotent Lie Algebras

Ghanam R^{1*}, Basim Mustafa B², Mustafa MT³ and Thompson G⁴

¹Department of Liberal Arts and Sciences, Virginia Commonwealth University in Qatar, Qatar

²Department of Mathematics, An-Najah National University, Palestine

³Department of Mathematics, Statistics and Physics, Qatar University, Qatar

⁴Department of Mathematics, University of Toledo, USA

Abstract

This paper is concerned with finding linear representations for seven-dimensional real, indecomposable nilpotent Lie algebras. We consider the first 39 algebras presented in Gong's classification which was based on the upper central series dimensions. For each algebra, we give a corresponding matrix Lie group, a representation of the Lie algebra in terms of left-invariant vector field and left-invariant one forms.

Keywords: Lie algebra; Lie group; Representation, Nilpotent

Introduction

Given a real Lie algebra \mathfrak{g} of dimension n a well known theorem due to Ado [1,2] asserts that \mathfrak{g} has a faithful representation as a subalgebra of $gl(p, R)$ for some p . The theorem does not give much information about the value of p but leads one to believe that p may be very large in relation to the size of n and consequently it seems to be of limited practical value. In previous work we have found linear representation for all indecomposable real Lie algebras in dimension six or less [3-5] In this paper we are concerned with finding linear representations for seven-dimensional nilpotent Lie algebras. Let \mathfrak{g} be a nilpotent Lie algebra, we shall assume throughout that \mathfrak{g} is indecomposable in the sense that \mathfrak{g} is not isomorphic to a direct sum of two proper nilpotent ideals. If \mathfrak{g} is decomposable and we have representations for both factors then we can easily find a representation for \mathfrak{g} . It is well known that a nilpotent Lie algebra has a non-trivial center and so finding representations is difficult because the adjoint representation is not faithful. In this article, we consider the list of seven-dimensional nilpotent Lie algebras over R classified by Gong [6], his list contained 147 indecomposable Lie algebras that are classified according to the upper central series. We consider the first 39 cases for which the dimensions of the upper central series are (27) (37) (247) (257) and (357). For each algebra \mathfrak{g} we give a corresponding Lie group that is a subgroup of $GL(7, R)$. The representation for the Lie algebra is then easily obtained by differentiating and evaluating at the identity. The structure of the paper is as follows: in Section 2, we give a brief history and background on the classification of the seven-dimensional nilpotent Lie algebras, In section 3, we give a matrix Lie group that corresponds to each Lie algebra. We also give a representation of the Lie algebra in terms of left-invariant one-forms and left-invariant vector fields. Throughout the paper we will use (p, q, r, x, y, z, w) as local coordinates on Lie groups.

Classifying Nilpotent Lie Algebras in Dimension Seven

Classification of solvable Lie algebras is not an easy problem, this is due to the fact the solvable Lie algebras are unlike semisimple Lie algebras. Semisimple Lie algebras are considered very beautiful since over the complex numbers we have the Killing form, Dynkin diagrams, root space decompositions, the Serre representation, the theory of highest weight representation, the Weyl character formula and much more [7-10]. On the other hand, we have Lie, Engel and Ado's theorems for solvable Lie algebras. There has been several attempts to classify the seven-dimensional nilpotent Lie algebras, we will mention the most recent ones that we are aware of. In 1993, Seeley [10] gave a

classification of over the field of complex numbers. His classification was based on the upper central series of the Lie algebras and knowledge of all lower dimensional nilpotent Lie algebras. His list contained 161 Lie algebras; 130 indecomposable and 31 decomposable. In 1998, Gong [8], presented a new list of seven-dimensional nilpotent Lie algebras. Gong's classification was based on the Skjelbred-Sund method [11]. Gong provided a classification of the seven-dimensional indecomposable algebras over Algebraically closed fields ($\chi \neq 2$) and another classification over the field of real numbers. Once again, he used the same labeling as Seeley; the dimensions of the upper central series.

Representations

In this section, we present a matrix Lie group representation for the first 39 seven-dimensional Lie algebras with upper central series (27) (37) (247) (257) and (357)

Algebras with upper central series dimensions (27)

$$1. (27A) : [x_1, x_2] = x_6, [x_1, x_4] = x_7, [x_3, x_5] = x_7;$$

A matrix Lie group is given by:

$$S = \begin{pmatrix} 1 & x & y & r & p & q \\ 0 & 1 & 0 & 0 & w & 0 \\ 0 & 0 & 1 & 0 & -z & 0 \\ 0 & 0 & 0 & 1 & 0 & x \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

Left invariant differential forms are given by:

$$F_1 = dp - wdx + zdy, \quad F_2 = -xdr + dq, \quad F_3 = dr, \quad F_4 = dx, \\ F_5 = dy, \quad F_6 = dz, \quad F_7 = dw$$

*Corresponding author: Ghanam R, Department of Liberal Arts and Sciences, Virginia Commonwealth University in Qatar, Qatar, Tel: +974 4402 0795; E-mail: raghanam@vcu.edu

Received September 15, 2015; Accepted January 12, 2016; Published January 18, 2016

Citation: Ghanam R, Basim Mustafa B, Mustafa MT, Thompson G (2016) Matrix Representation for Seven-Dimensional Nilpotent Lie Algebras. J Phys Math 7: 155. doi:10.4172/2090-0902.1000155

Copyright: © 2016 Ghanam R, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

The vector field representation is given by:

$$E_1 = -wD_p - D_x, \quad E_2 = -xD_q - D_r, \quad E_3 = D_z, \quad E_4 = D_w, \\ E_5 = zD_p - D_y, \quad E_6 = D_q, \quad E_7 = D_p$$

2. (27B) : $[x_1, x_4] = x_6, [x_2, x_5] = x_7, [x_3, x_4] = x_6 + x_7, [x_3, x_5] = x_6 + x_7$;

A matrix Lie group is given by:

$$S = \begin{bmatrix} 1 & y & x & r & p & q \\ 0 & 1 & 0 & 0 & w+z & z \\ 0 & 0 & 1 & 0 & z & z \\ 0 & 0 & 0 & 1 & 0 & x \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

Left invariant differential forms are given by:

$$F_1 = dp - zdx + (-w-z)dy, \quad F_2 = -xdr + dq - zdx - zdy, \quad F_3 = dr, \quad F_4 = dx \\ F_5 = dy, \quad F_6 = dz, \quad F_7 = dz + dw$$

Vector field representation is given by:

$$E_1 = D_w, \quad E_2 = -xD_q - D_r, \quad E_3 = D_z, \quad E_4 = (w+z)D_p + zD_q + D_x, \\ E_5 = zD_p + zD_q + D_x, \quad E_6 = D_p, \quad E_7 = D_q$$

Algebras with upper central series dimensions (37)

1. (37A) : $[x_1, x_2] = x_5, [x_2, x_3] = x_6, [x_2, x_4] = x_7$;

A matrix Lie group is given by:

$$S = \begin{bmatrix} 1 & 0 & 0 & w & 0 & 0 & q \\ 0 & 1 & 0 & 0 & w & 0 & -x \\ 0 & 0 & 1 & 0 & 0 & w & y \\ 0 & 0 & 0 & 1 & 0 & 0 & z \\ 0 & 0 & 0 & 0 & 1 & 0 & r \\ 0 & 0 & 0 & 0 & 0 & 1 & p \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

Left invariant differential forms are given by:

$$F_1 = dp, \quad F_2 = dq - zdw, \quad F_3 = dr, \quad F_4 = -dx - rdw \\ F_5 = dy - pdw, \quad F_6 = dz, \quad F_7 = dw$$

Vector field representation is given by:

$$E_1 = D_p, \quad E_2 = zD_q - rD_x + pD_y + D_w, \quad E_3 = D_r, \quad E_4 = D_z \\ E_5 = D_y, \quad E_6 = D_x, \quad E_7 = -D_q$$

2. (37B) : $[x_1, x_2] = x_5, [x_2, x_3] = x_6, [x_3, x_4] = x_7$;

A matrix Lie group is given by:

$$S = \begin{bmatrix} 1 & 0 & 0 & w & 0 & -z & q \\ 0 & 1 & 0 & 0 & w & 0 & x \\ 0 & 0 & 1 & 0 & 0 & w & y \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & -r \\ 0 & 0 & 0 & 0 & 0 & 1 & p \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

Left invariant differential forms are given by:

$$F_1 = dp, \quad F_2 = dq + pdz, \quad F_3 = dr, \quad F_4 = dx + rdw \\ F_5 = dy - pdw, \quad F_6 = dz, \quad F_7 = dw$$

Vector field representation is given by:

$$E_1 = -D_r, \quad E_2 = D_w - rD_x + pD_y, \quad E_3 = -D_p, \quad E_4 = D_z - pD_q, \\ E_5 = D_x, \quad E_6 = D_y, \quad E_7 = D_q$$

3. (37C) : $[x_1, x_2] = x_5, [x_2, x_3] = x_6, [x_2, x_4] = x_7, [x_3, x_4] = x_5$;

A matrix Lie group is given by:

$$S = \begin{bmatrix} 1 & 0 & 0 & w & 0 & r & -q \\ 0 & 1 & 0 & 0 & w & 0 & x \\ 0 & 0 & 1 & 0 & 0 & w & y \\ 0 & 0 & 0 & 1 & 0 & 0 & z \\ 0 & 0 & 0 & 0 & 1 & 0 & r \\ 0 & 0 & 0 & 0 & 0 & 1 & p \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

Left invariant differential forms are given by:

$$F_1 = dp, \quad F_2 = dr, \quad F_3 = dz, \quad F_4 = dw \\ F_5 = -pdr - dq - zdw, \quad F_6 = dx - rdw, \quad F_7 = dy - pdw$$

Vector field representation is given by:

$$E_1 = D_p, \quad E_2 = zD_q - rD_x + pD_y + D_w, \quad E_3 = D_r, \quad E_4 = D_z, \\ E_5 = D_y, \quad E_6 = D_x, \quad E_7 = -D_q$$

4. (37D) : $[x_1, x_2] = x_5, [x_1, x_3] = x_6, [x_2, x_4] = x_7, [x_3, x_4] = x_5$;

A matrix Lie group is given by:

$$S = \begin{bmatrix} 1 & 0 & 0 & w & -z & 0 & q \\ 0 & 1 & 0 & 0 & w & -z & x \\ 0 & 0 & 1 & 0 & 0 & w & y \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & r \\ 0 & 0 & 0 & 0 & 0 & 1 & p \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

Left invariant differential forms are given by:

$$F_1 = dp, \quad F_2 = dq + rdz, \quad F_3 = dr, \quad F_4 = dx + pdz - rdw \\ F_5 = dy - pdw, \quad F_6 = dz, \quad F_7 = dw$$

Vector field representation is given by:

$$E_1 = D_p, \quad E_2 = -rD_q - pD_x + D_z, \quad E_3 = rD_x + pD_y + D_w, \quad E_4 = D_r, \\ E_5 = -D_x, \quad E_6 = D_y, \quad E_7 = D_q$$

Algebras with upper central series dimensions (247)

1. (247A) : $[x_1, x_i] = x_{i+2}, i = 2, 3, 4, 5$;

A matrix Lie group is given by:

$$S = \begin{bmatrix} 1 & 0 & w & 0 & 1/2w^2 & 0 & q \\ 0 & 1 & 0 & w & 0 & 1/2w^2 & x \\ 0 & 0 & 1 & 0 & w & 0 & y \\ 0 & 0 & 0 & 1 & 0 & w & z \\ 0 & 0 & 0 & 0 & 1 & 0 & r \\ 0 & 0 & 0 & 0 & 0 & 1 & p \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

Left invariant differential forms are given by:

$$F_1 = dp, \quad F_2 = dq - ydw, \quad F_3 = dr, \quad F_4 = dx - zdw \\ F_5 = dy - rdw, \quad F_6 = dz - pdw, \quad F_7 = dw$$

Vector field representation is given by:

$$E_1 = -yD_q - zD_x - rD_y - pD_z - D_w, \quad E_2 = D_p, \quad E_3 = D_r, \quad E_4 = D_z \\ E_5 = D_y, \quad E_6 = D_x, \quad E_7 = D_q$$

2. (247B) : $[x_1, x_i] = x_{i+2}, i = 2, 3, 4, [x_3, x_5] = x_7;$

A matrix Lie group is given by:

$$S = \begin{bmatrix} 1 & 0 & 0 & p & 0 & -z + pw & q \\ 0 & 1 & w & 0 & 1/2w^2 & 0 & x \\ 0 & 0 & 1 & 0 & w & 0 & y \\ 0 & 0 & 0 & 1 & 0 & w & z \\ 0 & 0 & 0 & 0 & 1 & 0 & r \\ 0 & 0 & 0 & 0 & 0 & 1 & p \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

Left invariant differential forms are given by:

$$F_1 = dp, \quad F_2 = -zdp + dq + pdz - p^2dw, \quad F_3 = dr, \quad F_4 = dx - ydw \\ F_5 = dy - rdw, \quad F_6 = dz - pdw, \quad F_7 = dw$$

Vector field representation is given by:

$$E_1 = D_w + yD_x + rD_y + pD_z, \quad E_2 = D_r, \quad E_3 = D_p + zD_q, \quad E_4 = -D_y \\ E_5 = pD_q - D_z, \quad E_6 = D_x, \quad E_7 = 2D_q$$

3. (247C) : $[x_1, x_i] = x_{i+2}, i = 2, 3, 4, [x_1, x_5] = x_7, [x_3, x_5] = x_6;$

A matrix Lie group is given by:

$$S = \begin{bmatrix} 1 & 0 & w & p & 1/2w^2 & -z + pw & q \\ 0 & 1 & 0 & w & 0 & 1/2w^2 & x \\ 0 & 0 & 1 & 0 & w & 0 & y \\ 0 & 0 & 0 & 1 & 0 & w & z \\ 0 & 0 & 0 & 0 & 1 & 0 & r \\ 0 & 0 & 0 & 0 & 0 & 1 & p \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

Left invariant differential forms are given by:

$$F_1 = dp, \quad F_2 = -zdp + dq + pdz + (-p^2 - y)dw, \quad F_3 = dr, \quad F_4 = dx - zdw \\ F_5 = dy - rdw, \quad F_6 = dz - pdw, \quad F_7 = dw$$

Vector field representation is given by:

$$E_1 = D_w + yD_x + zD_y + rD_z + pD_z, \quad E_2 = 2D_r, \quad E_3 = D_p + zD_q, \quad E_4 = -2D_y \\ E_5 = pD_q - D_z, \quad E_6 = 2D_q, \quad E_7 = D_x$$

4. (247D) : $[x_1, x_i] = x_{i+2}, i = 2, 3, [x_1, x_4] = x_6, [x_2, x_5] = x_7, [x_3, x_4] = x_7;$

A matrix Lie group is given by:

$$S = \begin{bmatrix} 1 & 0 & 0 & r & 0 & rw - y & q \\ 0 & 1 & 0 & w & 0 & 1/2w^2 & x \\ 0 & 0 & 1 & 0 & w & 0 & y \\ 0 & 0 & 0 & 1 & 0 & w & z \\ 0 & 0 & 0 & 0 & 1 & 0 & r \\ 0 & 0 & 0 & 0 & 0 & 1 & p \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

Left invariant differential forms are given by:

$$F_1 = dp, \quad F_2 = -zdr + dq + pdy - prdw, \quad F_3 = dr, \quad F_4 = dx - zdw \\ F_5 = dy - rdw, \quad F_6 = dz - pdw, \quad F_7 = dw$$

Vector field representation is given by:

$$E_1 = zD_x + rD_y + pD_z + D_w, \quad E_2 = D_p, \quad E_3 = zD_q + D_r, \quad E_4 = -D_z \\ E_5 = pD_q - D_y, \quad E_6 = D_x, \quad E_7 = D_q$$

5. (247E) : $[x_1, x_i] = x_{i+2}, i = 2, 3, 4, [x_1, x_5] = x_6, [x_2, x_5] = x_7, [x_3, x_4] = x_7;$

A matrix Lie group is given by:

$$S = \begin{bmatrix} 1 & 0 & 0 & r & 0 & rw - y & q \\ 0 & 1 & w & w & 1/2w^2 & 1/2w^2 & x \\ 0 & 0 & 1 & 0 & w & 0 & y \\ 0 & 0 & 0 & 1 & 0 & w & z + pw \\ 0 & 0 & 0 & 0 & 1 & 0 & r \\ 0 & 0 & 0 & 0 & 0 & 1 & p \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

Left invariant differential forms are given by:

$$F_1 = dp, \quad F_2 = (-pw - z)dr + dq + pdy - prdw, \quad F_3 = dr, \quad F_4 = dx + (-pw - z - y)dw \\ F_5 = dy - rdw, \quad F_6 = wdp + dz, \quad F_7 = dw$$

Vector field representation is given by:

$$E_1 = (pw + z + y)D_x + rD_y + D_w, \quad E_2 = (z + pw)D_q + D_r, \quad E_3 = D_p - wD_z, \quad E_4 = pD_q - D_y \\ E_5 = -D_z, \quad E_6 = D_x, \quad E_7 = D_q$$

6. (247F) : $[x_1, x_i] = x_{i+2}, i = 2, 3, [x_2, x_4] = x_6, [x_2, x_5] = x_7;$

A matrix Lie group is given by:

$$S = \begin{bmatrix} 1 & 0 & 0 & w & 0 & r + 1/2w^2 & q \\ 0 & 1 & r & 0 & -y + rw & 0 & x \\ 0 & 0 & 1 & 0 & w & 0 & y \\ 0 & 0 & 0 & 1 & 0 & w & z \\ 0 & 0 & 0 & 0 & 1 & 0 & r \\ 0 & 0 & 0 & 0 & 0 & 1 & p \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

Left invariant differential forms are given by:

$$F_1 = dp, \quad F_2 = -pdr + dq - zdw, \quad F_3 = dr, \quad F_4 = -ydr + dx + rdy - r^2dw \\ F_5 = dy - rdw, \quad F_6 = dz - pdw, \quad F_7 = dw$$

Vector field representation is given by:

$$E_1 = rD_y + pD_z + D_w, \quad E_2 = D_r + yD_x, \quad E_3 = -D_p - zD_q, \quad E_4 = rD_x - D_y \\ E_5 = -pD_q + D_z, \quad E_6 = 2D_x, \quad E_7 = 2D_q$$

7. (247G) : $[x_1, x_i] = x_{i+2}, i = 2, 3, [x_1, x_4] = x_7, [x_2, x_4] = x_6, [x_3, x_5] = x_7;$

A matrix Lie group is given by:

$$S = \begin{bmatrix} 1 & 0 & w & p & 1/2w^2 & -z + pw & q \\ 0 & 1 & r & 0 & -y + rw & 0 & x \\ 0 & 0 & 1 & 0 & w & 0 & y \\ 0 & 0 & 0 & 1 & 0 & w & z \\ 0 & 0 & 0 & 0 & 1 & 0 & r \\ 0 & 0 & 0 & 0 & 0 & 1 & p \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

Left invariant differential forms are given by:

$$F_1 = dp, \quad F_2 = -zdp + dq + pdz + (-p^2 - y)dw, \quad F_3 = dr, \quad F_4 = -ydr + dx + rdy - r^2dw \\ F_5 = dy - rdw, \quad F_6 = dz - pdw, \quad F_7 = dw$$

Vector field representation is given by:

$$E_1 = yD_q + rD_y + pD_z + D_w, \quad E_2 = 2D_r + 2yD_x, \quad E_3 = -D_p - zD_q, \quad E_4 = 2rD_x - 2D_y \\ E_5 = -pD_q + D_z, \quad E_6 = 8D_x, \quad E_7 = 2D_q$$

8. (247H) :

$$[x_1, x_i] = x_{i+2}, i = 2, 3, [x_1, x_4] = x_7, [x_1, x_5] = x_6, [x_2, x_4] = x_6, [x_3, x_5] = x_7;$$

A matrix Lie group is given by:

$$S = \begin{bmatrix} 1 & 0 & w & p & 1/2w^2 & pw-z & q \\ 0 & 1 & r & w & rw-y & 1/2w^2 & x \\ 0 & 0 & 1 & 0 & w & 0 & y \\ 0 & 0 & 0 & 1 & 0 & w & z \\ 0 & 0 & 0 & 0 & 1 & 0 & r \\ 0 & 0 & 0 & 0 & 0 & 1 & p \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

Left invariant differential forms are given by:

$$F_1 = dp, \quad F_2 = -zdp + dq + pdz + (-p^2 - y)dw, \quad F_3 = dr, \quad F_4 = -ydr + dx + rdy + (-r^2 - z)dw \\ F_5 = dy - rdw, \quad F_6 = dz - pdw, \quad F_7 = dw$$

Vector field representation is given by:

$$E_1 = 2yD_q + 2zD_x + 2rD_y + 2pD_z + 2D_w, \quad E_2 = D_r + yD_x, \quad E_3 = D_p + zD_q, \quad E_4 = 2rD_x - 2D_y \\ E_5 = 2pD_q - 2D_z, \quad E_6 = 4D_x, \quad E_7 = 4D_q$$

9. (247I) : $[x_1, x_i] = x_{i+2}, i = 2, 3, [x_2, x_5] = x_6, [x_3, x_4] = x_6, [x_3, x_5] = x_7;$

A matrix Lie group is given by:

$$S = \begin{bmatrix} 1 & 0 & 0 & r & 0 & rw-y & q \\ 0 & 1 & 0 & p & 0 & -z & x \\ 0 & 0 & 1 & 0 & w & 0 & y \\ 0 & 0 & 0 & 1 & 0 & w & z+pw \\ 0 & 0 & 0 & 0 & 1 & 0 & r \\ 0 & 0 & 0 & 0 & 0 & 1 & p \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

Left invariant differential forms are given by:

$$F_1 = dp, \quad F_2 = (-z - pw)dr + dq + pdy - prdw, \quad F_3 = dr, \quad F_4 = zdp + dx + pdz \\ F_5 = dy - rdw, \quad F_6 = wdp + dz, \quad F_7 = dw$$

Vector field representation is given by:

$$E_1 = rD_x + D_w, \quad E_2 = (2z + 2pw)D_q + 2D_r, \quad E_3 = -D_p + (-z - pw)D_x + wD_z, \quad E_4 = 2pD_q - 2D_y \\ E_5 = -pD_x + D_z, \quad E_6 = -2D_q, \quad E_7 = 2D_x$$

10. (247J) : $[x_1, x_i] = x_{i+2}, i = 2, 3, 4, [x_2, x_5] = x_6, [x_3, x_4] = x_6, [x_3, x_5] = x_7;$

A matrix Lie group is given by:

$$S = \begin{bmatrix} 1 & 0 & 0 & p & 0 & pw-z & q \\ 0 & 1 & w & r & 1/2w^2 & rw-y & x \\ 0 & 0 & 1 & 0 & w & 0 & y \\ 0 & 0 & 0 & 1 & 0 & w & z \\ 0 & 0 & 0 & 0 & 1 & 0 & r \\ 0 & 0 & 0 & 0 & 0 & 1 & p \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

Left invariant differential forms are given by:

$$F_1 = dp, \quad F_2 = -zdp + dq + pdz - p^2dw, \quad F_3 = dr, \quad F_4 = -zdr + dx + pdy + (-pr - y)dw \\ F_5 = dy - rdw, \quad F_6 = dz - pdw, \quad F_7 = dw$$

Vector field representation is given by:

$$E_1 = yD_x + rD_y + pD_z + D_w, \quad E_2 = D_r + zD_x, \quad E_3 = D_p + zD_q, \quad E_4 = pD_x - D_y \\ E_5 = pD_q - D_z, \quad E_6 = D_x, \quad E_7 = 2D_q$$

11. (247K) : $[x_1, x_i] = x_{i+2}, i = 2, 3, 4, [x_2, x_5] = x_7, [x_3, x_4] = x_7, [x_3, x_5] = x_6;$

A matrix Lie group is given by:

$$S = \begin{bmatrix} 1 & 0 & 0 & r & 0 & rw-y & q \\ 0 & 1 & w & p & 1/2w^2 & -z & x \\ 0 & 0 & 1 & 0 & w & 0 & y \\ 0 & 0 & 0 & 1 & 0 & w & z+pw \\ 0 & 0 & 0 & 0 & 1 & 0 & r \\ 0 & 0 & 0 & 0 & 0 & 1 & p \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

Left invariant differential forms are given by:

$$F_1 = dp, \quad F_2 = (-z - pw)dr + dq + pdy - prdw, \quad F_3 = dr, \quad F_4 = -zdp + dx + pdz - ydw \\ F_5 = dy - rdw, \quad F_6 = wdp + dz, \quad F_7 = dw$$

Vector field representation is given by:

$$E_1 = yD_x + rD_y + D_w, \quad E_2 = (2z + 2pw)D_q + 2D_r, \quad E_3 = D_p + (z + pw)D_x - wD_z, \quad E_4 = 2pD_q - 2D_y \\ E_5 = pD_x - D_z, \quad E_6 = 2D_x, \quad E_7 = 2D_q$$

12. (247L) : $[x_1, x_i] = x_{i+2}, i = 2, 3, 4, 5, [x_2, x_3] = x_6$

A matrix Lie group is given by:

$$S = \begin{bmatrix} 1 & 0 & w & 0 & 1/2w^2 & r & q \\ 0 & 1 & 0 & w & 0 & 1/2w^2 & x \\ 0 & 0 & 1 & 0 & w & 0 & y \\ 0 & 0 & 0 & 1 & 0 & w & z \\ 0 & 0 & 0 & 0 & 1 & 0 & r \\ 0 & 0 & 0 & 0 & 0 & 1 & p \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

Left invariant differential forms are given by:

$$F_1 = dp, \quad F_2 = -pdr + dq - ydw, \quad F_3 = dr, \quad F_4 = dx - zdw \\ F_5 = dy - rdw, \quad F_6 = dz - pdw, \quad F_7 = dw$$

Vector field representation is given by:

$$E_1 = yD_q + zD_x + rD_y + pD_z + D_w, \quad E_2 = -pD_q - D_r, \quad E_3 = -D_p, \quad E_4 = D_y \\ E_5 = D_z, \quad E_6 = -D_q, \quad E_7 = -D_x$$

13. (247M) : $[x_1, x_i] = x_{i+2}, i = 2, 3, 4, [x_2, x_3] = x_6, [x_3, x_5] = x_7;$

A matrix Lie group is given by:

$$S = \begin{bmatrix} 1 & 0 & 0 & w & 0 & r+1/2w^2 & q \\ 0 & 1 & r & 0 & -y+rw & 0 & x \\ 0 & 0 & 1 & 0 & w & 0 & y \\ 0 & 0 & 0 & 1 & 0 & w & z \\ 0 & 0 & 0 & 0 & 1 & 0 & r \\ 0 & 0 & 0 & 0 & 0 & 1 & p \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

Left invariant differential forms are given by:

$$F_1 = dp, \quad F_2 = -pdr + dq - zdw, \quad F_3 = dr, \quad F_4 = -ydr + dx + rdy - r^2dw \\ F_5 = dy - rdw, \quad F_6 = dz - pdw, \quad F_7 = dw$$

Vector field representation is given by:

$$E_1 = -zD_q - rD_y - pD_z - D_w, \quad E_2 = D_p, \quad E_3 = pD_q + D_r + yD_x, \quad E_4 = D_z \\ E_5 = -rD_x + D_y, \quad E_6 = D_q, \quad E_7 = -2D_x$$

14. (247N) : $[x_1, x_i] = x_{i+2}, i = 2, 3, [x_1, x_5] = x_6, [x_2, x_3] = x_7, [x_2, x_4] = x_6;$

A matrix Lie group is given by:

$$S = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & r & q \\ 0 & 1 & r & w & rw-y & 1/2w^2 & x \\ 0 & 0 & 1 & 0 & w & 0 & y \\ 0 & 0 & 0 & 1 & 0 & w & z \\ 0 & 0 & 0 & 0 & 1 & 0 & r \\ 0 & 0 & 0 & 0 & 0 & 1 & p \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

Left invariant differential forms are given by:

$$F_1 = dp, \quad F_2 = -pdr + dq, \quad F_3 = dr, \quad F_4 = -ydr + dx + rdy + (-r^2 - z)dw \\ F_5 = dy - rdw, \quad F_6 = dz - pdw, \quad F_7 = dw$$

Vector field representation is given by:

$$E_1 = zD_x + rD_y + pD_z + D_w, \quad E_2 = pD_q + D_r + yD_s, \quad E_3 = 2D_p, \quad E_4 = rD_x - D_y \\ E_5 = -2D_z, \quad E_6 = 2D_x, \quad E_7 = -2D_q$$

15. (247O) : $[x_1, x_i] = x_{i+2}, i = 2, 3, 4, [x_1, x_5] = x_7, [x_2, x_3] = x_7, [x_3, x_5] = x_6;$

A matrix Lie group is given by:

$$S = \begin{bmatrix} 1 & 0 & w & 0 & 1/2w^2 & r & q \\ 0 & 1 & r & w & -y + rw & 1/2w^2 & x \\ 0 & 0 & 1 & 0 & w & 0 & y \\ 0 & 0 & 0 & 1 & 0 & w & z \\ 0 & 0 & 0 & 0 & 1 & 0 & r \\ 0 & 0 & 0 & 0 & 0 & 1 & p \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

Left invariant differential forms are given by:

$$F_1 = dp, \quad F_2 = -pdr + dq - ydw, \quad F_3 = dr, \quad F_4 = -ydr + dx + rdy + (-r^2 - z)*w \\ F_5 = dy - rdw, \quad F_6 = dz - pdw, \quad F_7 = dw$$

Vector field representation is given by:

$$E_1 = 2yD_q + 2zD_x + 2rD_y + 2pD_z + 2D_w, \quad E_2 = 4D_p, \quad E_3 = 2pD_q + 2D_r + 2yD_s, \quad E_4 = -8D_z \\ E_5 = 4rD_x - 4D_y, \quad E_6 = 16D_x, \quad E_7 = 8D_q$$

16. (247P) : $[x_1, x_i] = x_{i+2}, i = 2, 3, [x_2, x_3] = x_6, [x_2, x_5] = x_7, [x_3, x_4] = x_7;$

A matrix Lie group is given by:

$$S = \begin{bmatrix} 1 & 0 & 0 & 0 & p & 0 & q \\ 0 & 1 & 0 & r & 0 & rw-y & x \\ 0 & 0 & 1 & 0 & w & 0 & y \\ 0 & 0 & 0 & 1 & 0 & w & z \\ 0 & 0 & 0 & 0 & 1 & 0 & r \\ 0 & 0 & 0 & 0 & 0 & 1 & p \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

Left invariant differential forms are given by:

$$F_1 = dp, \quad F_2 = -rdp + dq, \quad F_3 = dr, \quad F_4 = -zdr + dx + pdy - prdw \\ F_5 = dy - rdw, \quad F_6 = dz - pdw, \quad F_7 = dw$$

Vector field representation is given by:

$$E_1 = rD_y + pD_z + D_w, \quad E_2 = -D_r - zD_x, \quad E_3 = -D_p - rD_q, \quad E_4 = -pD_x + D_y \\ E_5 = D_z, \quad E_6 = D_q, \quad E_7 = D_x$$

17. (247Q) : $[x_1, x_i] = x_{i+2}, i = 2, 3, 4, [x_2, x_3] = x_6, [x_2, x_5] = x_7, [x_3, x_4] = x_7;$

A matrix Lie group is given by:

$$S = \begin{bmatrix} 1 & 0 & 0 & w & 0 & r+1/2w^2 & q \\ 0 & 1 & p & r & pw-z & -y+rw & x \\ 0 & 0 & 1 & 0 & w & 0 & y \\ 0 & 0 & 0 & 1 & 0 & w & z \\ 0 & 0 & 0 & 0 & 1 & 0 & r \\ 0 & 0 & 0 & 0 & 0 & 1 & p \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

Left invariant differential forms are given by:

$$F_1 = dp, \quad F_2 = -pdr + dq - zdw, \quad F_3 = dr, \quad F_4 = -zdr - ydp + dx + pdy + rdz - 2prdw \\ F_5 = dy - rdw, \quad F_6 = dz - pdw, \quad F_7 = dw$$

Vector field representation is given by:

$$E_1 = 2zD_x + 2rD_y + 2pD_z + 2D_w, \quad E_2 = -2D_r - 2yD_s, \quad E_3 = 4pD_q + 4D_r + 4zD_z, \quad E_4 = -4rD_x + 4D_y \\ E_5 = 8pD_x - 8D_y, \quad E_6 = -8D_q, \quad E_7 = -32D_x$$

18. (247R) :

$$[x_1, x_i] = x_{i+2}, i = 2, 3, 4, [x_1, x_5] = x_6, [x_2, x_3] = x_6, [x_2, x_5] = x_7, [x_3, x_4] = x_7;$$

A matrix Lie group is given by:

$$S = \begin{bmatrix} 1 & 0 & w & w & p+1/2w^2 & 1/2w^2 & q \\ 0 & 1 & 0 & r & 0 & rw-y & x \\ 0 & 0 & 1 & 0 & w & 0 & y \\ 0 & 0 & 0 & 1 & 0 & w & z \\ 0 & 0 & 0 & 0 & 1 & 0 & r \\ 0 & 0 & 0 & 0 & 0 & 1 & p \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

Left invariant differential forms are given by:

$$F_1 = dp, \quad F_2 = -rdp + dq + (-z-y)dw, \quad F_3 = dr, \quad F_4 = -zdr + dx + pdy - rpdw \\ F_5 = dy - rdw, \quad F_6 = dz - pdw, \quad F_7 = dw$$

The Vector field representation is given by:

$$E_1 = (z+y)D_q + rD_y + pD_z + D_w, \quad E_2 = D_r + zD_x, \quad E_3 = D_p + rD_q, \quad E_4 = pD_x - D_y \\ E_5 = -D_z, \quad E_6 = D_q, \quad E_7 = D_x$$

Algebras with upper central series dimensions (257)

1. (257A) : $[x_1, x_2] = x_3, [x_1, x_3] = x_6, [x_1, x_5] = x_7, [x_2, x_4] = x_6;$

A matrix Lie group is given by:

$$S = \begin{bmatrix} 1 & x & -y & -r & p & q \\ 0 & 1 & x & 0 & w & 0 \\ 0 & 0 & 1 & 0 & z & 0 \\ 0 & 0 & 0 & 1 & 0 & x \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

Left invariant differential forms are given by:

$$F_1 = dp + (-w+zx)dx + zdy, \quad F_2 = xdr + dq, \quad F_3 = dr, \quad F_4 = dx, \\ F_5 = xdx + dy, \quad F_6 = dz, \quad F_7 = -zdx + dw$$

Vector field representation is given by:

$$E_1 = -D_x - wD_p + xD_y - zD_w, \quad E_2 = D_z, \quad E_3 = D_w, \quad E_4 = -D_y + zD_p \\ E_5 = D_r - xD_q, \quad E_6 = D_p, \quad E_7 = D_q$$

2. (257B) : $[x_1, x_2] = x_3, [x_1, x_3] = x_6, [x_1, x_4] = x_7, [x_2, x_5] = x_7;$

A matrix Lie group is given by:

$$S = \begin{bmatrix} 1 & 0 & -w & -w & 0 & 1/2w^2 & q \\ 0 & 1 & 0 & 0 & -w & z & -x \\ 0 & 0 & 1 & 0 & 0 & -w & -y \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & r \\ 0 & 0 & 0 & 0 & 0 & 1 & p \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

Left invariant differential forms are given by:

$$F_1 = dp, \quad F_2 = dq - ydw, \quad F_3 = dr, \quad F_4 = dx + pdz - rdw, \\ F_5 = dy - pdw, \quad F_6 = dz, \quad F_7 = dw$$

Vector field representation is given by:

$$E_1 = -yD_q - rD_x - pD_y - D_w, \quad E_2 = D_p, \quad E_3 = D_y, \quad E_4 = D_r, \\ E_5 = pD_x - D_z, \quad E_6 = D_q, \quad E_7 = D_x$$

$$3. (257C) : [x_1, x_2] = x_3, [x_1, x_3] = x_6, [x_2, x_4] = x_6, [x_2, x_5] = x_7;$$

A matrix Lie group is given by:

$$S = \begin{bmatrix} 1 & x & -y & -r & p & q \\ 0 & 1 & z & 0 & w & 0 \\ 0 & 0 & 1 & 0 & z & 0 \\ 0 & 0 & 0 & 1 & 0 & x \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

Left invariant differential forms are given by:

$$F_1 = dp + (-w + z^2)dx + zdy, \quad F_2 = xdr + dq, \quad F_3 = dr, \quad F_4 = dx, \\ F_5 = zdx + dy, \quad F_6 = dz, \quad F_7 = -zdz + dw$$

Vector field representation is given by:

$$E_1 = D_z + zD_w, \quad E_2 = wD_p + D_x - zD_y, \quad E_3 = zD_p - D_y, \quad E_4 = -D_w, \\ E_5 = xD_q - D_r, \quad E_6 = D_p, \quad E_7 = D_q$$

$$4. (257D) : [x_1, x_2] = x_3, [x_1, x_3] = x_6, [x_1, x_4] = x_7, [x_2, x_4] = x_6, [x_2, x_5] = x_7;$$

A matrix Lie group is given by:

$$S = \begin{bmatrix} 1 & 0 & -w & -w & p & 1/2w^2 & q \\ 0 & 1 & 0 & 0 & -w & z & -x \\ 0 & 0 & 1 & 0 & 0 & -w & -y \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & r \\ 0 & 0 & 0 & 0 & 0 & 1 & p \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

Left invariant differential forms are given by:

$$F_1 = dp, \quad F_2 = -rdp + dq - ydw, \quad F_3 = dr, \quad F_4 = dx + pdz - rdw, \\ F_5 = dy - pdw, \quad F_6 = dz, \quad F_7 = dw$$

Vector field representation is given by:

$$E_1 = D_w + yD_q + rD_x + pD_y, \quad E_2 = D_p + rD_q, \quad E_3 = -D_y, \quad E_4 = -D_r, \\ E_5 = -D_z + pD_x, \quad E_6 = D_q, \quad E_7 = D_x$$

$$5. (257E) : [x_1, x_2] = x_3, [x_1, x_3] = x_6, [x_2, x_4] = x_7, [x_4, x_5] = x_6;$$

A matrix Lie group is given by:

$$S = \begin{bmatrix} 1 & 0 & -w & -w & z & 1/2w^2 & q \\ 0 & 1 & 0 & 0 & 0 & z & -x \\ 0 & 0 & 1 & 0 & 0 & -w & -y \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & r \\ 0 & 0 & 0 & 0 & 0 & 1 & p \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

Left invariant differential forms are given by:

$$F_1 = dp, \quad F_2 = dq - rdz - ydw, \quad F_3 = dr, \quad F_4 = dx + pdz, \\ F_5 = dy - pdw, \quad F_6 = dz, \quad F_7 = dw$$

Vector field representation is given by:

$$E_1 = yD_q + pD_y + D_w, \quad E_2 = D_p, \quad E_3 = -D_y, \quad E_4 = -rD_q + pD_x - D_z, \\ E_5 = D_r, \quad E_6 = D_q, \quad E_7 = D_x$$

$$6. (257F) : \begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

A matrix Lie group is given by:

$$S = \begin{bmatrix} 1 & 0 & -w & -w & z & 1/2w^2 & q \\ 0 & 1 & 0 & 0 & -w & 0 & -x \\ 0 & 0 & 1 & 0 & 0 & -w & -y \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & r \\ 0 & 0 & 0 & 0 & 0 & 1 & p \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

Left invariant differential forms are given by:

$$F_1 = dp, \quad F_2 = dq - rdz - ydw, \quad F_3 = dr, \quad F_4 = dx - rdw, \\ F_5 = dy - pdw, \quad F_6 = dz, \quad F_7 = dw$$

Vector field representation is given by:

$$E_1 = -D_p, \quad E_2 = -yD_q - rD_x - pD_y - D_w, \quad E_3 = D_y, \quad E_4 = D_r, \\ E_5 = rD_q + D_z, \quad E_6 = D_q, \quad E_7 = D_x$$

$$7. (257G) : [x_1, x_2] = x_3, [x_1, x_3] = x_6, [x_1, x_5] = x_7, [x_2, x_4] = x_7, [x_4, x_5] = x_6;$$

A matrix Lie group is given by:

$$S = \begin{bmatrix} 1 & 0 & -w & -w & z & 1/2w^2 & q \\ 0 & 1 & 0 & 0 & -w & z & -x \\ 0 & 0 & 1 & 0 & 0 & -w & -y \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & r \\ 0 & 0 & 0 & 0 & 0 & 1 & p \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

Left invariant differential forms are given by:

$$F_1 = dp, \quad F_2 = dq - rdz - ydw, \quad F_3 = dr, \quad F_4 = dx + pdz - rdw, \\ F_5 = dy - pdw, \quad F_6 = dz, \quad F_7 = dw$$

Vector field representation is given by:

$$E_1 = -yD_q - rD_x - pD_y - D_w, \quad E_2 = D_p, \quad E_3 = D_y, \quad E_4 = -rD_q + pD_x - D_z, \\ E_5 = D_r, \quad E_6 = D_q, \quad E_7 = D_x$$

$$8. (257H) : [x_1, x_2] = x_3, [x_1, x_3] = x_6, [x_2, x_4] = x_6, [x_4, x_5] = x_7;$$

A matrix Lie group is given by:

$$S = \begin{bmatrix} 1 & x & -y & -r & p & q \\ 0 & 1 & z & 0 & w+1/2z^2 & 0 \\ 0 & 0 & 1 & 0 & z & 0 \\ 0 & 0 & 0 & 1 & 0 & w \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

Left invariant differential forms are given by:

$$F_1 = dp + (-w + (1/2)z^2)dx + zdy, \quad F_2 = wdr + dq, \quad F_3 = dr, \quad F_4 = dx, \\ F_5 = zdx + dy, \quad F_6 = dz, \quad F_7 = dw$$

Vector field representation is given by:

$$E_1 = -D_z, \quad E_2 = D_x + (w + (1/2)z^2)D_p - zD_y, \quad E_3 = -zD_p + D_y, \quad E_4 = -D_w, \\ E_5 = -wD_q + D_r, \quad E_6 = D_p, \quad E_7 = D_q$$

9. (257I) : $[x_1, x_2] = x_3, [x_1, x_3] = x_6, [x_1, x_4] = x_6, [x_1, x_5] = x_7, [x_2, x_3] = x_7;$

A matrix Lie group is given by:

$$S = \begin{bmatrix} 1 & 0 & -w & -w & 0 & 1/2w^2 & q \\ 0 & 1 & p & 0 & -w & y & -x \\ 0 & 0 & 1 & 0 & 0 & -w & -y - pw \\ 0 & 0 & 0 & 1 & 0 & 0 & z \\ 0 & 0 & 0 & 0 & 1 & 0 & r \\ 0 & 0 & 0 & 0 & 0 & 1 & p \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

Left invariant differential forms are given by:

$$F_1 = dp, \quad F_2 = dq + (-y - pw + z)dw, \quad F_3 = dr, \quad F_4 = ydp - dx - pdy + rdw, \\ F_5 = wdp + dy, \quad F_6 = dz, \quad F_7 = dw$$

Vector field representation is given by:

10. (257J) : $[x_1, x_2] = x_3, [x_1, x_3] = x_6, [x_1, x_5] = x_7, [x_2, x_3] = x_7, [x_2, x_4] = x_6;$

A matrix Lie group is given by:

$$S = \begin{bmatrix} 1 & 0 & -w & -w & 0 & 1/2w^2 + z & q \\ 0 & 1 & p & 0 & -w & y & -x \\ 0 & 0 & 1 & 0 & 0 & -w & -y - pw \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & r \\ 0 & 0 & 0 & 0 & 0 & 1 & p \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

Left invariant differential forms are given by:

$$F_1 = dp, \quad F_2 = dq - pdz + (-y - pw)dw, \quad F_3 = dr, \quad F_4 = ydp - dx - pdy + rdw, \\ F_5 = wdp + dy, \quad F_6 = dz, \quad F_7 = dw$$

Vector field representation is given by:

$$E_1 = (y + pw)D_q + rD_x + D_w, \quad E_2 = D_p + (y + pw)D_x - wD_y, \quad E_3 = pD_x - D_y, \quad E_4 = pD_q + D_z, \\ E_5 = -2D_r, \quad E_6 = D_q, \quad E_7 = 2D_x$$

11. (257K) : $[x_1, x_2] = x_3, [x_1, x_3] = x_6, [x_2, x_3] = x_7, [x_4, x_5] = x_7;$

A matrix Lie group is given by:

$$S = \begin{bmatrix} 1 & 0 & -z & y - wz & r & 0 & q \\ 0 & 1 & w & 1/2w^2 & 0 & 0 & x \\ 0 & 0 & 1 & w & 0 & 0 & 2y \\ 0 & 0 & 0 & 1 & 0 & 0 & 2z \\ 0 & 0 & 0 & 0 & 1 & 0 & p \\ 0 & 0 & 0 & 0 & 0 & 1 & r \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

Left invariant differential forms are given by:

$$F_1 = dp, \quad F_2 = -pdr + dq - 2zdy + 2ydz + 2z^2dw, \quad F_3 = dr, \quad F_4 = dx - 2ydw, \\ F_5 = dy - zdw, \quad F_6 = dz, \quad F_7 = dw$$

Vector field representation is given by:

$$E_1 = D_w + 2yD_x + zD_y, \quad E_2 = D_z - 2yD_q, \quad E_3 = -D_y - 2zD_q, \quad E_4 = -4D_p, \\ E_5 = D_r + pD_q, \quad E_6 = 2D_x, \quad E_7 = -4D_q$$

12. (257L) : $[x_1, x_2] = x_3, [x_1, x_3] = x_6, [x_2, x_3] = x_7, [x_2, x_4] = x_6, [x_4, x_5] = x_7;$

A matrix Lie group is given by:

$$S = \begin{bmatrix} 1 & 0 & -z & y - wz & r & 0 & q \\ 0 & 1 & w & 1/2w^2 & 0 & z & x \\ 0 & 0 & 1 & w & 0 & 0 & 2y \\ 0 & 0 & 0 & 1 & 0 & 0 & 2z \\ 0 & 0 & 0 & 0 & 1 & 0 & p \\ 0 & 0 & 0 & 0 & 0 & 1 & r \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

Left invariant differential forms are given by:

$$F_1 = dp, \quad F_2 = -pdr + dq - 2zdy + 2ydz + 2z^2dw, \quad F_3 = dr, \quad F_4 = dx - rdz - 2ydw, \\ F_5 = dy - zdw, \quad F_6 = dz, \quad F_7 = dw$$

Vector field representation is given by:

$$E_1 = D_w + 2yD_x + zD_y, \quad E_2 = D_z - 2yD_q + rD_x, \quad E_3 = -D_y - 2zD_q, \quad E_4 = -2D_r - 2pD_q, \\ E_5 = -2D_p, \quad E_6 = 2D_x, \quad E_7 = -4D_q$$

Algebras with upper central series dimensions (357)

1. (357A) : $[x_1, x_2] = x_3, [x_1, x_3] = x_5, [x_1, x_4] = x_7, [x_2, x_4] = x_6;$

A matrix Lie group is given by:

$$S = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & r & -q \\ 0 & 1 & 0 & w & 0 & 1/2w^2 & x \\ 0 & 0 & 1 & 0 & w & 0 & y \\ 0 & 0 & 0 & 1 & 0 & w & z \\ 0 & 0 & 0 & 0 & 1 & 0 & r \\ 0 & 0 & 0 & 0 & 0 & 1 & p \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

Left invariant differential forms are given by:

$$F_1 = dp, \quad F_2 = -pdr - dq, \quad F_3 = dr, \quad F_4 = dx - zdw, \\ F_5 = dy - rdw, \quad F_6 = dz - pdw, \quad F_7 = dw$$

Vector field representation is given by:

$$E_1 = -zD_x + rD_y - pD_z - D_w, \quad E_2 = D_p, \quad E_3 = D_z, \quad E_4 = -pD_q + D_r, \\ E_5 = D_x, \quad E_6 = -D_q, \quad E_7 = -D_y$$

2. (357B) : $[x_1, x_2] = x_3, [x_1, x_3] = x_5, [x_1, x_4] = x_7, [x_2, x_3] = x_6;$

A matrix Lie group is given by:

$$S = \begin{bmatrix} 1 & 0 & 0 & p & 0 & -z + pw & 2q \\ 0 & 1 & 0 & w & 0 & 1/2w^2 & x \\ 0 & 0 & 1 & 0 & w & 0 & y \\ 0 & 0 & 0 & 1 & 0 & w & z \\ 0 & 0 & 0 & 0 & 1 & 0 & r \\ 0 & 0 & 0 & 0 & 0 & 1 & p \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

Left invariant differential forms are given by:

$$F_1 = dp, \quad F_2 = -zdp + 2dq + pdz - p^2 dw, \quad F_3 = dr, \quad F_4 = dx - zdw, \\ F_5 = dy - rdw, \quad F_6 = dz - pdw, \quad F_7 = dw$$

The vector field representation is given by:

$$E_1 = zD_x + rD_y + pD_z + D_w, \quad E_2 = D_p + z/2D_q, \quad E_3 = p/2D_q - D_z, \quad E_4 = -D_r, \\ E_5 = D_x, \quad E_6 = D_q, \quad E_7 = D_y$$

$$3. (357C) : [x_1, x_2] = x_3, [x_1, x_3] = x_5, [x_1, x_4] = x_7, [x_2, x_3] = x_6, [x_2, x_4] = x_5;$$

A matrix Lie group is given by:

$$S = \begin{bmatrix} 1 & 0 & 0 & p & 0 & -z + pw & 2q \\ 0 & 1 & 0 & w & p & 1/2w^2 & x \\ 0 & 0 & 1 & 0 & w & 0 & y \\ 0 & 0 & 0 & 1 & 0 & w & z \\ 0 & 0 & 0 & 0 & 1 & 0 & r \\ 0 & 0 & 0 & 0 & 0 & 1 & p \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

Left invariant differential forms are given by:

$$F_1 = dp, \quad F_2 = -zdp + 2dq + pdz - p^2 dw, \quad F_3 = dr, \quad F_4 = -rdp + dx - zdw, \\ F_5 = dy - rdw, \quad F_6 = dz - pdw, \quad F_7 = dw$$

Vector field representation is given by:

$$E_1 = D_x + zD_z + rD_y + pD_z, \quad E_2 = D_p + ((1/2)z)D_q + rD_z, \quad E_3 = -D_z + ((1/2)p)D_q, \quad E_4 = -D_r, \\ E_5 = D_x, \quad E_6 = D_q, \quad E_7 = D_y$$

Conclusion

The Explanation regarding finding linear representations for seven-dimensional real, indecomposable nilpotent Lie algebras is done.

References

1. Ado ID (1935) Note on the representation of finite continuous groups by means of linear substitution. Izv Fiz-Mat Obsch (Kazan) 7: 01-43.
2. Ado ID (1947) The representation of Lie algebras by matrices (in Russian). Akademiya Nauk SSSR i Moskovskoe Matematicheskoe Obshchestvo. Uspekhi Matematicheskikh Nauk 2: 159-173.
3. Ghanam R, Thompson G, Tonon S (2006) Representations for six-dimensional nilpotent Lie algebras. Hadronic J 29: 299-317.
4. Ghanam R, Strugar I, Thompson G (2005) Matrix representations for low dimensional Lie algebras. Extracta Math 20: 151-184.
5. Ghanam R, Thompson G, Miller E (2004) Variationality of four-dimensional Lie group connections. J Lie Theory 14: 395-425.
6. Gong MP (1998) Classification of Nilpotent Lie algebras of dimension 7 (Over algebraically closed Fields and R). University of Waterloo.
7. Jacobson N (1962) Lie Algebras. Tracts in Pure and Applied Math Interscience Publishers, Newyork.
8. Humphreys JE (1972) Introduction to Lie algebras and representation theory. Graduate Text in Math Springer.
9. Seeley C. Degeneration of 6-dimensional Nilpotent Lie Algebras over C. Communications in Algebra 18: 3493-3505.
10. Seeley C (1993) 7-dimensional Nilpotent Lie Algebra. Trans Amer Math Soc 335: 479-496.
11. Skjelbred T, Sund T (1977). Classification of Nilpotent Lie Algebras in dimension six. University of Oslo.

Citation: Ghanam R, Basim Mustafa B, Mustafa MT, Thompson G (2016) Matrix Representation for Seven-Dimensional Nilpotent Lie Algebras. J Phys Math 7: 155. doi:10.4172/2090-0902.1000155

OMICS International: Publication Benefits & Features

Unique features:

- Increased global visibility of articles through worldwide distribution and indexing
- Showcasing recent research output in a timely and updated manner
- Special issues on the current trends of scientific research

Special features:

- 700 Open Access Journals
- 50,000 editorial team
- Rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at PubMed (partial), Scopus, EBSCO, Index Copernicus and Google Scholar etc
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://omicsonline.com/open-access/physical-mathematics.php>