

EINSTEIN SPACES OF POSITIVE SCALAR CURVATURE

YOSIO MUTŌ

1. Let M be an n -dimensional compact orientable Einstein space with positive scalar curvature K . Then the concircular curvature tensor Z_{kjih} defined by

$$(1) \quad Z_{kjih} = K_{kjih} - \frac{K}{n(n-1)}(g_{ji}g_{kh} - g_{ki}g_{jh})$$

satisfies

$$(2) \quad Z_{kjih}g^{ji} = 0,$$

because of

$$K_{ji} = \frac{K}{n}g_{ji}.$$

The purpose of the present paper is to prove the

Theorem. *If the concircular curvature tensor satisfies the inequality*

$$(3) \quad \frac{1}{K} |Z_{kjih}A^k B^j C^i D^h| < \frac{2}{5n^7}$$

at every point of M for any set of unit vectors A, B, C, D , then the Einstein space M is a space of constant curvature.

Roughly speaking, this theorem tells that, if M_0 is a space of positive constant curvature, there exist no Einstein spaces other than M_0 in a sufficiently small neighborhood of M_0 . The inequality (3) is a quite rough and modest estimation. We can get a better estimation by a more elaborate calculation.

2. In an Einstein space we have $\nabla_k K_{ji} = 0$, $\nabla_k K = 0$, and therefore $\nabla_i Z_{kjih} = \nabla_i K_{kjih}$. Thus by using Green's theorem and the second identity of Bianchi, we get

$$\begin{aligned} - \int_M (\nabla^i Z^{kjih})(\nabla_i Z_{kjih})dV &= \int_M Z^{kjih} \nabla^l \nabla_l K_{kjih} dV \\ &= 2 \int_M Z^{kjih} \nabla^l \nabla_l K_{lji h} dV, \end{aligned}$$

where dV is the volume element of M , and the last step becomes, by virtue of the Ricci identity and $\nabla^l K_{ljih} = 0$,

$$\begin{aligned} & 2 \int_M Z^{kjih} \left(\frac{K}{n} K_{kjih} - K^l{}_{kj}{}^m K_{lmih} - K^l{}_{ki}{}^m K_{ljmh} - K^l{}_{kh}{}^m K_{ljim} \right) dV \\ &= \frac{2K}{n} \int_M Z^{kjih} K_{kjih} dV + \int_M Z^{kjih} (K_{kj}{}^{lm} K_{lmih} - 4K^l{}_{kh}{}^m K_{jlm i}) dV, \end{aligned}$$

where we have used

$$K^l{}_{kj}{}^m - K^l{}_{jk}{}^m = -K_{kj}{}^{lm}.$$

We also obtain, in consequence of (1),

$$\begin{aligned} & \int_M Z^{kjih} K_{kjih} dV = \int_M Z^{kjih} Z_{kjih} dV, \\ & \int_M Z^{kjih} (K_{kj}{}^{lm} K_{lmih} - 4K^l{}_{kh}{}^m K_{jlm i}) dV \\ &= \int_M (Z^{kjih} Z_{kj}{}^{lm} Z_{lmih} - 4Z^{kjih} Z^l{}_{kh}{}^m Z_{jlm i}) dV \\ &+ \frac{K}{n(n-1)} \int_M Z^{kjih} (Z_{kjhi} - Z_{kjih} + Z_{jkih} - Z_{kjih}) dV \\ &+ \frac{4K}{n(n-1)} \int_M Z^{kjih} (Z_{ikhj} + Z_{jhki}) dV. \end{aligned}$$

In the last step the second and the third terms are cancelled with each other by virtue of the first identity of Bianchi, so that we have

$$\begin{aligned} & - \int_M (\nabla^l Z^{kjih}) (\nabla_i Z_{kjih}) dV \\ (4) \quad &= \int_M \left(\frac{2K}{n} Z^{kjih} Z_{kjih} + Z^{kjih} Z_{kj}{}^{lm} Z_{lmih} - 4Z^{kjih} Z^l{}_{kh}{}^m Z_{jlm i} \right) dV. \end{aligned}$$

3. At each point P of M let us take an orthonormal frame and consider all components of Z_{kjih} with respect to this frame. By defining $m(P)$ by

$$m(P) = \max \left(\frac{1}{K} |Z_{kjih}| \right),$$

we obtain

$$\begin{aligned} \int_M Z^{kjih} Z_{kjih} dV &\geq K^2 \int_M m^2 dV, \\ \int_M Z^{kjih} Z_{kj}{}^{lm} Z_{lmih} dV &\geq -n^6 K^3 \int_M m^3 dV, \\ - \int_M Z^{kjih} Z^l{}_{kh}{}^m Z_{jlm_i} dV &\geq -n^6 K^3 \int_M m^3 dV. \end{aligned}$$

Hence we have the inequality

$$(5) \quad \int_M m^2 \left(1 - \frac{5}{2} n^7 m\right) dV \leq 0.$$

If (3) holds, then m satisfies

$$m < \frac{2}{5n^7}$$

on M , and we have

$$\int_M m^2 \left(1 - \frac{5}{2} n^7 m\right) dV \geq 0,$$

which and (5) imply that in this case m must be identically zero, and therefore

$$Z_{kjih} = 0.$$

Hence the theorem is proved.

