

VOLUME ESTIMATES FOR KÄHLER-EINSTEIN METRICS: THE THREE-DIMENSIONAL CASE

X.-X. CHEN & S.K. DONALDSON

Abstract

We obtain an estimate for the volumes of neighborhoods of sets of large curvature in three-dimensional Kähler-Einstein manifolds. The key technical step is to prove that a version of monotonicity for L^2 energy holds as long as the underlying region does not “carry homology” (in the sense that the normalized energy in a ball controls the normalized energy in an interior ball).

1. Introduction

This is the first of a series of papers in which we obtain estimates for the volume of certain subsets of Kähler-Einstein manifolds. These estimates form the main analytical input for an approach to general existence questions [9]. Let (M, g) be any compact Riemannian manifold and $r > 0$. We let $K_r \subset M$ be the set of points x where $|\text{Riem}| \geq r^{-2}$ and write Z_r for the r -neighborhood of K_r . Thus any point of the complement $M \setminus Z_r$ is the centre of a metric r -ball on which the curvature is bounded by r^{-2} . Rescaling lengths by a factor r^{-1} , this becomes, in the rescaled metric, a unit ball on which the curvature is bounded by 1. If in addition (as will be the case in our situation), we have control of the local injectivity radius, we can say that, at the length scale r , the complement $M \setminus Z_r$ consists of “good points” with neighborhoods of bounded geometry. Our aim is to derive estimates for the *volume* of the “bad” set Z_r : the set where the geometry need not be standard at this scale.

Let g be a Kähler-Einstein metric so $\text{Ric} = \lambda g$ for constant λ . Then one can find in each complex dimension n constants a_n, b_n , depending only on n , so that pointwise on the manifold,

$$(1) \quad |\text{Riem}|^2 d\mu = (a_n c_1(\text{Riem})^2 + b_n c_2(\text{Riem})) \wedge \omega^{n-2},$$

where we write $c_i(\text{Riem})$ for the standard integrand defining the Chern classes in Chern-Weil theory. (These constants do not depend on λ . As

Received 4/26/2011.

usual we write ω for the 2-form corresponding to g .) Thus on a compact manifold M ,

$$(2) \quad \int_M |\text{Riem}|^2 = \langle a_n c_1^2 \cup \omega^{n-2} + b_n c_2 \cup \omega^{n-2}, M \rangle.$$

The right-hand side is a topological invariant, determined by the Chern classes of M and the Kähler class, which we will denote for brevity by $E(M)$. This identity shows that the curvature of the Kähler-Einstein metric cannot be very large on a set of large volume in M . More precisely, we have an obvious estimate for the volume of K_r ,

$$\text{Vol}(K_r) \leq E(M)r^4.$$

The goal of our work is to estimate the volume of Z_r rather than K_r . We consider Kähler-Einstein metrics with nonnegative Ricci curvature and in this first paper we restrict to complex dimension 3. We consider a compact Kähler-Einstein 3-fold (M, g) with $\text{Ric} = \lambda g, \lambda \geq 0$. We suppose that the metric satisfies the condition that

$$(3) \quad \text{Vol}M \geq \kappa \text{Diam}(M)^6$$

for some $\kappa > 0$. The Bishop-Gromov comparison theorem implies that for all metric balls $B(x, r)$ with $r \leq \text{Diam}(M)$,

$$(4) \quad \text{Vol}B(x, r) \geq \kappa r^6.$$

Our main result is the following.

Theorem 1. *In this situation,*

$$\text{Vol}(Z_r) \leq C (E(M)r^4 + b_2(M)r^6)$$

where C depends only on κ and $b_2(M)$ is the second Betti number of M .

Notice that the statement here is scale invariant (as of course it has to be). When $c_1 > 0$ (which is the case we have mainly in mind), the bound (3) follows from Myers' Theorem, with a constant κ determined by topological data. If we fix the scale by requiring that $\text{Ric} = g$, then the content of the theorem is the bound

$$(5) \quad \text{Vol}(Z_r) \leq Cr^4.$$

We will also establish a small extension, which is probably not optimal.

Theorem 2. *With notation as in Theorem 1, there is a constant C' such that for all r there is a connected open subset $\Omega' \subset M \setminus Z_r$ with*

$$\text{Vol}(M \setminus \Omega') \leq C(E(M)r^{18/5} + b_2(M)r^6).$$

(In applications such as in [9], any bound $O(r^\mu)$ with $\mu > 2$ will suffice.)

We can consider the same questions for Kähler-Einstein metrics in any complex dimension n . When $n = 2$, the proofs are much easier because of the *scale invariance* of the L^2 norm of the curvature in that dimension. In general for a ball $B(x, r)$ (with $r \leq \text{Diam}(M)$), we define the “normalized energy”

$$(6) \quad E(x, r) = r^{4-2n} \int_{B(x,r)} |\text{Riem}|^2 d\mu,$$

which is scale invariant. (When we want to emphasise the dependence on the metric, we write $E(x, r, g)$.) Notice that in our situation we have

$$\kappa r^{2n} \leq \text{Vol}(B(x, r)) \leq \omega_{2n} r^{2n}$$

where ω_{2n} is the volume of the unit ball in \mathbf{R}^{2n} . So it is essentially the same to normalise by the appropriate power of the volume of the ball. Normalized energy functionals of this kind appear in many other contexts in differential geometry, for example the theories of harmonic maps and Yang-Mills fields. In these two theories a crucial *monotonicity* property holds. This is the statement that with a fixed centre the normalized energy is a decreasing function of r . If this monotonicity property held in our situation for Kähler-Einstein metrics, the proof of our theorem would be relatively straightforward. (Of course when $n = 2$, the monotonicity is obviously true.) The main work in this paper is to establish a result which can be seen as an “approximate monotonicity” property. To state this cleanly, let us say that an open subset $U \subset M$ carries homology if the inclusion map $H_2(\partial U, \mathbf{R}) \rightarrow H_2(U, \mathbf{R})$ is not surjective. Note that if $V \subset U$ carries homology, then so does U , and that if $U_1, U_2, \dots, U_p \subset M$ are disjoint domains which each carry homology, then p cannot exceed the second Betti number of M .

Theorem 3. *With the same hypotheses as Theorem 1, for each $\epsilon > 0$ there is a $\delta > 0$ such that if $B(x, r) \subset M$ is a metric ball ($r \leq \text{Diam}(M)$) which does not carry homology and $E(x, r) \leq \delta$, then for any $y \in B(x, r/2)$ and $r' \leq r/2$ we have $E(y, r') \leq \epsilon$.*

The function $\delta(\epsilon)$ depends only on the non-collapsing constant κ .

In Section 2 of this paper we prove Theorem 3. The proof is an application of the extensive theory, due to Anderson, Cheeger, Colding, and Tian, of Gromov-Hausdorff limits of Riemannian manifolds with lower bounds on Ricci curvature. In particular we make use of deep results of Cheeger, Colding, and Tian on codimension-4 singularities and of Cheeger and Colding on tangent cones at infinity. Given Theorem 3, the deduction of Theorem 1, which we show in Section 3, is fairly straightforward. We will first prove a “small energy” result, as follows.

Proposition 1. *With the same hypotheses as Theorem 1, there are δ_0, K such that if $B(x, r) \subset M$ (with $r \leq \text{Diam}(M)$) is a ball which does not carry homology and $E(x, r) \leq \delta_0$, then $|\text{Riem}| \leq Kr^{-2}$ on the interior ball $B(x, r/3)$.*

Theorem 1 follows from a straightforward covering argument. In Section 4 we conclude with some remarks and discussion.

In the sequel to this paper we will extend the results to all dimensions using a rather different argument, making more use of the complex structure and developing ideas of Tian in [13]. (Tian has informed us that, using these ideas, he obtained related results some time ago.) This argument also gives another approach to the three-dimensional case here. But it appears to us worthwhile to write down both proofs.

The authors have had this paper in draft form since early 2010. Recently Cheeger and Naber have posted a preprint [6] establishing these volume estimates in all dimensions, and including results in a more general Riemannian geometry setting. Their approach is somewhat different and it seems valuable to have this variety of arguments in the literature.

Acknowledgments. The second author was partially supported by European Research Council award No. 247331.

2. Proof of Theorem 3

2.1. Cones with small energy. One foundation of our proof of Theorem 3 is a result of Cheeger, Colding, and Tian which states, roughly speaking, that the formation of codimension-4 singularities requires a definite amount of energy. Let $G \subset U(2)$ be a finite group acting freely on S^3 and $n \geq 2$. Consider the unit ball B in the metric product $\mathbf{C}^{n-2} \times (\mathbf{C}^2/G)$ centred at $(0, 0)$. Let V be a Kähler manifold of complex dimension n with nonnegative Ricci curvature and B' be a unit ball centred at $p \in V$. Write $d_{GH}(B, B')$ for the based Gromov-Hausdorff distance.

Proposition 2 ([5], Theorem 8.1). *There are $\alpha_n, \eta_n > 0$ such that if $d_{GH}(B, B') \leq \alpha_n$ and G is non-trivial, then*

$$\int_{B'} |\text{Riem}|^2 \geq \eta_n.$$

Using this, we now prove a result about metric cones in complex dimension 3 with small energy.

Proposition 3. *Let Y be a metric cone with vertex O . Suppose that the based space (Y, O) is the Gromov-Hausdorff limit of based Kähler-Einstein manifolds (of complex dimension 3) (X_n, g_n, O_n) with $\text{Ric}(g_n) = \lambda_n g_n$, where $\lambda_n \geq 0$ and $\lambda_n \rightarrow 0$. Suppose that the X_n satisfy a non-collapsing condition (4), with fixed $\kappa > 0$. Then there is a $\theta > 0$ such that if $E(O_n, 2, g_n) \leq \theta$ for all n , then Y is smooth away from the vertex.*

To see this, we argue by contradiction. Suppose there is a sequence of such examples Y_m , with fixed κ and with $\theta_m \rightarrow 0$. Taking a subsequence, we can suppose these have a based Gromov-Hausdorff limit Y_∞ , and by a diagonal argument this is the limit of a sequence of smooth based manifolds X_n , as above, with $E(O_n, 2, g_n) \rightarrow 0$. The non-collapsing condition means that, according to the Cheeger-Colding-Tian theory, Y_∞ is a smooth Ricci-flat Kähler manifold, of real dimension 6, outside a set $\mathcal{S} \subset Y_\infty$ of Hausdorff dimension at most 2. Further, because of the complex structures present, if the dimension of \mathcal{S} is strictly less than 2, it must be 0, and this means that the singular set contains at most the vertex O , since it is invariant under the dilation action on the cone. So it suffices to show that $\dim \mathcal{S} < 2$. If, on the contrary, the dimension is 2 then, again by the general theory, there is some point with a tangent cone of the form $\mathbf{C} \times \mathbf{C}^2/G$ for a non-trivial G . (In fact this is true for almost all points of the singular set, with respect to Hausdorff 2-measure.) By the invariance under the dilation action, we can suppose this point is at distance 1, say, from the vertex. By the definition of tangent cone and of Gromov-Hausdorff convergence we can find a fixed small ρ with the property that, for all large n , there is a point $p_n \in X_n$ such that the distance in X_n from p_n to O_n is approximately 1 and the Gromov-Hausdorff distance between the ρ -ball in X_n centred at p_n and the ρ -ball in the model space $\mathbf{C} \times \mathbf{C}^2/G$ is less than $\alpha_3\rho$. But, rescaling the result above, this implies that the integral of $|\text{Riem}|^2$ over this ρ -ball in X_n is at least $\eta_3\rho^2$, which gives a contradiction.

Proposition 4. *With the same hypotheses as in Proposition 2, for any σ, L we can choose θ so that for all large n and points x in X_n with $L^{-1} \leq d(x, O_n) \leq L$, we have*

$$|\text{Riem}| \leq \sigma d(x, O_n)^{-2}.$$

This follows from the same argument as above, and general theory. (By results of Anderson [1], at points where the Gromov-Hausdorff limit is smooth, the metrics converge in C^∞ .)

Proposition 5. *With the same hypotheses as in Proposition 2, we can choose θ so that the cone Y must have the form \mathbf{C}^3/Γ for some $\Gamma \subset U(3)$ acting freely on S^5 .*

We know that Y has a Kähler metric with zero Ricci curvature, so this follows from the *rigidity* of \mathbf{C}^3/Γ , among such cones. To give a direct argument, we use the fact that the curvature tensor of a Kähler-Einstein metric satisfies an identity of the form

$$(7) \quad \nabla^* \nabla \text{Riem} = \text{Riem} * \text{Riem},$$

for a certain bilinear algebraic expression $*$. With suitable normalizations, this gives a differential inequality

$$(8) \quad \Delta |\text{Riem}| \geq -|\text{Riem}|^2.$$

(We use the “analysts convention” for the sign of Δ .) Now apply this to our cone Y and set $f = |\text{Riem}|$. Write r for the radial function on the cone. Clearly f is homogeneous of degree -2 , so (in an obvious notation)

$$\frac{\partial f}{\partial r} = -\frac{2}{r}f \quad \frac{\partial^2 f}{\partial r^2} = \frac{6}{r^2}f.$$

Consider the restriction of f to the cross-section $r = 1$ and a point p where f attains its maximal value: m , say. If we write Δ_Σ for the Laplacian on the cross-section, we have the usual formula

$$\Delta = r^{-5} \frac{\partial}{\partial r} \left(r^5 \frac{\partial}{\partial r} \right) + r^{-2} \Delta_\Sigma.$$

By the homogeneity of f we have, at the point p ,

$$\Delta f = -4f + \Delta_\Sigma f,$$

and $\Delta_\Sigma f \leq 0$ at p by the maximum principle. So we deduce that $4m \leq m^2$, and if $m < 4$, we must have $m = 0$.

2.2. The main argument. The second foundation of our proof is the existence, due to Cheeger and Colding [3], of “tangent cones at infinity,” under suitable hypotheses, as metric cones. It is convenient to first state an alternative form of Theorem 3.

Proposition 6. *With the same hypotheses as in Theorem 3. for all $\epsilon > 0$ there is an $A(\epsilon)$ such that if the ball $B(x, r) \subset M$ does not carry homology and $E(x, r) \leq \epsilon A(\epsilon)^{-2}$, then for all $\rho \leq r$ we have $E(x, \rho) \leq \epsilon$.*

To see that this statement implies Theorem 3, let $B(y, r') \subset B(x, r)$ be a ball of the kind considered there. The $r/2$ ball $B(y, r/2)$ centred at y lies in $B(x, r)$ and the corresponding normalized energy is at most 2^2 times $E(x, r)$. It suffices to take $\delta(\epsilon) = \epsilon(2A(\epsilon))^{-2}$ and apply Proposition 6 to $B(y, r/2)$.

Now we start the proof of Proposition 6. Notice that in proving the result, we can obviously assume that ϵ is as small as we please, and we will always suppose that ϵ is less than the constant θ of Proposition 5. First fix ϵ, A and suppose that the statement is false for these parameters, so we have a point x in a manifold M and radii $\rho < r$ with $E(x, \rho) \geq \epsilon, E(x, r) \leq \epsilon A^{-2}$. This implies that $\rho < A^{-1}r$. Choose the largest possible value of ρ , so $E(x, s) < \epsilon$ for $s \in (\rho, r]$ but $E(x, \rho) = \epsilon$. Still keeping ϵ fixed, we suppose that we have such violating examples for a sequence $A_n \rightarrow \infty$, and data $M_n, g_n, x_n, \rho_n, r_n$. Rescale the (Riemannian) metrics by the factor $\rho_n^{-1/2}$: without loss of generality there is a pointed Gromov-Hausdorff limit

$$(M_n, \rho_n^{-1/2} g_n, x_n) \rightarrow (Z, O).$$

Here Z may be singular. Let R_m be any sequence with $R_m \rightarrow \infty$. Then we can rescale the metric on Z by factors R_m^{-1} , and the general

theory of [3] (using the noncollapsing condition (4)) tells us that a subsequence converges to a “tangent cone at infinity” Y which is a metric cone. The convergence is again in the sense of pointed Gromov-Hausdorff limits.

Now go back to the smooth Riemannian manifolds M_n and consider the rescalings $(R_m \rho_n)^{-1/2} g_n$, indexed by n, m . If we choose a suitable function $n(m)$ which increases sufficiently rapidly, then the corresponding sequence of Riemannian manifolds converges to the cone Y . Call these based manifold M_m, \tilde{g}_m, x_m . We can also suppose $n(m)$ increases so rapidly that $A_{n(m)}/R_m$ tends to infinity with m . The choice of parameters means that $E(x_m, s, \tilde{g}_m) < \epsilon < \theta$ if $s \leq A_n/R_m$ and $s > R_m^{-1}$. In particular this holds for $s = 2$, and we are in the position considered in Theorem 2. Thus we deduce that the tangent cone Y is smooth (away from its vertex) and of the form \mathbf{C}^3/Γ . It follows easily that Z is smooth outside a compact set and that the curvature of Z satisfies a bound

$$(9) \quad |\text{Riem}| \leq \sigma r_Z^{-2},$$

(outside a compact set) where r_Z denotes the distance in Z to the base point and the constant σ can be made as small as we like by choosing ϵ small.

The further result that we need is that the curvature actually decays faster.

Proposition 7. *There is a $\gamma > 3$ and $K > 0$ such that outside a compact subset of Z .*

$$|\text{Riem}| \leq K r_Z^{-\gamma}.$$

In fact our proof will establish the result for any $\gamma < 4$. The decay rate r^{-4} arises as that of the Green’s function in real dimension 6. We postpone the proof of Proposition 7, which is somewhat standard, and move on to complete the proof of Theorem 3, assuming this.

Consider the general situation of a domain $U \subset M$ with smooth boundary ∂U and a Kähler metric on M . Suppose that

- The cohomology class of ω in $H^2(\partial U)$ is zero.
- U does not carry homology.
- $H^1(\partial U) = 0$.

The first two conditions imply that the class of ω in $H^2(U)$ also vanishes, so we can write $\omega = da$ for a 1-form a on U . Now let p be the invariant polynomial corresponding to the characteristic class $a_3 c_1^2 + b_3 c_2$, as discussed in the Introduction. Thus $p(\text{Riem})$ is a closed 4-form on U and if the metric is Kähler-Einstein, we have

$$\int_U |\text{Riem}|^2 = \int_U p(\text{Riem}) \wedge \omega.$$

Then

$$(10) \quad \int_U p(\text{Riem}) \wedge \omega = \int_{\partial U} p(\text{Riem}) \wedge a.$$

Further, the third condition implies that if \tilde{a} is *any* 1-form on ∂U with $d\tilde{a} = \omega|_{\partial U}$, then

$$(11) \quad \int_{\partial U} p(\text{Riem}) \wedge a = \int_{\partial U} p(\text{Riem}) \wedge \tilde{a}.$$

Thus the integral of $p(\text{Riem}) \wedge \omega$ over U is determined by data on the boundary, and if the metric is Kähler-Einstein, this coincides with the integral of $|\text{Riem}|^2$ over U . In particular we get an inequality

$$(12) \quad \int_U |\text{Riem}|^2 \leq c \max_{\partial U} |\text{Riem}|^2 \int_{\partial U} |a|,$$

for some fixed constant c .

To apply this, consider first the flat cone \mathbf{C}^3/Γ and let Σ_r be the cross-section at radius r . Then the real cohomology of Σ_r vanishes in dimensions 1 and 2 and we can obviously write

$$\omega|_{\Sigma_r} = da_r$$

with $|a_r| = O(r)$. In fact we can take a_r to be one-half the contraction of ω with the vector field $r \frac{\partial}{\partial r}$, so $|a_r| = r$. Let $V_\Gamma r^5$ be the 5-volume of Σ_r .

Now choose R_0 so large that $cK^2 R_0^{6-2\gamma} V_\Gamma < \epsilon/2$. By our discussion of the cone at infinity in Z , we can choose $R \geq R_0$ such that near the level set $r_Z = R$ in Z there is a hypersurface Σ' with the property that the geometry of ω_Z restricted to Σ' is very close to that of the cone metric restricted to Σ_R , in an obvious sense. In particular we can suppose that $\omega_Z|_{\Sigma'} = da'$ where

$$\int_{\Sigma'} |a'| \leq \frac{3}{2} V_\Gamma R^6,$$

say. The bound on the curvature in Proposition 7 implies that

$$(13) \quad c \max_{\Sigma'} |\text{Riem}|^2 \int_{\Sigma'} |a'| \leq \frac{3\epsilon}{4}.$$

From now on R is fixed. We go back to the manifolds M_n with rescaled metrics $\rho_n^{-1/2} g_n$ converging to Z . We choose n so large that there is hypersurface $\Sigma'' \subset M_n$ on which the geometry is close to that of Σ' . Then Σ'' is the boundary of an open set $U \subset M_n$ which does not carry homology by hypothesis. By taking n large, we can make the boundary term in (10) as close as we like to that estimated in (12), (13). So we can suppose that

$$\int_U |\text{Riem}|^2 \leq \frac{5\epsilon}{6},$$

say. But the U contains the unit ball centred at x_n over which the integral of $|\text{Riem}|^2$ is ϵ by construction. This gives the desired contradiction.

2.3. Curvature decay. Here we prove Proposition 7. The proof exploits the differential inequality

$$\Delta|\text{Riem}| \geq -|\text{Riem}|^2 \geq -\sigma r^{-2}|\text{Riem}|.$$

To explain the argument, consider first a slightly different problem in which we work on a cone Y (of real dimension 6) with radial function r , and we have a smooth positive function f on the set $r \geq 1$ with $\Delta f \leq \sigma r^{-2}f$ and $f \leq \sigma r^{-2}$. We have

$$(14) \quad \left(\Delta + \frac{\sigma}{r^2}\right)r^\lambda = (\lambda(\lambda + 4) + \sigma)r^{\lambda-2}.$$

Let $\alpha = -2 - \sqrt{4 - \sigma}, \beta = -2 + \sqrt{4 - \sigma}$: the roots of the equation $\lambda(\lambda + 4) + \sigma = 0$. We suppose σ is small, so α is close to -4 and β is close to 0 . Then any linear combination

$$g = Ar^\alpha + Br^\beta$$

satisfies the equation $(\Delta + \frac{\sigma}{r^2})g = 0$. Fix $R > 1$ and let m_1, m_R be the maximum values of f on the cross-sections $r = 1, r = R$ respectively. We choose constants A, B so that $g(1) \geq m_1, g_R \geq m_R$. If we solve for the case of equality, we get

$$A_0 = \frac{R^\beta m_1 - m_R}{R^\beta - R^\alpha}, \quad B_0 = \frac{m_R - R^\alpha m_1}{R^\beta - R^\alpha}.$$

Since $R^\beta > R^\alpha$ it certainly suffices to take

$$A = \frac{R^\beta m_1}{R^\beta - R^\alpha}, \quad B = \frac{m_R}{R^\beta - R^\alpha}.$$

Then set $u = f - g$ so that $u \leq 0$ when $r = 1$ or $r = R$ and $(\Delta + \sigma r^{-2})u \geq 0$. We claim that $u \leq 0$ throughout the region $1 \leq r \leq R$. For if we write $u = hr^{-2}$, calculation gives

$$(15) \quad \left(\Delta + \frac{\sigma}{r^2}\right)u = r^{-2}\left(\Delta h - 2r^{-1}\frac{\partial h}{\partial r} - (4 - \sigma)r^{-2}h\right)$$

and (since $4 - \sigma > 0$) we see that there can be no interior positive maximum of h .

We see then that for $1 < r < R$ we have

$$f \leq Ar^\alpha + Br^\beta$$

where A, B are given by the formulae above. Using the information $m_1 \leq \theta, m_R \leq \sigma R^{-2}$, we get

$$Ar^\alpha \leq cr^\alpha, Br^\beta \leq cR^{-\beta+2}r^\beta$$

for some fixed c . Taking R very large compared to r , we get $f \leq 2cr^\alpha$, say.

We want to adapt this argument to the function $f = |\text{Riem}|$ on Z . Recall that Z has base point O and r_Z is the distance to O . We write

$$\mathcal{A}(r_1, r_2) = \{z \in Z : r_1 < r_Z(z) < r_2\}$$

with $r_2 = \infty$ allowed.

There are several complications. One minor difficulty is that f may not be smooth, but this is handled by standard approximation arguments. The second is that the manifold is not exactly a cone, even at large distances, and the radius function r_Z need not be smooth.

Lemma 1. *For any $\tau > 0$ we can find an $R_\tau > 0$ and a smooth function \underline{r} on the region $\mathcal{A}(R_\tau, \infty) \subset Z$ such that*

- $|\frac{\underline{r}}{r_Z} - 1| \leq \tau$,
- $||\nabla \underline{r}| - 1| \leq \tau$,
- $|\Delta \underline{r}^2 - 12| \leq \tau$.

We only sketch the proof. It is clear that we can choose R_τ so that we can find such a function on any annulus $\mathcal{A}(2^p R_\tau, 2^{p+1} R_\tau)$ for $p = 1, 2, \dots$. This just uses the convergence of the rescaled metric to the cone. If we extend these annuli slightly, we get a sequence of overlapping annuli and a function defined on each. To construct \underline{r} we glue these together using cut-off functions. Notice that we only have to glue adjacent terms so that the gluing errors do not accumulate.

Now think of σ and τ as fixed small numbers. (It will be clear from the discussion below that one could calculate appropriate values explicitly: for example $\sigma = \tau = 1/100$ will do.) We want to adapt the preceding argument to prove Proposition 7. There is no loss in supposing that in fact $R_\tau = 1/2$ and that $|\text{Riem}| \leq \sigma |\text{Riem}|^2$ on $\underline{r} \geq 1$. Let α, β be the roots as above and choose α', β' with α' slightly greater than α and β' slightly less than β . Then if τ is small, we will have

$$(16) \quad (\Delta + \sigma \underline{r}^{-2}) \underline{r}^{\alpha'}, (\Delta + \sigma \underline{r}^{-2}) \underline{r}^{\beta'} \leq 0.$$

We want to choose A, B such that $g(\underline{r}) = A \underline{r}^{\alpha'} + B \underline{r}^{\beta'}$ has $\Delta + \sigma \underline{r}^{-2} \leq 0$ and $g(1) \geq m_1, g(R) \geq m_R$ where m_1, m_R are the maxima of f on $\underline{r} = 1, \underline{r} = R$ respectively. We take

$$A = \frac{m_1 R^{\beta'}}{R^{\beta'} - R^{\alpha'}}, \quad B = \frac{m_R}{R^{\beta'} - R^{\alpha'}}.$$

Then $A, B \geq 0$, so the differential inequality follows from (16). Now consider $u = f - g$ and argue as before to show that u has no interior maximum. We have

$$\Delta \underline{r}^{-2} = \underline{r}^{-4} (8 |\nabla \underline{r}|^2 - \Delta \underline{r}^2) \leq (-4 + 9\delta) \underline{r}^{-4},$$

and the same argument goes through.

3. Proof of Theorem 1

We begin with a standard result.

Proposition 8. *Given $\kappa > 0$ there is a $\chi > 0$ such that if B is any unit ball in a Kähler-Einstein manifold (of real dimension 6) and*

- 1) $|\text{Riem}| = 1$ at the center of B ;
- 2) $|\text{Riem}| \leq 4$ throughout B ;
- 3) $\text{Vol}(B) \geq \kappa$,

then $\int_B |\text{Riem}|^2 \geq \chi$.

One way to prove this is to apply the Moser iteration technique to $|\text{Riem}|$, using the differential inequality (8) and the fact that in this situation the Sobolev constant is bounded. Another method is to use elliptic estimates in harmonic coordinates.

Proposition 9. *Suppose M is a Kähler-Einstein manifold as considered in Theorem 1. There is an $\epsilon > 0$ with the property that, if U is any domain such that the normalized energy of any ball contained in U is less than ϵ , then $|\text{Riem}| \leq 4d^{-2}$, where d denotes the distance to the boundary of U .*

To see this, we let S be the maximum value of $d^{-2}|\text{Riem}|$ over U and suppose that this is attained at p . If $S > 4$, then $d(q) \geq d(p)/2$ for any point q in the ball of radius $d(p)S^{-1/2}$ centred at p . Rescale this ball to unit size and we are in the situation considered in the preceding proposition. If we take ϵ to be the constant χ appearing there, we get a contradiction, so in fact $S \leq 4$.

We can now prove Proposition 1. Take ϵ as above and let $\delta(\epsilon)$ be the value given by Theorem 3. Suppose that $B(x, r)$ is a ball of normalized energy less than δ and let U be the half-sized ball. Then Theorem 3 tells us that the normalized energy of any ball in U is at most ϵ and we can apply the result above to see that $|\text{Riem}| \leq 4.6^2 r^{-2}$ in $B(x, r/3)$.

Of course some of the constants appearing in the statements above (such as the value 4 in Proposition 8) are rather arbitrary. The essential point is that there is a definite threshold value, so that if the normalized energy is below this threshold, on a ball which does not carry homology, then we get complete control of the metric on interior regions.

Now we prove Theorem 1. Given r and a Kähler-Einstein metric (M, g) as considered there, we pick a maximal collection of point $x_a, a \in I$ in M such that the distance between any pair is at least r . Then the r -balls $B(x_a, r)$ cover M . Consider a ball $B(x_a, 12r)$. If this ball does not carry homology and its normalized energy is less than δ , then by the result above we have $|\text{Riem}| \leq 1$ on $B(x_a, 2r)$. Thus no point in Z_r can lie in $B(x_a, r)$, and thus Z_r is covered by balls $B(x_a, r)$ where either $E(x_a, 12r) > \delta$ or $B(x_a, 12r)$ carries homology. Let $I' \subset I$ denote the indices of the first kind and $I'' \subset I$ those of the second.

Suppose that N balls of the form $B(x_a, 12r)$ have a non-empty common intersection. Let q be a point in the intersection, so the N centres x_a all lie in the $12r$ ball centred at q . By construction the balls $B(x_a, r/2)$ are disjoint and have volume at least $(\kappa/2^6)r^6$. Since the volume of $B(q, 12.5r)$ is bounded above by a fixed multiple of r^6 , this gives a fixed bound on N , independent of r . Thus

$$\sum_{a \in I'} \int_{B(x_a, 12r)} |\text{Riem}|^2 \leq N \int_M |\text{Riem}|^2.$$

On the other hand, by definition, $E(x_a, 12r) \geq \delta$ for $a \in I'$, so we see that the number of elements of I' is at most

$$|I'| \leq \frac{N}{\delta(12r)^2} \int_M |\text{Riem}|^2 = C_1 r^{-2},$$

say.

By a similar argument there is a fixed upper bound N' on the number of balls $B(x_a, 12r)$ which can meet any given one. It follows that the number of these balls which carry homology is bounded by N' times the second Betti number of M . So $|I''|$ is bounded by a fixed number. Then

$$\text{Vol}(Z_r) \leq \text{constant } r^6(|I'| + |I''|) \leq C_1 r^4 + C_2 r^6.$$

Now we turn to Theorem 2. Recall that this states that we can find a *connected* open subset $\Omega' \subset M \setminus Z_r$ so that the volume of the complement of Ω' is bounded by a multiple of $r^{18/5}$.

To prove this, we recall that in our situation there is a bound on the isoperimetric constant, due to Croke [8]. If $H \subset M$ is a rectifiable hypersurface dividing M into two components M_1, M_2 with $\text{Vol}(M_1) \leq \text{Vol}(M_2)$, then

$$(17) \quad \text{Vol}(M_1)^{1/6} \leq k \text{Vol}(H)^{1/5}$$

for a fixed constant k . By the construction in the proof of Theorem 1 above, the set Z_r is contained in W , which is a union of P balls of radius r with $P \leq C(r^{-2} + 1)$. By the Bishop comparison theorem the 5-volume of the boundary of one of these balls is bounded by Cr^5 . The boundary ∂W of W is a rectifiable set of 5-volume at most the sum of the boundaries of the balls, thus $\text{Vol}(\partial W) \leq C(r^3 + r^5)$. We can normalise so that the volume of M is 1 and then, without loss of generality, suppose that r is so small that $\text{Vol}(W) \leq 1/10$ and $k^5 \text{Vol}(\partial W) \leq 9/10$. Let Ω_i be the connected components of $M \setminus W$. If a component Ω_1 has volume greater than $1/2$ (i.e. one half the volume of M), then by (14) its complement has volume less than $k^6 \text{Vol}(\partial W)^{6/5} = k^6 C^{6/5} r^{18/5}$ and we can take $\Omega' = \Omega_1$. So suppose that all components Ω_i have volume less than $1/2$. Then it is clear that

$$\sum_i \text{Vol}(\partial \Omega_i) = \text{Vol}(\partial W)$$

while

$$\sum_i \text{Vol}(\Omega_i) \geq 9/10.$$

The second equation implies that $\sum_i \text{Vol}(\Omega_i)^{5/6} \geq 9/10$ and then (17) gives

$$\sum_i \text{Vol}(\partial\Omega_i) \geq 9k^{-5}/10,$$

so

$$\text{Vol}(\partial W) \geq 9k^{-5}/10,$$

contrary to our assumptions.

4. Discussion

- 1) Proposition 5 and the ensuing arguments in 2.2 above are closely related to a result of Cheeger, Colding, and Tian ([5], Theorem 9.26). Let X be a complete, noncompact Ricci-flat Kähler manifold of complex dimension n with base point p . Suppose that $\text{Vol}(B(p, R)) \geq \kappa R^{2n}$ and

$$R^{4-2n} \int_{B(p,R)} |\text{Riem}|^2 \rightarrow 0$$

as $R \rightarrow \infty$. Tian conjectured in [12] that in this situation X is an ALE manifold, with tangent cone at infinity of the form \mathbf{C}^n/Γ . When $n = 3$, this conjecture was established in the result quoted above. Our statements are a little different since we establish a definite “small asymptotic energy threshold” which implies that the manifold is ALE.

- 2) In our situation the tangent cone Y of Z at infinity is unique. In general, positive Ricci curvature does not imply uniqueness of tangent cones, even when $|\text{Riem}| \leq cr^{-2}$. See the discussion in [4] of examples, including an unpublished example due to Perelman.
- 3) Theorem 3 becomes false if we omit the condition that the ball does not carry homology. To see this, one can consider for example the quotient $\underline{M} = T^6/\Gamma$ of a complex torus by a group Γ of order 3, acting with isolated fixed points. Then \underline{M} has a resolution M with $c_1(M) = 0$. According to Joyce [11], there is a family of ALE metrics on the resolution of \mathbf{C}^3/Γ , parametrised by the Kähler class. For suitable Kähler classes on M , the Calabi-Yau metric is approximately given by gluing rescaled versions of these ALE metrics to the flat metric on \underline{M} , just as in the familiar picture of the Kummer construction for K3 surfaces. For any $\delta > 0$, we can use this scaling to find a Ricci-flat metric on M , a unit ball $B(x, 1) \subset M$ on which the normalized energy is less than δ , and an interior ball $B(x, \rho)$ on which the normalized energy exceeds c , for some fixed c . The arguments in our forthcoming paper will

show that this is essentially the only way in which approximate monotonicity can fail; see also [13].

- 4) It is interesting to ask if similar results to those proved above can be established for constant scalar curvature and extremal Kähler metrics. For this it might be sensible to assume a bound on the Sobolev constant. Perhaps some of the techniques used in [14], [7] can be applied to this problem.
- 5) Another question is whether a result like Theorem 3 holds for general 6-dimensional Einstein metrics (with nonnegative Ricci curvature and a non-collapsing condition). It might be that a different topological side condition is appropriate.
- 6) The function $\delta(\epsilon)$ in Theorem 3 depends only on the collapsing constant κ . It would be interesting to determine the function effectively, but our method cannot do this. One suspects that, in reality, it may be possible to take $\delta(\epsilon)$ not much smaller than ϵ , and that the constant C in Theorem 1 can (in reality) be taken not too large. Some results on the numerical analysis of Kähler-Einstein, and more generally extremal, metrics seem to give evidence for this suspicion ([2], [10]) but a theoretical derivation of realistic estimates seems a long way off.

References

- [1] M.T. Anderson, *Convergence and rigidity of metrics under Ricci curvature bounds*, Invent. Math. **102** 429–445 (1990), MR 1074481 Zbl 0711.53038.
- [2] R.S. Bunch & S.K. Donaldson, *Numerical approximations to extremal metrics on toric surfaces*, In: Handbook of Geometric Analysis No. 1 1–28 Adv. Lect. Math. Int. Press (2008), MR 2483361.
- [3] J. Cheeger & T.H. Colding, *Lower bounds on the Ricci curvature and the almost rigidity of warped products*, Annals of Math. **144** 189–237 (1996) MR 1405949 Zbl 0865.53037.
- [4] J. Cheeger & T.H. Colding, *On the structure of spaces with Ricci curvature bounded below, I*, Jour. Differential Geometry, **46** 406–480 (1997), MR 1484888 Zbl 0902.53034.
- [5] J. Cheeger, T.H. Colding & G. Tian, *On the singularities of spaces with bounded Ricci curvature*, Geometric and Functional Analysis **12** 873–914 (2002), MR 1937830 Zbl 1030.53046.
- [6] J. Cheeger & A. Naber, *Lower bounds on Ricci curvature and quantitative behaviour of singular sets*, arxiv.1103.1819 (To appear in Inventiones Math.).
- [7] X.-X. Chen & B. Weber, *Moduli spaces of critical Riemannian metrics with $L^{n/2}$ norm of curvature bounded*, Adv. Math. **226** No. 2 1307–1370 (2011), MR 2737786 Zbl 1205.53074.
- [8] C.B. Croke, *Some isoperimetric inequalities and eigenvalue estimates*, Ann. Sc. Ec. Norm. Sup. Paris **13** 419–435 (1980), MR 0608287 Zbl 0465.53032.
- [9] S. Donaldson, *Stability, birational transformations and the Kähler-Einstein problem*, arxiv.1007.4220 (To appear in Surveys in Differential Geometry Vol. 17.)

- [10] M.R. Douglas, R.L. Karp, S. Lukic & R. Reinbacher, *Numerical Calabi-Yau metrics*, J. Math. Phys. **49** No. 3 032302 (2008), MR 2406784 Zbl 1153.81351.
- [11] D. Joyce, *Compact manifolds with special holonomy*, Oxford Math. Monographs Oxford UP 2000 MR 1787733 Zbl 1027.53052.
- [12] G. Tian, *Kähler-Einstein metrics on algebraic manifolds*, Proceedings Int. Cong. Math. 1990, Math. Soc. Japan 1991 587–598 MR 1159246.
- [13] G. Tian, *Compactness theorems for Kähler-Einstein manifolds of dimension 3 and up*, Jour. Differential Geometry **35** 535–558 (1992) MR 1163448 Zbl 0766.53054.
- [14] G. Tian & J. Viaclovsky, *Moduli spaces of critical Riemannian metrics in dimension 4*, Adv. Math. **196** No. 2 346–372 (2005)MR 2166311 Zbl pre02213018.

DEPT OF MATHEMATICS
STONY BROOK UNIVERSITY
STONY BROOK, NY 11794
E-mail address: xiu@math.sunysb.edu

DEPARTMENT OF MATHEMATICS
IMPERIAL COLLEGE
QUEEN'S GATE
LONDON SW7 2AZ
E-mail address: s.donaldson@imperial.ac.uk