

## DISPERSIONLESS INTEGRABLE SYSTEMS IN 3D AND EINSTEIN–WEYL GEOMETRY

EUGENE V. FERAPONTOV & BORIS S. KRUGLIKOV

### Abstract

For several classes of second-order dispersionless PDEs, we show that the symbols of their formal linearizations define conformal structures that must be Einstein–Weyl in 3D (or self-dual in 4D) if and only if the PDE is integrable by the method of hydrodynamic reductions. This demonstrates that the integrability of these dispersionless PDEs can be seen from the geometry of their formal linearizations.

### 1. Introduction

Let

$$(1) \quad F(x^i, u, u_{x^i}, u_{x^i x^j}, \dots) = 0$$

be a partial differential equation (PDE), where  $u$  is a (scalar) function of the independent variables  $x^1, \dots, x^n$ . The formal linearization of (1) results upon setting  $u \rightarrow u + \epsilon v$  and keeping terms of the order  $\epsilon$ . This leads to a linear PDE for  $v$ ,

$$(2) \quad \ell_F(v) = 0,$$

where  $\ell_F$  is the operator of formal linearization,

$$\ell_F = F_u + F_{u_{x^i}} \mathcal{D}_{x^i} + F_{u_{x^i x^j}} \mathcal{D}_{x^i} \mathcal{D}_{x^j} + \dots$$

(here  $\mathcal{D}_{x^i}$  is the operator of total differentiation by  $x^i$ ). Note that, for nonlinear  $F$ , the formal linearization depends on the solution  $u$ . For instance, the linearization of the dispersionless Kadomtsev–Petviashvili (dKP) equation,  $u_{xt} - (uu_x)_x - u_{yy} = 0$ , reads as  $v_{xt} - (uv)_{xx} - v_{yy} = 0$ .

The linearized equation (2) appears in a wide range of constructions and applications:

- *Stability analysis* of a solution  $u$  is based on the investigation of the spectrum of the linearized operator; this goes back to Lyapunov.
- *Classical, contact, and higher symmetries* of equation (1) correspond to solutions of the linearized equation of the form  $v = v(x^i, u, u_{x^i}, u_{x^i x^j}, \dots)$ , which are required to satisfy (2) identically modulo (1); this goes back to Lie [41, 44, 55].

---

Received 10/30/2012.

- *Contact invariants* of scalar ordinary differential equations can be obtained from the Wilczynski invariants of the corresponding linearized equations [14, 15].
- *Generalized Laplace invariants* of PDEs appearing in the context of Darboux integrability can be obtained from the Laplace invariants of linearized equations [71, 2, 43, 46, 45].
- *Algebraic integrability* of ordinary differential equations can be seen from the structure of the differential Galois group of linearized equations: it must be Abelian; see [52] and references therein.

In general, coefficients of a linear PDE have differential-geometric meaning. In particular, its symbol can be interpreted as a symmetric tensor field. The following natural question arises:

*Can one read integrability of a given PDE off the geometry of its formal linearization?*

In this paper we answer this question in the affirmative for the four particularly interesting classes of PDEs in 3D, namely

#### Equations of type I

$$(3) \quad (a(u))_{xx} + (b(u))_{yy} + (c(u))_{tt} + 2(p(u))_{xy} + 2(q(u))_{xt} + 2(r(u))_{yt} = 0.$$

This class was introduced in [18] in the context of the “central quadric ansatz”; see the remark at the end of Section 3. The corresponding integrability aspects were discussed in [32].

#### Equations of type II

$$(4) \quad f_{11}u_{xx} + f_{22}u_{yy} + f_{33}u_{tt} + 2f_{12}u_{xy} + 2f_{13}u_{xt} + 2f_{23}u_{yt} = 0.$$

Here the coefficients  $f_{ij}$  depend on the first-order derivatives  $u_x, u_y, u_t$  only. Equations of this type can be called quasilinear wave equations, their integrability was analyzed in [8, and references therein].

#### Equations of type III

$$(5) \quad F(u_{xx}, u_{xy}, u_{yy}, u_{xt}, u_{yt}, u_{tt}) = 0.$$

Equations of this form are known as the dispersionless Hirota type or Hessian type equations. Their integrability was studied in [56, 29].

#### Equations of type IV

$$(6) \quad A(u)u_x + B(u)u_y + C(u)u_t = 0.$$

Here  $u$  is a two-component column vector, and  $A, B, C$  are  $2 \times 2$  matrices. Equations of this form are known as systems of hydrodynamic type. Their integrability was investigated in [27]. It was demonstrated in [54] that coefficients of the generic integrable equations of the types (4), (5), and (6) can be parametrized by generalized hypergeometric functions.

Equations (3)–(6) belong to the class of dispersionless PDEs. They arise in a wide range of applications in mathematical physics, general relativity, differential geometry, and the theory of integrable systems (as

dispersionless limits of integrable soliton equations of the KP/Toda type [70]; see Section 9). In the dispersionless limit, the familiar “solitonic” integrability (based on Lax pairs, algebro-geometric techniques, etc.) requires a modification. An adequate approach is provided by the method of hydrodynamic reductions [26], which is based on the requirement of the existence of special multi-phase solutions that can be viewed as dispersionless analogues of multi-soliton/multi-gap solutions. To make this paper as self-contained as possible, in Section 9 we included an appendix with a brief overview of this approach, and further references.

Solutions to equations (3)–(6) carry a canonical conformal structure that can be defined as follows: The symbol of formal linearization is a symmetric (2,0)-tensor  $g^\sharp \in \Gamma(S^2TM)$  on the base manifold  $M$  with coordinates  $(x^1, x^2, x^3) = (x, y, t)$ , which depends on a finite jet of the solution  $u$  (in the case (6) we use the dispersion relation). This tensor is only defined up to multiplication by a non-zero factor, which makes our theory conformal. All considerations are local, and  $M$  will be identified with an open domain of  $\mathbb{R}^3$ . We will always assume  $g^\sharp$  to be non-degenerate; in this case, the inverse (0,2)-tensor  $g \in \Gamma(S^2T^*M)$  defines a metric. In coordinates,  $g = g_{ij} dx^i dx^j$  where  $g_{ij}$  is the inverse of the matrix of  $g^\sharp$ . We will assume the Lorentzian signature of  $g$ ; this is equivalent to the requirement of hyperbolicity of the corresponding PDE (see Section 9 for a discussion of the elliptic case). The conformal class  $[g]$  of the metric  $g$  is the key invariant responsible for the linearizability/integrability of the equations under study. Our main results can be summarized as follows:

- Equations (3)–(6) are linearizable (by a transformation from the natural equivalence group specified in each particular case below) if and only if the corresponding conformal structures  $g$  are conformally flat on every solution (an extra condition is needed in the case (6)). This provides a simple linearizability test based on the vanishing of the corresponding Cotton tensor.
- Equations (3)–(6) are integrable by the method of hydrodynamic reductions if and only if the corresponding conformal structures  $g$  are Einstein–Weyl on every solution (again, an extra condition is needed for (6)). Recall that an Einstein–Weyl structure consists of a symmetric connection  $\mathbb{D}$  and a conformal structure  $g$  such that (a) the connection  $\mathbb{D}$  preserves the conformal class,  $\mathbb{D}[g] = 0$ , and (b) the trace-free part of the symmetrized Ricci tensor of  $\mathbb{D}$  vanishes.

In coordinates, this gives

$$(7) \quad \mathbb{D}_k g_{ij} = \omega_k g_{ij}, \quad R_{(ij)} = \Lambda g_{ij},$$

where  $\omega = \omega_k dx^k$  is a covector,  $R_{(ij)}$  is the symmetrized Ricci tensor of  $\mathbb{D}$ , and  $\Lambda$  is some function [11]. In fact, one needs to specify

$g$  and  $\omega$  only; then the first set of equations uniquely determines  $\mathbb{D}$ . We point out that for all examples considered in this paper, the covector  $\omega$  is expressed in terms of  $g$  by the universal explicit formula

$$(8) \quad \omega_k = 2g_{kj}\mathcal{D}_{x^s}(g^{js}) + \mathcal{D}_{x^k}(\ln \det g_{ij}).$$

Note that in 3D this formula is invariant under the transformation  $g \rightarrow \lambda g$ ,  $\omega \rightarrow \omega + d \ln \lambda$ , which is characteristic of the Einstein–Weyl geometry.

We recall that the Einstein–Weyl equations (7) are integrable by twistor-theoretic methods [39]. Thus, solutions of integrable PDEs carry “integrable” geometry. Equivalently, one can say that second-order dispersionless integrable systems in 3D (having non-degenerate symbol) can be viewed as reductions of the Einstein–Weyl conditions, which therefore play the role of a universal “master-equation” [69]. Let us mention that relations of dispersionless integrable systems to the Einstein–Weyl geometry have been discussed in [68, 9, 10, 47, 20, 22, 23, 19, 36].

Given a class of integrable PDEs such as (4)–(6), the verification of the Einstein–Weyl conditions (7) can be a formidable task, primarily due to a rather intricate structure of the integrability conditions. A way to bypass computational difficulties is to use the result of Cartan [11], which says that the Einstein–Weyl property of a triple  $(\mathbb{D}, g, \omega)$  is equivalent to the existence of a two-parameter family of surfaces that are null with respect to the conformal structure  $g$  (that is, tangential to the null cones of  $g$ ) and totally geodesic in the Weyl connection  $\mathbb{D}$ . In the context of dispersionless integrable systems, such surfaces are provided by the corresponding dispersionless Lax pairs: these consist of  $\lambda$ -dependent vector fields  $X, Y$  that are required to commute modulo the equation, identically in the “spectral parameter”  $\lambda$  (for all classes of PDEs discussed in this paper, the existence of such Lax pairs is equivalent to the integrability by the method of hydrodynamic reductions). Note that  $X$  and  $Y$  may contain derivatives with respect to  $\lambda$ .

Taking integral surfaces of the distribution spanned by  $X, Y$  in the extended four-space with coordinates  $x, y, t, \lambda$ , and projecting them down to the space of independent variables  $x, y, t$ , we obtain the required two-parameter family of null totally geodesic surfaces. Computationally, this approach has an advantage, allowing one to avoid working with the full set of integrability conditions (the problem is to prove the existence of a Lax pair, and this is where the integrability conditions are needed). In this approach, the key object within the triple  $(\mathbb{D}, g, \omega)$  is the connection  $\mathbb{D}$ , which is uniquely specified by the given two-parameter family of null totally geodesic surfaces. Both Einstein–Weyl conditions (7) will be satisfied automatically.

Our main results relating linearizability/integrability to geometry of formal linearizations are proved in Sections 3–6 (we find it more convenient to treat the above four classes separately: explicit forms of the corresponding linearizability/integrability conditions are rather different). Known integrable equations of types (3)–(6) provide an abundance of Einstein–Weyl structures parametrized by elementary functions, elliptic functions, modular forms, and Painlevé transcendents. Some further examples are collected in Section 7. Most of the examples of Einstein–Weyl structures exhibited in this paper are apparently new (otherwise, a reference is given).

In Section 8 we discuss geometric aspects of integrability of second-order dispersionless PDEs in 4D, indicating that the associated conformal structures must be self-dual. This is in agreement with the fundamental fact that the Einstein–Weyl equations are reductions of the equations of self-duality [42, 9, 10].

The method of hydrodynamic reductions, providing an efficient approach to the integrability of equations (3)–(6), is summarized in Section 9. Hydrodynamic reductions of elliptic PDEs are also discussed.

In calculations of the Cotton tensor and the Einstein–Weyl conditions, we used symbolic packages of Maple. We shall omit unnecessary lengthy formulae from the text. All relevant programs, including the integrability conditions of equations (3)–(6) and details of proofs from Sections 3–6, are available from arXiv:1208.2728v3.

Since the involved functions are rational differential expressions with integer coefficients, the results are exact and the proofs are rigorous.

**Conventions.** All our considerations are micro-local, i.e., local for a solution, with the size of neighbourhood also depending on (jet of) the solution. We work either in the real smooth category or in the complex holomorphic category. In the latter case we only assume non-degeneracy of the symbol, while in the former we assume that the symbol is hyperbolic (see the next section for precise definition). As our approach to the dispersionless integrability is based on the method of hydrodynamic reductions, which generally refers to hyperbolic systems, we assume the Lorentzian signature of  $g$  in 3D, and the neutral signature in 4D, but this requirement can be removed without restricting the generality if the PDE in question is analytic. In fact, one can treat the elliptic case by the complexification approach; see Section 9 for more details.

Although results of this paper are local, we perceive that global versions may be available through the twistor theory.

**Acknowledgments.** We would like to thank D. Calderbank, M. Dunajski, K. Khusnutdinova, and M. Pavlov for useful comments and clarifying discussions. We also thank the referee for numerous suggestions, which helped to improve the presentation. We acknowledge financial

support from the LMS (BK) and the University of Tromsø (EVF) making this collaboration possible. Special thanks go to the University Hospital of North Norway, where part of this work was completed. The research of EVF was partially supported by the European Research Council Advanced Grant FroM-PDE.

## 2. Preliminaries

In this section we discuss the necessary background material. For simplicity we restrict to the case of a single partial differential equation  $\mathcal{E}$  of the second order. A scalar second-order differential operator is a function  $F \in C^\infty(J^2M)$  on the space of 2-jets of the base manifold  $M$ , and the equation  $\mathcal{E} = \{F = 0\}$  is a submanifold in  $J^2M$ . We shall assume, as is customary, that the projection  $p_{2,1} : \mathcal{E} \rightarrow J^1M$  is a submersion. Then  $\pi_{\mathcal{E}} : \mathcal{E} \rightarrow M$  is also a submersion.

As described in the Introduction, the linearization operator  $\ell_F$  is a second-order linear differential operator defined modulo the equation  $\mathcal{E}$ , which means that its coefficients, being functions of the finite-order jets of  $u$ , are subject to this PDE. The important property of formal linearization is its contact invariance, which ensures that contact transformations lift naturally to the tangent bundle (covering) of the equation. To the best of our knowledge, this invariance was explored for the first time in [2].

In this paper we need a simpler fact that the symbol  $\sigma_F$  of the linearization operator  $\ell_F$ , also called the symbol of  $F$ , is contact invariant. The symbol is a bi-vector (section)  $\sigma_F \in C^\infty(\mathcal{E}, \pi_{\mathcal{E}}^* S^2 T^* M)$  depending on the 2-jet  $[u]_x^2 \in \mathcal{E}$ ; in local coordinates,  $\sigma_F = F_{u_{x^i x^j}} \partial_{x^i} \partial_{x^j}$ . Let us briefly indicate the proof.

A contact transformation  $\Phi : J^1M \rightarrow J^1M$  lifts naturally to a transformation  $\Phi^{(1)} : J^2M \rightarrow J^2M$  (the latter is usually defined on an open dense subset of  $J^2M$  due to mixing of dependent and independent variables). The fibers of the projection  $\pi_{2,1} : J^2M \rightarrow J^1M$  are affine spaces associated to the fibers of  $S^2 T^* M$ . The prolongation  $\Phi^{(1)}$  is fiberwise projective on them. Its symbol  $\sigma_{\Phi^{(1)}} : S^2 T^* M \rightarrow S^2 T^* M$  (here and below the bundles are pulled back to  $\mathcal{E}$  via the natural projection  $\pi_{\mathcal{E}}$ ) is the differential of this projective transformation. Using the properties of the symbol (cf. [44, 64]), we conclude

$$\sigma_{(\Phi^{(1)})^*(F)} = \sigma_{F \circ \Phi^{(1)}} = \sigma_F \circ \sigma_{\Phi^{(1)}}.$$

This is the required contact invariance of the symbol  $\sigma_F : S^2 T^* M \rightarrow \mathbb{R}$ .

A non-zero covector  $p \in T_x^* M$  is called characteristic for  $F$  at  $[u]_x^2 \in \mathcal{E}$  if  $\sigma_F([u]_x^2)(p, p) = 0$ . The projectivized (complexified) set of characteristic covectors is called the (complex) characteristic variety  $\text{Char}(\mathcal{E})$

at  $[u]_x^2$ . The equation  $\mathcal{E}$  is hyperbolic if its complex characteristic variety  $\text{Char}(\mathcal{E})$  is the complexification of a real variety. The characteristic variety is invariant under contact transformations. Indeed, let  $g_{\mathcal{E}} = \text{Ker}(\sigma_F) = T\mathcal{E} \cap \text{Ker} d\pi_{2,1} \subset S^2T^*M$  be the symbol of  $\mathcal{E}$ . Then the claim follows from the fact [64] that a covector  $p$  is characteristic whenever  $p^2 = p \cdot p \in g_{\mathcal{E}}$ .

The basic object of our study is the bi-vector  $\sigma_F = g^{\sharp}$ , and when this bi-vector is non-degenerate, we consider the dual (conformal) metric  $g \in S^2T_x^*M$ , which depends only on  $[u]_x^2 \in \mathcal{E}$  (we point out that the projectivized null cone of  $g$  is dual to the characteristic variety). Together with the 1-form  $\omega$ , the metric  $g$  uniquely defines the connection  $\mathbb{D}$  by the first equation of (7), and the triple  $(\mathbb{D}, g, \omega)$  defines an Einstein–Weyl structure if the second equation of (7) is satisfied.

We say that a certain tensor (Cotton, Einstein–Weyl, etc.), which depends on higher-order jets of  $u$ , vanishes on every solution, if it vanishes modulo the equation  $\mathcal{E}$ , meaning again that jets of  $u$  are constrained by the equation, and a finite number of its differential consequences. In practice we eliminate, say, all higher-order derivatives of  $u$  containing differentiation by  $t$  more than once (i.e.,  $u_{tt}, u_{ttt}, u_{xtt}$ , etc.), and equate to zero terms at the remaining higher-order derivatives. Since all objects depend on a finite-order jet of  $u$ , we do not rely upon smooth solvability of the equation, and carry out calculations formally using the geometric theory of PDEs.

**Proposition 1.** *For an equation  $\mathcal{E}$ , the properties for the conformal metric  $g$  to have the Cotton tensor zero on every solution, or to satisfy the Einstein–Weyl conditions on every solution, are contact invariant.*

*Proof.* Let  $F \circ \Phi^{(1)} = \tilde{F}$  be the transformed operator. Then the prolonged contact transformation maps  $\sigma_{\tilde{F}}$  to  $\sigma_F$ , and consequently the metric  $g$  is mapped to the corresponding metric  $\tilde{g}$ . Denote by  $\tilde{\omega}$  the pullback of  $\omega$  (higher prolongations of  $\Phi$  are used at this step). The contact map  $\Phi$  sends a solution  $\tilde{S}$  of  $\tilde{\mathcal{E}} = \{\tilde{F} = 0\}$ , considered as a Legendrian submanifold in  $J^1M$ , to a solution  $S$  of  $\mathcal{E}$ , and the vanishing of the Cotton tensor of  $\tilde{g}$  on  $\tilde{S}$  is equivalent to the same condition for  $g$  on  $S$ . Similarly, the Einstein–Weyl property for  $(\tilde{g}, \tilde{\omega})$  on  $\tilde{S}$  is equivalent to the same property for  $(g, \omega)$  on  $S$ .

Classical solutions (projecting diffeomorphically onto  $M$ ) may be mapped to multi-valued solutions (Legendrian submanifolds), but, locally most of them are mapped to classical solutions. More precisely, if we consider  $(k + 1)$ -jets of solutions, then the prolongation  $\Phi^{(k)}$  is defined on an open dense subset thereof. Thus, the vanishing of the Cotton tensor, and the trace-freeness for the symmetrized Ricci tensor on every solution of  $\mathcal{E}$ , implies similar properties for almost all solutions of  $\tilde{\mathcal{E}}$ . The latter we quantify to hold on an open dense set of the prolonged

equation  $\tilde{\mathcal{E}}^{(k-1)} \subset J^{k+1}M$ , and this by continuity implies the required property for all solutions. q.e.d.

Thus, we obtain a covariant approach to integrability. Notice, however, that the integrability by the method of hydrodynamic reductions, as well as the explicit form (8) of the covector  $\omega$ , are coordinate-dependent. More about this will be said in the concluding remarks.

The above discussion covers integrable PDE of types (3)–(5). For systems of first-order PDEs, such as (6), the theory is similar: the only difference is that, by virtue of the Lie–Bäcklund theorem, contact transformations should be changed to point transformations, i.e., diffeomorphisms of  $J^0(M, \mathbb{R}^2) = M \times \mathbb{R}^2$ .

### 3. Equations of type I

In this section we consider equations of the form (3),

$$(a(u))_{xx} + (b(u))_{yy} + (c(u))_{tt} + 2(p(u))_{xy} + 2(q(u))_{xt} + 2(r(u))_{yt} = 0.$$

Their integrability was investigated in [32] based on the method of hydrodynamic reductions. This boils down to the requirement of the existence of an infinity of multi-phase solutions, which imposes strong constraints on the coefficients of the equation and provides an efficient classification criterion (see Section 9 for a brief summary of the method). To formulate the classification result, we introduce the symmetric matrix

$$V(u) = \begin{pmatrix} a' & p' & q' \\ p' & b' & r' \\ q' & r' & c' \end{pmatrix},$$

where prime denotes differentiation by  $u$ . The classification is performed modulo (complex) linear changes of the independent variables  $x, y, t$ , as well as transformations  $u \rightarrow \varphi(u)$ , which constitute the equivalence group of our problem.

**Theorem 1 ([32]).** *Equation (3) is integrable by the method of hydrodynamic reductions if and only if the matrix  $V(u)$  satisfies the constraint*

$$(9) \quad V'' = (\ln \det V)'V' + kV,$$

for some scalar function  $k = k(u)$ . Modulo equivalence transformations, this leads to the five canonical forms of nonlinear integrable models:

$$u_{xx} + u_{yy} - (\ln(e^u - 1))_{yy} - (\ln(e^u - 1))_{tt} = 0,$$

$$u_{xx} + u_{yy} - (e^u)_{tt} = 0,$$

$$(e^u - u)_{xx} + 2u_{xy} + (e^u)_{tt} = 0,$$

$$u_{xt} - (uu_x)_x - u_{yy} = 0,$$

$$(u^2)_{xy} + u_{yy} + 2u_{xt} = 0.$$

*Examples 2 and 4 are the familiar Boyer–Finley (BF) and the dKP equations.*

We point out that the constraint (9), which implies  $V'' \in \text{span}\{V, V'\}$ , means that the “curve”  $V(u)$  lies in a two-dimensional linear subspace of the space of  $3 \times 3$  symmetric matrices. The classification of normal forms of such linear subspaces, which is equivalent to the classification of pencils of conics, leads to the five canonical forms of Theorem 1. It was pointed out by D. Calderbank that equations of the form (3) are related to generalized Nahm equations with the gauge group  $SDiff(\Sigma^2)$ . In this language, the five canonical forms of Theorem 1 correspond to the five types of generalized Nahm equations obtained in [9]. For the dKP equation, this correspondence was explicitly demonstrated in [18].

The linearized equation (3) is

$$a'(u)v_{xx} + b'(u)v_{yy} + c'(u)v_{tt} + 2p'(u)v_{xy} + 2q'(u)v_{xt} + 2r'(u)v_{yt} + \dots = 0,$$

where dots denote terms with lower-order derivatives of  $v$ . Its symbol defines a conformal structure  $g = g_{ij}(u)dx^i dx^j$  where  $(x^1, x^2, x^3) = (x, y, t)$ , and the matrix  $g_{ij}$  is the inverse of  $V$ . Our first result is as follows.

**Theorem 2.** *Equation (3) is linearizable by a transformation from the equivalence group if and only if the conformal structure  $g$  is conformally flat on every solution.*

*Proof.* The condition responsible for conformal flatness in three dimensions is the vanishing of the Cotton tensor,

$$(10) \quad \nabla_r(R_{pq} - \frac{1}{4}Rg_{pq}) = \nabla_q(R_{pr} - \frac{1}{4}Rg_{pr}),$$

where  $R_{pq}$  is the Ricci tensor,  $R$  is the scalar curvature, and  $\nabla$  denotes covariant differentiation in the Levi–Civita connection of  $g$ . Calculating (10) and using (3) and its differential consequences to eliminate all higher-order partial derivatives of  $u$  containing differentiation by  $t$  more than once, we obtain expressions that have to vanish identically in the remaining higher-order derivatives of  $u$  (without loss of generality, we will assume that  $c(u) = u$ ; this can be achieved by a transformation from the equivalence group). Requiring the vanishing of coefficients at the remaining derivatives of  $u$ , we obtain that all entries of the matrix  $V$  must be constant, which leads to linear equations. q.e.d.

It turns out that conformal structures corresponding to all five integrable models from Theorem 1 satisfy the Einstein–Weyl property. In fact, this follows from the construction of [9], which provides Einstein–Weyl structures from solutions of the gauge field equations with the gauge group  $SDiff(\Sigma^2)$  modeled on Riccati spaces; our goal here is to present the explicit formulae. In addition, for all equations from Theorem 1 we present dispersionless Lax pairs in the form  $[X, Y] = 0$ , where

$X$  and  $Y$  are  $\lambda$ -dependent vector fields which commute modulo the equation. Projecting integral surfaces of the distribution spanned by  $X$  and  $Y$  from the extended space of coordinates  $x, y, t, \lambda$  down to the space of independent variables  $x, y, t$ , we obtain two-parameter families of null totally geodesic surfaces of the corresponding Einstein–Weyl structures.

Here and in what follows we choose the “simplest” representative metric  $g$  within the conformal class  $[g]$ . In all cases it is given by either the inverse or the cofactor matrix of  $g^\sharp$ .

**Equation 1:**  $u_{xx} + u_{yy} - (\ln(e^u - 1))_{yy} - (\ln(e^u - 1))_{tt} = 0$ .

Conformal structure:  $g = dx^2 + (1 - e^u)dy^2 + (e^{-u} - 1)dt^2$ .

Covector:  $\omega = \frac{e^u + 1}{e^u - 1}u_x dx - u_y dy + u_t dt$ .

Lax pair:

$$X = \partial_y + \sqrt{e^u - 1} \sin \varphi \partial_x + \left( \frac{e^u}{e^u - 1} u_t - \frac{\cos \varphi}{\sqrt{1 - e^{-u}}} u_x \right) \partial_\lambda,$$

$$Y = \partial_t + \sqrt{1 - e^{-u}} \cos \varphi \partial_x + \left( \frac{1}{1 - e^u} u_y + \frac{\sin \varphi}{\sqrt{e^u - 1}} u_x \right) \partial_\lambda,$$

here  $\varphi = -\arctan(e^{-u/2} \tan \lambda/2)$ .

**Equation 2:**  $u_{xx} + u_{yy} - (e^u)_{tt} = 0$  (BF equation).

Conformal structure:  $g = dx^2 + dy^2 - e^{-u} dt^2$ .

Covector:  $\omega = -u_x dx - u_y dy + u_t dt$ .

This Einstein–Weyl structure was obtained in [68]; see also [47].

Lax pair:

$$X = \partial_y - e^{u/2} \sin \lambda \partial_t - \frac{1}{2}(u_x + e^{u/2} u_t \cos \lambda) \partial_\lambda,$$

$$Y = \partial_x - e^{u/2} \cos \lambda \partial_t + \frac{1}{2}(u_y + e^{u/2} u_t \sin \lambda) \partial_\lambda.$$

**Equation 3:**  $(e^u - u)_{xx} + 2u_{xy} + (e^u)_{tt} = 0$ .

Conformal structure:  $g = 2dxdy + (1 - e^u)dy^2 + e^{-u} dt^2$ .

Covector:  $\omega = -u_x dx + (2e^u u_x - u_y) dy + u_t dt$ .

Lax pair:

$$X = \partial_t - \lambda \partial_x + (\lambda^2 + 1)u_x \partial_\lambda,$$

$$Y = \partial_y + \frac{1}{2}(e^u(\lambda^2 + 1) - 1)\partial_x - \frac{1}{2}e^u(u_t + \lambda u_x)(\lambda^2 + 1)\partial_\lambda.$$

**Equation 4:**  $u_{xt} - (uu_x)_x - u_{yy} = 0$  (dKP equation).

Conformal structure:  $g = 4dxdt - dy^2 + 4udt^2$ .

Covector:  $\omega = -4u_x dt$ .

This Einstein–Weyl structure was obtained in [20].

Lax pair:

$$X = \partial_y - \lambda \partial_x + u_x \partial_\lambda, \quad Y = \partial_t - (\lambda^2 + u)\partial_x + (u_x \lambda + u_y)\partial_\lambda.$$

**Equation 5:**  $(u^2)_{xy} + u_{yy} + 2u_{xt} = 0$ .

Conformal structure:  $g = 2dxdt + dy^2 - 2udydt + u^2 dt^2$ .

Covector:  $\omega = 2u_x dy + 2(u_y - uu_x) dt$ .

Lax pair:

$$X = \partial_y - \lambda \partial_x + 2u_x \lambda \partial_\lambda, \quad Y = \partial_t + \left(\frac{1}{2} \lambda^2 + u \lambda\right) \partial_x - (u_x \lambda + u_y + 2uu_x) \lambda \partial_\lambda.$$

All of the above conformal structures  $g$  and covectors  $\omega$  can be represented in terms of the matrix  $V = V(u)$  as follows (the symbol  $\mathfrak{t}$  denotes the transposition):

$$(11) \quad \begin{aligned} g &= (dx \ dy \ dt) V^{-1} (dx \ dy \ dt)^{\mathfrak{t}}, \\ \omega &= 2(dx \ dy \ dt) V^{-1} V'(u_x, u_y, u_t)^{\mathfrak{t}} - d(\ln \det V). \end{aligned}$$

They satisfy the Einstein–Weyl equations (7) if and only if  $V(u)$  satisfies the integrability condition (9). Setting  $(x, y, t) = (x^1, x^2, x^3)$ , one can represent the components of  $\omega = \omega_k dx^k$  by the formula (8). It turns out that exactly the same formula holds for all other classes of dispersionless PDEs discussed in this paper. The covector  $\omega$  is related to the symbol of formal linearization via the identity

$$g^{ij} v_{x^i x^j} = \nabla^i \nabla_i v - \frac{1}{2} \omega^i \nabla_i v,$$

where  $\nabla^i = g^{ik} \nabla_k$ ,  $\omega^i = g^{ik} \omega_k$ , and  $\nabla$  denotes covariant differentiation in the Levi–Civita connection of the metric  $g_{ij}$ . Note that the right-hand side of this identity can be interpreted as a special case of the Weyl wave operator of weight 0 [22]. The second main result of this section is as follows.

**Theorem 3.** *Equation (3) is integrable by the method of hydrodynamic reductions if and only if the corresponding conformal structure  $g$  is Einstein–Weyl on every solution, with the covector  $\omega$  given by (8).*

*Proof.* Given an equation of the form (3), the conformal structure of its formal linearization is  $g = (dx \ dy \ dt) V^{-1} (dx \ dy \ dt)^{\mathfrak{t}}$ ; see (11). We will seek a covector  $\omega$  in the form  $\omega = (dx \ dy \ dt) T(u_x \ u_y \ u_t)^{\mathfrak{t}}$  where  $T(u)$  is an unknown  $3 \times 3$  matrix depending on  $u$ . Imposing the Einstein–Weyl equations (7) and using (3) to eliminate the second-order derivative  $u_{tt}$  (no higher-order derivatives of  $u$  will occur in this calculation), we obtain a set of relations that have to vanish identically in the remaining partial derivatives of  $u$ . Thus, equating to zero terms at the second-order derivatives of  $u$ , we obtain  $T$  in terms of  $V$ ,

$$T = 2V^{-1} V' - (\ln \det V)' E,$$

where  $E$  is the  $3 \times 3$  identity matrix. This is equivalent to the formulae (11) and (8) for  $\omega$ . The remaining terms vanish identically if and only if  $V$  satisfies the constraint (9). q.e.d.

REMARK 1. It was assumed in the proof of Theorem 3 that  $\omega$  depends linearly on the first-order derivatives of  $u$ . One can show that

this assumption is unnecessary: even a rather general requirement that  $\omega$  depends on some finite-order jets of  $u$  is already sufficiently restrictive and leads to the same formula (8) for  $\omega$ .

REMARK 2. Equations from Theorem 1 possess implicit solutions  $u(x, y, t)$  of the form

$$(12) \quad (x, y, t)M(u)(x, y, t)^t = 1,$$

where  $M(u)$  is a  $3 \times 3$  symmetric matrix of  $u$ . The level surfaces of such solutions,  $u = \text{const}$ , are central quadrics in the space of independent variables  $x, y, t$ . This construction is known as the central quadric ansatz [65, 18]. The equation for  $M(u)$  is

$$M' = sMVM/\sqrt{\det M},$$

$s = \text{const}$ . It was demonstrated in [65, 18] that in the cases of BF and dKP, this equation reduces to Painlevé transcendents P3-P1. It was shown in [32] that other integrable models from Theorem 1 lead to the remaining Painlevé equations P6-P4, with the full P6 corresponding to the first equation. Thus, we obtain a whole variety of Einstein–Weyl structures parametrized by Painlevé transcendents.

#### 4. Type II: quasilinear wave equations

In this section we discuss geometric aspects of quasilinear wave equations (4),

$$f_{11}u_{xx} + f_{22}u_{yy} + f_{33}u_{tt} + 2f_{12}u_{xy} + 2f_{13}u_{xt} + 2f_{23}u_{yt} = 0;$$

here the coefficients  $f_{ij} = f_{ij}(u_x, u_y, u_t)$  are functions of the first-order derivatives only. PDEs of this type were investigated in [8] based on their correspondence with conformal structures in projective space. It was pointed out that the moduli space of integrable equations is 20-dimensional. It was shown in [54] that coefficients of the generic integrable equations of the form (4) can be parametrized by generalized hypergeometric functions. We recall that the class of quasilinear wave equations is invariant under the group  $SL(4)$  of linear transformations of the variables  $x^i, u$ , where  $(x^1, x^2, x^3) = (x, y, t)$ . These transformations constitute the natural equivalence group of the problem.

The linearized equation has the form

$$f_{ij}v_{x^i x^j} + \dots = 0,$$

and it defines the conformal structure  $g = g_{ij}dx^i dx^j$  where the matrix of  $g_{ij}$  is the inverse of  $f_{ij}$ . Here dots denote terms with lower-order derivatives of  $v$ . Our first result is as follows.

**Theorem 4.** *Equation (4) is linearizable by a transformation from the equivalence group  $SL(4)$  if and only if the conformal structure  $g$  is conformally flat on every solution.*

*Proof.* Let us first recall, following [8], the linearizability conditions for equations of the form (4). Since the structure of these conditions is the same in any dimension, we will consider the general  $n$ -dimensional case,

$$f_{ij}u_{x^i x^j} = 0,$$

where  $f_{ij}$  are functions of the first-order derivatives  $u_{x^k}$  only,  $i, j, k = 1, \dots, n$ . Setting  $p_k = u_{x^k}$ , we will write down a system of differential constraints for  $f_{ij}(\mathbf{p})$  which are necessary and sufficient for the linearizability of the equation under study by a transformation from the equivalence group  $SL(n+1)$ . Let us introduce the object

$$a_{ijk} = \partial_{p_k} f_{ij} - (c_k + 2s_k)f_{ij} - s_i f_{kj} - s_j f_{ki},$$

where

$$s_k = \frac{f^{ij}}{(n+2)(1-n)} (\partial_{p_k} f_{ij} - n\partial_{p_j} f_{ik}),$$

$$c_k = \frac{f^{ij}}{(n+2)(n-1)} ((n+3)\partial_{p_k} f_{ij} - 2(n+1)\partial_{p_j} f_{ik}).$$

Then the linearizability is equivalent to the following two conditions:

- $a_{ijk} = 0$ ,
- $\partial_{p_j} s_i - s_i s_j = 0$ .

Geometrically, these conditions are equivalent to the existence of a flat connection  $\nabla$  (in  $\mathbf{p}$ -coordinates) with Christoffel symbols  $\Gamma_{jk}^i = s_j \delta_k^i + s_k \delta_j^i$  such that  $\nabla_k f_{ij} = c_k f_{ij}$ ; see [8] for more details.

Let us now require that the conformal structure  $g = g_{ij} dx^i dx^j$  is conformally flat on every solution. Calculating the Cotton tensor (10) (now we set  $n = 3$ ), and using (4) and its differential consequences to eliminate all higher-order partial derivatives of  $u$  that contain differentiation by  $t$  more than once, we obtain a complicated expression, which has to vanish identically in the remaining higher-order derivatives of  $u$ . In particular, requiring that coefficients at the remaining fourth-order derivatives vanish identically (no higher-order derivatives of  $u$  will occur in this calculation), we obtain the first linearizability condition  $a_{ijk} = 0$ .

There are two ways to proceed: Collecting terms at the lower-order derivatives of  $u$  one can obtain the second set of linearizability conditions (this, however, leads to quite complicated calculations). Another way is to point out that the condition  $a_{ijk} = 0$  alone is already sufficiently restrictive [8, 61, 1] and implies that the PDE in question is either linearizable or reducible to the equation for minimal hypersurfaces in a (pseudo) Euclidean space,

$$[(\nabla u)^2 - 1]\Delta u - (\nabla u)H(\nabla u)^t = 0,$$

where  $\nabla u = (u_{x^1}, \dots, u_{x^n})$  is the gradient of  $u$ ,  $\Delta$  is the Laplacian, and  $H$  is the Hessian matrix of  $u$ . In the three-dimensional case we arrive

at the equation

$$(u_y^2 + u_t^2 - 1)u_{xx} + (u_x^2 + u_t^2 - 1)u_{yy} + (u_x^2 + u_y^2 - 1)u_{tt} \\ - 2(u_x u_y u_{xy} + u_x u_t u_{xt} + u_y u_t u_{yt}) = 0.$$

To complete the proof, it remains to point out that the corresponding conformal structure  $g$  is not conformally flat on generic solutions; furthermore, the equation itself is not linearizable (in fact, not even integrable for  $n \geq 3$ ). q.e.d.

**REMARK 3.** This proof, and the proof of Theorem 2, generalize to any dimension  $n > 3$ , with the only change that one needs the Weyl tensor of conformal curvature instead of the Cotton tensor. For non-linearizable differential equations, the requirement of conformal flatness singles out a subclass of exact solutions of a given PDE; see [49].

Our next goal is to prove that conformal structures corresponding to formal linearizations of *integrable* equations of the form (4) give rise to the Einstein–Weyl geometry. Let us begin with examples of known integrable PDEs.

**Example 1.** The equation  $u_x u_{yt} + u_y u_{xt} + u_t u_{xy} = 0$  constitutes the Euler–Lagrange equation for the Lagrangian density  $u_x u_y u_t$  that was obtained in [30] in the classification of first-order integrable Lagrangians. Conformal structure:

$$g = (u_x dx + u_y dy + u_t dt)^2 - 2u_x^2 dx^2 - 2u_y^2 dy^2 - 2u_t^2 dt^2.$$

$$\text{Covector: } \omega = -4 \frac{u_x u_{yt}}{u_y u_t} dx - 4 \frac{u_y u_{tx}}{u_t u_x} dy - 4 \frac{u_t u_{xy}}{u_x u_y} dt.$$

**Example 2.** The equation

$$(u_y p(u_t))_x + (u_x p(u_t))_y + (u_x u_y p'(u_t))_t = 0$$

constitutes the Euler–Lagrange equation for the Lagrangian density  $u_x u_y p(u_t)$ , which can be viewed as a deformation of Example 1. In this case, the integrability conditions reduce to a single fourth-order ODE for  $p$ ,

$$p''''(p^2 p'' - 2pp'^2) - p^2 p'''' + 2pp'p''p''' + 8p'^3 p''' - 9p'^2 p''^2 = 0.$$

It was shown in [31] that the general solution to this ODE is a modular form of weight 1 and level 3 known as the Eisenstein series  $E_{1,3}$ .

Conformal structure:

$$g = (p' u_x dx + p' u_y dy + p dt)^2 - 2p'^2 u_x^2 dx^2 \\ - 2p'^2 u_y^2 dy^2 - 2p^2 dt^2 - 2pp'' u_x u_y dx dy.$$

Covector:

$$\begin{aligned} \omega = 2 & \left( \frac{pp''^2 - pp'p''' + p'^2p''}{p(pp'' - 2p'^2)} u_x u_{tt} - 2 \frac{p'u_x}{p u_y} u_{ty} \right) dx \\ & + 2 \left( \frac{pp''^2 - pp'p''' + p'^2p''}{p(pp'' - 2p'^2)} u_y u_{tt} - 2 \frac{p'u_y}{p u_x} u_{tx} \right) dy \\ & + 2 \left( \frac{pp''' - 3p'p''}{pp'' - 2p'^2} u_{tt} + 2 \left( \frac{u_{tx}}{u_x} + \frac{u_{ty}}{u_y} \right) \right) dt. \end{aligned}$$

This structure is Einstein–Weyl if and only if  $p$  satisfies the above fourth-order ODE.

The second main result of this section is as follows:

**Theorem 5.** *Equation (4) is integrable by the method of hydrodynamic reductions if and only if the corresponding conformal structure  $g$  is Einstein–Weyl on every solution, with the covector  $\omega$  given by (8).*

*Proof.* We will give two proofs of this result. The first one is computational, based on the explicit calculation of the Einstein–Weyl constraints and the integrability conditions as derived in [8]. The second proof utilises the fact that any integrable PDE of the form (4) possesses a dispersionless Lax pair [8]. We demonstrate that the existence of a Lax pair implies that the Weyl connection  $\mathbb{D}$ , specified by the conformal structure  $g = g_{ij} dx^i dx^j$  and the covector (8), possesses a two-parameter family of null totally geodesic surfaces, the property known to be characteristic of the Einstein–Weyl geometry [11].

The first proof can be summarized as follows. Given an equation of the form (4), the conformal structure of its formal linearization is  $g = g_{ij} dx^i dx^j$ , where  $g_{ij}$  is the inverse of  $f_{ij}$ . We will seek a covector  $\omega = \omega_k dx^k$  in the form  $\omega_k = T_k^{ij} u_{x^i x^j}$ , where  $T_k^{ij}$  are certain functions of the first-order derivatives of  $u$ . Imposing the Einstein–Weyl equations (7) and using (4) and its differential consequences to eliminate all higher-order derivatives of  $u$  that contain differentiation by  $t$  more than once (maximum third-order derivatives of  $u$  will occur in this calculation), we obtain a set of relations that have to vanish identically in the remaining partial derivatives of  $u$ . Thus, equating to zero terms at the third-order derivatives of  $u$ , we obtain the expression (8) for  $\omega$ . The remaining terms vanish identically if and only if the coefficients  $f_{ij}$  satisfy the set of integrability conditions as derived in [8]. This finishes the first proof of Theorem 5.

Let us now give a somewhat more conceptual (as well as less computational) demonstration that the integrability is equivalent to the Einstein–Weyl property. It is based on the fact that any integrable equation of the form (4) possesses a dispersionless Lax pair of the form

$$(13) \quad S_t = f(S_x, u_x, u_y, u_t), \quad S_y = g(S_x, u_x, u_y, u_t).$$

This means that the consistency condition,  $S_{ty} = S_{yt}$ , is equivalent to the equation (4). Lax pairs of this form are known to arise as dispersionless limits of solitonic Lax pairs in 2+1 dimensions [70]. Let us first outline the general construction that leads from the Lax pair (13) to totally geodesic null surfaces of the Weyl connection  $\mathbb{D}$ . Differentiating (13) by  $x$  and setting  $S_x = \lambda$ ,  $u_x = a$ ,  $u_y = b$ ,  $u_t = c$ , we obtain

$$(14) \quad \lambda_t = f_\lambda \lambda_x + f_a a_x + f_b b_x + f_c c_x, \quad \lambda_y = g_\lambda \lambda_x + g_a a_x + g_b b_x + g_c c_x.$$

With this system we associate the vector fields

$$\begin{aligned} X &= \frac{\partial}{\partial t} - f_\lambda \frac{\partial}{\partial x} + (f_a a_x + f_b b_x + f_c c_x) \frac{\partial}{\partial \lambda}, \\ Y &= \frac{\partial}{\partial y} - g_\lambda \frac{\partial}{\partial x} + (g_a a_x + g_b b_x + g_c c_x) \frac{\partial}{\partial \lambda}, \end{aligned}$$

which live in the extended four-dimensional space with coordinates  $x, y, t, \lambda$ . Note that the compatibility condition  $\lambda_{ty} = \lambda_{yt}$  is equivalent to the commutativity of these vector fields:  $[X, Y] = 0$ . The geometry behind this construction is as follows. Let us consider the cotangent bundle  $Z$  of the solution  $u(x, y, t)$ , with local coordinates  $(x, y, t, S_x, S_y, S_t)$ . Equations (13) specify a four-dimensional submanifold  $M^4 \subset Z$  parametrized by  $x, y, t$  and  $\lambda$ . The compatibility of the equations (13) indicates that this submanifold is coisotropic. The vector fields  $X, Y$  generate the kernel of the restriction to  $M^4$  of the symplectic form  $dS_x \wedge dx + dS_y \wedge dy + dS_t \wedge dt$ . Equations (14) mean that the vectors  $X, Y$  are tangential to the hypersurface of  $M^4$  defined by the equation  $\lambda = \lambda(x, y, t)$ .

Projecting the two-parameter family of integral surfaces of the distribution spanned by  $X, Y$  to the space of independent variables  $x, y, t$ , we obtain a two-parameter family of null totally geodesic surfaces of the Weyl connection  $\mathbb{D}$ . To see this we first project  $X$  and  $Y$ . This gives two vector fields

$$\hat{X} = \frac{\partial}{\partial t} - f_\lambda \frac{\partial}{\partial x}, \quad \hat{Y} = \frac{\partial}{\partial y} - g_\lambda \frac{\partial}{\partial x},$$

which commute if and only if  $\lambda$  satisfies the equations (14). It remains to show that  $\hat{X}$  and  $\hat{Y}$  form a null distribution (i.e., tangential to the null cones of  $g$ ), and that the covariant derivatives  $\mathbb{D}_{\hat{X}} \hat{X}$ ,  $\mathbb{D}_{\hat{X}} \hat{Y}$ ,  $\mathbb{D}_{\hat{Y}} \hat{X}$ ,  $\mathbb{D}_{\hat{Y}} \hat{Y}$  belong to the span of  $\hat{X}, \hat{Y}$ . Equivalently, one can introduce the covector  $\theta = dx + g_\lambda dy + f_\lambda dt$  that annihilates  $\hat{X}, \hat{Y}$ , and verify that  $\theta$  is null, and that  $\mathbb{D}_{\hat{X}} \theta \wedge \theta = \mathbb{D}_{\hat{Y}} \theta \wedge \theta = 0$ . This follows from the equations satisfied by the functions  $f(\lambda, a, b, c)$  and  $g(\lambda, a, b, c)$  as derived in [8]:

$$(15) \quad \begin{aligned} f_a &= 2kf_{12} + g_\lambda k f_{22} + f_\lambda (k f_{23} - p), & g_b &= -k f_{23} + p, \\ g_a &= -2kf_{13} - f_\lambda k f_{33} - g_\lambda (k f_{23} + p), & f_c &= k f_{23} + p, \\ f_b &= k f_{22}, & g_c &= -k f_{33}, \end{aligned}$$

where  $p(\lambda, a, b, c)$  and  $k(\lambda, a, b, c)$  are yet another two auxiliary functions. Furthermore,  $f_\lambda$  and  $g_\lambda$  satisfy the relation

$$(16) \quad f_{11} + f_{22}g_\lambda^2 + f_{33}f_\lambda^2 + 2f_{12}g_\lambda + 2f_{13}f_\lambda + 2f_{23}f_\lambda g_\lambda = 0,$$

which means that the covector  $\theta$  is null. To close the system (15)–(16), one proceeds as follows. Calculating the consistency conditions for equations (15),  $f_{ab} = f_{ba}$ ,  $g_{ab} = g_{ba}$ , etc., six conditions altogether, and differentiating the relation (16) by  $a, b, c$ , and  $\lambda$ , one obtains ten relations that can be solved for  $f_{\lambda\lambda}, g_{\lambda\lambda}$  and the first-order derivatives of  $k$  and  $p$ . Modulo these relations, the conditions  $\mathbb{D}_{\hat{X}}\theta \wedge \theta = \mathbb{D}_{\hat{Y}}\theta \wedge \theta = 0$  are satisfied identically. This finishes the second proof of Theorem 5. q.e.d.

### 5. Type III: dispersionless Hirota equations

In this section we discuss geometric aspects of PDEs (5) of the dispersionless Hirota type,

$$F(u_{xx}, u_{xy}, u_{yy}, u_{xt}, u_{yt}, u_{tt}) = 0,$$

which were investigated in [56, 29], revealing a remarkable correspondence with hypersurfaces of the Lagrangian Grassmanian. Geometric aspects of  $GL(2)$  structures associated with such equations were studied in [63]. It was shown in [54] that the generic integrable equation of the form (5) can be parametrized by generalized hypergeometric functions. Recall that equations of the form (5) are invariant under the group  $Sp(6)$  of linear symplectic transformations of the variables  $x^i, u_{x^i}$ , where  $(x^1, x^2, x^3) = (x, y, t)$ . These transformations constitute the natural equivalence group of the problem.

The linearized equation has the form

$$F_{ij}v_{x^i x^j} = 0$$

and defines a conformal structure  $g = g_{ij}dx^i dx^j$  where the matrix of  $g_{ij}$  is the inverse of  $F_{ij} = \partial F / \partial u_{x^i x^j}$ . Our first result is as follows.

**Theorem 6.** *Equation (5) is linearizable by a transformation from the equivalence group  $Sp(6)$  if and only if the conformal structure  $g$  is conformally flat on every solution.*

*Proof.* Solving for  $u_{tt}$  (we can always bring our equation into the form with a non-trivial dependence on  $u_{tt}$  due to the presence of a large equivalence group), we can rewrite (5) in the form

$$(17) \quad u_{tt} = f(u_{xx}, u_{xy}, u_{yy}, u_{xt}, u_{yt}).$$

The corresponding linearized equation is

$$v_{tt} = f_{u_{xx}}v_{xx} + f_{u_{xy}}v_{xy} + f_{u_{yy}}v_{yy} + f_{u_{xt}}v_{xt} + f_{u_{yt}}v_{yt},$$

with the associated conformal structure  $g^\sharp$  (with upper indices) defined by the matrix

$$(18) \quad P = \begin{pmatrix} f_{u_{xx}} & \frac{1}{2}f_{u_{xy}} & \frac{1}{2}f_{u_{xt}} \\ \frac{1}{2}f_{u_{xy}} & f_{u_{yy}} & \frac{1}{2}f_{u_{yt}} \\ \frac{1}{2}f_{u_{xt}} & \frac{1}{2}f_{u_{yt}} & -1 \end{pmatrix}.$$

We require this structure to be conformally flat for every background solution  $u(x, y, t)$ . Calculating the Cotton tensor (10) and using (17) and its differential consequences to eliminate all higher-order derivatives of  $u$  containing differentiation by  $t$  more than once, we obtain a complicated expression that has to vanish identically in the remaining higher-order derivatives of  $u$  (maximum fifth-order derivatives will appear in this calculation). In particular, requiring that coefficients at the remaining fifth-order derivatives of  $u$  vanish identically, we obtain nine second-order differential constraints for  $f$ :

$$(19) \quad \begin{aligned} f_{u_{xx}}f_{u_{xt}u_{xt}} + f_{u_{xx}u_{xx}} &= 0, & f_{u_{yy}}f_{u_{yt}u_{yt}} + f_{u_{yy}u_{yy}} &= 0, \\ f_{u_{xt}}f_{u_{xt}u_{xt}} + 2f_{u_{xt}u_{xx}} &= 0, & f_{u_{yt}}f_{u_{yt}u_{yt}} + 2f_{u_{yt}u_{yy}} &= 0, \\ f_{u_{yt}}f_{u_{xt}u_{xt}} + 2(f_{u_{xt}}f_{u_{xt}u_{yt}} + f_{u_{xt}u_{xy}} + f_{u_{yt}u_{xx}}) &= 0, \\ f_{u_{xt}}f_{u_{yt}u_{yt}} + 2(f_{u_{yt}}f_{u_{xt}u_{yt}} + f_{u_{yt}u_{xy}} + f_{u_{xt}u_{yy}}) &= 0, \\ f_{u_{xy}}f_{u_{xt}u_{xt}} + 2f_{u_{xx}}f_{u_{xt}u_{yt}} + 2f_{u_{xx}u_{xy}} &= 0, \\ f_{u_{xy}}f_{u_{yt}u_{yt}} + 2f_{u_{yy}}f_{u_{xt}u_{yt}} + 2f_{u_{yy}u_{xy}} &= 0, \\ f_{u_{yy}}f_{u_{xt}u_{xt}} + f_{u_{xx}}f_{u_{yt}u_{yt}} + 2f_{u_{xy}}f_{u_{xt}u_{yt}} + 2f_{u_{xx}u_{yy}} + f_{u_{xy}u_{xy}} &= 0. \end{aligned}$$

It was shown in [60, 6, 13] that these relations characterize symplectic Monge–Ampère equations; i.e., PDEs (5) such that the left-hand side  $F$  can be represented as a linear combination of all possible minors of the Hessian matrix of  $u$ ,

$$U = \begin{pmatrix} u_{xx} & u_{xy} & u_{xt} \\ u_{xy} & u_{yy} & u_{yt} \\ u_{xt} & u_{yt} & u_{tt} \end{pmatrix}.$$

Symplectic Monge–Ampère equations and differential constraints (19) have a clear geometric interpretation. Let us consider the Lagrangian Grassmannian  $\Lambda^6$  that can be (locally) parametrized by  $3 \times 3$  symmetric matrices  $U$ . Minors of  $U$  define the Plücker embedding of  $\Lambda^6$  into projective space  $P^{13}$ . We will identify  $\Lambda^6$  with the image of this projective embedding. Symplectic Monge–Ampère equations can be viewed as hyperplane sections  $M^5$  of  $\Lambda^6 \subset P^{13}$ . We point out that differential

constraints (19) can be represented in compact form as

$$(20) \quad \begin{aligned} d^2 f &= 2a_0(df du_{xy} - du_{xt} du_{yt}) + 2a_1(df du_{yy} - (du_{yt})^2) \\ &+ 2a_2(df du_{xx} - (du_{xt})^2) + 2b_0(du_{xx} du_{yy} - (du_{xy})^2) \\ &+ 2b_1(du_{xt} du_{xy} - du_{yt} du_{xx}) + 2b_2(du_{yt} du_{xy} - du_{xt} du_{yy}). \end{aligned}$$

Indeed, they follow from (20) by elimination of the coefficients  $a_i, b_i$ . Here  $d^2 f$  is the symmetric differential of  $f$ . Notice that  $d^2 f$  and the six quadratic expressions on the right-hand side of (20) are nothing but second fundamental forms of the submanifold  $M^5 \subset \Lambda^6 \subset P^{13}$  defined by (17). Furthermore, the six fundamental forms on the right-hand side of (20) are the restrictions to  $M^5$  of the second fundamental forms of  $\Lambda^6 \subset P^{13}$ . Thus, (20) says that  $M^5$  has no nontrivial second fundamental forms “of its own”; i.e., all its second fundamental forms can be obtained as restriction of the second fundamental forms of  $\Lambda^6$ . This is an obvious necessary condition for a submanifold  $M^5 \subset \Lambda^6 \subset P^{13}$  to be a hyperplane section. In the present case, it is also sufficient. Calculating consistency conditions for (20), we obtain the system

$$(21) \quad \begin{aligned} da_0 &= a_0 \varphi - 2s du_{12}, & db_0 &= b_0 \varphi + s df, \\ da_1 &= a_1 \varphi + s du_{11}, & db_1 &= b_1 \varphi + 2s du_{02}, \\ da_2 &= a_2 \varphi + s du_{22}, & db_2 &= b_2 \varphi + 2s du_{01}, & ds &= s \varphi. \end{aligned}$$

Here  $\varphi = a_0 du_{xy} + a_1 du_{yy} + a_2 du_{xx}$ , and  $s$  is yet another auxiliary function. One can verify that  $d\varphi = 0$ . Equations (20) and (21) constitute an involutive differential system for  $f$  that characterizes symplectic Monge–Ampère equations.

Once we know that our PDE is of symplectic Monge–Ampère type, there are two ways to proceed. The first one is to use the fact that linearizable Monge–Ampère equations correspond to special hyperplane sections of  $\Lambda^6$  such that the corresponding hyperplane is tangential to  $\Lambda^6$ , i.e., belongs to the dual variety [29, 16]. Written in differential form, this simple geometric property gives just one extra condition that can be used to express  $s$  (via a cumbersome formula reflecting the fact the the dual variety of  $\Lambda^6$  is a complicated quartic hypersurface):

$$\begin{aligned} s &= \frac{-1}{4 \det P} \left( a_0^2 f_{u_{xx}} f_{u_{yy}} + a_1^2 f_{u_{xx}}^2 + a_2^2 f_{u_{yy}}^2 \right. \\ &- a_0 a_1 f_{u_{xx}} f_{u_{xy}} - a_0 a_2 f_{u_{xy}} f_{u_{yy}} + a_1 a_2 (f_{u_{xy}}^2 - 2 f_{u_{xx}} f_{u_{yy}}) \\ &- a_0 b_0 (f_{u_{xt}} f_{u_{yt}} + f_{u_{xy}}) - a_0 b_1 f_{u_{xt}} f_{u_{yy}} - a_0 b_2 f_{u_{yt}} f_{u_{xx}} \\ &+ a_1 b_0 (f_{u_{xt}}^2 + 2 f_{u_{xx}}) + a_1 b_1 (f_{u_{xt}} f_{u_{xy}} - f_{u_{yt}} f_{u_{xx}}) + a_1 b_2 f_{u_{xt}} f_{u_{xx}} \\ &+ a_2 b_0 (f_{u_{yt}}^2 + 2 f_{u_{yy}}) + a_2 b_1 f_{u_{yt}} f_{u_{yy}} + a_2 b_2 (f_{u_{yt}} f_{u_{xy}} - f_{u_{xt}} f_{u_{yy}}) \\ &\left. + b_0^2 - b_1^2 f_{u_{yy}} - b_2^2 f_{u_{xx}} - b_0 b_1 f_{u_{yt}} - b_0 b_2 f_{u_{xt}} - b_1 b_2 f_{u_{xy}} \right), \end{aligned}$$

where the matrix  $P$  is given by (18).

A direct calculation shows that the remaining coefficients of the Cotton tensor vanish if and only if the above linearizability condition holds. This, however, is a rather complicated calculation.

Another way is to use the fact that, in three dimensions, any non-degenerate symplectic Monge–Ampère equation is either linearizable or  $Sp(6)$ -equivalent to one of the three canonical forms [48, 3]

$$(22) \quad \text{Hess } u = 1, \quad \text{Hess } u = u_{xx} + u_{yy} + u_{tt}, \quad \text{Hess } u = u_{xx} + u_{yy} - u_{tt}.$$

The first of these equations governs improper affine hyperspheres, while the last two describe special Lagrangian 3-folds. One can verify by a direct calculation that formal linearizations of the equations (22) are not conformally flat for generic solutions. Thus, once again conformal flatness proves to be equivalent to the linearizability. q.e.d.

As in the previous cases, formal linearizations of *integrable* equations of the form (5) give rise to conformal structures satisfying the Einstein–Weyl property.

**Example 1.** The equation  $e^{utt} = e^{u_{xx}} + e^{u_{yy}}$  appeared in [29] in the classification of integrable PDEs of the form  $F(u_{xx}, u_{yy}, u_{tt}) = 0$ .  
Conformal structure:  $g = e^{-u_{xx}} dx^2 + e^{-u_{yy}} dy^2 - e^{-u_{tt}} dt^2$ .

Covector:  $\omega = 2(u_{ttt}dt + u_{xxx}dx + u_{yyy}dy) - d(u_{tt} + u_{xx} + u_{yy})$ .

**Example 2.** The equation  $u_{tt} = \frac{u_{xy}}{u_{xt}} + \frac{1}{6}\eta(u_{xx})u_{xt}^2$  appeared in [56] in the classification of integrable hydrodynamic chains. Here the integrability conditions reduce to the Chazy equation for  $\eta$ ,  $\eta''' + 2\eta\eta'' = 3(\eta')^2$ .

Conformal structure:

$$g = 4u_{xt}dx dy - \left(\frac{2}{3}\eta' u_{xt}^4 + s^2\right) dy^2 + 2sdy dt - dt^2, \quad s = \frac{1}{3}\eta u_{xt}^2 - \frac{u_{xy}}{u_{xt}}.$$

Covector:

$$\begin{aligned} \omega = & \left[ \left( \frac{1}{9} u_{tx}^3 \eta \eta' + \frac{2}{3} u_{tx}^3 \eta'' - \frac{1}{3} u_{xy} \eta' \right) u_{xxx} + (u_{xy} u_{tx}^{-3} - \frac{1}{3} \eta) u_{xxy} \right. \\ & + \left( \frac{2}{9} u_{tx}^2 \eta^2 + \frac{8}{3} u_{tx}^2 \eta' - u_{xy}^2 u_{tx}^{-4} - \frac{1}{3} u_{xy} u_{tx}^{-1} \eta \right) u_{txx} \\ & \left. - 2 u_{tx}^{-1} u_{txy} + \left( \frac{2}{3} u_{tx} \eta + 4 u_{xy} u_{tx}^{-2} \right) u_{ttx} \right] dy \\ & - \left[ \frac{1}{3} \eta' u_{tx} u_{xxx} - u_{t,x}^{-2} u_{xxy} + (u_{xy} u_{tx}^{-3} + \frac{2}{3} \eta) u_{txx} - 2 u_{tx}^{-1} u_{ttx} \right] dt. \end{aligned}$$

This structure is Einstein–Weyl if and only if  $\eta$  solves the Chazy equation.

The second main result of this section is as follows:

**Theorem 7.** *Equation (5) is integrable by the method of hydrodynamic reductions if and only if the corresponding conformal structure  $g$  is Einstein–Weyl on every solution, with the covector  $\omega$  given by (8).*

*Proof.* Given an equation of the form (5), the conformal structure of its formal linearization is  $g = g_{ij}dx^i dx^j$ , where  $g_{ij}$  is the inverse of  $F_{ij}$ . We will seek a covector  $\omega = \omega_k dx^k$  in the form  $\omega_k = T_k^{ijl} u_{x^i x^j x^l}$ , where  $T_k^{ijl}$  are certain functions of the second-order derivatives of  $u$ . Imposing the Einstein–Weyl equations (7) and using (5) and its differential consequences to eliminate all higher-order derivatives of  $u$  that contain differentiation by  $t$  more than once (we use the representation (17); maximum fourth-order derivatives of  $u$  will occur in this calculation), we obtain a set of relations which have to vanish identically in the remaining partial derivatives of  $u$ . Thus, equating to zero coefficients at the fourth-order derivatives of  $u$ , we obtain the expression (8) for  $\omega$ .

With this expression,  $R - \Lambda g$  is purely quadratic in the third-order derivatives  $u_{x^i x^j x^l}$ . Choosing  $\Lambda$  in such a way that  $dt^2$  term disappears, we get  $5 \cdot 28 = 140$  coefficients at these quadratic terms that are third-order differential polynomials in  $f$ . Note that these coefficients are linear in the third-order derivatives of  $f$ . Their vanishing is equivalent to some 35 identities constituting an involutive closed system of third-order PDEs for  $f$ . These are precisely the integrability conditions as derived in [29]. q.e.d.

## 6. Type IV: systems of hydrodynamic type

In this section we discuss geometric aspects of integrable systems of hydrodynamic type (6),

$$A(u)u_x + B(u)u_y + C(u)u_t = 0,$$

where  $u = (u^1, u^2)^t$  is a two-component column vector of the dependent variables, and  $A(u), B(u), C(u)$  are  $2 \times 2$  matrices. It will be assumed that there is a matrix in the span of  $A, B, C$  that is hyperbolic. This class is invariant under arbitrary changes of variables  $u^1, u^2$ , as well as linear transformations of  $x, y, t$ , which constitute the natural equivalence group of the problem. If  $C$  is non-degenerate, the multiplication by  $C^{-1}$  brings the system into evolutionary form. The integrability of systems of hydrodynamic type was investigated in [27]. It was shown in [54] that generic integrable system of the form (6) can be parametrized by generalized hypergeometric functions. The linearized system has the form

$$A(u)v_x + B(u)v_y + C(u)v_t + \dots = 0,$$

where dots denote terms which do not contain derivatives of  $v$ . The corresponding dispersion relation (which coincides with the equation for characteristic covectors) is given by the formula

$$\det(\lambda^1 A(u) + \lambda^2 B(u) + \lambda^3 C(u)) = 0.$$

The left-hand side is a quadratic form  $Q$  in  $\lambda = (\lambda^1, \lambda^2, \lambda^3)$ :  $Q(\lambda) = \lambda \cdot D \cdot \lambda^t$ , where  $D = D(u)$  is a  $3 \times 3$  symmetric matrix. It defines the

conformal structure  $g = g_{ij}dx^i dx^j$ , where the matrix of  $g_{ij}$  is the inverse of  $D$ . Recall that  $(x^1, x^2, x^3) = (x, y, t)$ . Our first result is as follows:

**Theorem 8.** *System (6) is linearizable by a transformation from the equivalence group if and only if it is integrable and the conformal structure  $g$  is conformally flat on every solution (note that conformal flatness alone is no longer sufficient for the linearizability).*

*Proof.* Without any loss of generality, one can assume that system (6) is represented in the form

$$(23) \quad \begin{pmatrix} u^1 \\ u^2 \end{pmatrix}_t + \begin{pmatrix} a & 0 \\ 0 & b \end{pmatrix} \begin{pmatrix} u^1 \\ u^2 \end{pmatrix}_x + \begin{pmatrix} p & q \\ r & s \end{pmatrix} \begin{pmatrix} u^1 \\ u^2 \end{pmatrix}_y = 0$$

(multiply by  $C^{-1}$  and use a change of variables  $u^1, u^2$  to make the matrix  $C^{-1}A$  diagonal: such diagonalization is always possible in the two-component hyperbolic situation). Here the matrix elements  $a, b, p, q, r, s$  are functions of  $u^1, u^2$ . The corresponding dispersion relation takes the form

$$(24) \quad (\lambda^1 a + \lambda^2 p + \lambda^3)(\lambda^1 b + \lambda^2 s + \lambda^3) - qr(\lambda^2)^2 = 0,$$

with the associated matrix

$$D(u) = \begin{pmatrix} ab & \frac{as+bp}{2} & \frac{a+b}{2} \\ \frac{as+bp}{2} & ps - qr & \frac{p+s}{2} \\ \frac{a+b}{2} & \frac{p+s}{2} & 1 \end{pmatrix}.$$

This gives rise to the conformal structure  $g = g_{ij}dx^i dx^j$ , where the matrix of  $g_{ij}$  is the inverse of  $D$ . We require  $g$  to be conformally flat for every background solution  $u$ . Calculating the Cotton tensor (10) and using (23) and its differential consequences to eliminate all higher-order derivatives of  $u$  that contain differentiation by  $t$ , we obtain the expression which has to vanish identically in the remaining higher-order derivatives of  $u$  (maximum third-order derivatives will appear in this calculation). In particular, requiring that coefficients at the remaining third-order derivatives of  $u$  vanish identically, we obtain that  $a, b, p, s$ , and  $qr$  must be constant (thus, all coefficients of the dispersion relation are constants). In this case, the rest of the Cotton tensor vanishes identically. By a transformation from the equivalence group any such system can be brought to the form

$$u_t^1 = qu_y^2, \quad u_y^1 = qu_x^2,$$

where  $q$  is still an arbitrary function of  $u^1, u^2$ . For generic  $q$ , such systems are neither linearizable nor integrable. Imposing the integrability conditions as derived in [27], we obtain just one extra constraint,

$(\ln q)_{u^1 u^2} = 0$ , which is equivalent to the existence of a change of variables  $u^1 \rightarrow \varphi^1(u^1)$ ,  $u^2 \rightarrow \varphi^2(u^2)$  bringing system (23) to a constant coefficient form. q.e.d.

Formal linearizations of *integrable* systems of the form (6) give rise to conformal structures satisfying the Einstein–Weyl property.

**Example 1.** Let us consider the system

$$\begin{aligned} u_t^1 + \frac{u^1 u_x^1 + u^2 u_x^2}{(u^1)^2 + (u^2)^2} + \frac{u^1 u_y^2 - u^2 u_y^1}{(u^1)^2 + (u^2)^2} &= 0, \\ u_t^2 + \frac{u^2 u_x^1 - u^1 u_x^2}{(u^1)^2 + (u^2)^2} + \frac{u^1 u_y^1 + u^2 u_y^2}{(u^1)^2 + (u^2)^2} &= 0, \end{aligned}$$

which was obtained in [26] as the first-order form of the Boyer–Finley equation: indeed, the expression  $\rho = (u^1)^2 + (u^2)^2$  satisfies the equation

$$\rho_{tt} = \Delta \ln \rho,$$

$\Delta = \partial_x^2 + \partial_y^2$ . In this case, the dispersion relation is

$$\frac{(\lambda^1)^2 + (\lambda^2)^2}{\rho} - (\lambda^3)^2 = 0.$$

The corresponding Einstein–Weyl structure takes the familiar form [68]:

Conformal structure:  $g = \rho(dx^2 + dy^2) - dt^2$ .

Covector:  $\omega = 2\frac{\rho_t}{\rho} dt$ .

**Example 2.** A class of Hamiltonian systems of hydrodynamic type can be represented in the form

$$(25) \quad v_t + (H_v)_y = 0, \quad w_x + (H_w)_y = 0;$$

here  $H(v, w)$  is the Legendre transform of the Hamiltonian density. The dispersion relation,

$$(\lambda^2 H_{vv} + \lambda^3)(\lambda^2 H_{ww} + \lambda^1) - (\lambda^2 H_{vw})^2 = 0,$$

gives rise to the conformal structure

$$g = (dy - H_{ww} dx - H_{vw} dt)^2 - 4H_{vw}^2 dx dt.$$

The corresponding covector  $\omega$  is given by the formula (8),

$$\begin{aligned} H_{vw}^2 \omega &= \left( H_{ww} \Delta_y + 2H_{vw}^2 (H_{ww})_y + H_{ww} ((H_{vv})_x + (H_{ww})_t) \right) dx \\ &\quad - \left( \Delta_y + (H_{vv})_x + (H_{ww})_t \right) dy \\ &\quad + \left( H_{vv} \Delta_y + 2H_{vw}^2 (H_{vv})_y + H_{vv} ((H_{vv})_x + (H_{ww})_t) \right) dt; \end{aligned}$$

here  $\Delta = H_{vv} H_{ww} - 2H_{vw}^2$ . One can verify that  $g, \omega$  satisfy the Einstein–Weyl constraints if and only if the potential  $H(v, w)$  satisfies the set

of integrability conditions as derived in [26] based on the method of hydrodynamic reductions:

$$\begin{aligned}
 H_{vw}H_{vvv} &= 2H_{vv}H_{vvw}, \\
 H_{vw}H_{vvv} &= 2H_{vv}H_{vww}, \\
 (26) \quad H_{vw}H_{vvvw} &= H_{vv}H_{vww} + H_{vv}H_{wvw}, \\
 H_{vw}H_{vww} &= 2H_{vv}H_{wvw}, \\
 H_{vw}H_{wvw} &= 2H_{vw}H_{wvw}.
 \end{aligned}$$

This system is in involution. It was shown in [33] that its generic solution is given by the formula

$$(27) \quad H(v, w) = Z(v + w) + \epsilon Z(v + \epsilon w) + \epsilon^2 Z(v + \epsilon^2 w);$$

here  $\epsilon = e^{2\pi i/3}$ , and  $Z''(s) = \zeta(s)$ , where  $\zeta$  is the Weierstrass zeta-function:  $\zeta' = -\wp$ ,  $(\wp')^2 = 4\wp^3 - g_3$  (equianharmonic case  $g_2 = 0$ ). Degenerations of this solution correspond to

$$H(v, w) = \frac{1}{2}v^2\zeta(w), \quad H(v, w) = (v + w) \ln(v + w),$$

as well as the following polynomial potentials:

$$H(v, w) = v^2w^2, \quad H(v, w) = vw^2 + \frac{\alpha}{5}w^5, \quad H(v, w) = vw + \frac{1}{6}w^3.$$

The second main result of this section is as follows:

**Theorem 9.** *System (6) with non-constant dispersion relation is integrable by the method of hydrodynamic reductions if and only if the corresponding conformal structure  $g$  is Einstein–Weyl on every solution, with the covector  $\omega$  given by (8).*

*Proof.* We will give two proofs of this result. The first one is computational, based on the explicit calculation of the Einstein–Weyl constraints and the integrability conditions as derived in [27]. The second proof utilizes the fact that any two-component integrable system of hydrodynamic type possesses a dispersionless Lax pair [27].

The first proof can be summarized as follows. We consider system (6) represented in the form (23). The associated conformal structure is  $g = g_{ij}dx^i dx^j$ , where  $g_{ij}$  is the inverse of  $D$ . We will seek a covector  $\omega = \omega_k dx^k$  in the form  $\omega_k = T_{kj}^i u_{x^i}^j$ , where  $T_{kj}^i$  are certain functions of  $u^1, u^2$ . Imposing the Einstein–Weyl equations (7) and using (23) and its differential consequences to eliminate all higher-order derivatives of  $u$  that contain differentiation by  $t$  (maximum second-order derivatives of  $u$  will occur in this calculation), we obtain a set of relations that have to vanish identically in the remaining partial derivatives of  $u$ . Thus, equating to zero coefficients at the second-order derivatives of  $u$ , we obtain the expression (8) for  $\omega$ . Equating to zero the remaining terms, we obtain 15 relations providing all second-order partial derivatives (in  $u^1, u^2$ ) of

the coefficients of the dispersion relation, i.e., of  $a, b, p, s$ , and  $qr$ . These relations constitute part of the total set of 16 integrability conditions as derived in [27]. To recover the missing condition, we calculate the compatibility conditions for the 15 relations at hand. This leads to the following two cases:

**Case 1:** All coefficients of the dispersion relation are constant, in this case all of the 15 relations are satisfied identically. Any such system can be brought to the form  $u_t^1 = qu_y^2$ ,  $u_y^1 = qu_x^2$ , see the proof of Theorem 8. Although the Einstein–Weyl property is trivially satisfied, the system does not need to be integrable for generic  $q(u^1, u^2)$ . This is why we need to eliminate the case of constant dispersion relation from the statement of Theorem 9.

**Case 2:** One of the 15 relations—namely, the expression for the mixed derivative  $(qr)_{u^1u^2}$ , splits into two separate expressions for  $q_{u^1u^2}$  and  $r_{u^1u^2}$ . This gives all of the 16 integrability conditions as derived in [27] based on the method of hydrodynamic reductions, thus finishing the first proof.

As in Theorem 5, there exists another demonstration that the integrability is equivalent to the Einstein–Weyl property. It is based on the fact that any integrable system of hydrodynamic type, which we again assume represented in the form (23), possesses a dispersionless Lax pair

$$(28) \quad S_t = f(S_y, u^1, u^2), \quad S_x = g(S_y, u^1, u^2).$$

This means that the consistency condition  $S_{tx} = S_{xt}$  is equivalent to the system (23). Differentiating (28) by  $y$  and setting  $S_y = \lambda$ , we obtain

$$(29) \quad \lambda_t = f_\lambda \lambda_y + f_{u^1} u_y^1 + f_{u^2} u_y^2, \quad \lambda_x = g_\lambda \lambda_y + g_{u^1} u_y^1 + g_{u^2} u_y^2.$$

With this system we associate the pair of vector fields

$$X = \frac{\partial}{\partial t} - f_\lambda \frac{\partial}{\partial y} + (f_{u^1} u_y^1 + f_{u^2} u_y^2) \frac{\partial}{\partial \lambda}, \quad Y = \frac{\partial}{\partial x} - g_\lambda \frac{\partial}{\partial y} + (g_{u^1} u_y^1 + g_{u^2} u_y^2) \frac{\partial}{\partial \lambda},$$

which live in the extended four-dimensional space with coordinates  $x, y, t, \lambda$ . Note that the compatibility condition  $\lambda_{tx} = \lambda_{xt}$  is equivalent to the commutativity of these vector fields. Projecting the two-parameter family of integral surfaces of the distribution spanned by  $X, Y$  to the space of independent variables  $x, y, t$ , we obtain a two-parameter family of null totally geodesic surfaces of the Weyl connection  $\mathbb{D}$ . This can be seen as follows. Equations (29) mean that the vectors  $X, Y$  are tangential to the hypersurface  $\lambda = \lambda(x, y, t)$ . Projecting  $X$  and  $Y$  to the space of independent variables  $x, y, t$ , we obtain two vector fields

$$\hat{X} = \frac{\partial}{\partial t} - f_\lambda \frac{\partial}{\partial y}, \quad \hat{Y} = \frac{\partial}{\partial x} - g_\lambda \frac{\partial}{\partial y},$$

which commute if and only if  $\lambda$  satisfies the equations (29). It remains to show that  $\hat{X}$  and  $\hat{Y}$  form a null distribution (i.e., tangential to the null

cones of the conformal structure  $g$ ), and that the covariant derivatives  $\mathbb{D}_{\hat{X}}\hat{X}$ ,  $\mathbb{D}_{\hat{X}}\hat{Y}$ ,  $\mathbb{D}_{\hat{Y}}\hat{X}$ ,  $\mathbb{D}_{\hat{Y}}\hat{Y}$  belong to the span of  $\hat{X}, \hat{Y}$ . Equivalently, one can introduce the covector  $\theta = g_\lambda dx + dy + f_\lambda dt$  that annihilates  $\hat{X}, \hat{Y}$  and verify that  $\theta$  is null, and that  $\mathbb{D}_{\hat{X}}\theta \wedge \theta = \mathbb{D}_{\hat{Y}}\theta \wedge \theta = 0$ . This follows from the equations satisfied by the functions  $f(\lambda, u^1, u^2)$  and  $g(\lambda, u^1, u^2)$  as derived in [27]:

$$\begin{aligned} f_{u^1} &= -a g_{u^1}, & f_{u^2} &= -b g_{u^2}, \\ f_\lambda &= \frac{b(p+r\frac{g_{u^2}}{g_{u^1}}) - a(s+q\frac{g_{u^1}}{g_{u^2}})}{a-b}, & g_\lambda &= \frac{s+q\frac{g_{u^1}}{g_{u^2}} - p-r\frac{g_{u^2}}{g_{u^1}}}{a-b}, \\ g_{u^1 u^1} &= \frac{g_{u^1}(g_{u^2}^2(r(b_{u^1}-a_{u^1})+(a-b)r_{u^1})+g_{u^1}g_{u^2}((a-b)p_{u^1}+(s-p)a_{u^1}-ra_{u^2})+qa_{u^1}g_{u^2}^2)}{(a-b)rg_{u^2}^2}, \\ g_{u^1 u^2} &= \frac{1}{b-a}(a_{u^2}g_{u^1} - b_{u^1}g_{u^2}), \\ g_{u^2 u^2} &= \frac{g_{u^2}(g_{u^1}^2(q(a_{u^2}-b_{u^2})+(b-a)q_{u^2})+g_{u^1}g_{u^2}((b-a)s_{u^2}+(p-s)b_{u^2}-qb_{u^1})+rb_{u^2}g_{u^1}^2)}{(b-a)qg_{u^1}^2}. \end{aligned}$$

Note that  $f_\lambda$  and  $g_\lambda$  satisfy the relation

$$(a g_\lambda + p + f_\lambda)(b g_\lambda + s + f_\lambda) - qr = 0,$$

which means that the covector  $\theta$  is null in the conformal structure defined by the dispersion relation (24). This finishes the second proof of Theorem 9. q.e.d.

## 7. Further integrable examples in 3D

In this section we collect miscellaneous examples of dispersionless integrable systems in 3D related to Einstein–Weyl geometry, which do not fit into the classes discussed above.

**Example 1.** The following system was derived by Manakov and Santini [51] as a two-component generalization of the dKP equation:

$$u_{xt} - u_{yy} + (uu_x)_x + v_x u_{xy} - v_y u_{xx} = 0, \quad v_{xt} - v_{yy} + uv_{xx} + v_x v_{xy} - v_y v_{xx} = 0.$$

Its formal linearization results upon setting  $u \rightarrow u + \epsilon u_1$ ,  $v \rightarrow v + \epsilon v_1$ , which gives

$$Lu_1 + \dots = 0, \quad Lv_1 + \dots = 0;$$

here  $L = \partial_x \partial_t - \partial_y^2 + (u - v_y) \partial_x^2 + v_x \partial_x \partial_y$ , and dots denote terms that do not contain second-order derivatives of  $u_1$  and  $v_1$ . The symbol of  $L$  gives rise to the conformal structure

$$g = (dy - v_x dt)^2 - 4(dx - (u - v_y) dt) dt,$$

which satisfies the Einstein–Weyl equations with the corresponding covector  $\omega$  given by the formula (8):

$$\omega = -v_{xx} dy + (4u_x - 2v_{xy} + v_x v_{xx}) dt.$$

This Einstein–Weyl structure was obtained in [19].

**Example 2.** The following system was proposed by Bogdanov [4] as a two-component generalization of the BF equation:

$$(e^{-\phi})_{tt} = m_t \phi_{xy} - m_x \phi_{yt}, \quad m_{tt} e^{-\phi} = m_x m_{yt} - m_t m_{xy}.$$

Its formal linearization results upon setting  $\phi \rightarrow \phi + \epsilon \phi_1$ ,  $m \rightarrow m + \epsilon m_1$ , which gives

$$L\phi_1 + \dots = 0, \quad Lm_1 + \dots = 0;$$

here  $L = e^{-\phi} \partial_t^2 - m_x \partial_y \partial_t + m_t \partial_x \partial_y$ , and dots denote terms that do not contain second-order derivatives of  $\phi_1$  and  $m_1$ . The symbol of  $L$  gives rise to the conformal structure

$$g = (m_x dx + m_t dt)^2 + 4e^{-\phi} m_t dx dy,$$

which satisfies the Einstein–Weyl equations with the corresponding covector  $\omega$  given by the formula (8):

$$\omega = \left( \frac{m_{tt}}{m_t^2} - 2 \frac{\phi_t}{m_t} \right) (m_x dx + m_t dt) + 2 \frac{m_{yt}}{m_t} dy.$$

**Example 3.** The equation

$$(30) \quad m_t^\rho m_{tt} = m_x m_{yt} - m_t m_{xy}$$

was obtained in [4] as a reduction of the two-component BF system from Example 2: set  $\phi = -\rho \ln m_t$ . Since this equation fits into the class of quasilinear wave equations discussed in Section 4, it gives rise to the Einstein–Weyl geometry with the corresponding conformal structure

$$g = (m_x dx + m_t dt)^2 + 4m_t^{\rho+1} dx dy,$$

and the covector  $\omega$  given by (8):

$$\omega = (2\rho + 1) \frac{m_{tt}}{m_t^2} (m_x dx + m_t dt) + 2 \frac{m_{yt}}{m_t} dy.$$

Note that these  $g$  and  $\omega$  result, via the substitution  $\phi = -\rho \ln m_t$ , from the formulae of Example 2. It was pointed out in [5] that, applying to this equation the contact transformation defined as

$$(x, y, t, m, m_x, m_y, m_t) \rightarrow (x, y, \tau, F, F_x, F_y, F_\tau),$$

where  $\tau = \frac{1}{2} \ln m_t$  and

$$F = \frac{m - tm_t}{\sqrt{m_t}}, \quad F_x = \frac{m_x}{\sqrt{m_t}}, \quad F_y = \frac{m_y}{\sqrt{m_t}}, \quad F_\tau = -2t\sqrt{m_t} - \frac{m - tm_t}{\sqrt{m_t}},$$

one obtains the equation derived by Dunajski and Tod in the context hyper-Kähler metrics with conformal symmetry [22],

$$(31) \quad (F_y + F_{y\tau})(F_x - F_{x\tau}) - (F - F_{\tau\tau})F_{xy} = 4e^{2\rho\tau}.$$

It was shown in [22] that this equation gives rise to the Einstein–Weyl structure

$$g = (F d\tau + F_x dx - F_y dy - dF_\tau)^2 + 16e^{2\rho\tau} dx dy,$$

$$\omega = 4\rho d\tau + \frac{(2 + 4\rho)(F_x - F_{x\tau})dx + (2 - 4\rho)(F_y + F_{y\tau})dy}{F - F_{\tau\tau}}.$$

One can verify that the Einstein–Weyl structure of the equation (31) can be obtained from that of the equation (30) by applying the above contact transformation (and an appropriate rescaling). Note that in the last case the covector  $\omega$  is no longer given by our formula (8) (unless  $\rho = 0$ , in which case the equation becomes translationally invariant). The reason for this is that, although the Einstein–Weyl property is clearly contact-invariant, this is not the case for our formula for  $\omega$ . We recall that, given a conformal structure  $g$ , the problem of reconstruction of the corresponding covector  $\omega$  from the Einstein–Weyl constraints is far from trivial [24]. What simplifies this problem in our case is that  $\omega$  should be given by a “universal” formula depending on finite-order jets of a solution  $u$ . In any case, a fully contact-invariant approach to dispersionless integrable systems in 3D is yet to be developed.

## 8. Integrability in 4D and self-duality

There are very few classification results in 4D. Here we consider the case of symplectic Monge–Ampère equations, i.e., equations represented as linear combinations of minors of the Hessian matrix of a function  $u(x^1, x^2, x^3, x^4)$ , which constitute a subclass of equations of type III (in 4D). Below,  $u_{ij}$  denotes  $u_{x^i x^j}$ .

**Theorem 10** ([16]). *Over the field of complex numbers, any integrable non-degenerate symplectic Monge–Ampère equation is  $Sp(8)$ -equivalent to one of the following normal forms:*

- 1)  $u_{11} - u_{22} - u_{33} - u_{44} = 0$  (linear wave equation);
- 2)  $u_{13} + u_{24} + u_{11}u_{22} - u_{12}^2 = 0$  (second heavenly equation [59]);
- 3)  $u_{13} = u_{12}u_{44} - u_{14}u_{24}$  (modified heavenly equation);
- 4)  $u_{13}u_{24} - u_{14}u_{23} = 1$  (first heavenly equation [59]);
- 5)  $u_{11} + u_{22} + u_{13}u_{24} - u_{14}u_{23} = 0$  (Husain equation [40]);
- 6)  $\alpha u_{12}u_{34} + \beta u_{13}u_{24} + \gamma u_{14}u_{23} = 0$  (general heavenly equation),  
 $\alpha + \beta + \gamma = 0$ .

These heavenly-type equations are known to possess Lax pairs of the form  $[X_1, X_2] = 0$ , where  $X_1, X_2$  are parameter-dependent vector fields that commute modulo the corresponding equations [59, 40, 16, 50]:

**Second heavenly equation**  $u_{13} + u_{24} + u_{11}u_{22} - u_{12}^2 = 0$ :

$$X_1 = \partial_4 + u_{11}\partial_2 - (u_{12} - \lambda)\partial_1, \quad X_2 = \partial_3 - (u_{12} + \lambda)\partial_2 + u_{22}\partial_1.$$

**Modified heavenly equation**  $u_{13} = u_{12}u_{44} - u_{14}u_{24}$ :

$$X_1 = u_{14}\partial_2 - u_{12}\partial_4 + \lambda\partial_1, \quad X_2 = u_{44}\partial_2 - (u_{24} - \lambda)\partial_4 - \partial_3.$$

**First heavenly equation**  $u_{13}u_{24} - u_{14}u_{23} = 1$ :

$$X_1 = u_{13}\partial_4 - u_{14}\partial_3 + \lambda\partial_1, \quad X_2 = -u_{23}\partial_4 + u_{24}\partial_3 - \lambda\partial_2.$$

**Husain equation**  $u_{11} + u_{22} + u_{13}u_{24} - u_{14}u_{23} = 0$ :

$$X_1 = \partial_2 + u_{13}\partial_4 - u_{14}\partial_3 + \lambda\partial_1, \quad X_2 = \partial_1 - u_{23}\partial_4 + u_{24}\partial_3 - \lambda\partial_2.$$

**General heavenly equation**  $\alpha u_{12}u_{34} + \beta u_{13}u_{24} + \gamma u_{14}u_{23} = 0$ :

$$X_1 = \partial_1 - \frac{u_{13}}{u_{34}}\partial_4 + \gamma\lambda\left(\partial_1 - \frac{u_{14}}{u_{34}}\partial_3\right), \quad X_2 = \frac{u_{23}}{u_{34}}\partial_4 - \partial_2 + \beta\lambda\left(\partial_2 - \frac{u_{24}}{u_{34}}\partial_3\right).$$

One can show that the distribution spanned by  $X_1$  and  $X_2$  is totally null with respect to the conformal structure provided by the symbol of formal linearization. The integrability of this distribution for any value of the spectral parameter  $\lambda$  implies the existence of a three-parameter family of null surfaces ( $\alpha$ -surfaces), the property known to be equivalent to self-duality. Another way to see this is to notice that the above Lax pairs are linear in  $\lambda$ :

$$X_1 = X + \lambda Y, \quad X_2 = P + \lambda Q.$$

Furthermore, the corresponding formal linearizations can be written in the form

$$(XQ - YP)v + \dots = 0,$$

where the expression  $XQ - YP$  is understood as a second-order differential operator and dots denote terms containing first-order derivatives of  $v$ . According to [21, 37], this means that the symbols of formal linearizations must be self-dual (in fact, hypercomplex, but this property is known to imply self-duality). These and other examples support the following conjectures relating the linearizability/integrability of four-dimensional PDEs to conformal geometry of symbols of their formal linearizations:

- A 4D second-order dispersionless PDE is linearizable (by a transformation from the appropriate equivalence group) if and only if the corresponding conformal structure is conformally flat on every solution.
- A 4D second-order dispersionless PDE is integrable by the method of hydrodynamic reductions if and only if the corresponding conformal structure is (anti) self-dual on every solution. Since the equations of self-duality are known to be integrable by the twistor construction [57], this supports the evidence that solutions to integrable PDEs must carry integrable geometry.

A convenient approach to the integrability of four-dimensional PDEs is based on the requirement that all their traveling wave reductions to 3D must be integrable. This necessary condition turns out to be very strong indeed, and in many cases is already sufficient for integrability [16]. Since the integrability in 3D is related to the Einstein–Weyl condition, this gives yet another confirmation of the well-known fact that symmetry reductions of the self-duality equations lead to Einstein–Weyl geometry [42, 9, 10].

### 9. Appendix: the method of hydrodynamic reductions

As proposed in [26], the method of hydrodynamic reductions applies to first-order quasilinear systems of the form

$$(32) \quad A(\mathbf{u})\mathbf{u}_x + B(\mathbf{u})\mathbf{u}_y + C(\mathbf{u})\mathbf{u}_t = 0,$$

or equations transformable into this form by a suitable change of variables. Here  $\mathbf{u} = (u^1, \dots, u^m)^\mathfrak{t}$  is an  $m$ -component column vector of the dependent variables and  $A, B, C$  are  $l \times m$  matrices where  $l$ , the number of equations, is allowed to exceed the number of the unknowns,  $m$ . The system (32) will be assumed involutive with the general solution depending on a certain number of arbitrary functions of *two* variables. Systems of this type are referred to as 3D dispersionless PDEs. Typically, they arise as dispersionless limits of integrable soliton equations: the canonical example is the KP equation,  $u_t - uu_x + \epsilon^2 u_{xxx} - w_y = 0$ ,  $w_x = u_y$ , which assumes the form (32) in the limit  $\epsilon \rightarrow 0$ .

It will be demonstrated below that equations of types (3)–(5) are indeed within the class (32). The method of hydrodynamic reductions consists of seeking  $N$ -phase solutions in the form

$$(33) \quad \mathbf{u} = \mathbf{u}(R^1, \dots, R^N),$$

where every “phase”  $R^i(x, y, t)$ ,  $1 \leq i \leq N$ , is required to satisfy a pair of consistent equations (no summation)

$$(34) \quad R_{yt}^i = \mu^i(R)R_x^i, \quad R_t^i = \lambda^i(R)R_x^i.$$

The variables  $R^i$  are also known as *Riemann invariants*. The compatibility conditions  $R_{yt}^i = R_{ty}^i$  imply the following restrictions for the characteristic speeds  $\mu^i$  and  $\lambda^i$ :

$$(35) \quad \frac{\partial_j \mu^i}{\mu^j - \mu^i} = \frac{\partial_j \lambda^i}{\lambda^j - \lambda^i}, \quad i \neq j, \quad \partial_i = \frac{\partial}{\partial R^i}.$$

Commuting systems of the form (34) can be solved by the generalized hodograph method [67]. Equations of the form (34) are known as  $N$ -component *systems of hydrodynamic type*; their Hamiltonian and geometric aspects have been thoroughly investigated in [17, 67].

We require that *every* solution of (34) gives rise to a solution of the original system (32) via (33). In this case, equations (34) are said to constitute an  $N$ -component *hydrodynamic reduction* of the original system. Each hydrodynamic reduction can be viewed as a decomposition of the original  $m$ -component 3D system into a pair of commuting  $N$ -component 2D systems. It is remarkable that there exist 3D systems possessing an infinity of such reductions.

**Definition** ([26]). *The system (32) is said to be integrable by the method of hydrodynamic reductions if, for any  $N$ , it possesses infinitely many  $N$ -component hydrodynamic reductions parametrized by  $N$  arbitrary functions of one variable.*

One can show that the existence of 3-component reductions depending on three arbitrary functions of one variable is already sufficiently restrictive, and implies the integrability. This is reminiscent of the well-known 3-soliton condition in the Hirota bilinear approach. For a particular reduction, the corresponding solutions constitute only a very narrow subclass of solutions of the original system (32). As  $N$  varies, solutions coming from  $N$ -component hydrodynamic reductions form an ever growing subset of solutions of the system (32) that is, in a sense, locally dense (i.e., these solutions do not satisfy any finite-order differential constraints other than the equation itself and its differential consequences). Multi-phase solutions of this form originate from gas dynamics [62, 7, 58, 38, see also references therein]. They are sometimes referred to as *non-linear interactions of planar simple waves*. For  $N = 1, 2$ , they are called *simple waves* and *double waves*, respectively, and belong to the class of solutions with a degenerate hodograph.

The above definition provides an efficient classification criterion. In general, one proceeds as follows: substituting (33) into (32) and using (34), one obtains

$$(36) \quad (A + \mu^i B + \lambda^i C) \partial_i \mathbf{u} = 0.$$

The condition of the non-trivial solvability of this linear system provides the dispersion relation for the characteristic speeds  $\mu^i$  and  $\lambda^i$  for any  $i = 1, \dots, N$ . Namely, the pair  $\lambda, \mu$  (more precisely, the covector  $dx + \mu dy + \lambda dt$ ) is called characteristic if

$$rk(A + \mu B + \lambda C) < m.$$

We will assume that the dispersion relation defines an irreducible algebraic curve. For instance, in the case  $l = m$  this gives an algebraic curve of degree  $m$ ,

$$det(A + \mu B + \lambda C) = 0.$$

In the language of PDEs, this is the part of the characteristic variety of our system [44] lying in an affine chart.

For all examples discussed in this paper, the dispersion relation reduces to a non-degenerate conic determined by the symbol of formal linearization. Equations (35) and (36) form an overdetermined system for the functions  $\mathbf{u}(R)$  and the characteristic speeds  $\mu^i(R), \lambda^i(R)$ . The requirement that they are consistent, with the general solution depending on  $N$  arbitrary functions of one variable, leads to the integrability conditions in terms of the original matrices  $A, B, C$ . Let us illustrate

the method of hydrodynamic reductions using the example of the dKP equation, in which case all calculations can be verified by hand.

**Example 1: hydrodynamic reductions of the dKP equation.** Let us write the dKP equation  $u_{xt} - (uu_x)_x - u_{yy} = 0$  in the form (32):

$$(37) \quad u_t - uu_x = w_y, \quad u_y = w_x.$$

Looking for  $N$ -phase solutions,  $u = u(R^1, \dots, R^N)$ ,  $w = w(R^1, \dots, R^N)$ , where the phases  $R^i$  satisfy Equations (34), one obtains the relations

$$(38) \quad \partial_i w = \mu^i \partial_i u, \quad \lambda^i = u + (\mu^i)^2,$$

(the second one is the dispersion relation). The compatibility condition  $\partial_i \partial_j w = \partial_j \partial_i w$  implies the equations ( $1 \leq i, j \leq N$ )

$$(39) \quad \partial_i \partial_j u = \frac{\partial_j \mu^i}{\mu^j - \mu^i} \partial_i u + \frac{\partial_i \mu^j}{\mu^i - \mu^j} \partial_j u, \quad i \neq j,$$

which, along with the commutativity conditions (35), result in the equations for  $u(R)$  and  $\mu^i(R)$ , known as the Gibbons–Tsarev system,

$$(40) \quad \partial_j \mu^i = \frac{\partial_j u}{\mu^j - \mu^i}, \quad \partial_i \partial_j u = 2 \frac{\partial_i u \partial_j u}{(\mu^j - \mu^i)^2}, \quad i \neq j,$$

first derived in [35] in the context of hydrodynamic reductions of the Benney moment equations; see also [34].

For any solution  $\mu^i, u$  of the system (40), one can reconstruct  $\lambda^i$  and  $w$  by virtue of (38). The system (40) is compatible and its general solution depends, modulo transformations  $R^i \rightarrow f^i(R^i)$ , on  $N$  arbitrary functions of one variable. This gives the required family of  $N$ -component hydrodynamic reductions parametrized by  $N$  arbitrary functions of one variable and establishes the integrability of dKP. We point out that the compatibility conditions,  $\partial_k \partial_j \mu^i = \partial_j \partial_k \mu^i$  and  $\partial_k \partial_i \partial_j u = \partial_j \partial_i \partial_k u$ , involve *triples* of indices  $i \neq j \neq k$  only. Thus, the consistency of the system (40) for  $N = 3$  implies its consistency for arbitrary  $N$ . This turns out to be a general phenomenon: as mentioned above, the existence of 3-component reductions is already sufficient for the integrability.

Let us demonstrate that equations (3)–(6) can be brought into the first-order quasilinear form (32).

### Equations of type I:

$$a_{xx} + b_{yy} + c_{tt} + 2p_{xy} + 2q_{xt} + 2r_{yt} = 0.$$

Recall that  $a, b, c, p, q, r$  are functions of one and the same dependent variable  $u$ . Consider the first-order system

$$a_x + p_y + q_t = -\varphi_t, \quad b_y + p_x + r_t = \psi_t, \quad c_t + q_x + r_y = \varphi_x - \psi_y,$$

which implies the above second-order PDE on elimination of the auxiliary potentials  $\varphi, \psi$ . This system is invariant under gauge transformations  $\varphi \rightarrow \varphi + \eta_y$ ,  $\psi \rightarrow \psi + \eta_x$ , where  $\eta$  is an arbitrary function of  $x, y$ .

One can show that this gauge freedom can be eliminated by imposing a compatible differential constraint

$$a'\psi_x + p'\psi_y + q'\psi_t + p'\varphi_x + b'\varphi_y + r'\varphi_t = 0.$$

This results in a system of the form (32), with  $\mathbf{u} = (u, \varphi, \psi)$  and  $(l, m) = (4, 3)$ . It was investigated by the method of hydrodynamic reductions in [32] (see Theorem 1 for the integrability conditions).

#### Equations of type II:

$$f_{11}u_{xx} + f_{22}u_{yy} + f_{33}u_{tt} + 2f_{12}u_{xy} + 2f_{13}u_{xt} + 2f_{23}u_{yt} = 0.$$

Recall that  $f_{ij}$  are functions of the first-order derivatives of  $u$  only. Setting  $u_x = a$ ,  $u_y = b$ ,  $u_t = c$ , one obtains an equivalent first-order representation,

$$\begin{aligned} a_y = b_x, \quad a_t = c_x, \quad b_t = c_y, \\ f_{11}a_x + f_{22}b_y + f_{33}c_t + 2f_{12}a_y + 2f_{13}a_t + 2f_{23}b_t = 0, \end{aligned}$$

where  $f_{ij}$  are now viewed as functions of  $a, b, c$ . This is again of the form (32) with  $\mathbf{u} = (a, b, c)$  and  $(l, m) = (4, 3)$ . This type of equations was investigated by the method of hydrodynamic reductions in [8].

#### Equations of type III:

$$F(u_{xx}, u_{xy}, u_{yy}, u_{xt}, u_{yt}, u_{tt}) = 0.$$

Rewriting the equation in explicit form,

$$u_{tt} = f(u_{xx}, u_{xy}, u_{yy}, u_{xt}, u_{yt}),$$

and setting  $u_{xx} = a$ ,  $u_{xy} = b$ ,  $u_{yy} = c$ ,  $u_{xt} = p$ ,  $u_{yt} = q$ ,  $u_{tt} = f(a, b, c, p, q)$ , one obtains a first-order quasilinear system by writing out all possible consistency conditions among the second-order derivatives,

$$\begin{aligned} a_y = b_x, \quad a_t = p_x, \quad b_y = c_x, \quad b_t = p_y = q_x, \quad c_t = q_y, \\ p_t = f(a, b, c, p, q)_x, \quad q_t = f(a, b, c, p, q)_y. \end{aligned}$$

This system is of the form (32), with  $\mathbf{u} = (a, b, c, p, q)$  and  $(l, m) = (8, 5)$ . This type of equations was investigated by the method of hydrodynamic reductions in [29].

#### Equations of type IV:

$$A(u)u_x + B(u)u_y + C(u)u_t = 0.$$

This system is of the form (32), with  $(l, m) = (2, 2)$ . It was investigated by the method of hydrodynamic reductions in [27].

**REMARK 4 (hydrodynamic reductions of elliptic PDEs).** Although the method of hydrodynamic reductions was primarily designed to deal with hyperbolic systems, it can be extended to elliptic PDEs. The only difference is that, in the elliptic case, the dispersion curve (characteristic variety) has no real points, so that the characteristic

speeds  $\mu^i, \lambda^i$ , as well as the variables  $R^i$ , must be complex. We will say that an elliptic system (32) with real-analytic coefficients is integrable by the method of hydrodynamic reductions if it possesses “infinitely many” complex-valued solutions (33) where the complex phases  $R^i$  satisfy the commuting systems (34). In this context,  $\mathbf{u}, \mu^i, \lambda^i$  are viewed as complex-analytic functions of  $R^i$ . We would like to stress that the integrability conditions for elliptic PDEs, as well as the procedure of their derivation, are identically the same as in the hyperbolic case. Since the integrability conditions are all we need to prove the Einstein–Weyl property of the symbol of formal linearization, it will still hold for elliptic integrable systems. On imposing appropriate “reality” constraints, the above scheme can lead to elliptic reductions with *real* coefficients. Let us demonstrate this using again the example of dKP.

**Example 2: elliptic reduction of the dKP equation.** Two-phase solutions of the system (37) have the form  $u = u(R^1, R^2)$ ,  $w = w(R^1, R^2)$  where the phases  $R^1, R^2$  satisfy Equations (34). As shown in Example 1, one obtains the relations (38) for  $w$  and  $\lambda^i$ , along with the Gibbons–Tsarev equations (40) for  $\mu^i$  and  $u$ :

$$\partial_2 \mu^1 = \frac{\partial_2 u}{\mu^2 - \mu^1}, \quad \partial_1 \mu^2 = \frac{\partial_1 u}{\mu^1 - \mu^2}, \quad \partial_1 \partial_2 u = 2 \frac{\partial_1 u \partial_2 u}{(\mu^2 - \mu^1)^2}.$$

A particular solution of these equations (the so-called shallow water reduction) is given by

$$\mu^1 = R^2 + 3R^1, \quad \mu^2 = R^1 + 3R^2, \quad u = (R^1 - R^2)^2,$$

so that relations (38) imply

$$w = 2(R^1 - R^2)^2(R^1 + R^2), \\ \lambda^1 = 2(R^1 + R^2)^2 + 8(R^1)^2, \quad \lambda^2 = 2(R^1 + R^2)^2 + 8(R^2)^2.$$

Let us now allow  $R^1$  and  $R^2$  to be complex, such that  $R^2 = \bar{R}^1$ . Setting  $R^1 = p + iq$ ,  $R^2 = p - iq$ , we obtain  $u = -4q^2$ ,  $w = -16pq^2$  (both real!), while equations (34) result, on separating real and imaginary parts, in the following pair of two-component commuting elliptic systems with real coefficients:

$$\begin{pmatrix} p \\ q \end{pmatrix}_y = \begin{pmatrix} 4p & -2q \\ 2q & 4p \end{pmatrix} \begin{pmatrix} p \\ q \end{pmatrix}_x, \\ \begin{pmatrix} p \\ q \end{pmatrix}_t = \begin{pmatrix} 16p^2 - 8q^2 & -16pq \\ 16pq & 16p^2 - 8q^2 \end{pmatrix} \begin{pmatrix} p \\ q \end{pmatrix}_x.$$

Thus, we obtain a two-component elliptic reduction of dKP.

**REMARK 5 (hydrodynamic reductions in higher dimensions).** The method of hydrodynamic reductions generalizes to higher-dimensional quasilinear systems, say,

$$A(\mathbf{u})\mathbf{u}_x + B(\mathbf{u})\mathbf{u}_y + C(\mathbf{u})\mathbf{u}_z + D(\mathbf{u})\mathbf{u}_t = 0.$$

One again looks at  $N$ -phase solutions in the form

$$\mathbf{u} = \mathbf{u}(R^1, \dots, R^N),$$

where the phases  $R^i(x, y, t)$  satisfy a triple of consistent equations

$$R_y^i = \mu^i(R)R_x^i, \quad R_t^i = \lambda^i(R)R_x^i, \quad R_z^i = \eta^i(R)R_x^i;$$

see [25, 28] for further details.

## 10. Concluding remarks

(a) We have characterized the integrability of several classes of dispersionless PDEs in 3D by the Einstein–Weyl property of their formal linearizations. It should be emphasized that the four types of equations under consideration, although invariant under certain equivalence groups, are clearly not contact/point invariant. On the other hand, the property for a second-order PDE to have Einstein–Weyl symbol of formal linearization, is invariant under arbitrary contact (and more generally Lie–Bäcklund type) transformations. Although the covector  $\omega$  will no longer be given by the simple formula (8), it will still depend on finite-order jets of the transformed equation. This suggests a contact-invariant approach to the dispersionless integrability unifying all known examples. The class of Monge–Ampère equations in 3D would be the natural venue to develop a fully contact-invariant theory. We should however, warn the reader that, in general, the Einstein–Weyl property alone may not be sufficient for the dispersionless integrability: as we saw in the proof of Theorem 8, the system  $u_t^1 = qu_y^2$ ,  $u_y^1 = qu_x^2$  has a conformally flat symbol but is not integrable for generic  $q(u^1, u^2)$ .

(b) Solutions of dispersionless PDEs discussed in this paper, and the induced Einstein–Weyl structures, are generally defined on open domains in  $\mathbb{R}^3$ . One cannot guarantee more, since the construction applies to *all* solutions. We believe, however, that, for some particular solutions, global aspects/singularities may come into play. As an illustration, let us consider equations (5) of the dispersionless Hirota type. Geometrically, an equation of this form specifies a hypersurface  $M^5$  in the Lagrangian Grassmannian  $\Lambda^6$  [29]. For some particular integrable equations, such as, e.g., the potential dKP equation  $u_{xt} - u_{xx}^2 - u_{yy} = 0$ , this hypersurface is algebraic. Solutions of the equation correspond to Lagrangian submanifolds in the six-dimensional symplectic space whose Gaussian image belongs to  $M^5$ . Some of them may give rise to algebraic 3-folds in  $M^5$ , with non-trivial global properties. The investigation of algebraic solutions and the induced Einstein–Weyl structures is one of the interesting problems left outside the scope of this paper.

(c) In the case of higher-order dispersionless PDEs in 3D, the symbol of formal linearization defines a generalized conformal structure that supplies each solution with a field of algebraic null cones. Furthermore, the

Lax pair provides a two-parameter family of null surfaces. This suggests that, in the spirit of [11], one can define the “generalized Einstein–Weyl” property responsible for the integrability of higher-order PDEs. We expect that our results will carry over to this more general context.

(d) Attempts to extend our results to dimensions higher than four meet an immediate obstacle: all known integrable (non-linearizable) examples in dimensions five and higher have degenerate symbols. Thus, their solutions do not carry any conventional “geometry.” One possible way to proceed is to require that non-degenerate traveling wave reductions of such PDEs to 3D/4D give rise to Einstein–Weyl/self-dual geometries.

(e) Einstein–Weyl structures are known to be related to third-order ODEs satisfying the so-called Wünschmann and Cartan relations [12, 66, 53]. Thus, every solution of a dispersionless integrable system has a third-order ODE naturally associated with it (to be precise, a point equivalence class of ODEs). The (complexified) space of the dependent and independent variables of this ODE, which can be identified with the space of null totally geodesic surfaces of the Einstein–Weyl structure, is known as the “minitwistor” space [39, 65]. Solutions of this ODE correspond to curves of totally geodesic surfaces passing through a fixed point: these form a three-parameter family, and are known as “twistor curves.” It would be of interest to develop an efficient procedure to explicitly calculate this ODE for multi-phase solutions provided by the method of hydrodynamic reductions.

## References

- [1] M.A. Akivis & V.V. Goldberg, *Projective differential geometry of submanifolds*, Elsevier Science Publishers, Amsterdam (1993), MR 1234487, Zbl 0865.53013.
- [2] I.M. Anderson & N. Kamran, *The variational bicomplex for hyperbolic second-order scalar partial differential equations in the plane*, Duke Math. J. **87**, no. 2 (1997), 265–319, MR 1443529, Zbl 0881.35069.
- [3] B. Banos, *On symplectic classification of effective 3-forms and Monge–Ampère equations*, Diff. Geom. Appl. **19** (2003), 147–166, MR 2002657, Zbl 1035.34004.
- [4] L.V. Bogdanov, *Non-Hamiltonian generalizations of the dispersionless 2DTL hierarchy*, J. Phys. A **43**, no. 43 (2010), 434008, MR 2727782, Zbl 1208.37040.
- [5] L. V. Bogdanov, *Dunajski-Tod equation and reductions of the generalized dispersionless 2DTL hierarchy*, Phys. Lett. A **376**, no. 45 (2012), 2894–2898, MR 2997171.
- [6] G. Boillat, *Sur l'équation générale de Monge–Ampère à plusieurs variables*, C. R. Acad. Sci. Paris Sér. I Math. **313**, no. 11 (1991), 805–808, MR 1139843, Zbl 0753.35107.
- [7] M. Burnat, *The method of characteristics and Riemann's invariants for multidimensional hyperbolic systems*, Sibirsk. Mat. Z. **11** (1970), 279–309, MR 0287163, Zbl 0213.37303; *The method of Riemann invariants and its applications to the theory of plasticity*. I, II. Arch. Mech. (Arch. Mech. Stos.) **23** (1971), 817–838; *ibid.* **24** (1972), 3–26, MR 0321393, Zbl 0238.73026.

- [8] P.A. Burovskii, E.V. Ferapontov & S.P. Tsarev, *Second-order quasilinear PDEs and conformal structures in projective space*, International J. Math. **21**, no. 6 (2010), 799–841, MR 2658410, Zbl 1198.35104.
- [9] David M. J. Calderbank, *Integrable background geometries*, preprint, <http://people.bath.ac.uk/dmjc20/mpapers.html> (2002).
- [10] D. M. J. Calderbank & H. Pedersen, *Selfdual spaces with complex structures, Einstein–Weyl geometry and geodesics*, Ann. Inst. Fourier (Grenoble) **50**, no. 3 (2000), 921–963, MR 1779900, Zbl 0970.53027.
- [11] E. Cartan, *Sur une classe d’espaces de Weyl*, Ann. Sci. École Norm. Sup. (3) **60** (1943), 1–16, MR 0014292, Zbl 0028.30802.
- [12] E. Cartan, *The geometry of differential equations of third order*, Revista Mat. Hisp.-Amer. **4** (1941), 3–33, MR 0004537, Zbl 0063.01955.
- [13] A. Donato, U. Ramgulum & C. Rogers, *The (3+1)-dimensional Monge–Ampère equation in discontinuity wave theory: Application of a reciprocal transformation*, Meccanica **27** (1992), 257–262, Zbl 0764.35066.
- [14] B. Doubrov, *On a class of contact invariants of systems of ordinary differential equations*, Russian Math. (Iz. VUZ) **50**, no. 1 (2006), 73–74, MR 2227551, Zbl 1176.53023.
- [15] B. Doubrov, *Generalized Wilczynski invariants for non-linear ordinary differential equations*, in M. Eastwood and W. Miller (eds.), *Symmetries and Overdetermined Systems of Partial Differential Equations*, IMA Vol. Math. Appl. **144** Springer, NY (2008), 25–40, MR 2384704, Zbl 1149.34026.
- [16] B. Doubrov & E.V. Ferapontov, *On the integrability of symplectic Monge–Ampère equations*, J. Geom. Phys. **60** (2010), 1604–1616, MR 2661158, Zbl 1195.35109.
- [17] B.A. Dubrovin & S.P. Novikov, *Hydrodynamics of weakly deformed soliton lattices: differential geometry and Hamiltonian theory*, Russian Math. Surveys, **44**, no. 6 (1989), 35–124, MR 1037010, Zbl 0712.58032.
- [18] M. Dunajski & P. Tod, *Einstein–Weyl spaces and dispersionless Kadomtsev–Petviashvili equation from Painlevé I and II*, Physics Letters A **303** (2002), 253–264, MR 1937101, Zbl 0999.35083.
- [19] M. Dunajski, *An interpolating dispersionless integrable system*, J. Phys. A **4**, no. 31 (2008), 315202, 9 pp, MR 2425804, Zbl 1234.37044.
- [20] M. Dunajski, L.J. Mason & P. Tod, *Einstein–Weyl geometry, the dKP equation and twistor theory*, J. Geom. Phys. **37**, no. 1–2 (2001), 63–93, MR 1807082, Zbl 0990.53052.
- [21] M. Dunajski, *The twisted photon associated to hyper-Hermitian four-manifolds*, J. Geom. Phys. **30**, no. 3 (1999), 266–281, MR 1692233, Zbl 0959.53022.
- [22] M. Dunajski & P. Tod, *Einstein–Weyl structures from hyper-Kähler metrics with conformal Killing vectors*, Differential Geom. Appl. **14**, no. 1 (2001), 39–55, MR 1811563, Zbl 0997.53036.
- [23] M. Dunajski, *A class of Einstein–Weyl spaces associated to an integrable system of hydrodynamic type*, J. Geom. Phys. **51**, no. 1 (2004), 126–137, MR 2078688, Zbl 1110.53032.
- [24] M.G. Eastwood & K.P. Tod, *Local constraints on Einstein–Weyl geometries: the 3-dimensional case*, Ann. Global Anal. Geom. **18**, no. 1 (2000), 1–27, MR 1739522, Zbl 0957.53015.

- [25] E.V. Ferapontov & M.V. Pavlov, *Hydrodynamic reductions of the heavenly equation*, *Class. Quantum Grav.* **20** (2003), 2429–2441, MR 1984982, Zbl 1029.83022.
- [26] E.V. Ferapontov & K.R. Khusnutdinova, *On the integrability of (2+1)-dimensional quasilinear systems*, *Comm. Math. Phys.* **248** (2004), 187–206, MR 2104610, Zbl 1070.37047.
- [27] E.V. Ferapontov & K.R. Khusnutdinova, *The characterization of 2-component (2+1)-dimensional integrable systems of hydrodynamic type*, *J. Phys. A: Math. Gen.* **37**, no. 8 (2004), 2949–2963, MR 2042985, Zbl 1040.35042.
- [28] E.V. Ferapontov & K.R. Khusnutdinova, *Hydrodynamic reductions of multi-dimensional dispersionless PDEs: the test for integrability*, *J. Math. Phys.* **45**, no. 6 (2004), 2365–2377, MR 2059699, Zbl 1071.35118.
- [29] E.V. Ferapontov, L. Hadjikos & K.R. Khusnutdinova, *Integrable equations of the dispersionless Hirota type and hypersurfaces in the Lagrangian Grassmannian*, *Int. Math. Res. Notices* (2010), no. 3, 496–535; arXiv: 0705.1774 (2007), MR 2587572, Zbl 1203.35215.
- [30] E.V. Ferapontov, K.R. Khusnutdinova & S.P. Tsarev, *On a class of three-dimensional integrable Lagrangians*, *Comm. Math. Phys.* **261**, no. 1 (2006), 225–243, MR 2193210, Zbl 1108.37045.
- [31] E.V. Ferapontov & A.V. Odesskii, *Integrable Lagrangians and modular forms*, *J. Geom. Phys.* **60**, no. 6–8 (2010), 896–906, MR 2647291, Zbl 1188.35161.
- [32] E.V. Ferapontov, B. Huard & A. Zhang, *On the central quadric ansatz: integrable models and Painlevé reductions*, *J. Phys. A: Math. Theor.* **45** (2012), 195–204, MR 2924500, Zbl 1252.35017.
- [33] E.V. Ferapontov, A. Moro & V. V. Sokolov, *Hamiltonian systems of hydrodynamic type in 2+1 dimensions*, *Comm. Math. Phys.* **285**, no. 1 (2009), 31–65, MR 2453590, Zbl 1228.37041.
- [34] J. Gibbons & Y. Kodama, *A method for solving the dispersionless KP hierarchy and its exact solutions. II*, *Phys. Lett. A* **135**, no. 3 (1989), 167–170, MR 0985513.
- [35] J. Gibbons & S.P. Tsarev, *Reductions of the Benney equations*, *Phys. Lett. A* **211** (1996), 19–24, MR 1372470, Zbl 1072.35588; *Conformal maps and reductions of the Benney equations*, *Phys. Lett. A* **258** (1999), 263–270, MR 1710008, Zbl 0936.35184.
- [36] M. Godlinski & P. Nurowski, *On three-dimensional Weyl structures with reduced holonomy*, *Classical Quantum Gravity* **23**, no. 3 (2006), 603–608, MR 2202439, Zbl 1087.53043.
- [37] J.D.E. Grant & I.A.B. Strachan, *Hypercomplex integrable systems*, *Nonlinearity* **12**, no. 5 (1999), 1247–1261, MR 1710101, Zbl 0934.35181.
- [38] A. Grundland & R. Zelazny, *Simple waves in quasilinear hyperbolic systems. I, II. Riemann invariants for the problem of simple wave interactions*, *J. Math. Phys.* **24**, no. 9 (1983), 2305–2328, MR 0715404, Zbl 0555.35080.
- [39] N.J. Hitchin, *Complex manifolds and Einstein's equations*, *Twistor geometry and nonlinear systems* (Primorsko, 1980), 73–99, *Lecture Notes in Math.* **970**, Springer, Berlin/New York (1982), MR 0699802, Zbl 0507.53025.
- [40] V. Husain, *Self-dual gravity as a two-dimensional theory and conservation laws*, *Classical Quantum Gravity* **11**, no. 4 (1994), 927–937, MR 1277228.
- [41] N.Kh. Ibragimov, *Transformation groups applied to mathematical physics* (translated from the Russian) *Mathematics and its Applications* (Soviet Series), D. Reidel Publishing Co., Dordrecht, 1985, MR 0785566, Zbl 0558.53040.

- [42] P.E. Jones & K.P. Tod, *Minitwistor spaces and Einstein–Weyl spaces*, Classical Quantum Gravity **2**, no. 4 (1985), 565–577, MR 0795102, Zbl 0575.53042.
- [43] M. Juras & I. M. Anderson, *Generalized Laplace invariants and the method of Darboux*, Duke Math. J. **89**, no. 2 (1997), 351–375, MR 1460626, Zbl 0885.35075.
- [44] I.S. Krasil’shchik, V.V. Lychagin & A.M. Vinogradov, *Geometry of jet spaces and nonlinear partial differential equations*, Advanced Studies in Contemporary Mathematics, 1. Gordon and Breach Science Publishers, New York, 1986, MR 0861121, Zbl 0722.35001.
- [45] B. Kruglikov, *Laplace transformation of Lie class  $\omega = 1$  overdetermined systems*, J. Nonlin. Math. Phys. **18**, no. 4 (2011), 583–611, MR 2864633, Zbl 1243.35007.
- [46] A. Kushner, *Classification of Monge–Ampère equations*, in B. Kruglikov, V. Lychagin, E. Straume (eds.), *Differential Equations: Geometry, Symmetries and Integrability: The Abel Symposium 2008*, *Abel Symposia* **5**, Springer-Verlag, Berlin (2009), MR 2562576, Zbl 1233.35005.
- [47] C. LeBrun, *Explicit self-dual metrics on  $CP_2 \# \dots \# CP_2$* , J. Differential Geom. **34**, no. 1 (1991), 223–253, MR 1114461, Zbl 0725.53067.
- [48] V.V. Lychagin, V.N. Rubtsov & I.V. Chekalov, *A classification of Monge–Ampère equations*, Ann. Sci. Ecole Norm. Sup. **26**, no. 4 (1993), 281–308, MR 1222276, Zbl 0789.58078.
- [49] V.V. Lychagin & V. Yumaguzhin, *Minkowski metrics on solutions of the Khokhlov–Zabolotskaya equation*, Lobachevskii J. Math. **30**, no. 4 (2009), 333–336, MR 2587855, Zbl 1225.35011.
- [50] A.A. Malykh & M.B. Sheftel, *General heavenly equation governs anti-self-dual gravity*, J. Phys. A **44**, no. 15 (2011), 155201, MR 2783639, Zbl 1215.83026.
- [51] S.V. Manakov & P.M. Santini, *The Cauchy problem on the plane for the dispersionless Kadomtsev–Petviashvili equation*, JETP Lett. **83** (2006), 462–466.
- [52] J.J. Morales Ruiz, *Differential Galois theory and non-integrability of Hamiltonian systems*, Progress in Mathematics **179**, Birkhäuser Verlag, Basel (1999), MR 1713573, Zbl 0934.12003.
- [53] P. Nurowski, *Differential equations and conformal structures*, J. Geom. Phys. **55**, no. 1 (2005), 19–49, MR 2157414, Zbl 1082.53024.
- [54] A. Odesskii & V. Sokolov, *Integrable  $(2+1)$ -dimensional systems of hydrodynamic type*, Theor. Math. Phys. **163**, no. 2 (2010), 549–586, Zbl 06102200.
- [55] P.J. Olver, *Applications of Lie Groups to Differential Equations. Graduate Texts in Mathematics*, **107**, Springer-Verlag, New York, (1986), MR 0836734, Zbl 0588.22001.
- [56] M.V. Pavlov, *New integrable  $(2+1)$ -equations of hydrodynamic type*, Russian Math. Surv., **58**, no. 2 (2003), 384–385, MR 1992575, Zbl 1056.37082; *Classifying integrable Egoroff hydrodynamic chains*, Theor. Math. Phys. **138** (2004), 45–58, MR 2061092, Zbl 1178.37087.
- [57] R. Penrose, *Nonlinear gravitons and curved twistor theory*, General Relativity and Gravitation **7**, no. 1 (1976), 31–52, MR 0439004, Zbl 0354.53025.
- [58] Z. Peradzyński, *Nonlinear plane  $k$ -waves and Riemann invariants*, Bull. Acad. Polon. Sci. Sr. Sci. Tech. **19** (1971), 625–632, MR 0387827, Zbl 0224.35036; *Riemann invariants for the nonplanar  $k$ -waves*, Bull. Acad. Polon. Sci. Sr. Sci. Tech. **19** (1971), 717–724, MR 0387828, Zbl 0244.35059.
- [59] J.F. Plebański, *Some solutions of complex Einstein equations*, J. Math. Phys. **16** (1975), 2395–2402, MR 0386609.

- [60] T. Ruggeri, *Su una naturale estensione a tre variabili dell' equazione di Monge–Ampere*, Accad. Naz. Lincei, LV (1973), 445–449, MR 0380066, Zbl 0294.35013.
- [61] L.P. Safaryan, *Certain classes of manifolds of cones of order two in  $P_n$* , Akad. Nauk Armjan. SSR Dokl. **50** (1970), 83–90, MR 0286004, Zbl 0219.50013.
- [62] A.F. Sidorov, V.P. Shapeev & N.N. Yanenko, *The method of differential constraints and its applications in gas dynamics*, ‘Nauka’, Novosibirsk (1984), MR 0780937, Zbl 0604.76062.
- [63] A.D. Smith, *Integrable  $GL(2)$  geometry and hydrodynamic partial differential equations*, Communications in Analysis and Geometry **18**, no. 4 (2010), 743–790, MR 2765729, Zbl 1239.58020.
- [64] D.C. Spencer, *Overdetermined systems of linear partial differential equations*, Bull. Amer. Math. Soc. **75** (1969), 179–239, MR 0242200, Zbl 0185.33801.
- [65] K.P. Tod, *Scalar-flat Kähler and hyper-Kähler metrics from Painlevé-III*, Class. Quantum Grav. **12** (1995), 1535–1547, MR 1344288, Zbl 0828.53061.
- [66] K.P. Tod, *Einstein–Weyl spaces and third-order differential equations*, J. Math. Phys. **41**, no. 8 (2000), 5572–5581, MR 1770973, Zbl 0979.53050.
- [67] S.P. Tsarev, *Geometry of hamiltonian systems of hydrodynamic type. Generalized hodograph method*, Izvestija AN USSR Math. **54**, no. 5 (1990), 1048–1068, MR 1086085, Zbl 0796.76014.
- [68] R.S. Ward, *Einstein–Weyl spaces and  $SU(\infty)$  Toda fields*, Class. Quantum Grav. **7**, no. 4 (1990), L95–L98, MR 1045295, Zbl 0687.53044.
- [69] R.S. Ward, *Integrable and solvable systems, and relations among them*, Philos. Trans. Roy. Soc. London Ser. A **315**, no. 1533 (1985), 451–457, MR 0836745, Zbl 0579.35078.
- [70] V.E. Zakharov, *Dispersionless limit of integrable systems in 2+1 dimensions*, in N.M. Ercolani et al. (eds.), *Singular Limits of Dispersive Waves*, Plenum Press, NY (1994), 165–174, MR 1321203, Zbl 0852.35130.
- [71] A.V. Zhiber & V.V. Sokolov, *Exactly integrable hyperbolic equations of Liouville type*, Russian Math. Surveys **56**, no. 1 (2001), 61–101, MR 1845643, Zbl 1003.35093.

DEPARTMENT OF MATHEMATICAL SCIENCES  
 LOUGHBOROUGH UNIVERSITY  
 LOUGHBOROUGH, LEICESTERSHIRE LE11 3TU, UNITED KINGDOM  
*E-mail address:* E.V.Ferapontov@lboro.ac.uk

INSTITUTE OF MATHEMATICS AND STATISTICS  
 NT-FACULTY  
 UNIVERSITY OF TROMSØ  
 TROMSØ 90-37, NORWAY  
*E-mail address:* Boris.Kruglikov@uit.no