



TORIC KÄHLER METRICS: COHOMOGENEITY ONE EXAMPLES OF CONSTANT SCALAR CURVATURE IN ACTION-ANGLE COORDINATES

MIGUEL ABREU

Communicated by Izu Vaisman

Abstract. In these notes of the mini-course given by the author at the XI-th Conference on Geometry, Integrability and Quantization, Varna, Bulgaria, 2009 after the introduction of the toric Kähler geometry, we present Calabi's family of $U(n)$ -invariant extremal Kähler metrics in symplectic action-angle coordinates and show that it actually contains, as particular cases, many interesting cohomogeneity one examples of constant scalar curvature.

1. Introduction

In 1982 Calabi [7] constructed, using local complex coordinates, a general four-parameter family of $U(n)$ -invariant extremal Kähler metrics, which he used to put an extremal Kähler metric on

$$H_m^n := \mathbb{P}(\mathcal{O}(-m) \oplus \mathbb{C}) \rightarrow \mathbb{P}^{n-1}$$

for all $n, m \in \mathbb{N}$ and any possible Kähler cohomology class. In particular, when $n = 2$, on all Hirzebruch surfaces.

The main goal of these notes is to present Calabi's general family in local symplectic action-angle coordinates, using the set-up of [1, 2] for toric Kähler geometry, and show that it actually contains other interesting cohomogeneity one Kähler metrics as particular cases (see also [22]). These include:

- the Fubini-Study, flat and Bergman Kähler-Einstein metrics of constant holomorphic sectional curvature (positive, zero and negative, respectively).
- the complete Ricci flat Kähler metric on the total space of

$$\mathcal{O}(-n) \rightarrow \mathbb{P}^{n-1}$$

for all $n \in \mathbb{N}$ and any possible Kähler cohomology class, constructed by Calabi [6] in 1979.

- the complete scalar flat Kähler metric on the total space of

$$\mathcal{O}(-m) \rightarrow \mathbb{P}^{n-1}$$

for all $m, n \in \mathbb{N}$ and any possible Kähler cohomology class, constructed for $n = 2$ by LeBrun [18] in 1988 and for $n > 2$ by Pedersen & Poon [21] in 1991 (see also Simanca [24]).

- the complete Kähler-Einstein metric with negative scalar curvature on the total space of the open disc bundle

$$\mathcal{D}(-m) \subset \mathcal{O}(-m) \longrightarrow \mathbb{P}^{n-1}$$

for all $n < m \in \mathbb{N}$ and any possible Kähler cohomology class, constructed by Pedersen & Poon [21].

- the complete constant negative scalar curvature Kähler metric on the total space of the open disc bundle

$$\mathcal{D}(-m) \subset \mathcal{O}(-m) \longrightarrow \mathbb{P}^{n-1}$$

for all $n, m \in \mathbb{N}$ and any possible Kähler cohomology class, also constructed by Pedersen & Poon [21].

Calabi's general family contains many other interesting cohomogeneity one special Kähler metrics. Besides the Bochner-Kähler orbifold examples presented in [3], it contains for example a family of singular Kähler-Einstein metrics on certain H_m^n that are directly related to the Sasaki-Einstein metrics constructed by Gauntlett-Martelli-Sparks-Waldram [13, 14] in 2004 - see [4].

These notes are organized as follows. In Section 2 we give a basic introduction to symplectic geometry and discuss some fundamental features of toric symplectic manifolds. Section 3 is devoted to the toric Kähler metrics. After some relevant linear algebra preliminaries, we explain how these can be parametrized in action-angle coordinates via symplectic potentials for the associated toric compatible complex structures and discuss some important properties of these symplectic potentials. In Section 4 we write down the symplectic potentials that give rise to the toric constant (scalar) curvature metrics on real two dimensional manifold and identify the underlying toric symplectic surfaces. This is a warm-up for Section 5 where we discuss in detail the above higher dimensional examples that arise in Calabi's general family of local $U(n)$ -invariant extremal Kähler metrics.

2. Toric Symplectic Manifolds

In this section we give a basic introduction to symplectic geometry and discuss some fundamental features of toric symplectic manifolds.

2.1. Symplectic Manifolds

Definition 1. A **symplectic manifold** is a pair (B, ω) where B is a smooth manifold and ω is a closed and non-degenerate two-form, i.e.,

- i) $\omega \in \Omega^2(B)$ is such that $d\omega = 0$ and
- ii) for any $p \in B$ and $0 \neq X \in T_p B$, there exists $Y \in T_p B$ such that $\omega_p(X, Y) \neq 0$.

The non-degeneracy condition ii) implies that a symplectic manifold is always even dimensional. If B has dimension $2n$, the non-degeneracy condition ii) is equivalent to requiring that

$$\omega^n \equiv \omega \wedge \cdots \wedge \omega \in \Omega^{2n}(B) \text{ is a volume form.}$$

Hence, a symplectic manifold (B, ω) is always oriented.

Example 2. The most basic example is \mathbb{R}^{2n} with linear coordinates $(u_1, \dots, u_n, v_1, \dots, v_n)$ and symplectic form

$$\omega_{\text{st}} = du \wedge dv := \sum_{j=1}^n du_j \wedge dv_j.$$

Example 3. Any two-dimensional surface equipped with an area form is a symplectic manifold. For example, the sphere S^2 or any other compact orientable surface Σ_g of genus g .

Example 4. If (B_1, ω_1) and (B_2, ω_2) are symplectic manifolds, then

$$(B = B_1 \times B_2, \quad \omega = \omega_1 \times \omega_2)$$

is also a symplectic manifold. Here, $\omega_1 \times \omega_2$ means the sum of the pullbacks of the symplectic forms ω_1 and ω_2 from the factors B_1 and B_2 .

Example 5. *The imaginary part of the Hermitian metric on any Kähler manifold is a symplectic form. Hence, any Kähler manifold is a symplectic manifold. In particular, the complex projective space \mathbb{P}^n equipped with its Fubini-Study form ω_{FS} is a symplectic manifold.*

When (B, ω) is a compact symplectic manifold we have that

$$\omega^n = \text{volume form} \Rightarrow 0 \neq [\omega^n] \in H^{2n}(B, \mathbb{R}) \Rightarrow 0 \neq [\omega] \in H^2(B, \mathbb{R}).$$

In particular, the spheres \mathbb{S}^{2n} have no symplectic form when $n > 1$, since

$$H^2(\mathbb{S}^{2n}, \mathbb{R}) = 0 \quad \text{when } n > 1.$$

2.2. Symplectomorphisms and Darboux's Theorem

Definition 6. *Let (B, ω) be a symplectic manifold. A **symplectomorphism** of B is a diffeomorphism $\varphi : B \rightarrow B$ such that $\varphi^*(\omega) = \omega$. These form the **symplectomorphism group**, a subgroup of $\text{Diff}(B)$ that will be denoted by $\text{Diff}(B, \omega)$.*

Example 7. *Consider the symplectic manifold (B, ω) and let $h : B \rightarrow \mathbb{R}$ be a smooth function on B . The non-degeneracy of ω implies that there exists a unique vector field $X_h \in \mathcal{X}(B)$ such that $X_h \lrcorner \omega = dh$. This vector field X_h is called the **Hamiltonian vector field** of the function h and has the following fundamental property:*

the flow $\varphi_t \equiv (X_h)_t : B \rightarrow B$ consists of symplectomorphisms of B .

This can be proved by using Cartan's formula to compute

$$\mathcal{L}_{X_h} \omega = X_h \lrcorner d\omega + d(X_h \lrcorner \omega) = X_h \lrcorner 0 + d(dh) = 0.$$

Hence, on a symplectic manifold (B, ω) any smooth function $h \in C^\infty(B)$ gives rise, through the flow of the corresponding Hamiltonian vector field $X_h \in \mathcal{X}(B)$, to a local one-parameter group of symplectomorphisms.

One can use the symplectomorphisms constructed in the previous example to prove that

- i) the symplectomorphism group $\text{Diff}(B, \omega)$ is always infinite-dimensional
- ii) the action of $\text{Diff}(B, \omega)$ on the manifold B is always k -transitive, for any $k \in \mathbb{N}$

iii) in particular, any point of a symplectic manifold (B, ω) looks locally like any other point of (B, ω) .

This last statement is made more precise in the following

Theorem 8 (Darboux) *Let (B, ω) be a symplectic manifold of dimension $2n$. Then, any point $p \in B$ has a neighborhood $U \subset B$ symplectomorphic to a neighborhood V of the origin in $(\mathbb{R}^{2n}, \omega_{\text{st}})$, i.e., there exists a diffeomorphism*

$$\phi : U \subset B \rightarrow V \subset \mathbb{R}^{2n} \text{ such that } \phi(p) = 0 \text{ and } \phi^*(\omega_{\text{st}}) = \omega.$$

In other words,

there are no local invariants in symplectic geometry

which is in sharp contrast with what happens, for example, in Riemannian geometry.

2.3. Symplectic and Hamiltonian Vector Fields

The Lie algebra of the symplectomorphism group $\text{Diff}(B, \omega)$, viewed as an infinite-dimensional Lie group, is naturally identified with the vector space $\mathcal{X}(B, \omega)$ of *symplectic vector fields*, i.e., vector fields $X \in \mathcal{X}(B)$ such that $\mathcal{L}_X \omega = 0$, with Lie bracket $[\cdot, \cdot]$ given by the usual Lie bracket of vector fields. As before, we can use Cartan's formula to obtain

$$\mathcal{L}_X \omega = X \lrcorner d\omega + d(X \lrcorner \omega) = X \lrcorner 0 + d(X \lrcorner \omega) = d(X \lrcorner \omega).$$

Hence, the vector space of symplectic vector fields is given by

$$\mathcal{X}(B, \omega) = \{X \in \mathcal{X}(B); \text{ the one-form } X \lrcorner \omega \text{ is closed}\}$$

while its subspace of *Hamiltonian vector fields* is given by

$$\mathcal{X}_H(B, \omega) = \{X \in \mathcal{X}(B); \text{ the one-form } X \lrcorner \omega \text{ is exact}\}.$$

In fact, as the following theorem shows, $\mathcal{X}_H(B, \omega)$ is a Lie subalgebra of $\mathcal{X}(B, \omega)$.

Theorem 9. *If $X, Y \in \mathcal{X}(B, \omega)$ are symplectic vector fields, then $[X, Y]$ is the Hamiltonian vector field of the function $\omega(Y, X) : B \rightarrow \mathbb{R}$, i.e.,*

$$[X, Y] = X_{\omega(Y, X)} \in \mathcal{X}_H(B, \omega).$$

Proof: It suffices to compute $[X, Y] \lrcorner \omega$, using the standard formulas from differential geometry and the defining properties of X, Y and ω

$$\begin{aligned} [X, Y] \lrcorner \omega &= \mathcal{L}_X(Y \lrcorner \omega) - Y \lrcorner (\mathcal{L}_X \omega) \\ &= d(X \lrcorner (Y \lrcorner \omega)) + X \lrcorner (d(Y \lrcorner \omega)) - Y \lrcorner (d(X \lrcorner \omega)) - Y \lrcorner (X \lrcorner d\omega) \\ &= d(\omega(Y, X)). \quad \blacksquare \end{aligned}$$

Remark 10. $\mathcal{X}_H(B, \omega)$ is the Lie algebra of the fundamental subgroup of the symplectomorphism group, namely the subgroup $\text{Ham}(B, \omega) \subset \text{Diff}(B, \omega)$ of Hamiltonian symplectomorphisms of (B, ω) . It follows from Theorem 9 that this Lie algebra can be naturally identified with the vector space $C^\infty(B)/\mathbb{R}$, i.e., the smooth functions on B modulo constants, equipped with a bracket $\{\cdot, \cdot\}$ known as the Poisson bracket

$$\{f, g\} \equiv \omega(X_f, X_g).$$

Note also that when $H^1(B, \mathbb{R}) = 0$ we have that $\mathcal{X}_H(B, \omega) = \mathcal{X}(B, \omega)$.

2.4. Hamiltonian Torus Actions

Let (B, ω) be a symplectic manifold equipped with a symplectic action of

$$\mathbb{T}^m \equiv \mathbb{R}^m / 2\pi\mathbb{Z}^m \equiv \mathbb{R} / 2\pi\mathbb{Z} \times \cdots \times \mathbb{R} / 2\pi\mathbb{Z} \equiv \mathbb{S}^1 \times \cdots \times \mathbb{S}^1$$

i.e., with a homomorphism $\mathbb{T}^m \rightarrow \text{Diff}(B, \omega)$. Let $X_1, \dots, X_m \in \mathcal{X}(B)$ be the vector fields generating the action of each individual \mathbb{S}^1 -factor. Then, since the action is symplectic, we have that

$$\mathcal{L}_{X_k} \omega = 0 \Leftrightarrow X_k \lrcorner d\omega + d(X_k \lrcorner \omega) = 0 \Leftrightarrow d(X_k \lrcorner \omega) = 0$$

i.e.,

$$X_k \in \mathcal{X}(B, \omega) \quad \text{for all } k \in \{1, \dots, m\}.$$

Definition 11. A symplectic \mathbb{T}^m -action on a symplectic manifold (B, ω) is said to be **Hamiltonian** if for every $k \in \{1, \dots, m\}$ there exists a function $h_k : B \rightarrow \mathbb{R}$ such that $X_k \lrcorner \omega = dh_k$, i.e., $X_k \equiv X_{h_k} \in \mathcal{X}_H(B, \omega)$ is the Hamiltonian vector field of h_k . In this case, the map $\mu : B \rightarrow \mathbb{R}^m$ defined by

$$\mu(p) = (h_1(p), \dots, h_m(p)) \quad \text{for all } p \in B$$

which will be called a **moment map** for the action.

Remark 12. Suppose $\mu : B \rightarrow \mathbb{R}^m$ is a moment map for a Hamiltonian \mathbb{T}^m -action on (B, ω) . Then $\mu + c$, for any given constant $c \in \mathbb{R}^m$, is also a moment map for that same action.

Remark 13. The orbits of a Hamiltonian \mathbb{T}^m -action on a symplectic manifold (B, ω) are always **isotropic**, i.e.,

$$\omega|_{\text{orbit}} \equiv 0.$$

In fact, the tangent space to an orbit is generated by the Hamiltonian vector fields X_{h_k} , $k \in \{1, \dots, m\}$. Using Theorem 9 and the fact that the torus \mathbb{T}^m is abelian, we have that

$$X_{\omega(X_{h_k}, X_{h_l})} = -[X_{h_k}, X_{h_l}] \equiv 0 \Rightarrow \omega(X_{h_k}, X_{h_l}) \equiv \text{const}, \quad k, l \in \{1, \dots, m\}.$$

Since \mathbb{T}^m is compact, there is for each $k \in \{1, \dots, m\}$ and on each \mathbb{T}^m -orbit a point p_k where the function $h_k|_{\text{orbit}}$ attains its maximum. Then

$$\omega(X_{h_k}, X_{h_l}) = (dh_k)_{p_k}(X_{h_l}) = 0.$$

Hence, the above constant is actually zero and each \mathbb{T}^m -orbit is indeed isotropic. This fact will be used below, in the proof of Proposition 17.

Example 14. Consider $(\mathbb{R}^{2n}, \omega_{\text{st}})$, where

$$\omega_{\text{st}} = du \wedge dv := \sum_{j=1}^n du_j \wedge dv_j$$

as in Example 2, and its usual identification with \mathbb{C}^n given by

$$z_j = u_j + iv_j, \quad j = 1, \dots, n. \quad (1)$$

The standard \mathbb{T}^n -action τ_{st} on \mathbb{R}^{2n} given by

$$(y_1, \dots, y_n) \cdot (z_1, \dots, z_n) = (e^{-iy_1} z_1, \dots, e^{-iy_n} z_n)$$

is Hamiltonian with a moment map $\mu_{\text{st}} : \mathbb{R}^{2n} \rightarrow \mathbb{R}^n$ given by

$$\mu_{\text{st}}(u_1, \dots, u_n, v_1, \dots, v_n) = \frac{1}{2}(u_1^2 + v_1^2, \dots, u_n^2 + v_n^2).$$

Example 15. Consider the projective space $(\mathbb{P}^n, \omega_{\text{FS}})$ equipped with the homogeneous coordinates $[z_0; z_1; \dots; z_n]$.

The \mathbb{T}^n -action τ_{FS} on \mathbb{P}^n given by

$$(y_1, \dots, y_n) \cdot [z_0; z_1; \dots; z_n] = [z_0; e^{-iy_1} z_1; \dots; e^{-iy_n} z_n]$$

is Hamiltonian, with a moment map $\mu_{\text{FS}} : \mathbb{P}^n \rightarrow \mathbb{R}^n$ given by

$$\mu_{\text{FS}}[z_0; z_1; \dots; z_n] = \frac{1}{\|z\|^2} (\|z_1\|^2, \dots, \|z_n\|^2).$$

Note that the image of μ_{FS} is the convex hull of the images of the $n+1$ fixed points of the action, i.e., the standard simplex in \mathbb{R}^n .

Atiyah [5] and Guillemin-Sternberg [16] proved in 1982 the following Convexity Theorem.

Theorem 16. Let (B, ω) be a compact, connected, symplectic manifold, equipped with a Hamiltonian \mathbb{T}^m -action with a moment map $\mu : B \rightarrow \mathbb{R}^m$. Then

- i) the level sets $\mu^{-1}(\lambda)$ of the moment map are connected (for any $\lambda \in \mathbb{R}^m$)
- ii) the image $\mu(B) \subset \mathbb{R}^m$ of the moment map is the convex hull of the images of the fixed points of the action.

2.5. Toric Symplectic Manifolds

The following proposition motivates the definition of a toric symplectic manifold.

Proposition 17. If a symplectic manifold (B, ω) has an effective Hamiltonian \mathbb{T}^m -action, then $m \leq (\dim B)/2$.

Proof:

$$\begin{aligned} \text{Effective action} &\Rightarrow \text{there exist } m\text{-dimensional orbits} \\ \text{Hamiltonian } \mathbb{T}^m\text{-action} &\Rightarrow \text{orbits are isotropic (see Remark 13)} \\ \text{Linear Algebra} &\Rightarrow \dim(\text{isotropic orbit}) \leq \frac{1}{2} \dim B. \end{aligned}$$

■

Definition 18. A toric symplectic manifold is a connected symplectic manifold (B^{2n}, ω) , equipped with an effective Hamiltonian action of the n -torus

$$\tau : \mathbb{T}^n \cong \mathbb{R}^n / 2\pi\mathbb{Z}^n \hookrightarrow \text{Ham}(B, \omega)$$

such that the corresponding moment map

$$\mu : B \rightarrow \mathbb{R}^n$$

which is well-defined up to a constant, is proper onto its convex image $P = \mu(B) \subset \mathbb{R}^n$.

Remark 19. *The requirement that the moment map be “proper onto its convex image”, something that is automatic for compact manifolds, makes the theory of non-compact toric symplectic manifolds analogous to the compact one (see [17]).*

Example 20. $(\mathbb{R}^{2n}, \omega_{\text{st}})$, equipped with the standard Hamiltonian \mathbb{T}^n -action described in Example 14, is a non-compact toric symplectic manifold.

Example 21. $(\mathbb{P}^n, \omega_{\text{FS}})$, equipped with the Hamiltonian \mathbb{T}^n -action described in Example 15, is a compact toric symplectic manifold.

2.6. Classification Theorem and Action-Angle Coordinates

Any toric symplectic manifold has an associated convex set, the image of the moment map of the torus action. The convex sets that arise in this way are characterized in the following definition.

Definition 22. *A convex polyhedral set P in \mathbb{R}^n is called simple and integral if*

- 1) *there are n edges meeting at each vertex p*
- 2) *the edges meeting at the vertex p are rational, i.e., each edge is of the form $p + tv_i$, $0 \leq t \leq \infty$, where $v_i \in \mathbb{Z}^n$*
- 3) *the v_1, \dots, v_n in (2) can be chosen to be a \mathbb{Z} -basis of the lattice \mathbb{Z}^n .*

A facet is a face of P of codimension one.

A Delzant set is a simple and integral convex polyhedral set $P \subset \mathbb{R}^n$. A Delzant polytope is a compact Delzant set.

*Two Delzant sets are termed to be **isomorphic** if one can be mapped to the other by a translation.*

In 1988 Delzant [10] showed that any Delzant polytope determines a unique compact toric symplectic manifold. More precisely, if two compact toric symplectic

manifolds have the same Delzant polytope, then there exists an equivariant symplectomorphism between them. This result can be generalized to the possibly non-compact setting of Definition 18 (see [17]).

Theorem 23. *Let (B, ω, τ) be a toric symplectic manifold, with a moment map $\mu : B \rightarrow \mathbb{R}^n$. Then $P \equiv \mu(B)$ is a Delzant set.*

Two toric symplectic manifolds are equivariant symplectomorphic (with respect to a fixed torus acting on both) if and only if their associated Delzant sets are isomorphic. Moreover, every Delzant set arises from some toric symplectic manifold.

Remark 24. *One can use the results of Lerman and Tolman [19] to generalize Theorem 23 to orbifolds. The outcome is a classification of symplectic toric orbifolds via **labeled Delzant sets**, i.e., convex polyhedral sets, as in Definition 22, with “ \mathbb{Z} -basis” in (3) replaced by “ \mathbb{Q} -basis” and with a positive integer label attached to each facet.*

Each facet F of a labeled Delzant set $P \subset \mathbb{R}^n$ determines a unique lattice vector $\nu_F \in \mathbb{Z}^n$: the primitive inward pointing normal lattice vector. A convenient way of thinking about a positive integer label $m_F \in \mathbb{N}$ associated to F is by dropping the primitive requirement from this lattice vector: consider $m_F \nu_F$ instead of ν_F .

In other words, a labeled Delzant set can be defined as a rational simple polyhedral set $P \subset \mathbb{R}^n$ with an inward pointing normal lattice vector associated to each of its facets.

The proof gives an explicit symplectic reduction construction of a canonical model for each toric symplectic manifold, i.e., it associates to each Delzant set P an explicit toric symplectic manifold (B_P, ω_P, τ_P) with moment map $\mu_P : B_P \rightarrow P$. One can use these canonical models to derive general properties of toric symplectic manifolds. For example, let \check{P} denote the interior of P , and consider $\check{B}_P \subset B_P$ defined by $\check{B}_P = \mu_P^{-1}(\check{P})$. One easily checks that \check{B}_P is an open dense subset of B_P , consisting of all points where the \mathbb{T}^n -action is free. It can be described as

$$\check{B}_P \cong \check{P} \times \mathbb{T}^n = \left\{ (x, y); x \in \check{P} \subset \mathbb{R}^n, y \in \mathbb{R}^n / 2\pi\mathbb{Z}^n \right\}$$

where (x, y) are symplectic **action-angle** coordinates for ω_P , i.e.,

$$\omega_P|_{\check{B}} = dx \wedge dy = \sum_{j=1}^n dx_j \wedge dy_j.$$

Hence, one has a global equivariant Darboux’s Theorem in the toric context. Note that in these action-angle coordinates the moment map is given simply by

$$\mu_P(x, y) = x$$

i.e., projection in the action coordinates.

3. Toric Kähler Metrics

3.1. Linear Compatible Complex Structures

Definition 25. A compatible complex structure on the symplectic vector space (V, ω) is a complex structure J on V , i.e., $J \in \text{End}(V)$ with $J^2 = -\text{Id}$, such that

$$\langle \cdot, \cdot \rangle_J := \omega(\cdot, J\cdot)$$

is an inner product on V . This is equivalent to

$$\omega(J\cdot, J\cdot) = \omega(\cdot, \cdot) \quad \text{and} \quad \omega(v, Jv) > 0, \quad \text{for all } v \neq 0 \in V.$$

The set of all compatible complex structures on a symplectic vector space (V, ω) will be denoted by $\mathcal{J}(V, \omega)$.

The symplectic linear group $\text{Sp}(V, \omega)$ acts on $\mathcal{J}(V, \omega)$ by conjugation

$$\begin{aligned} \text{Sp}(V, \omega) \times \mathcal{J}(V, \omega) &\rightarrow \mathcal{J}(V, \omega) \\ (\Phi, J) &\mapsto \Phi J \Phi^{-1}. \end{aligned}$$

This action can be easily seen to be transitive and, if we fix $J_0 \in \mathcal{J}(V, \omega)$ and corresponding inner product $\langle \cdot, \cdot \rangle_0$, we have that

$$\mathcal{J}(V, \omega) = \text{Sp}(V, \omega) / \text{U}(V, \omega, \langle \cdot, \cdot \rangle_0)$$

where $\text{U}(V, \omega, \langle \cdot, \cdot \rangle_0) = \text{Sp}(V, \omega) \cap \text{O}(V, \langle \cdot, \cdot \rangle_0)$ is the unitary group. $\mathcal{J}(V, \omega)$ is a symmetric space and admits a beautiful explicit description due to Siegel [23].

Definition 26. The Siegel upper half space \mathcal{S}_n is the open contractible subset of the complex vector space of complex symmetric matrices defined by

$$\begin{aligned} \mathcal{S}_n := \{ Z = R + iS; R \text{ and } S \text{ are real symmetric } (n \times n) \text{ matrices} \\ \text{and } S \text{ is positive definite} \}. \end{aligned}$$

Choose a symplectic basis for (V, ω) , i.e., an isomorphism $(V, \omega) \cong (\mathbb{R}^{2n}, \omega_{\text{st}})$, and identify the symplectic linear group $\text{Sp}(V, \omega)$ with the matrix group $\text{Sp}(2n, \mathbb{R})$ consisting of $(2n \times 2n)$ real matrices Φ such that

$$\Phi^t \cdot \omega_0 \cdot \Phi = \omega_0$$

where

$$\omega_0 = \begin{bmatrix} 0 & \vdots & \text{Id} \\ \dots\dots\dots & & \\ -\text{Id} & \vdots & 0 \end{bmatrix}$$

is the matrix form of ω_{st} written in $(n \times n)$ blocks. We will also write any $\Phi \in \text{Sp}(2n, \mathbb{R})$ in $(n \times n)$ blocks

$$\Phi = \begin{bmatrix} A & \vdots & B \\ \dots\dots\dots & & \\ C & \vdots & D \end{bmatrix} .$$

The following proposition is proved in [23].

Proposition 27. $\text{Sp}(2n, \mathbb{R})$ acts on \mathcal{S}_n by linear fractional transformations

$$\begin{aligned} \text{Sp}(2n, \mathbb{R}) \times \mathcal{S}_n &\rightarrow \mathcal{S}_n \\ (\Phi, Z) &\mapsto \Phi(Z) := (AZ + B) \cdot (CZ + D)^{-1} . \end{aligned}$$

This action is transitive and the isotropy group of $i\text{Id} \in \mathcal{S}_n$ is $\text{U}(n) \subset \text{Sp}(2n, \mathbb{R})$:

$$\text{U}(n) = \text{Sp}(2n, \mathbb{R}) \cap \text{O}(2n) = \{ \Phi \in \text{Sp}(2n, \mathbb{R}) ; \Phi^t \cdot \Phi = \text{Id} \} .$$

Hence,

$$\mathcal{S}_n \cong \text{Sp}(2n, \mathbb{R}) / \text{U}(n) .$$

Given $Z = R + iS \in \mathcal{S}_n$, define

$$\Phi_Z := \begin{bmatrix} S^{1/2} & \vdots & RS^{-1/2} \\ \dots\dots\dots & & \\ 0 & \vdots & S^{-1/2} \end{bmatrix} \in \text{Sp}(2n, \mathbb{R}) .$$

Under the action of $\text{Sp}(2n, \mathbb{R})$ on \mathcal{S}_n , we have that

$$\Phi_Z(i\text{Id}) = Z .$$

Let $J_0 \in \mathcal{J}(\mathbb{R}^{2n}, \omega_{\text{st}})$ be given by

$$J_0 = \begin{bmatrix} 0 & \vdots & -\text{Id} \\ \dots\dots\dots & & \\ \text{Id} & \vdots & 0 \end{bmatrix} . \tag{2}$$

For each $Z \in \mathcal{S}_n$, define $J_Z \in \mathcal{J}(\mathbb{R}^{2n}, \omega_{\text{st}})$ by

$$J_Z := (J_0 \cdot \Phi_Z) \cdot J_0 \cdot (J_0 \cdot \Phi_Z)^{-1} = \begin{bmatrix} -S^{-1}R & \vdots & -S^{-1} \\ \dots\dots\dots & & \\ RS^{-1}R + S & \vdots & RS^{-1} \end{bmatrix}.$$

This defines a bijection

$$\begin{aligned} \mathcal{S}_n &\rightarrow \mathcal{J}(\mathbb{R}^{2n}, \omega_{\text{st}}) \\ Z &\mapsto J_Z \end{aligned}$$

which, up to J_0 -conjugation, is equivariant with respect to the $\text{Sp}(2n, \mathbb{R})$ -action on both spaces. More precisely, if $\Phi \in \text{Sp}(2n, \mathbb{R})$ then

$$\Phi \cdot J_Z \cdot \Phi^{-1} = J_{Z'} \Leftrightarrow Z' = (J_0^{-1} \cdot \Phi \cdot J_0)(Z).$$

In particular, $\mathcal{J}(\mathbb{R}^{2n}, \omega_{\text{st}})$ is a contractible space and, for any symmetric $(n \times n)$ real matrix U , we have that

$$\Phi = \begin{bmatrix} I & \vdots & 0 \\ \dots\dots\dots & & \\ U & \vdots & I \end{bmatrix} \in \text{Sp}(2n, \mathbb{R}) \quad \text{and} \quad \Phi \cdot J_Z \cdot \Phi^{-1} = J_{(Z-U)}. \quad (3)$$

This will be relevant below.

3.2. Toric Compatible Complex Structures

Definition 28. A compatible almost complex structure on a symplectic manifold (B, ω) is an almost complex structure J on B , i.e., $J \in \Gamma(\text{End}(TB))$ with $J^2 = -\text{Id}$, such that

$$\langle \cdot, \cdot \rangle_J := \omega(\cdot, J\cdot)$$

is a Riemannian metric on B . This is equivalent to $\omega(J\cdot, J\cdot) = \omega(\cdot, \cdot)$ and $\omega(X, JX) > 0$, for all $X \neq 0, X \in TB$.

The space of all compatible almost complex structures on a symplectic manifold (B, ω) will be denoted by $\mathcal{J}(B, \omega)$.

Remark 29.

- i) The fact that $\mathcal{J}(\mathbb{R}^{2n}, \omega_{\text{st}})$ is contractible implies that $\mathcal{J}(B, \omega)$ is a non-empty, infinite-dimensional and contractible, for any symplectic manifold (B, ω) .

- ii) A Kähler manifold is a symplectic manifold (B, ω) with an integrable compatible complex structure J , i.e., one that is locally isomorphic to the standard complex structure J_0 on \mathbb{R}^{2n} . Note that (1) gives the standard isomorphism $(\mathbb{R}^{2n}, J_0) \cong \mathbb{C}^n$.
- iii) The space of integrable compatible complex structures on the symplectic manifold (B, ω) will be denoted by $\mathcal{I}(B, \omega) \subset \mathcal{J}(B, \omega)$.
- iv) In general, $\mathcal{I}(B, \omega)$ can be empty.

Definition 30. A toric compatible complex structure on a toric symplectic manifold (B^{2n}, ω, τ) is a \mathbb{T}^n -invariant $J \in \mathcal{I}(B, \omega) \subset \mathcal{J}(B, \omega)$. The space of all such structures will be denoted by $\mathcal{I}^{\mathbb{T}^n}(B, \omega) \subset \mathcal{J}^{\mathbb{T}^n}(B, \omega)$.

Remark 31. It follows from the classification in Theorem 23, more precisely from the explicit symplectic reduction construction of the canonical model for any compact toric symplectic manifold (B^{2n}, ω, τ) , that $\mathcal{I}^{\mathbb{T}^n}(B, \omega)$ is always non-empty.

3.3. Local Form of Toric Compatible Complex Structures

It follows from the above bijection between $\mathcal{J}(\mathbb{R}^{2n}, \omega_{\text{st}})$ and the Siegel upper half space \mathcal{S}_n that any $J \in \mathcal{J}^{\mathbb{T}^n}(\check{B}, \omega|_{\check{B}})$ can be written in action-angle coordinates (x, y) on $\check{B} \cong \check{P} \times \mathbb{T}^n$ as

$$J = \begin{bmatrix} -S^{-1}R & \vdots & -S^{-1} \\ \dots\dots\dots & & \\ RS^{-1}R + S & \vdots & RS^{-1} \end{bmatrix}$$

where $R = R(x)$ and $S = S(x)$ are real symmetric $(n \times n)$ matrices, with S positive definite.

For *integrable* toric compatible complex structures we have that

$$J \in \mathcal{I}^{\mathbb{T}^n} \subset \mathcal{J}^{\mathbb{T}^n} \Leftrightarrow \frac{\partial Z_{ij}}{\partial x_k} = \frac{\partial Z_{ik}}{\partial x_j}$$

and there exists $f : \check{P} \rightarrow \mathbb{C}$, $f(x) = r(x) + \text{is}(x)$, such that

$$Z_{ij} = \frac{\partial^2 f}{\partial x_i \partial x_j} = \frac{\partial^2 r}{\partial x_i \partial x_j} + \text{i} \frac{\partial^2 s}{\partial x_i \partial x_j} = R_{ij} + \text{i}S_{ij}.$$

Any real function $h : \check{P} \rightarrow \mathbb{R}$ is the Hamiltonian of one-parameter family

$$\phi_t : \check{B} \rightarrow \check{B}$$

of \mathbb{T}^n -equivariant symplectomorphisms. These are given in action-angle coordinates (x, y) on $\check{B} \cong \check{P} \times \mathbb{T}^n$ by

$$\phi_t(x, y) = \left(x, y - t \frac{\partial h}{\partial x}\right).$$

Hence, it follows from (3) that the natural action of ϕ_t on $\mathcal{J}^{\mathbb{T}^n}$, given by

$$\phi_t \cdot J = (d\phi_t) \circ J \circ (d\phi_t)^{-1}$$

corresponds in the Siegel upper half space parametrization to

$$\phi_t \cdot (Z = R + iS) = (R + tH) + iS$$

where

$$H = (h_{ij}) = \left(\frac{\partial^2 h}{\partial x_i \partial x_j} \right).$$

This implies that, for any integrable $J \in \mathcal{I}^{\mathbb{T}^n}$, there exist action-angle coordinates (x, y) on $\check{B} \cong \check{P} \times \mathbb{T}^n$ such that $R \equiv 0$, i.e.,

$$J = \begin{bmatrix} 0 & \vdots & -S^{-1} \\ \dots\dots\dots & & \\ S & \vdots & 0 \end{bmatrix}$$

with

$$S = S(x) = (s_{ij}(x)) = \left(\frac{\partial^2 s}{\partial x_i \partial x_j} \right)$$

for some real potential function $s : \check{P} \rightarrow \mathbb{R}$.

The holomorphic coordinates for J are given in this case by

$$z(x, y) = u(x, y) + iv(x, y) = \frac{\partial s}{\partial x}(x) + iy \tag{4}$$

and the corresponding Riemannian (Kähler) metric

$$\langle \cdot, \cdot \rangle_J := \omega(\cdot, J\cdot)$$

on $\check{B} \cong \check{P} \times \mathbb{T}^n$ can be written in matrix form as

$$\omega_0 \cdot J = \begin{bmatrix} 0 & \vdots & \text{Id} \\ \dots\dots\dots & & \\ -\text{Id} & \vdots & 0 \end{bmatrix} \cdot \begin{bmatrix} 0 & \vdots & -S^{-1} \\ \dots\dots\dots & & \\ S & \vdots & 0 \end{bmatrix} = \begin{bmatrix} S & \vdots & 0 \\ \dots\dots\dots & & \\ 0 & \vdots & S^{-1} \end{bmatrix} \quad (5)$$

with

$$S = \left(\frac{\partial^2 s}{\partial x_i \partial x_j} \right).$$

Definition 32. We will call such a potential function

$$s : \check{P} \rightarrow \mathbb{R}$$

the **symplectic potential** of both the complex structure J and the metric $\langle \cdot, \cdot \rangle_J$.

Remark 33. This particular way to arrive at the above local form for any compatible structure $J \in \mathcal{I}^{\mathbb{T}^n}$ is due to Donaldson [12], and illustrates a small part of his formal general framework for the action of the symplectomorphism group of a symplectic manifold on its space of compatible complex structures (cf. [11]).

Example 34. Consider the standard linear complex structure $J_0 \in \mathcal{I}^{\mathbb{T}^n}(\mathbb{R}^{2n}, \omega_{\text{st}})$ given by (2). In action-angle coordinates (x, y) on

$$\check{\mathbb{R}}^{2n} = (\mathbb{R}^2 \setminus \{(0, 0)\})^n \cong (\mathbb{R}^+)^n \times \mathbb{T}^n = \check{P} \times \mathbb{T}^n$$

its symplectic potential is given by

$$s : \check{P} = (\mathbb{R}^+)^n \longrightarrow \mathbb{R}$$

$$x = (x_1, \dots, x_n) \longmapsto s(x) = \frac{1}{2} \sum_{i=1}^n x_i \log(x_i).$$

Hence, in these action-angle coordinates, the standard complex structure has the matrix form

$$J_0 = \begin{bmatrix} 0 & \vdots & \text{diag}(-2x_i) \\ \dots\dots\dots & & \\ \text{diag}(1/2x_i) & \vdots & 0 \end{bmatrix}$$

while the standard flat Euclidean metric becomes

$$\begin{bmatrix} \text{diag}(1/2x_i) & \vdots & 0 \\ \dots\dots\dots & & \\ 0 & \vdots & \text{diag}(2x_i) \end{bmatrix}.$$

3.4. Symplectic Potentials for Compact Toric Symplectic Manifolds

The proof of Theorem 23 associates to each Delzant set $P \subset \mathbb{R}^n$, via an explicit symplectic reduction construction, a canonical Kähler toric manifold

$$(B_P^{2n}, \omega_P, \tau_P, \mu_P, J_P) \quad \text{such that} \quad \mu_P(B_P) = P.$$

In [15] Guillemin gave an explicit formula for the symplectic potential of this canonical Kähler metric. To write it down one needs some simple combinatorial data that can be easily obtained directly from the polytope P .

Let F_i denote the i -th facet (codimension-one face) of the polytope P . The affine defining function of F_i is the function

$$\begin{aligned} \ell_i : \mathbb{R}^n &\longrightarrow \mathbb{R} \\ x &\longmapsto \ell_i(x) = \langle x, \nu_i \rangle - \lambda_i \end{aligned}$$

where $\nu_i \in \mathbb{Z}^n$ is a primitive inward pointing normal to F_i and $\lambda_i \in \mathbb{R}$ is such that $\ell_i|_{F_i} \equiv 0$. Note that $\ell_i|_{\check{P}} > 0$.

Theorem 35. *In appropriate action-angle coordinates (x, y) , the canonical symplectic potential $s_P : \check{P} \rightarrow \mathbb{R}$ for $J_P|_{\check{P}}$ is given by*

$$s_P(x) = \frac{1}{2} \sum_{i=1}^d \ell_i(x) \log \ell_i(x)$$

where d is the number of the facets of P .

Example 36. *The symplectic potential presented in Example 34 for the standard flat Euclidean metric on \mathbb{R}^{2n} is the canonical symplectic potential of the corresponding Delzant set $P = (\mathbb{R}_0^+)^n \subset \mathbb{R}^n$.*

Example 37. *For projective space \mathbb{P}^n the polytope $P \subset \mathbb{R}^n$ can be taken to be the standard simplex, with defining affine functions*

$$\ell_i(x) = x_i, \quad i = 1, \dots, n, \quad \text{and} \quad \ell_{n+1}(x) = 1 - r$$

where $r = \sum_i x_i$.

The canonical symplectic potential $s_P : \check{P} \rightarrow \mathbb{R}$, given by

$$s_P(x) = \frac{1}{2} \sum_{i=1}^n x_i \log x_i + \frac{1}{2} (1 - r) \log(1 - r)$$

defines the standard complex structure J_{FS} and Fubini-Study metric on \mathbb{P}^n .

Theorem 39 below provides the symplectic version of the $\partial\bar{\partial}$ -lemma in the toric context, characterizing the symplectic potentials that correspond to the toric compatible complex structures on a toric symplectic manifold. It is an immediate extension to our possibly non-compact setting of the compact version proved in [2]. To properly state it we need the following definition.

Definition 38. *Let (B, ω, τ) be a symplectic toric manifold and let us denote by*

$$Y_1, \dots, Y_n \in \mathcal{X}_H(B, \omega)$$

the Hamiltonian vector fields generating the torus action. A toric compatible complex structure $J \in \mathcal{I}^T(B, \omega)$ is said to be complete if the J -holomorphic vector fields

$$JY_1, \dots, JY_n \in \mathcal{X}(B)$$

are complete. The space of all complete toric compatible complex structures on (B, ω, τ) will be denoted by $\mathcal{I}_c^T(B, \omega)$.

Theorem 39. *Let J be any complete compatible toric complex structure on the symplectic toric manifold (B_P, ω_P, τ_P) . Then, in suitable action-angle (x, y) -coordinates on $\check{B}_P \cong \check{P} \times \mathbb{T}^n$, J is given by a symplectic potential $s \in C^\infty(\check{P})$ of the form*

$$s(x) = s_P(x) + h(x)$$

where s_P is given by Theorem 35, h is smooth on the whole P , and the matrix $S = \text{Hess}_x(s)$ is positive definite on \check{P} and has determinant of the form

$$\text{Det}(S) = \left(\delta \prod_{r=1}^d \ell_r \right)^{-1}$$

with δ being a smooth and strictly positive function on the whole P .

Conversely, any such potential s determines a (not necessarily complete) complex structure on $\check{B}_P \cong \check{P} \times \mathbb{T}^n$, that extends uniquely to a well-defined compatible toric complex structure J on the toric symplectic manifold (B_P, ω_P, τ_P) .

Remark 40. *If one takes into account Remark 24, the word “manifold” can be replaced by “orbifold” in Theorem 35 and Theorem 39 (see [3]).*

Remark 41. *There is no immediate relation between completeness of a toric compatible complex structure and completeness of the associated toric Kähler metric. See Remark 43.*

3.5. Scalar Curvature

We now recall from [1] a particular formula for the scalar curvature in action-angle (x, y) -coordinates. A Kähler metric of the form (5) has scalar curvature Sc given by¹

$$Sc = - \sum_{j,k} \frac{\partial}{\partial x_j} \left(g^{jk} \frac{\partial \log \text{Det}(S)}{\partial x_k} \right)$$

which after some algebraic manipulations becomes more compact

$$Sc = - \sum_{j,k} \frac{\partial^2 s^{jk}}{\partial x_j \partial x_k} \tag{6}$$

where the s^{jk} , $1 \leq j, k \leq n$, are the entries of the inverse of the matrix $S = \text{Hess}_x(s)$, $s \equiv$ symplectic potential. See [11] for an appropriate interpretation of this formula for the scalar curvature.

3.6. Symplectic Potentials and Affine Transformations

Because the Delzant set $P \subset \mathbb{R}^n$ of a symplectic toric manifold is well defined up to translations (i.e., additions of constants to the moment map) and $\text{SL}(n, \mathbb{Z})$ transformations (i.e., changes of basis of the torus $\mathbb{T}^n = \mathbb{R}^n / 2\pi\mathbb{Z}^n$), symplectic potentials should transform naturally under these type of maps. While the effect of translations is trivial to analyse, the effect of $\text{SL}(n, \mathbb{Z})$ transformations is more interesting. In fact

symplectic potentials transform quite naturally
under any $\text{GL}(n, \mathbb{R})$ linear transformation.

Let $T \in \text{GL}(n, \mathbb{R})$ and consider the linear symplectic change of action-angle coordinates

$$x := T^{-1}x' \quad \text{and} \quad y := T^t y'.$$

Then

$$P' = \bigcap_{a=1}^d \{x' \in \mathbb{R}^n; \ell'_a(x') := \langle x', \nu'_a \rangle + \lambda'_a \geq 0\}$$

¹The normalization for the value of the scalar curvature we are using here differs from the one used in [1, 2] by a factor of 1/2.

becomes

$$P := T^{-1}(P') = \bigcap_{a=1}^d \{x \in \mathbb{R}^n; \ell_a(x) := \langle x, \nu_a \rangle + \lambda_a \geq 0\}$$

with

$$\nu_a = T^t \nu'_a \quad \text{and} \quad \lambda_a = \lambda'_a$$

and symplectic potentials transform by

$$s = s' \circ T \quad (\text{in particular, } s_P = s'_P \circ T).$$

The corresponding Hessians are related by

$$S = T^t(S' \circ T)T$$

and

$$S_C = S'_C \circ T.$$

Example 42. Fig. 1 illustrates two equivalent descriptions of a toric symplectic rational ruled four-manifold or, equivalently, of a Hirzebruch surface

$$H_m^2 := \mathbb{P}(\mathcal{O}(-m) \oplus \mathbb{C}) \rightarrow \mathbb{P}^1, \quad m \in \mathbb{N}.$$

The linear map $T \in \text{GL}(2, \mathbb{R})$ relating the two is given by

$$T = \begin{bmatrix} m & -1 \\ 0 & 1 \end{bmatrix}.$$

The inward pointing normal that should be considered for each facet is specified. The polytope on the right is a standard Delzant polytope for the Hirzebruch surface H_m^2 . The polytope on the left is very useful for the Kähler metric constructions of Section 5 and was implicitly used by Calabi in [7].

4. Toric Constant Curvature Metrics on the Real Two Dimensional Manifolds

Any real two-dimensional orientable Riemannian manifold is Kähler since its area form is a symplectic form and oriented rotation by $\pi/2$ on each tangent plane is a compatible complex structure. In this section we write down the symplectic potentials that give rise to toric constant (scalar) curvature metrics of real two




Figure 1. Hirzebruch surfaces.

dimensional manifold and identify the underlying toric symplectic manifolds B . This will be a warm-up for the higher dimensional examples that will be presented in Section 5.

According to the formula (6) for the scalar curvature, we are looking for symplectic potentials $s : \check{P} \subset \mathbb{R} \rightarrow \mathbb{R}$ such that $s'' > 0$ and

$$-\left(\frac{1}{s''(x)}\right)'' = 2k$$

where $k \in \mathbb{R}$ denotes the Gauss curvature. This implies that

$$s''(x) = -\frac{1}{kx^2 - 2bx - c}, \quad b, c \in \mathbb{R}$$

where $x \in \check{P} \subset \mathbb{R}$ is such that $s''(x) > 0$.

4.1. Cylinders

Suppose that $k = b = 0$. Then

$$s''(x) = \frac{1}{c} > 0 \Rightarrow c > 0 \quad \text{and} \quad x \in \mathbb{R}.$$

This means that $P = \mathbb{R}$ and $s : \check{P} = \mathbb{R} \rightarrow \mathbb{R}$ can be written as

$$s(x) = \frac{x^2}{2c}.$$

Hence

$$B = \check{B} = \check{P} \times \mathbb{T}^1 = \mathbb{R} \times S^1 = \{(x, y); x \in \mathbb{R}, y \in \mathbb{R}/2\pi\mathbb{Z}\}$$

and the metric is given in matrix form by

$$\begin{bmatrix} 1/c & 0 \\ 0 & c \end{bmatrix}$$

i.e., we get a flat cylinder of radius \sqrt{c} .

4.2. Cones

Suppose that $k = 0$ and $b \neq 0$. Then, modulo a translation and possible sign change in the action variable x , we can assume that $c = 0$, $b > 0$ and

$$s''(x) = \frac{1}{2bx} > 0 \quad \text{and therefore} \quad x > 0.$$

This means that $P = [0, +\infty[$ and $s : \check{P} =]0, +\infty[\rightarrow \mathbb{R}$ can be written as

$$s(x) = \frac{1}{b} \cdot \frac{1}{2} x \log x.$$

If $b = 1$ this is the canonical symplectic potential giving the flat Euclidean metric on \mathbb{R}^2 (cf. Example 34). In general, as explained in [3], this is the symplectic potential of a cone metric of angle πb on \mathbb{R}^2 , given in matrix form by

$$\begin{bmatrix} \frac{1}{2bx} & 0 \\ 0 & 2bx \end{bmatrix}.$$

When $b = 1/p$ with $p \in \mathbb{N}$, this corresponds to an orbifold flat metric on $\mathbb{R}^2/\mathbb{Z}_p$ (see [3]).

4.3. Footballs

Suppose that $k > 0$. Then, modulo a translation in the action variable x , we can assume that $b = 0$ and

$$s''(x) = \frac{1}{c - kx^2} > 0 \Rightarrow c > 0 \quad \text{and} \quad -\sqrt{c/k} < x < \sqrt{c/k}.$$

This means that $P = [-\sqrt{c/k}, \sqrt{c/k}]$ and $s : \check{P} =]-\sqrt{c/k}, \sqrt{c/k}[\rightarrow \mathbb{R}$ can be written as

$$s(x) = \frac{1}{\sqrt{ck}} \cdot \frac{1}{2} \left[(\sqrt{c/k} + x) \log(\sqrt{c/k} + x) + (\sqrt{c/k} - x) \log(\sqrt{c/k} - x) \right].$$

If $c = 1/k$ this is the canonical symplectic potential giving the smooth round european football metric of total area $4\pi/k$ and constant Gauss curvature k on $\mathbb{P}^1 \cong \mathbb{S}^2$ (cf. Example 37). In general, this is the symplectic potential of a singular american football metric of angle $\pi\sqrt{ck}$ at the ‘‘poles’’.

4.4. Hyperbolic Metrics

Suppose that $k < 0$. Then, modulo a translation in the action variable x , we can assume that $b = 0$ and

$$s''(x) = \frac{1}{c - kx^2}.$$

If $c > 0$ then $s''(x) > 0$, for all $x \in \mathbb{R}$, which means that $P = \mathbb{R}$ and $s : \check{P} = \mathbb{R} \rightarrow \mathbb{R}$ can be written as

$$s(x) = \sqrt{\frac{-1}{ck}} \arctan \left(\sqrt{\frac{-k}{c}} x \right).$$

This is the symplectic potential of a metric of constant Gauss curvature $k < 0$ on the manifold

$$B = \check{B} = \check{P} \times \mathbb{T}^1 = \mathbb{R} \times S^1 = \{(x, y); x \in \mathbb{R}, y \in \mathbb{R}/2\pi\mathbb{Z}\}$$

which is obviously the standart hyperboloid.

If $c < 0$ then

$$s''(x) > 0 \Rightarrow x \in]-\infty, -\sqrt{c/k} [\cup] \sqrt{c/k}, +\infty [.$$

Hence, up to a sign change in the action variable x , we may assume that $P =]\sqrt{c/k}, +\infty [$ and $s : \check{P} =]\sqrt{c/k}, +\infty [\rightarrow \mathbb{R}$ can be written as

$$s(x) = \frac{1}{\sqrt{ck}} \cdot \frac{1}{2} \left[(x - \sqrt{c/k}) \log(x - \sqrt{c/k}) - (x + \sqrt{c/k}) \log(x + \sqrt{c/k}) \right].$$

If $c = 1/k$ this is the symplectic potential of the hyperbolic metric of constant Gauss curvature $k < 0$ on \mathbb{R}^2 . In other words, in the action-angle coordinates (x, y) of this symplectic model, the hyperbolic metric is given in matrix form by

$$\begin{bmatrix} \frac{-k}{(kx)^2-1} & 0 \\ 0 & \frac{(kx)^2-1}{-k} \end{bmatrix}.$$

More generally, i.e., when $c \neq 1/k$, we get singular hyperbolic metrics on \mathbb{R}^2 , with a cone singularity of angle $\pi\sqrt{ck}$ at the origin.

Remark 43. *This case illustrates the fact that there is no immediate relation between completeness of a toric compatible complex structure and completeness of the associated toric Kähler metric. Here the metric is complete but the complex structure is not. In fact, it easily follows from (4) that \mathbb{R}^2 with this complex structure is biholomorphic to an open bounded disc $D \subset \mathbb{C}$.*

If $c = 0$ we have that

$$s''(x) = -\frac{1}{kx^2} > 0, \quad \text{for all } x \neq 0$$

and $s : \check{P} =]0, +\infty[\rightarrow \mathbb{R}$ can be written as

$$s(x) = \frac{1}{k} \log(x).$$

This is the symplectic potential of a complete hyperbolic cusp metric on $\check{B} =]0, +\infty[\times S^1$, given in matrix form by

$$\begin{bmatrix} -\frac{1}{kx^2} & 0 \\ 0 & -kx^2 \end{bmatrix}.$$

5. Calabi's Family of Extremal Kähler Metrics

In [7], Calabi introduced the notion of extremal Kähler metrics. These are defined, for a fixed closed complex manifold (M, J) , as critical points of the square of the L^2 -norm of the scalar curvature, considered as a functional on the space of all symplectic Kähler forms ω in a fixed Kähler class $\Omega \in H^2(M, \mathbb{R})$. The extremal Euler-Lagrange equation is equivalent to the gradient of the scalar curvature being a holomorphic vector field (see [6]), and so these metrics generalize constant scalar curvature Kähler metrics. Moreover, Calabi showed in [8] that extremal Kähler metrics are always invariant under a maximal compact subgroup of the group of holomorphic transformations of (M, J) . Hence, on a toric manifold, extremal Kähler metrics are automatically toric Kähler metrics, and one should be able to write them down using the previous action-angle coordinates framework.

In this section, following [1], we will do that for the four-parameter family of $U(n)$ -invariant extremal Kähler metrics constructed by Calabi in [7]. Calabi used this family to put extremal Kähler metrics on

$$\mathbb{P}(\mathcal{O}(-m) \oplus \mathbb{C}) \longrightarrow \mathbb{P}^{n-1}, \quad n, m \in \mathbb{N}$$

for any Kähler class. In particular, when $n = 2$, on all Hirzebruch surfaces (cf. Example 42). As we will see here, this family can be used to write down many other interesting extremal Kähler metrics, including the non-compact, cohomogeneity one, constant scalar curvature examples that were later constructed by LeBrun [18], Pedersen-Poon [21] and Simanca [24]. Using the action-angle coordinates set-up for toric Sasaki geometry developed in [20], one can show [4] that Calabi's family also contains a family of Kähler-Einstein metrics directly related to the Sasaki-Einstein metrics constructed in 2004 by Gauntlett-Martelli-Sparks-Waldram [13, 14].

5.1. Calabi's Family in Action-Angle Coordinates

Let us take symplectic potentials $s : \check{P} \subset (\mathbb{R}^+)^n \rightarrow \mathbb{R}$ of the form

$$s(x) = \frac{1}{2} \left(\sum_{i=1}^n x_i \log x_i + h(r) \right) \quad (7)$$

where

$$r = x_1 + \cdots + x_n$$

and \check{P} will be determined in each of the particular cases that we will consider. A simple computation shows that

$$\text{Det}(S) = \frac{1 + rh''(r)}{2^n x_1 \cdots x_n} \quad \text{and} \quad S^{-1} = (s^{ij} = 2(\delta_{ij}x_i - x_i x_j f'(r)))$$

where $f = h''/(1 + rh'')$. Then, (6) implies that the scalar curvature of the corresponding toric Kähler metric is given by

$$Sc(x) = Sc(r) = 2r^2 f''(r) + 4(n+1)rf'(r) + 2n(n+1)f(r). \quad (8)$$

The Euler-Lagrange equation defining an extremal Kähler metric can be shown to be equivalent to

$$\frac{\partial Sc}{\partial x_j} \equiv \text{constant}, \quad j = 1, \dots, n \quad (9)$$

i.e., the metric is extremal if and only if its scalar curvature Sc is an affine function of x (see [1]).

Requiring that the scalar curvature $Sc \equiv Sc(r)$, given by (8), is an affine function of r is easily seen to be equivalent to

$$h''(r) = -\frac{1}{r} + \frac{r^{n-1}}{r^n - A - Br - Cr^{n+1} - Dr^{n+2}} \quad (10)$$

where $A, B, C, D \in \mathbb{R}$ are the four parameters describing the family.

As shown by Calabi in [7], one can determine explicit values for the constants $A, B, C, D \in \mathbb{R}$ so that the corresponding symplectic potential, given by (7), gives rise to an extremal Kähler metric on

$$H_m^n := \mathbb{P}(\mathcal{O}(-m) \oplus \mathbb{C}) \longrightarrow \mathbb{P}^{n-1}, \quad n, m \in \mathbb{N}.$$

In our framework, this can be seen as follows. Up to a $\text{GL}(n, \mathbb{R})$ transformation, generalizing to higher dimensions the one considered in Example 42, H_m^n is determined by a moment polytope $P_m^n(a, b) \subset \mathbb{R}^n$ with defining affine functions

$$\ell_i(x) = x_i, \quad i = 1, \dots, n, \quad \ell_{n+1}(x) = \frac{1}{m}(r - a) \quad \text{and} \quad \ell_{n+2}(x) = \frac{1}{m}(b - r)$$

where the real numbers $0 < a < b$ determine the Kähler class, i.e., the cohomology class of the symplectic form $\omega_{a,b}$. Hence, it follows from Theorem 39 that, to determine a toric compatible complex structure on $(H_m^n, \omega_{a,b})$, the symplectic potential has to be of the form

$$2s(x) = \sum_{i=1}^n x_i \log x_i + \frac{1}{m} ((r-a) \log(r-a) + (b-r) \log(b-r)) + \tilde{h}(r) \quad (11)$$

where \tilde{h} is smooth on $P_m^n(a, b) \subset \mathbb{R}^n$. Comparing (7), (10) and (11), one concludes that we must have

$$\frac{r^{n-1}}{r^n - A - Br - Cr^{n+1} - Dr^{n+2}} = \frac{1}{m} \left(\frac{1}{r-a} + \frac{1}{b-r} \right) + R(r) \quad (12)$$

where $R(r)$ is a smooth function on $P_m^n(a, b) \subset \mathbb{R}^n$. This gives rise to a system of four linear equations in the four unknowns $A, B, C, D \in \mathbb{R}$, which admits a unique explicit solution for any $n, m \in \mathbb{N}$ and $a, b \in \mathbb{R}$ such that $0 < a < b$ (see page 285 of [7] or [22]).

5.2. Particular Cases

By construction, all Kähler metrics in Calabi's four-parameter family are extremal. A simple computation shows that their scalar curvature is given by

$$Sc(r) = 2(n+1)((2+n)Dr + nC).$$

Hence, these metrics have

$$\text{constant scalar curvature iff } D = 0$$

and are

$$\text{scalar-flat iff } C = D = 0.$$

Moreover, one can show that these metrics are

$$\text{Kähler-Einstein iff } B = D = 0$$

and

$$\text{Ricci-flat iff } B = C = D = 0.$$

We will now analyse in more detail these constant scalar curvature particular cases. Note that when $A = B = C = D = 0$ we have

$$s(x) = \frac{1}{2} \sum_{i=1}^n x_i \log x_i$$

which is the standard symplectic potential of the Delzant set $P = (\mathbb{R}_0^+)^n$ and determines the standard flat Euclidean metric on \mathbb{R}^{2n} (cf. Example 34).

5.3. Ricci-Flat Metrics

Assume that $B = C = D = 0$ and $A = a^n$ with $0 < a \in \mathbb{R}$. Then

$$\begin{aligned} h''(r) &= -\frac{1}{r} + \frac{r^{n-1}}{r^n - a^n} \\ &= -\frac{1}{r} + \frac{r^{n-1}}{(r-a) \sum_{k=1}^n a^{k-1} r^{n-k}} \\ &= -\frac{1}{r} + \frac{1}{n} \cdot \frac{1}{r-a} + R(r) \end{aligned}$$

where $R(r)$ is a smooth function on the rational Delzant set $P^n(a) \subset \mathbb{R}^n$ with defining affine functions

$$\ell_i(x) = x_i, \quad \text{for all } i = 1, \dots, n \quad \text{and} \quad \ell_{n+1}(x) = \frac{1}{n}(r-a).$$

The symplectic potential can be written as

$$s(x) = \frac{1}{2} \left(\sum_{i=1}^n x_i \log x_i + \frac{1}{n}(r-a) \log(r-a) + \tilde{h}(r) \right)$$

where \tilde{h} is smooth on $P^n(a) \subset \mathbb{R}^n$. Hence, for each $a > 0$, it defines a Ricci-flat Kähler metric on the total space of the canonical line bundle

$$\mathcal{O}(-n) \longrightarrow \mathbb{P}^{n-1}$$

(as before, up to a $\text{GL}(n, \mathbb{R})$ transformation, the underlying non-compact toric symplectic manifold is determined by $P^n(a) \subset \mathbb{R}^n$). These are the metrics constructed by Calabi in [6].

5.4. Scalar-Flat Metrics

We will now show that Calabi's family also contains the complete scalar-flat Kähler metrics on the total space of the line bundles

$$\mathcal{O}(-m) \longrightarrow \mathbb{P}^{n-1}, \quad \text{for all } m \in \mathbb{N}$$

constructed by LeBrun [18] and Pedersen-Poon [21] (see also Simanca [24]).

Up to a $\text{GL}(n, \mathbb{R})$ transformation, these spaces are determined by the rational Delzant sets $P_m^n(a) \subset \mathbb{R}^n$, with $0 < a \in \mathbb{R}$ and defining affine functions

$$\ell_i(x) = x_i, \quad i = 1, \dots, n, \quad \text{and} \quad \ell_{n+1}(x) = \frac{1}{m}(r-a).$$

which means that the symplectic potential has to be of the form

$$s(x) = \frac{1}{2} \left(\sum_{i=1}^n x_i \log x_i + \frac{1}{m} (r-a) \log(r-a) + \tilde{h}(r) \right)$$

where \tilde{h} is smooth on $P_m^n(a)$. This implies that

$$h''(r) = -\frac{1}{r} + \frac{1}{m} \cdot \frac{1}{r-a} + R(r)$$

with $R(r)$ smooth on $P_m^n(a)$. Since the scalar-flat condition is equivalent to $C = D = 0$, we get that

$$-\frac{1}{r} + \frac{r^{n-1}}{r^n - A - Br} = -\frac{1}{r} + \frac{1}{m} \cdot \frac{1}{r-a} + R(r).$$

This relation gives rise to a system of two linear equations in the two unknowns $A, B \in \mathbb{R}$, which admits a unique solution for any $n, m \in \mathbb{N}$ and $0 < a \in \mathbb{R}$

$$A = a^n(1 - n + m) \quad \text{and} \quad B = (n - m)a^{n-1}.$$

Note that when $m = 1$ we get complete scalar-flat Kähler metrics on the total space of the line bundle

$$\mathcal{O}(-1) \longrightarrow \mathbb{P}^{n-1}$$

i.e., on \mathbb{C}^n blown-up at the origin. These were originally constructed by D. Burns (at least when $n = 2$).

5.5. Fubini-Study and Bergman Metrics

Assume that $A = B = D = 0$, which implies in particular that we are considering Kähler-Einstein metrics. Then

$$\begin{aligned} h''(r) &= -\frac{1}{r} + \frac{r^{n-1}}{r^n - Cr^{n+1}} \\ &= -\frac{1}{r} + \frac{1}{r(1 - Cr)} = \frac{1}{\frac{1}{C} - r} \end{aligned}$$

which implies that the symplectic potential can be written as

$$s(x) = \frac{1}{2} \left(\sum_{i=1}^n x_i \log x_i + \left| \frac{1}{C} - r \right| \log \left| \frac{1}{C} - r \right| \right).$$

When $C = 1$ we recover Example 37, i.e., the standard complex structure and Fubini-Study metric on \mathbb{P}^n . More generally, for any $C > 0$, this defines the standard complex structure and suitably scaled Fubini-Study metric on \mathbb{P}^n . The corresponding moment polytope is the simplex in \mathbb{R}^n with defining affine functions

$$\ell_i(x) = x_i, \quad \text{for all } i = 1, \dots, n, \quad \text{and} \quad \ell_{n+1}(x) = \frac{1}{C} - r.$$

When $C < 0$ it follows from Theorem 39 that the above symplectic potential determines a toric compatible complex structure J_C on the toric symplectic manifold $(\mathbb{R}^{2n}, \omega_{\text{st}})$ with Delzant set $P = (\mathbb{R}_0^+)^n \subset \mathbb{R}^n$. The corresponding Kähler metric is a $U(n)$ -invariant Kähler-Einstein metric of negative scalar curvature on the complex manifold (\mathbb{R}^{2n}, J_C) . Using the holomorphic coordinates given by (4), one easily concludes that (\mathbb{R}^{2n}, J_C) is biholomorphic to a ball $B \subset \mathbb{C}^n$, which implies in particular that J_C is not complete. Moreover, the Kähler metric $\langle \cdot, \cdot \rangle_C := \omega_{\text{st}}(\cdot, J_C \cdot)$ is, in fact, the well-known and complete Bergman metric.

5.6. Other Kähler-Einstein Metrics

We will now show that Calabi's family also contains the complete Kähler-Einstein metrics with negative scalar curvature on the total space of the open disc bundles

$$\mathcal{D}(-m) \subset \mathcal{O}(-m) \longrightarrow \mathbb{P}^{n-1} \quad \text{for all } n < m \in \mathbb{N}$$

constructed by Pedersen-Poon [21].

As toric symplectic manifolds, and up to a $\text{GL}(n, \mathbb{R})$ transformation, these spaces are again determined by the rational Delzant sets $P_m^n(a) \subset \mathbb{R}^n$ with defining affine functions

$$\ell_i(x) = x_i, \quad \text{for all } i = 1, \dots, n, \quad \text{and} \quad \ell_{n+1}(x) = \frac{1}{m}(r - a)$$

which means that the symplectic potential has to be of the form

$$s(x) = \frac{1}{2} \left(\sum_{i=1}^n x_i \log x_i + \frac{1}{m}(r - a) \log(r - a) + \tilde{h}(r) \right)$$

where \tilde{h} is smooth on $P_m^n(a)$. This implies that

$$h''(r) = -\frac{1}{r} + \frac{1}{m} \cdot \frac{1}{r - a} + R(r)$$

with $R(r)$ smooth on $P_m^n(a)$. Since the Kähler-Einstein condition is equivalent to $B = D = 0$, we get that

$$-\frac{1}{r} + \frac{r^{n-1}}{r^n - A - Cr^{n+1}} = -\frac{1}{r} + \frac{1}{m} \cdot \frac{1}{r-a} + R(r).$$

This relation gives rise to a system of two linear equations in the two unknowns $A, C \in \mathbb{R}$, which admits a unique solution for any $n < m \in \mathbb{N}$ and $0 < a \in \mathbb{R}$

$$A = \frac{(m+1)a^n}{n+1} > 0 \quad \text{and} \quad C = \frac{n-m}{(n+1)a} < 0.$$

As remarked by Pedersen-Poon, these metrics are a superposition of Calabi's Ricci-flat metrics ($A > 0$) and Bergman metrics ($C < 0$). The analogous superposition of Calabi's Ricci-flat metrics ($A > 0$) with Fubini-Study metrics ($C > 0$) gives rise to Kähler-Einstein metrics on the projectivization of the above line bundles, with cone-like singularities in the normal directions to the zero and infinity sections. As explained in [4], these metrics are directly related to the smooth Sasaki-Einstein metrics constructed in 2004 by Gauntlett-Martelli-Sparks-Waldram [13, 14].

5.7. Other Constant Scalar Curvature Metrics

We will now show that Calabi's family also contains the complete constant negative scalar curvature Kähler metrics on the total space of the open disc bundles

$$\mathcal{D}(-m) \subset \mathcal{O}(-m) \longrightarrow \mathbb{P}^{n-1} \quad \text{for all } n, m \in \mathbb{N}$$

constructed again by Pedersen-Poon [21].

As before, we are interested in the rational Delzant sets $P_m^n(a) \subset \mathbb{R}^n$ and symplectic potentials of the form

$$s(x) = \frac{1}{2} \left(\sum_{i=1}^n x_i \log x_i + \frac{1}{m} (r-a) \log(r-a) + \tilde{h}(r) \right)$$

with \tilde{h} smooth on $P_m^n(a)$. Assuming $D = 0$ and $C = -1$, i.e., $Sc = -2n(n+1)$, this implies that

$$-\frac{1}{r} + \frac{r^{n-1}}{r^n - A - Br + r^{n+1}} = -\frac{1}{r} + \frac{1}{m} \cdot \frac{1}{r-a} + R(r)$$

with $R(r)$ smooth on $P_m^n(a)$. This relation gives rise to a system of two linear equations in the two unknowns $A, B \in \mathbb{R}$, which admits a unique solution for any $n, m \in \mathbb{N}$ and $0 < a \in \mathbb{R}$

$$A = (m-n+(1-n)a)a^n \quad \text{and} \quad B = (n-m+1+na)a^{n-1}.$$

Acknowledgements

I would like to thank to the organizers of the XI-th International Conference on Geometry, Integrability and Quantization held in Varna, Bulgaria, June 5–10, 2009, where this material was presented as part of a mini-course.

This work is partially supported by the Fundação para a Ciência e a Tecnologia (FCT/Portugal) and that one presented in Section 5 was carried out in January–June of 2001, while I was at the Fields Institute for Research in Mathematical Sciences in Toronto, Canada. It was presented during seminar talks given at the Workshop on Hamiltonian Group Actions and Quantization organized by the Fields Institute, June 4–13, 2001, and at Differentialgeometrie im Großen, Mathematisches Forschungsinstitut Oberwolfach, Germany, June 10–16, 2001. Here I want to express my thanks for the support and hospitality of the organizers of those meetings.

References

- [1] Abreu M., *Kähler Geometry of Toric Varieties and Extremal Metrics*, Int. J. Math. **9** (1998) 641–651.
- [2] Abreu M., *Kähler Geometry of Toric Manifolds in Symplectic Coordinates*, In: *Symplectic and Contact Topology: Interactions and Perspectives*, Y. Eliashberg, B. Khesin and F. Lalonde (Eds), Fields Institute Communications 35, American Mathematical Society 2003, pp 1–24.
- [3] Abreu M., *Kähler Metrics on Toric Orbifolds*, J. Diff. Geom. **58** (2001) 151–187.
- [4] Abreu M., *Kähler-Sasaki Geometry of Toric Symplectic Cones in Action-Angle Coordinates*, to appear in the Proceedings of the Geometry Summer School, Lisbon, July 13–17, 2009.
- [5] Atiyah M., *Convexity and Commuting Hamiltonians*, Bull. London Math. Soc. **14** (1982) 1–15.
- [6] Calabi E., *Métriques Kähleriennes et fibrés holomorphes*, Ann. Sci. Ec. Norm. Sup. **12** (1979) 269–294.
- [7] Calabi E., *Extremal Kähler Metrics*, In: *Seminar on Differential Geometry*, S.-T. Yau (Ed), Ann. Math. Studies vol.102, Princeton Univ. Press 1982, pp 259–290.

-
- [8] Calabi E., *Extremal Kähler Metrics II*, In: Differential Geometry and Complex Analysis, I. Chavel and H. Farkas (Eds), Springer, Berlin 1985, pp 95–114.
- [9] Calderbank D., David L. and Gauduchon P., *The Guillemin Formula and Kähler Metrics on Toric Symplectic Manifolds*, J. Symplectic Geom. **1** (2003) 767–784.
- [10] Delzant T., *Hamiltoniens périodiques et images convexes de l'application moment*, Bull. Soc. Math. France **116** (1988) 315–339.
- [11] Donaldson S., *Remarks on Gauge Theory, Complex Geometry and 4-manifold Topology*, In: Fields Medallists' Lectures, M. Atiyah and D. Iagolnitzer (Eds), World Scientific, Singapore 1997, pp 384–403.
- [12] Donaldson S., *Scalar Curvature and Stability of Toric Varieties*, J. Diff. Geom. **62** (2002) 289–349.
- [13] Gauntlett J., Martelli D., Sparks J. and Waldram D., *Sasaki-Einstein Metrics on $S^2 \times S^3$* , Adv. Theor. Math. Phys. **8** (2004) 711–734.
- [14] Gauntlett J., Martelli D., Sparks J. and Waldram D., *A New Infinite Class of Sasaki-Einstein Manifolds*, Adv. Theor. Math. Phys. **8** (2004) 987–1000.
- [15] Guillemin V., *Kähler Structures on Toric Varieties*, J. Diff. Geom. **40** (1994) 285–309.
- [16] Guillemin V. and Sternberg S., *Convexity Properties of the Moment Mapping*, Invent. Math. **67** (1982) 491–513.
- [17] Karshon Y. and Lerman E., *Non-Compact Symplectic Toric Manifolds*, arXiv:0907.2891.
- [18] LeBrun C., *Counter-Examples to the Generalized Positive Action Conjecture*, Comm. Math. Phys. **118** (1988) 591–596.
- [19] Lerman E. and Tolman S., *Hamiltonian Torus Actions on Symplectic Orbifolds and Toric Varieties*, Trans. Amer. Math. Soc. **349** (1997) 4201–4230.
- [20] Martelli D., Sparks J. and Yau S.-T., *The Geometric Dual of α -maximisation for Toric Sasaki-Einstein Manifolds*, Comm. Math. Phys. **268** (2006) 39–65.
- [21] Pedersen H. and Poon Y., *Hamiltonian Constructions of Kähler-Einstein Metrics and Kähler Metrics of Constant Scalar Curvature*, Comm. Math. Phys. **136** (1991) 309–326.
- [22] Raza A., *An Application of Guillemin-Abreu Theory to a Non-Abelian Group Action*, Diff. Geom. Appl. **25** (2007) 266–276.
- [23] Siegel C., *Symplectic Geometry*, Amer. J. Math. **65** (1943) 1–86.

-
- [24] Simanca S., *Kähler Metrics of Constant Scalar Curvature on Bundles Over $\mathbb{C}P^{n-1}$* , Math. Ann. **291** (1991) 239–246.

Miguel Abreu

Centro de Análise Matemática

Geometria e Sistemas Dinâmicos

Departamento de Matemática

Instituto Superior Técnico

Av. Rovisco Pais, 1049-001 Lisboa

PORTUGAL

E-mail address: mabreu@math.ist.utl.pt