

THE NAVIER-STOKES EQUATIONS AND WEAK HERZ SPACES

YOHEI TSUTSUI

Department of Mathematics, Osaka University, Japan

(Submitted by: Yoshikazu Giga)

Abstract. In this paper, we discuss the Cauchy problem for Navier-Stokes equations in homogeneous weak Herz spaces $WK_{p,q}^\alpha(\mathbb{R}^n)$. More precisely, we construct the solution in the class $L^\infty(0, T; WK_{p,q}^\alpha)$ with the initial data in $WK_{p,q}^\alpha$. Further, we consider the blow-up phenomena of time-local solutions and the uniqueness of global solutions with large initial data in $WK_{p,q}^\alpha$. Also, we give several embeddings of weak Herz spaces into homogeneous Besov spaces $\dot{B}_{p,\infty}^{-\alpha}(\mathbb{R}^n)$ ($\alpha > 0$), or $bmo^{-1}(\mathbb{R}^n)$.

1. INTRODUCTION

In this paper, we consider the Cauchy problem for the incompressible homogeneous Navier-Stokes equations on the whole space \mathbb{R}^n ,

$$(N-S) \quad \begin{cases} \partial_t u - \Delta u + (u \cdot \nabla)u + \nabla p = 0, \\ \operatorname{div} u = 0, \\ u(0) = u_0 \end{cases}$$

with the initial data u_0 in a homogeneous weak Herz space $WK_{p,q}^\alpha$. Here $u = (u^1, \dots, u^n)$ is the unknown velocity vector field, p is the unknown pressure scalar field and $u_0 = (u_0^1, \dots, u_0^n)$ is the given initial velocity with $\operatorname{div} u_0 = \nabla \cdot u_0 = 0$. Solving the Cauchy problem (N-S) can be reduced to finding a divergence free solution u of the integral equation

$$(I.E.) \quad \begin{aligned} u(t) &= e^{t\Delta} u_0 - \int_0^t e^{(t-s)\Delta} \mathbb{P}(u \cdot \nabla)u(s) ds \\ &= e^{t\Delta} u_0 - \int_0^t \nabla e^{(t-s)\Delta} \mathbb{P}(u \otimes u)(s) ds =: e^{t\Delta} u_0 - B(u, u)(t), \end{aligned}$$

where $u \otimes v := (u^i v^j)_{1 \leq i, j \leq n}$ is a matrix-valued function whose (i, j) component is $u^i v^j$, $e^{t\Delta}$ is the heat semigroup and $\mathbb{P} = \{\mathbb{P}_{i,j} := \delta_{i,j} + R_i R_j\}_{1 \leq i, j \leq n}$

Accepted for publication: April 2011.

AMS Subject Classifications: 35Q30, 76D05.

denotes the Leray-Hopf operator or the Weyl-Helmholtz projection which is the orthogonal projection on the solenoidal vector field, where $R_j = (-\Delta)^{-1/2}\partial_j$ is the j th Riesz transform. Of course, the operator $e^{t\Delta}$ is defined by the convolution

$$e^{t\Delta}f(x) := f * G_{\sqrt{t}}(x),$$

where G is the Gaussian $G(x) := \frac{1}{(4\pi)^{n/2}}e^{-|x|^2/4}$ and $G_t(x) := t^{-n}G(x/t)$.

Formally, if u is a divergence free solution of (I.E.) and

$$p := (-\Delta)^{-1} \sum_{j,k=1}^n \partial_j \partial_k (u^j u^k) = \sum_{j,k=1}^n R_j R_k (u^j u^k)$$

then (u, p) solves (N-S). We call the solution of (I.E.) the mild solution of (N-S). There are many papers which studied Navier-Stokes equations on several function spaces, for example, [2], [17], [9], [13], [14], [21], [23], [24], [27], [30], [33], [32], [38], [40], [44], [45] etc. In the present paper, we construct local and global mild solutions of (N-S) with the initial velocity u_0 belonging to the weak Herz space. We also discuss the uniqueness of our solutions and blow-up phenomena of our local solutions. Moreover, we investigate embeddings of weak Herz spaces into Besov spaces or bmo^{-1} .

Here we recall a remark of Cannone [7] to make clear the advantages of using the dyadic decomposition of a function in x -space and ξ -space, respectively. In [7], he proved that if the initial data $u_0 \in L^3(\mathbb{R}^3)$ is small in the sense of $\dot{B}_{p,\infty}^{-1+3/p}$ where $3 < p \leq 6$, then there exists a global mild solution of (N-S) in $C([0, \infty), L^3)$. Also, he remarked that, for any $f \in L^n(\mathbb{R}^n)$,

$$\lim_{|k| \rightarrow \infty} \|\omega_k f ; \dot{B}_{p,\infty}^{-1+n/p}\| = 0,$$

where $\omega_k(x) := e^{ikx}$ and $n < p < \infty$. In other words, for the initial data $u_0 \in L^n$, if u_0 oscillates sufficiently, then there exists a global mild solution $u \in C((0, \infty); L^n)$ to (N-S) with u_0 , even if $\|u_0 ; L^n\|$ is large. On the other hand, roughly speaking, Theorem 1.3 below allows us to deal with the following function f as the initial data:

$$f(x) := \sum_{k \in \mathbb{Z}} \frac{1}{|x - x_k|} \chi_k(x),$$

where $x_k := (\frac{3}{2}2^{k-1}, 0, \dots, 0) \in \mathbb{R}^n$ and χ_k is the characteristic function of $\{2^{k-1} \leq |x| < 2^k\}$. It is not hard to see that $f \in \dot{W}K_{n,\infty}^0$ and $f \notin \cup_{n < p < \infty} \dot{B}_{p,\infty}^{-1+n/p}$. For the latter, see Theorem 1.7. In other words, the

theorem says that even if the initial data u_0 has infinitely many singular points, if $\|u_0 ; W\dot{K}_{n,\infty}^0\|$ is sufficiently small, then there exists a global mild solution $u \in C((0, \infty); W\dot{K}_{n,\infty}^0)$ to (N-S) with u_0 . It is well known that $L^{n,\infty} \hookrightarrow \dot{B}_{p,\infty}^{-1+n/p} \hookrightarrow BMO^{-1}$ when $n < p < \infty$ and there exists a global mild solution $u \in L^\infty(0, \infty; BMO^{-1})$ to (N-S) with u_0 when $\|u_0 ; BMO^{-1}\|$ is sufficiently small, see [24]. Since we disprove $\dot{B}_{p,\infty}^{-1+n/p} \hookrightarrow W\dot{K}_{n,\infty}^0$ and Professor Akihiko Miyachi proved the inclusion $W\dot{K}_{n,\infty}^0 \hookrightarrow BMO^{-1}$ recently, our result are independent from Cannone's result and we have

$$\begin{aligned} L^n \hookrightarrow L^{n,\infty} \hookrightarrow \dot{B}_{p,\infty}^{-1+n/p} \hookrightarrow BMO^{-1}. \\ \hookrightarrow W\dot{K}_{n,\infty}^0 \hookrightarrow. \end{aligned}$$

Then, our initial data in Theorem 1.3 below were dealt with by Koch-Tataru in [24]. In [26], they introduced the Fourier-Besov spaces $FB_{p,q}^\alpha$ which are defined by $\|f ; FB_{p,q}^\alpha\| := \|\hat{f} ; \dot{K}_{p,q}^\alpha\|$, where \hat{f} is the Fourier transform of f ; see Definition 1.1 below for the definition of the norm $\|\cdot ; \dot{K}_{p,q}^\alpha\|$.

In the present paper, we use the symbol $\dot{K}_{p,q}^\alpha$, instead of $\dot{K}_q^{\alpha,p}$, because the parameter p was adopted as the one describing local regularity of functions in many papers studying Navier-Stokes equations. See below for the precise definition of $\dot{K}_{p,q}^\alpha$.

We explain about the Herz spaces. Nonhomogeneous Herz spaces $K_{p,1}^{n(1-\frac{1}{p})}$ with $1 < p < \infty$ are now called Beurling algebras which were introduced by Beurling [4]. Feichtinger [16] gave the different norms of Beurling algebras, which are equivalent to those in [4]. The space $\dot{K}_{p,q}^\alpha$ for all p, q and α was introduced by Herz [20] but his definition differs from our definition. In [8], the Hardy spaces associated to the Beurling algebras were introduced, then these spaces are generalized and are now called Herz-type Hardy spaces. The theories of the Herz space and the Herz-type Hardy space were developed by, for example, [10], [11], [34], [35], [36], [37], [41] etc.... These spaces are useful in the analysis of mapping properties of important operators. For example, Baernstein II and Sawyer [1] showed some multiplier theorems on the Hardy space $H^p(\mathbb{R}^n)$ by using a norm of the Herz space. Also, see [48] and the references there for applications of Herz spaces.

Firstly we recall the definitions of Herz spaces and weak Herz spaces.

Definition 1.1. *Let $0 < p, q \leq \infty$ and $\alpha \in \mathbb{R}$. One defines the homogeneous Herz space $\dot{K}_{p,q}^\alpha(\mathbb{R}^n)$ as $\dot{K}_{p,q}^\alpha(\mathbb{R}^n) := \{f \in L_{loc}^p(\mathbb{R}^n \setminus \{0\}) : \|f ; \dot{K}_{p,q}^\alpha\| < \infty\}$,*

where

$$\|f ; \dot{K}_{p,q}^\alpha\| := \left(\sum_{k \in \mathbb{Z}} 2^{k\alpha q} \|f ; L^p(A_k)\|^q \right)^{1/q},$$

with the usual modification in the case $q = \infty$, and $A_k := \{x \in \mathbb{R}^n : 2^{k-1} \leq |x| < 2^k\}$.

Definition 1.2. With the same exponents as above, one defines the weak Herz spaces $W\dot{K}_{p,q}^\alpha(\mathbb{R}^n)$ as the space of measurable functions f such that

$$\|f ; W\dot{K}_{p,q}^\alpha\| := \left(\sum_{k \in \mathbb{Z}} 2^{k\alpha q} \|f ; L^{p,\infty}(A_k)\|^q \right)^{1/q} < \infty,$$

with the usual modification in the case $q = \infty$, where

$$\|f ; L^{p,\infty}(A_k)\| := \sup_{\lambda > 0} \lambda |\{x \in A_k : |f(x)| > \lambda\}|^{1/p}$$

for $p < \infty$ and $L^{\infty,\infty}(A_k) = L^\infty(A_k)$.

A typical example of $W\dot{K}_{p,\infty}^\alpha$ is $\sum_{k \in \mathbb{Z}} \frac{2^{-k\alpha}}{|x-x_k|^{n/p}} \chi_k(x)$, where $x_k := (\frac{3}{2}2^{k-1}, 0, \dots, 0) \in \mathbb{R}^n$ and χ_k is the characteristic function of A_k . These functions are used in the proof of Theorem 1.7 below.

Here we remark that there is another type of weak Herz space, [25], defined by

$$\|f ; \dot{\mathcal{K}}_{(p,\infty),q}^\alpha\| := \sup_{\lambda > 0} \lambda \left(\sum_{k \in \mathbb{Z}} 2^{k\alpha q} |\{x \in A_k : |f(x)| > \lambda\}|^{p/q} \right)^{1/q}.$$

It is obvious $\|f ; \dot{\mathcal{K}}_{(p,\infty),q}^\alpha\| \leq \|f ; W\dot{K}_{p,q}^\alpha\|$; i.e., $W\dot{K}_{p,q}^\alpha \hookrightarrow \dot{\mathcal{K}}_{(p,\infty),q}^\alpha$.

Moreover, to state our theorems, we introduce the ‘‘smooth’’ weak Herz spaces

$$W\dot{\mathcal{K}}_{p,\infty}^\alpha := \{f \in W\dot{K}_{p,\infty}^\alpha ; e^{t\Delta} f \rightarrow f \text{ in } W\dot{K}_{p,\infty}^\alpha \text{ as } t \searrow 0\}.$$

Now we state our theorems. The first one concerns the local existence of a mild solution for large data.

Theorem 1.1 (local existence for large data). *Let $n \geq 2$, $n < p < \infty$ and $0 \leq \alpha < 1 - n/p$. Then for every $u_0 \in W\dot{K}_{p,\infty}^\alpha$ with $\operatorname{div} u_0 = 0$, there exist a positive constant $T > 0$ and a solution $u \in X_T \cap C((0, T); W\dot{K}_{p,\infty}^\alpha)$ of (I.E.) with u_0 such that*

$$\sup_{0 < t \leq T} t^{(\alpha+n/p)/2} \|u(t) ; L^\infty\| < \infty, \tag{1.1}$$

$$u(t) \in W\dot{\mathcal{K}}_{p,\infty}^\alpha \text{ for } t > 0, \tag{1.2}$$

$$u(t) \rightarrow u_0 \text{ in the weak* topology as } t \searrow 0, \tag{1.3}$$

$$u(\cdot) - e^{\Delta}u_0 \in L^\infty(0, T; \dot{K}_{p,\infty}^\alpha), \tag{1.4}$$

and

$$u(\cdot) - e^{\Delta}u_0 \text{ is right continuous in } \dot{K}_{p,\infty}^\alpha \text{ on } (0, T), \tag{1.5}$$

where $X_T = \{u \in L^\infty(0, T; W\dot{K}_{p,\infty}^\alpha) : \|u ; X_T\| < \infty, \operatorname{div} u = 0\}$,

$$\begin{aligned} \|u ; X_T\| &:= \sup_{0 < t \leq T} \|u(t) ; W\dot{K}_{p,\infty}^\alpha\| + \sup_{0 < t \leq T} t^{1/2} \|\nabla u(t) ; W\dot{K}_{p,\infty}^\alpha\| \\ &=: \|u ; X_{T,1}\| + \|u ; X_{T,2}\|. \end{aligned}$$

Moreover, if $u_0 \in W\dot{K}_{p,\infty}^\alpha$, then we can prove $u(t) \rightarrow u_0$ in $W\dot{K}_{p,\infty}^\alpha$ as $t \searrow 0$. As for uniqueness, u is the only solution of (I.E.) in the class $L^\infty(0, T; W\dot{K}_{p,\infty}^\alpha)$.

Remark 1.1. 1. The local solution above is in $\cap_{0 < T_1 < T_2 < T} L^\infty(\mathbb{R}^n \times (T_1, T_2))$. Hence, thanks to the proof of Proposition 15.1 in [31], we obtain the smoothness $u \in C^\infty(\mathbb{R}^n \times (0, T))$.

2. The life span T of the solution u in Theorem 1.1 is characterized by $T = \frac{C}{\|u_0 ; W\dot{K}_{p,\infty}^\alpha\|^{\frac{2}{1-n/p-\alpha}}}$, with a constant C depending on the exponents p, α and n only.

3. Because we show the inclusion $W\dot{K}_{p,\infty}^\alpha \hookrightarrow bmo^{-1}$ for $n < p \leq \infty$ and $0 \leq \alpha < 1 - n/p$, see Theorem 1.6, our initial data u_0 was dealt with by Koch and Tataru [24], where $bmo^{-1} = F_{\infty,2}^{-1}$ is defined by

$$\|f ; bmo^{-1}\| := \sup_{|Q| \leq 1} \left(\int_Q \int_0^{l(Q)^2} |e^{t\Delta} f(x)|^2 dt dx \right)^{1/2},$$

where the supremum is taken over all cubes Q with the volume $|Q| \leq 1$, $l(Q)$ is the side length of Q and the slashed integral $\int_Q f dx$ denotes the average $\frac{1}{|Q|} \int_Q f dx$. But Theorem 1.1 is not included in [24], because they used the condition

$$\sup_{0 < t \leq T} t^{1/2} \|u(t) ; L^\infty\| < \infty$$

to prove the uniqueness of local solutions. We show the uniqueness of our local solutions without such a condition.

4. In the case $u_0 \in \dot{K}_{p,\infty}^\alpha$, we can show Theorem 1.1 in which $W\dot{K}_{p,\infty}^\alpha$ is replaced with $\dot{K}_{p,\infty}^\alpha$. However we omit the details; this fact says that the assumption of Theorem 1.2 below is natural.

Also, we are interested in whether the local solution in Theorem 1.1 blows up at $t = T$ or can be continued beyond $t = T$. Such a problem was considered by Beale, Kato and Majda [3], Giga [12], Kozono and Taniuchi [29], Kozono, Ogawa and Taniuchi [27] and Kozono and Shimada [28]. We can establish an extension criterion on our local-in-time mild solutions by using the new function space $\dot{K}_{BMO,\infty}^0$ which is strictly larger than BMO by applying our previous work [47].

Definition 1.3. We define the space $\dot{K}_{BMO,\infty}^0$ as the space of locally integrable functions f on $\mathbb{R}^n \setminus \{0\}$ satisfying

$$\|f ; \dot{K}_{BMO,\infty}^0\| := \sup_{k \in \mathbb{Z}} \|f ; BMO(Q_k^*)\| := \sup_{k \in \mathbb{Z}} \sup_{Q \subset Q_k^*} \inf_{c \in \mathbb{C}} \int_Q |f - c| dy < \infty,$$

where $Q_k^* := Q_{k-1} \cup Q_k \cup Q_{k+1}$, $Q_k := (-2^k, 2^k)^n \setminus (-2^{k-1}, 2^{k-1})^n$ and the supremum $\sup_{Q \subset Q_k^*}$ is taken over all cubes Q contained in Q_k^* . Then, $\dot{K}_{BMO,\infty}^0/\mathbb{C}$ is a Banach space under the above norm $\|\cdot ; \dot{K}_{BMO,\infty}^0\|$.

Our extension criterion theorem is the following. Theorems 1.1 and 1.2 are proved in Section 3.

Theorem 1.2 (continuation principle for local mild solutions). *Let $n \geq 2$, $n < p < \infty$, $0 \leq \alpha < 1 - n/p$ and $0 < T^* < \infty$. Let $u_0 \in \dot{K}_{p,\infty}^\alpha$ with $\operatorname{div} u_0 = 0$ and u be a solution of (I.E.) in the class $L^\infty(0, T; \dot{K}_{p,\infty}^\alpha)$ for all $T \in (0, T^*)$. Suppose that u satisfies the condition*

$$\int_0^{T^*} \|\nabla u(t) ; \dot{K}_{BMO,\infty}^0\| dt < \infty;$$

then there exists $\tilde{T} > T^$ so that u can be extended to a solution of (I.E.) in the class $L^\infty(0, \tilde{T}; \dot{K}_{p,\infty}^\alpha)$.*

Remark 1.2. Kozono and Shimada [28] showed that if

$$\int_0^{T^*} \|u(t) ; \dot{F}_{\infty,\infty}^{-\alpha}\|^\gamma dt < \infty$$

for some $0 < \alpha < 1$ with $\gamma = 2(1 - \alpha)$, then a similar continuation principle of strong solutions holds. Here $\dot{F}_{\infty,\infty}^{-\alpha}$ is the homogeneous Triebel-Lizorkin space.

Although the Navier-Stokes equations are invariant with respect to the scaling $(u(x, t), p(x, t)) \rightarrow (\lambda u(\lambda x, \lambda^2 t), \lambda^2 p(\lambda x, \lambda^2 t))$, $(\lambda > 0)$ the weak Herz spaces appearing in Theorem 1.1 are not invariant with respect to such a

scaling. Roughly speaking, it is well known that if a function space X is invariant with respect to the scaling, i.e., $\|f(\lambda \cdot) ; X\| \approx \lambda^{-1} \|f ; X\|$, then the global existence of solutions for small data to (I.E.) is expected. It is not hard to show that $W\dot{K}_{p,q}^{1-n/p}$ and $\dot{K}_{p,q}^{1-n/p}$ have the described invariance, see Lemma 2.5 in Section 2. Thus the global existence of mild solutions for small data is expected on these spaces. Under the restriction $n \leq p \leq \infty$, since the largest space is $W\dot{K}_{n,\infty}^0$ among $\dot{K}_{p,q}^{1-n/p}$ and $W\dot{K}_{p,q}^{1-n/p}$, we state our global existence theorem for $W\dot{K}_{n,\infty}^0$ only. More precisely, we have the following theorem.

Theorem 1.3 (global existence for small data). *Let $n \geq 2$ and $n < p < \infty$. Then, there exists $\delta > 0$ such that for all $u_0 \in W\dot{K}_{n,\infty}^0$ with $\|u_0 ; W\dot{K}_{n,\infty}^0\| \leq \delta$ and $\operatorname{div} u_0 = 0$ there exists a solution $u \in X \cap C((0, \infty); W\dot{K}_{n,\infty}^0)$ of (I.E.) such that*

$$u(t) \in W\dot{K}_{n,\infty}^0 \text{ for } t > 0, \tag{1.6}$$

$$u(t) \rightarrow u_0 \text{ in weak* topology as } t \searrow 0, \tag{1.7}$$

$$u(\cdot) - e^{\cdot \Delta} u_0 \in L^\infty(0, \infty; \dot{K}_{n,\infty}^0), \tag{1.8}$$

$$u(\cdot) - e^{\cdot \Delta} u_0 \text{ is right continuous in } \dot{K}_{n,\infty}^0 \text{ on } (0, \infty), \tag{1.9}$$

where $X = \{u \in L^\infty(0, \infty; W\dot{K}_{n,\infty}^0) : \|u ; X\| < \infty, \operatorname{div} u = 0\}$,

$$\begin{aligned} \|u ; X\| &:= \sup_{t>0} \|u(t) ; W\dot{K}_{n,\infty}^0\| + \sup_{t>0} t^{(1-n/p)/2} \|u(t) ; \dot{K}_{p,\infty}^0\| \\ &\quad + \sup_{t>0} t^{1/2} \|u(t) ; L^\infty\| + \sup_{t>0} t^{1/2} \|\nabla u(t) ; W\dot{K}_{n,\infty}^0\| \\ &=: \|u ; X_1\| + \|u ; X_2\| + \|u ; X_3\| + \|u ; X_4\|. \end{aligned}$$

Moreover, if $u_0 \in W\dot{K}_{n,\infty}^0$, then we can show that

$$\begin{aligned} u(t) &\rightarrow u_0 \text{ in } W\dot{K}_{n,\infty}^0 \text{ as } t \searrow 0, \text{ and} \\ \lim_{t \searrow 0} t^{(1-n/p)/2} \|u(t) ; \dot{K}_{p,\infty}^0\| &= \lim_{t \searrow 0} t^{1/2} \|u(t) ; L^\infty\| = 0. \end{aligned}$$

Remark 1.3. 1. The solution above is in $\cap_{0 < T_1 < T_2 < T} L^\infty(\mathbb{R}^n \times (T_1, T_2))$ for every $T \in (0, \infty)$. Hence, thanks to the proof of Proposition 15.1 in [31], we obtain the smoothness $u \in C^\infty(\mathbb{R}^n \times (0, \infty))$.

2. We know the following inclusion relations from Theorem 1.6: for $2 \leq n < p < \sigma < \infty$,

$$W\dot{K}_{p,\infty}^{1-n/p} \hookrightarrow \dot{B}_{\sigma,\infty}^{-1+n/\sigma} \hookrightarrow BMO^{-1},$$

and for $q < \infty$

$$W\dot{K}_{n,q}^0 \hookrightarrow \dot{B}_{\sigma,\infty}^{-1+n/\sigma} \hookrightarrow BMO^{-1},$$

where $BMO^{-1} = \dot{F}_{\infty,2}^{-1}$ is defined by

$$\|f ; BMO^{-1}\| := \sup_Q \left(\int_Q \int_0^{l(Q)^2} |e^{t\Delta} f(x)|^2 dt dx \right)^{1/2},$$

where the supremum is taken over all cubes Q . Therefore if we replace $W\dot{K}_{n,\infty}^0$ with $W\dot{K}_{p,\infty}^{1-n/p}$ or $W\dot{K}_{n,q}^0$ in Theorem 1.3 where $p > n$ and $q < \infty$, then the initial data was dealt with in [24]. On the other hand, $W\dot{K}_{n,\infty}^0$ is not included in $\cup_{0 < p < \infty} \dot{B}_{p,\infty}^{-1+n/p}$, see Theorem 1.7.

As for uniqueness of global-in-time mild solutions, in the case $u_0 \in L^3(\mathbb{R}^3)$, it is well known that the solution is unique in the class $C([0, T]; L^3)$ even if initial data is large, see [18], [19] and [39]. Nevertheless, the existence of global solutions for large data in $L^3(\mathbb{R}^3)$ is an open problem. In this connection, Cannone [7] showed the existence of a global solution u in $C([0, \infty); L^3(\mathbb{R}^3))$ when an initial data $u_0 \in L^3(\mathbb{R}^3)$ is sufficiently small in the scale of the $\dot{B}_{p,\infty}^{-1+3/p}$ norm, where $3 < p \leq 6$. We can prove an analogous result for the uniqueness for large data in the weak Herz space setting.

Theorem 1.4 (uniqueness of global solution for large data). *Let $n \geq 3$. Let $u_0 \in W\dot{K}_{n,\infty}^0$ with $\operatorname{div} u_0 = 0$. If u and v are divergence free mild solutions of (I.E.) with initial data u_0 in the class $C([0, \infty); W\dot{K}_{n,\infty}^0)$ satisfying $u(t), v(t) \rightarrow u_0$ in $W\dot{K}_{n,\infty}^0$ as $t \searrow 0$, then $u = v$ on $[0, \infty)$.*

Also, as a corollary of the boundedness of B , see Remark 4.1, we can show a stability result of global solutions for small data, see [22].

Theorem 1.5 (stability of global solution for small data). *Let $n \geq 3$ and $u_0, v_0 \in W\dot{K}_{n,\infty}^0$. There exists $\delta > 0$ such that, if $\|u_0; W\dot{K}_{n,\infty}^0\|, \|v_0; W\dot{K}_{n,\infty}^0\| < \delta$, and*

$$\lim_{t \nearrow \infty} \|e^{t\Delta}(u_0 - v_0) ; W\dot{K}_{n,\infty}^0\| = 0,$$

then for the solutions u, v constructed in Theorem 1.3 with initial data u_0, v_0 , respectively, we have

$$\lim_{t \nearrow \infty} \|u(t) - v(t) ; W\dot{K}_{n,\infty}^0\| = 0.$$

Theorems 1.3, 1.4 and 1.5 are proved in Section 4. Furthermore, we discuss embeddings of weak Herz spaces into Besov spaces or bmo^{-1} . To prove a negative result, we use the characterization of homogeneous Besov spaces $\dot{B}_{p,\infty}^{-\alpha}$ ($\alpha > 0$), in terms of the heat semigroup $e^{t\Delta}$. The characterization we use is the following; for $1 \leq \sigma \leq \infty$, $\alpha > 0$ and $f \in \mathcal{S}'$:

$$\sup_{t>0} t^{\alpha/2} \|e^{t\Delta} f ; L^\sigma\| \approx \|f ; \dot{B}_{\sigma,\infty}^{-\alpha}\|. \tag{1.10}$$

See [7] or [46], for the details. Theorems 1.6 and 1.7 are proved in Section 5.

Theorem 1.6. (I) (The case $\sigma < \infty$)

(i) For $1 < p < \sigma < \infty$ and $0 < \alpha < n(1 - 1/p)$, we have

$$W\dot{K}_{p,\infty}^\alpha \hookrightarrow \dot{B}_{\sigma,\infty}^{-(\alpha+n(1/p-1/\sigma))}. \tag{1.11}$$

(ii) For $1 < p < \infty$ and $0 < \alpha < n(1 - 1/p)$, we have

$$\dot{K}_{p,\infty}^\alpha \hookrightarrow \dot{B}_{p,\infty}^{-\alpha}. \tag{1.12}$$

(iii) For $1 < p < \sigma < \infty$, one has

$$W\dot{K}_{p,\sigma}^0 \hookrightarrow \dot{B}_{\sigma,\infty}^{-n(1/p-1/\sigma)}. \tag{1.13}$$

(II) (The case $\sigma = \infty$)

(iv) For $1 < p < \infty$ and $0 \leq \alpha < n(1 - 1/p)$, we have

$$W\dot{K}_{p,\infty}^\alpha \hookrightarrow \dot{B}_{\infty,\infty}^{-(\alpha+n/p)}. \tag{1.14}$$

(v) For $0 \leq \alpha < n$, we have

$$\dot{K}_{\infty,\infty}^\alpha \hookrightarrow \dot{B}_{\infty,\infty}^{-\alpha}. \tag{1.15}$$

(vi) For $1 < p \leq \infty$ and $0 \leq \alpha \leq n(1 - 1/p)$, one has

$$W\dot{K}_{p,1}^\alpha \hookrightarrow \dot{B}_{\infty,\infty}^{-(\alpha+n/p)}. \tag{1.16}$$

(vii) We have

$$L^1 = \dot{K}_{1,1}^0 \hookrightarrow \dot{B}_{\infty,\infty}^{-n}. \tag{1.17}$$

(III)

(viii) For $n < p \leq \infty$ and $0 \leq \alpha < 1 - n/p$, one obtains

$$W\dot{K}_{p,\infty}^\alpha \hookrightarrow bmo^{-1}. \tag{1.18}$$

In (1.11) and (1.14), the lower bounds of α are sharp in the following sense.

Theorem 1.7. (i) (The case $\sigma < \infty$) For $1 < p < \sigma < \infty$ and $-n(1/p - 1/\sigma) < \alpha \leq 0$, the embedding

$$WK_{p,\infty}^\alpha \hookrightarrow \dot{B}_{\sigma,\infty}^{-(\alpha+n(1/p-1/\sigma))} \tag{1.19}$$

does not hold.

(ii) (The case $\sigma = \infty$) For $1 < p < \infty$ and $-n/p < \alpha < 0$, the embedding

$$WK_{p,\infty}^\alpha \hookrightarrow \dot{B}_{\infty,\infty}^{-(\alpha+n/p)} \tag{1.20}$$

does not hold.

Remark 1.4. 1. We give some examples of bmo^{-1} . For $n < p < \infty$, $0 \leq \alpha < 1 - n/p$ and $0 \leq \beta < 1$,

$$\sum_{k \in \mathbb{Z}} \frac{2^{-k\alpha}}{|x - x_k|^{n/p}} \chi_k(x), \frac{1}{|x|^\beta} \in \bigcup_{\substack{n < p < \infty \\ 0 \leq \alpha < 1 - n/p}} WK_{p,\infty}^\alpha,$$

where $x_k := \frac{3}{2}2^{k-1}e_1$ and $e_1 := (1, 0, \dots, 0)$. Then from (1.18), these functions belong to bmo^{-1} .

2. A special case of Theorem 1.7 is $WK_{p,\infty}^{1-n/p} \not\hookrightarrow \dot{B}_{\infty,\infty}^{-1}$, where $0 < p < n$. We remark that Bourgain and Pavlović [5] proved the ill posedness of the Navier-Stokes equations in $\dot{B}_{\infty,\infty}^{-1}$. Recently, in [49] he showed the ill posedness in $\dot{B}_{\infty,q}^{-1}$ for $q \in (2, \infty)$.

Many of our results above are consequences of the boundedness of convolution operators with some good functions on weak Herz spaces, see Propositions 2.1 and 2.2 in Section 2.

This paper is organized as follows. In Section 2 we recall the fundamental facts for Herz spaces and weak Herz spaces, and establish several estimates on weak Herz spaces, that is, the $WK_{p,q}^\alpha - WK_{\sigma,\delta}^\beta$ estimates of the heat semigroup $e^{t\Delta}$, the boundedness of a class of operators which contains the fractional integral operator I_α , and the critical estimates of the bilinear form B . In Section 3, we construct the unique local-in-time solution in weak Herz spaces and we prove the extension criterion Theorem 1.2. In Section 4, we prove the existence of global-in-time solutions on a scaling invariant space $WK_{n,\infty}^0$ for small data and the uniqueness of the solution in the class $C([0, \infty); WK_{n,\infty}^0)$ for smooth large data. Finally, in Section 5, we give the proof of Theorems 1.6 and 1.7.

2. PRELIMINARIES

Throughout this paper we use the following notation. \mathcal{S} and \mathcal{S}' denote the Schwartz spaces of rapidly decreasing smooth functions and tempered distributions, respectively. $A \lesssim B$ means $A \leq cB$ with a positive constant c . Also, $A \approx B$ means $c_1B \leq A \leq c_2B$ with positive constants c_1 and c_2 . In what follows, c denotes a constant that is independent of the functions involved, which may differ from line to line.

In this section, we recall the fundamental facts on Herz and weak Herz spaces and also establish some propositions, for example, the estimates of the heat semigroup and the boundedness of some operators on the weak Herz spaces. We begin with discussions for the relations between the Herz spaces or the weak Herz spaces and other function spaces.

Lebesgue spaces with power weights are special cases of Herz spaces; in the case $p < \infty$, $\dot{K}_{p,p}^\alpha = L^p(|x|^{\alpha p} dx)$, and $\|f ; \dot{K}_{\infty,\infty}^\alpha\| \approx \| |x|^\alpha f ; L^\infty \|$. Further, Herz spaces $\dot{K}_{p,\infty}^\alpha$ include Morrey spaces $\mathcal{M}_q^p(\mathbb{R}^n)$ and Lorentz spaces $L^{p,q}(\mathbb{R}^n)$; for $0 < q < p < \infty$, $L^{p,\infty} \hookrightarrow \mathcal{M}_q^p \hookrightarrow \dot{K}_{q,\infty}^{n(1/p-1/q)}$.

More precisely, we obtain the following.

Lemma 2.1. *Let $0 < q \leq p \leq \infty$. Then one has the embedding*

$$\mathcal{M}_q^p \hookrightarrow \dot{K}_{q,\infty}^{n(1/p-1/q)}.$$

Here, $\|f ; \mathcal{M}_q^p\| := \sup_Q |Q|^{1/p-1/q} \|f ; L^q(Q)\|$, where the supremum is taken over all cubes Q , and

$$\|f ; L^{p,q}\| := \left(\int_0^\infty t^{q/p} f^*(t)^q \frac{dt}{t} \right)^{1/q},$$

where f^* is the decreasing rearrangement of f . The usual modifications in the definitions above are made when $q = \infty$. The scaling invariant spaces $\dot{K}_{n,\infty}^0$ are not included in $L^{n,\infty}$. In fact, let $h(x) = 1$ if $2^k \leq x_i < 2^k + 1$ for some $k \in \mathbb{N}$ and all i , $h(x) = 0$ otherwise. Then, it is easy to see that $h \notin L^{n,\infty}$ and $h \in \dot{K}_{n,\infty}^0$.

On the other hand, $W\dot{K}_{n,\infty}^0$ is scaling invariant and also includes $L^{n,\infty}$. A typical member of $W\dot{K}_{n,\infty}^0$ is

$$f(x) := \sum_{k \in \mathbb{Z}} \frac{1}{|x - x_k|} \chi_{A_k}(x).$$

Then, we see that f does not belong to $\cup_{0 < p < \infty} \dot{B}_{p,\infty}^{-1+n/p}$, $\cup_{n < p \leq \infty} W\dot{K}_{p,\infty}^{1-n/p}$, $\dot{K}_{n,\infty}^0$ and $L^{n,\infty}$. Note that $L^{n,\infty} \hookrightarrow \cup_{n < p < \infty} \dot{B}_{p,\infty}^{-1+n/p}$. See Theorem 1.7 for

the proof of $f \notin \dot{B}_{p,\infty}^{-1+n/p}$. But the author knows no inclusion relations between Herz space and Morrey spaces, or weak Herz spaces and weak Morrey spaces which are scaling invariant, see [40] for the definition of weak Morrey spaces.

The following four lemmas are fundamental facts of homogeneous Herz spaces and weak Herz spaces. Because it is not hard to prove them, we omit the proofs.

Lemma 2.2. *Let $1 \leq p, q \leq \infty$ and $\alpha \in \mathbb{R}$.*

(i) $\mathcal{S} \hookrightarrow \dot{K}_{p,q}^\alpha \hookrightarrow W\dot{K}_{p,q}^\alpha$, provided that $q < \infty$ and $-n/p < \alpha < \infty$ or $q = \infty$ and $-n/p \leq \alpha < \infty$.

(ii) $\dot{K}_{p,q}^\alpha \hookrightarrow W\dot{K}_{p,q}^\alpha \hookrightarrow \mathcal{S}'$, provided that $q = 1$ and $-\infty < \alpha \leq n(1 - 1/p)$ or $1 < q \leq \infty$ and $-\infty < \alpha < n(1 - 1/p)$.

Lemma 2.3 ([41], Lemma 2.3). *Let $0 < p, q \leq \infty$, $\alpha \in \mathbb{R}$ and $f \in W\dot{K}_{p,q}^\alpha$. Then, $f \in L^r_{loc}(\mathbb{R}^n)$ for every r satisfying $\max(\alpha, 0)/n + 1/p < 1/r < \infty$.*

Lemma 2.4. *Let $0 < p_i, q_i \leq \infty$ and $\alpha_i \in \mathbb{R}$, ($i = 1, 2$). Then, if*

$$\frac{1}{p} = \frac{1}{p_1} + \frac{1}{p_2}, \quad \frac{1}{q} = \frac{1}{q_1} + \frac{1}{q_2} \quad \text{and} \quad \alpha = \alpha_1 + \alpha_2,$$

then

$$\|fg ; W\dot{K}_{p,q}^\alpha\| \leq c \|f ; W\dot{K}_{p_1,q_1}^{\alpha_1}\| \|g ; W\dot{K}_{p_2,q_2}^{\alpha_2}\|.$$

To construct the global solutions of (N-S), we need to know the parameters (p, q, α) for which $\dot{K}_{p,q}^\alpha$ and $W\dot{K}_{p,q}^\alpha$ are scaling invariant. From the lemma below, we see that the spaces $\dot{K}_{p,q}^{-1-n/p}$ and $W\dot{K}_{p,q}^{-1-n/p}$ are scaling invariant.

Lemma 2.5 ([41], Lemma 2.1). *For $0 < p, q \leq \infty$, $\alpha \in \mathbb{R}$ and $0 < \lambda < \infty$,*

$$\|f(\lambda \cdot) ; \dot{K}_{p,q}^\alpha\| \approx \lambda^{-(\alpha+n/p)} \|f ; \dot{K}_{p,q}^\alpha\|$$

and

$$\|f(\lambda \cdot) ; W\dot{K}_{p,q}^\alpha\| \approx \lambda^{-(\alpha+n/p)} \|f ; W\dot{K}_{p,q}^\alpha\|.$$

To establish the $W\dot{K}_{p,q}^\alpha - \dot{K}_{\sigma,q}^\beta$ or $W\dot{K}_{p,q}^\alpha - W\dot{K}_{\sigma,q}^\beta$ estimates of the heat semigroup $e^{t\Delta}$, we apply the following lemma to the Gaussian G , see Corollary 2.1 below.

Lemma 2.6. *Let $1 < p \leq \sigma \leq \infty$, $k, j \in \mathbb{Z}$, $1 + 1/\sigma = 1/p + 1/r$ and $\phi \in L^{r,1} \cap L^\infty$ with $|\phi(x)| \leq C_* |x|^{-m}$ for some $m \geq 0$ and $x \neq 0$.*

(i) *In the case $p < \sigma \leq \infty$, we have*

$$\|f_j * \phi ; L^\sigma(A_k)\| \leq c \|f_j ; L^{p,\infty}\|$$

$$\times \begin{cases} 2^{kn/\sigma} 2^{jn(1-1/p)} \min(2^{-km}, 1) =: R_1, & \text{if } j \leq k-2, \\ \min(2^{kn/\sigma} 2^{jn(1-1/p)}, 1) =: R_2, & \text{if } k-1 \leq j \leq k+1, \\ 2^{kn/\sigma} 2^{jn(1-1/p)} \min(2^{-jm}, 1) =: R_3, & \text{if } k+2 \leq j, \end{cases}$$

where $f_j := f\chi_{A_j}$ and a constant c depends on n, p, σ, ϕ and C_* only.

(ii) In the case $p = \sigma \leq \infty$, we have the same estimate for $\|f_j * \phi; L^{\sigma, \infty}(A_k)\|$ as above (i).

Proof. (i) In the case $j \leq k-2$, for $x \in A_k$ and $y \in A_j$, one has $|x-y| \gtrsim 2^k$. Then it is easy to see that

$$\|f_j * \phi; L^\sigma(A_k)\| \lesssim |A_k|^{1/\sigma} \min(2^{-km}, 1) \|f_j; L^1\|.$$

Since $\text{supp } f_j \subset A_j$, the inequality $\|f_j; L^1\| \leq p/(p-1)|A_j|^{1-1/p} \|f_j; L^{p, \infty}\|$ holds. Therefore we obtain the desired inequality in this case.

In the case $k-1 \leq j \leq k+1$, from the same argument as above, we have $\|f_j * \phi; L^\sigma(A_k)\| \lesssim 2^{kn/\sigma} 2^{jn(1-1/p)} \|f_j; L^{p, \infty}\|$. On the other hand, in the case $\sigma < \infty$, by using the stronger Young's inequality, see page 63 in [15], we have

$$\|f_j * \phi; L^\sigma(A_k)\| \leq \|f_j * \phi; L^\sigma(\mathbb{R}^n)\| \lesssim \|\phi; L^r\| \|f_j; L^{p, \infty}\|.$$

In the case $\sigma = \infty$, from the inequality of Hardy and Littlewood for rearrangements, [1], and the shift-invariance of the $L^{r,1}$ norm, we have

$$\|f_j * \phi; L^\infty(A_k)\| \lesssim \|\phi; L^{r,1}\| \|f_j; L^{p, \infty}\|.$$

The desired inequality in the case $k+2 \leq j$ can be showed by the same argument as the first case.

(ii) In this case, since the stronger Young's inequality is not available, the left-hand side is replaced by $L^{\sigma, \infty}$. □

The following proposition is a consequence of Lemma 2.6 and yields the $W\dot{K}_{p,q}^\alpha - \dot{K}_{p,q}^\alpha$ or $W\dot{K}_{p,q}^\alpha - W\dot{K}_{\sigma,q}^\beta$ estimates of the heat semigroup $e^{t\Delta}$ and the operator $\nabla e^{t\Delta}\mathbb{P}$, see below.

Proposition 2.1. *Let $1 < p \leq \sigma \leq \infty, 0 < q \leq \infty$ and $m \geq 0$. Suppose that $\phi \in L^{r,1} \cap L^\infty$, with $1 + 1/\sigma = 1/p + 1/r$, satisfies $|\phi(x)| \leq C_* |x|^{-m}$ for all $x \neq 0$. Then we obtain the following estimates.*

(i) In the case $p < \sigma$

$$\|f * \phi; \dot{K}_{\sigma,q}^\beta\| \leq c \|f; W\dot{K}_{p,q}^\alpha\|, \text{ and}$$

(ii) in the case $p = \sigma$

$$\|f * \phi; W\dot{K}_{p,q}^\beta\| \leq c \|f; W\dot{K}_{p,q}^\alpha\|$$

provided that one of the following cases holds:

- (1) $0 < q \leq 1$, $-n/\sigma < \beta \leq \alpha \leq n(1-1/p)$, $n(1-1/p) - \alpha + \beta + n/\sigma \leq m$ and $\beta + n/\sigma < m$,
 (2) $1 < q < \infty$, $-n/\sigma < \beta \leq \alpha < n(1-1/p)$ and $n(1-1/p) - \alpha + \beta + n/\sigma \leq m$,
 (3) $q = \infty$, $-n/\sigma \leq \beta \leq \alpha < n(1-1/p)$, $n(1-1/p) - \alpha + \beta + n/\sigma \leq m$ and $n(1-1/p) - \alpha < m$.

Proof. We prove the case (i) only. By using Lemma 2.6, we decompose

$$\|f * \phi ; \dot{K}_{\sigma,q}^\beta\| \leq c(\text{I} + \text{II} + \text{III}),$$

where

$$\begin{aligned} \text{I} &= \left(\sum_{k \in \mathbb{Z}} 2^{k\beta q} \left(\sum_{j=-\infty}^{k-2} \|f ; L^{p,\infty}(A_j)\| R_1 \right)^q \right)^{1/q}, \\ \text{II} &= \left(\sum_{k \in \mathbb{Z}} 2^{k\beta q} \left(\sum_{j=k-1}^{k+1} \|f ; L^{p,\infty}(A_j)\| R_2 \right)^q \right)^{1/q}, \\ \text{III} &= \left(\sum_{k \in \mathbb{Z}} 2^{k\beta q} \left(\sum_{j=k+2}^{\infty} \|f ; L^{p,\infty}(A_j)\| R_3 \right)^q \right)^{1/q}. \end{aligned}$$

We shall show the case (1). By the assumption $n(1-1/p) - \alpha + \beta + n/\sigma \leq m$ and $\beta \leq \alpha$, II can be estimated as follows:

$$\begin{aligned} \text{II} &\leq \left(\sum_{k \in \mathbb{Z}} 2^{k\beta q} \sum_{j=k-1}^{k+1} \|f ; L^{p,\infty}(A_j)\|^q R_2^q \right)^{1/q} \\ &\leq c \left(\sum_{j \in \mathbb{Z}} 2^{j\alpha q} \|f ; L^{p,\infty}(A_j)\|^q \min(2^{j(n(1-1/p) - \alpha + \beta + n/\sigma)q}, 2^{j(\beta - \alpha)q}) \right)^{1/q} \\ &\leq c \|f ; W\dot{K}_{p,q}^\alpha\|. \end{aligned}$$

To estimate I, we decompose again

$$\begin{aligned} \text{I} &\leq \left(\sum_{k=-\infty}^2 2^{k\beta q} \sum_{j=-\infty}^{k-2} \|f ; L^{p,\infty}(A_j)\|^q R_1^q \right)^{1/q} \\ &\quad + \left(\sum_{k=3}^{\infty} 2^{k\beta q} \sum_{j=-\infty}^0 \dots \right)^{1/q} + \left(\sum_{k=3}^{\infty} 2^{k\beta q} \sum_{j=1}^{k-2} \dots \right)^{1/q}. \end{aligned}$$

Because $-n/\sigma < \beta$ and $\alpha \leq n(1 - 1/p)$, we have

$$\begin{aligned} \text{1st term} &\leq \left(\sum_{j=-\infty}^0 2^{j\alpha q} \|f ; L^{p,\infty}(A_j)\|^q 2^{j(n(1-1/p)-\alpha)q} \sum_{k=-\infty}^2 2^{k(\beta+n/\sigma)q} \right) \\ &\leq c \left(\sum_{j=-\infty}^0 2^{j\alpha q} \|f ; L^{p,\infty}(A_j)\|^q 2^{j(n(1-1/p)-\alpha)q} \right)^{1/q} \leq c \|f ; W\dot{K}_{p,q}^\alpha\|. \end{aligned}$$

Also, since $\alpha \leq n(1 - 1/p)$ and $\beta + n/\sigma < m$, one has

$$\begin{aligned} \text{2nd term} &\leq \left(\sum_{k=3}^\infty 2^{k(\beta+n/\sigma-m)q} \right)^{\frac{1}{q}} \left(\sum_{j=-\infty}^0 2^{j\alpha q} \|f ; L^{p,\infty}(A_j)\|^q 2^{j(n(1-1/p)-\alpha)q} \right)^{\frac{1}{q}} \\ &\leq c \|f ; W\dot{K}_{p,q}^\alpha\|. \end{aligned}$$

Furthermore, since $n(1 - 1/p) - \alpha + \beta + n/\sigma \leq m$, we obtain

$$\begin{aligned} \text{3rd term} &= \left(\sum_{j=1}^\infty 2^{j\alpha q} \|f ; L^{p,\infty}(A_j)\|^q 2^{j(n(1-1/p)-\alpha)q} \sum_{k=j+2}^\infty 2^{k(\beta+n/\sigma-m)q} \right)^{1/q} \\ &\leq c \|f ; W\dot{K}_{p,q}^\alpha\|, \end{aligned}$$

which implies $I \leq c \|f ; W\dot{K}_{p,q}^\alpha\|$.

Going through a similar argument as above, one obtains

$$III \leq c \|f ; W\dot{K}_{p,q}^\alpha\|,$$

which completes the proof of the case (i).

Next we consider the case (3) with the same decompositions as above. By using $n(1 - 1/p) - \alpha + \beta + n/\sigma \geq 0$ and $\beta - \alpha \leq 0$, we have

$$\begin{aligned} II &= \sup_{k \in \mathbb{Z}} 2^{k\beta} \sum_{j=k-1}^{k+1} \|f ; L^{p,\infty}(A_j)\| \min(2^{kn/\sigma} 2^{jn(1-1/p)}, 1) \\ &\leq c \|f ; W\dot{K}_{p,\infty}^\alpha\| \sup_{k \in \mathbb{Z}} \min(2^{k(n(1-1/p)-\alpha+\beta+n/\sigma)}, 2^{k(\beta-\alpha)}) \leq c \|f ; W\dot{K}_{p,\infty}^\alpha\|. \end{aligned}$$

I is dominated by $\|f ; W\dot{K}_{p,\infty}^\alpha\|$ in the following way:

$$\begin{aligned} \text{1st term} &= \sup_{k \leq 2} 2^{k\beta} \sum_{j=-\infty}^{k-2} \|f ; L^{p,\infty}(A_j)\| 2^{kn/\sigma} 2^{jn(1-1/p)} \\ &\leq \|f ; W\dot{K}_{p,\infty}^\alpha\| \sup_{k \leq 2} 2^{k(\beta+n/\sigma)} \sum_{j=-\infty}^0 2^{j(n(1-1/p)-\alpha)} \leq c \|f ; W\dot{K}_{p,\infty}^\alpha\|, \end{aligned}$$

$$\begin{aligned} \text{2nd term} &= \sup_{k \geq 3} 2^{k\beta} \sum_{j=-\infty}^0 \|f ; L^{p,\infty}(A_j)\| 2^{k(n/\sigma-m)} 2^{jn(1-1/p)} \\ &\leq \|f ; \dot{W}K_{p,\infty}^\alpha\| \sup_{k \geq 3} 2^{k(\beta+n/\sigma-m)} \sum_{j=-\infty}^0 2^{j(n(1-1/p)-\alpha)} \leq c \|f ; \dot{W}K_{p,\infty}^\alpha\|, \end{aligned}$$

$$\begin{aligned} \text{3rd term} &= \sup_{k \geq 3} 2^{k\beta} \sum_{j=1}^{k-2} \|f ; L^{p,\infty}(A_j)\| 2^{k(n/\sigma-m)} 2^{jn(1-1/p)} \\ &\leq c \|f ; \dot{W}K_{p,\infty}^\alpha\| \sup_{k \geq 3} 2^{k(\beta+n/\sigma-m)} 2^{k(n(1-1/p)-\alpha)} \leq c \|f ; \dot{W}K_{p,\infty}^\alpha\|. \end{aligned}$$

Hence, we have $I \leq c \|f ; \dot{W}K_{p,\infty}^\alpha\|$. Using a similar argument as above, we can get the estimate

$$III \leq c \|f ; \dot{W}K_{p,\infty}^\alpha\|,$$

which completes the proof of the case (3).

The inequality in the case (2) follows from the interpolation. □

The next proposition says that if we restrict the condition of the weight parameters α and β , then we can improve the right-hand side of Proposition 2.1 with respect to the summation parameter q .

Proposition 2.2. *Let $1 < p \leq \sigma \leq \infty$, $0 < \delta < q \leq \infty$ and $m \geq 0$. Suppose that $\phi \in L^{r,1} \cap L^\infty$, with $1 + 1/\sigma = 1/p + 1/r$, satisfies $|\phi(x)| \leq C_* |x|^{-m}$ for all $x \neq 0$. If $-n/\sigma < \beta < \alpha < n(1 - 1/p) < m$, then we obtain the following estimates.*

(i) *In the case $p < \sigma$*

$$\|f * \phi ; \dot{K}_{\sigma,\delta}^\beta\| \leq c \|f ; \dot{W}K_{p,q}^\alpha\|, \text{ and}$$

(ii) *in the case $p = \sigma$*

$$\|f * \phi ; \dot{W}K_{p,\delta}^\beta\| \leq c \|f ; \dot{W}K_{p,q}^\alpha\|.$$

Proof. It suffices to prove the estimate in the cases $q = \infty$ and $0 < \delta < 1$. We omit the proof of this proposition, because it is not hard to complete the proof modifying the argument of the proof of Proposition 2.1. □

Using Propositions 2.1 and 2.2, we can investigate the behavior of the heat semigroup $e^{t\Delta}$ and $\nabla e^{t\Delta}\mathbb{P}$ in the weak Herz spaces. We use this corollary many times.

Corollary 2.1. *Let $1 < p \leq \sigma \leq \infty$, $0 < q, \delta \leq \infty$, $-n/\sigma \leq \beta \leq \alpha \leq n(1 - 1/p)$ and $j \in \{0, 1\}$.*

[I] *(In the case $q = \delta$) If the exponents satisfy the first two conditions of one of (1), (2) and (3) in Proposition 2.1, the following inequalities hold.*

(i) *In the case $p < \sigma$*

$$\|\nabla^j e^{t\Delta} f ; \dot{K}_{\sigma,q}^\beta\| \leq ct^{-j/2-(\alpha-\beta+n(1/p-1/\sigma))/2} \|f ; W\dot{K}_{p,q}^\alpha\|.$$

(ii) *In the case $p = \sigma$,*

$$\|\nabla^j e^{t\Delta} f ; W\dot{K}_{p,q}^\beta\| \leq ct^{-j/2-(\alpha-\beta)/2} \|f ; W\dot{K}_{p,q}^\alpha\|.$$

(iii) *In the case $p < \sigma$,*

$$\|\nabla^j e^{t\Delta} \mathbb{P}f ; \dot{K}_{\sigma,q}^\beta\| \leq ct^{-j/2-(\alpha-\beta+n(1/p-1/\sigma))/2} \|f ; W\dot{K}_{p,q}^\alpha\|.$$

(iv) *In the case $p = \sigma$, if $j = 1$ or $\beta < \alpha$, it follows that*

$$\|\nabla^j e^{t\Delta} \mathbb{P}f ; W\dot{K}_{p,q}^\beta\| \leq ct^{-j/2-(\alpha-\beta)/2} \|f ; W\dot{K}_{p,q}^\alpha\|.$$

[II] *In the case $q = \delta$, when $1 < p < \infty$ and $-n/p < \alpha < n(1 - 1/p)$, it follows that*

$$\|e^{t\Delta} \mathbb{P}f ; W\dot{K}_{p,q}^\alpha\| \leq c \|f ; W\dot{K}_{p,q}^\alpha\|.$$

[III] *In the case $\delta < q$, the following inequalities hold if $-n/\sigma < \beta < \alpha < n(1 - 1/p)$.*

(i) *In the case $p < \sigma$*

$$\|\nabla^j e^{t\Delta} f ; \dot{K}_{\sigma,\delta}^\beta\| \leq ct^{-j/2-(\alpha-\beta+n(1/p-1/\sigma))/2} \|f ; W\dot{K}_{p,q}^\alpha\|.$$

(ii) *In the case $p = \sigma$*

$$\|\nabla^j e^{t\Delta} f ; W\dot{K}_{p,\delta}^\beta\| \leq c t^{-j/2-(\alpha-\beta)/2} \|f ; W\dot{K}_{p,q}^\alpha\|.$$

(iii) *In the case $p < \sigma$*

$$\|\nabla^j e^{t\Delta} \mathbb{P}f ; \dot{K}_{\sigma,\delta}^\beta\| \leq c t^{-j/2-(\alpha-\beta+n(1/p-1/\sigma))/2} \|f ; W\dot{K}_{p,q}^\alpha\|.$$

(iv) *In the case $p = \sigma$*

$$\|\nabla^j e^{t\Delta} \mathbb{P}f ; W\dot{K}_{p,\delta}^\beta\| \leq c t^{-j/2-(\alpha-\beta)/2} \|f ; W\dot{K}_{p,q}^\alpha\|.$$

Proof. (I-i), (I-ii) The proofs of the inequalities are concluded from Lemma 2.5, Proposition 2.1 and the equation $e^{t\Delta} f = (f_{1/\sqrt{t}} * G)_{\sqrt{t}}$.

(I-iii) It is well known that the operator $e^{t\Delta} \mathbb{P}$ is a convolution operator with the Oseen kernel $\mathbb{K}_t(x) = t^{-n/2} \mathbb{K}(x/\sqrt{t})$ such that $|\nabla^m \mathbb{K}(x)| \leq c \langle x \rangle^{-(n+m)}$, see for example [31]. Then the inequality can be verified by applying Proposition 2.1 to \mathbb{K} .

(I-iv) If $j = 1$, then the inequality is showed by the same argument as above. We shall consider the case $\beta < \alpha$. When $p < \infty$ and $-n/p < \beta$, from Proposition 2.3 \mathbb{P} is bounded on $W\dot{K}_{p,q}^\beta$. Therefore, we have

$$\|e^{t\Delta}\mathbb{P}f ; W\dot{K}_{p,q}^\beta\| \leq c\|e^{t\Delta}f ; W\dot{K}_{p,q}^\beta\| \leq ct^{-(\alpha-\beta)/2}\|f ; W\dot{K}_{p,q}^\alpha\|.$$

When $p < \infty$ and $\beta = -n/p$, for two positive numbers ε and θ with $-n/p + \varepsilon < n(1 - 1/p)$ and $-n/p + \varepsilon + \theta = \alpha$, we have

$$\begin{aligned} \|e^{t\Delta}\mathbb{P}f ; W\dot{K}_{p,q}^{-n/p}\| &\leq ct^{-\varepsilon/2}\|e^{t/2\Delta}\mathbb{P}f ; W\dot{K}_{p,q}^{-n/p+\varepsilon}\| \\ &\leq ct^{-\varepsilon/2}\|e^{t/2\Delta}f ; W\dot{K}_{p,q}^{-n/p+\varepsilon}\| \leq ct^{-(\varepsilon+\theta)/2}\|f ; W\dot{K}_{p,q}^{-n/p+\varepsilon+\theta}\|. \end{aligned}$$

When $p = \infty$, there exists $r \in (1, \infty)$ with $-n/r < \beta < n(1 - 1/r)$. Hence, one obtains

$$\begin{aligned} \|e^{t\Delta}\mathbb{P}f ; W\dot{K}_{\infty,q}^\beta\| &\leq ct^{-n/(2r)}\|e^{t/2\Delta}\mathbb{P}f ; W\dot{K}_{r,q}^\beta\| \\ &\leq ct^{-n/(2r)}\|e^{t/2\Delta}f ; W\dot{K}_{r,q}^\beta\| \leq ct^{-n/(2r)}\|e^{t/2\Delta}f ; W\dot{K}_{\infty,q}^{\beta+n/r}\| \\ &\leq ct^{-(\alpha+\beta)/2}\|f ; W\dot{K}_{\infty,q}^\alpha\|. \end{aligned}$$

(II), (III) We omit the details. □

The following proposition is useful to establish estimates on the weak Herz spaces for many operators, for example, the Hardy-Littlewood maximal operator, Calderon-Zygmund operators and fractional integral operators.

Proposition 2.3. *Let $1 < p < n/\gamma$, $0 < q \leq \infty$, $0 \leq \gamma < n$ and $-n/p + \gamma < \alpha < n(1 - 1/p)$. Let $1/p - 1/r = \gamma/n$ and T_γ be a bounded operator from $L^{p,\infty}$ to $L^{r,\infty}$ with $\|T_\gamma ; L^{p,\infty} \rightarrow L^{r,\infty}\| \leq C_1$ satisfying*

$$|T_\gamma(f)(x)| \leq C_2 \int \frac{|f(y)|}{|x - y|^{n-\gamma}} dy$$

for $f \in L^1_{loc}$ with $x \notin \text{supp } f$. Then T_γ is also a bounded operator from $W\dot{K}_{p,q}^\alpha$ to $W\dot{K}_{r,q}^\alpha$ with $\|T_\gamma\| \leq c(C_1 + C_2)$.

Proof. We prove the case $q = \infty$ only:

$$\|T_\gamma(f) ; W\dot{K}_{r,\infty}^\alpha\| \leq c(C_1 + C_2)\|f ; W\dot{K}_{p,\infty}^\alpha\|.$$

We decompose

$$\|T_\gamma(f) ; W\dot{K}_{r,\infty}^\alpha\| \leq c \sup_{k \in \mathbb{Z}} 2^{k\alpha} \sup_{\lambda > 0} \lambda |\{x \in A_k ; \sum_{j=-\infty}^{k-2} |T_\gamma(|f\chi_j|)(x)| > \lambda\}|^{1/r}$$

$$\begin{aligned}
 &+ c \sup_{k \in \mathbb{Z}} 2^{k\alpha} \sup_{\lambda > 0} \lambda |\{x \in A_k; \sum_{j=k-1}^{k+1} |T_\gamma(|f\chi_j|)(x)| > \lambda\}|^{1/r} \\
 &+ c \sup_{k \in \mathbb{Z}} 2^{k\alpha} \sup_{\lambda > 0} \lambda |\{x \in A_k; \sum_{j=k+2}^{\infty} |T_\gamma(|f\chi_j|)(x)| > \lambda\}|^{1/r} \\
 &=: \text{I} + \text{II} + \text{III}.
 \end{aligned}$$

The boundedness of T_γ yields that $\text{II} \leq cC_1 \|f; W\dot{K}_{p,\infty}^\alpha\|$. Let $x \in A_k$. Since from the property of T_γ we have

$$\begin{aligned}
 \sum_{j=-\infty}^{k-2} T_\gamma(|f\chi_j|)(x) &\leq cC_2 2^{-k(n-\gamma)} \sum_{j=-\infty}^{k-2} \|f; L^1(A_j)\| \\
 &\leq cC_2 2^{-k(n-\gamma)} \sum_{j=-\infty}^{k-2} 2^{jn(1-1/p)} \|f; L^{(p,\infty)}(A_j)\| =: A,
 \end{aligned}$$

and hence, I is dominated by $C_2 \|f; W\dot{K}_{p,\infty}^\alpha\|$ in the following way:

$$\begin{aligned}
 \text{I} &= \sup_{k \in \mathbb{Z}} 2^{k\alpha} \sup_{0 < \lambda \leq A} \lambda |\{x \in A_k; \sum_{j=-\infty}^{k-2} T_\gamma(|f\chi_j|)(x) > \lambda\}|^{1/r} \\
 &\leq cC_2 \sup_{k \in \mathbb{Z}} 2^{k(\alpha+n/r-n+\gamma)} \sum_{j=-\infty}^{k-2} 2^{jn(1-1/p)} \|f; L^{(p,\infty)}(A_j)\| \leq cC_2 \|f; W\dot{K}_{p,\infty}^\alpha\|.
 \end{aligned}$$

It is not hard to show that $\text{III} \leq cC_2 \|f; W\dot{K}_{p,\infty}^\alpha\|$ by the same argument as above. □

The final proposition is a generalization of [31, Lemma 27.6] in the setting of the weak Herz spaces and the proof is based on that of [31, Lemma 27.6]. The proposition is applied in the proof of Theorems 1.4 and 1.5 in Section 4.

Proposition 2.4. *Let $n \geq 2$, $1 < p < n$, $1 \leq q \leq \infty$ and $1 - n/p < \alpha < n(1 - 1/p)$. For a positive $\beta > 0$ satisfying*

$$0 < \beta < \min\left(\frac{1}{2}, \frac{n}{2}\left(\frac{1}{p} - \frac{1}{n}\right), \frac{n}{2}\left(1 - \frac{1}{p} + \frac{1}{n}\right), \frac{1}{2}\left(\alpha - \left(1 - \frac{n}{p}\right)\right)\right),$$

suppose that $1/r_1 = 1/r + 2\beta/n$, $1/r_2 = 1/r - 2\beta/n$ and $1/p - 1/r = 1/n$. Then, we have that, for every $j, k \in \{1, \dots, n\}$

$$\left\| \int_0^\infty (-\Delta)^{1/2} e^{t\Delta} \mathbb{P}_{j,k} f(t) dt ; (W\dot{K}_{r_1,q}^\alpha, W\dot{K}_{r_2,q}^\alpha)_{1/2,\infty} \right\|$$

$$\leq c \|f ; L^\infty(0, \infty; W\dot{K}_{p,q}^\alpha)\|,$$

with a constant c independent of f . In particular, in the case $q = \infty$, we have

$$\| \int_0^\infty (-\Delta)^{1/2} e^{t\Delta} \mathbb{P}_{j,k} f(t) dt ; W\dot{K}_{r,\infty}^\alpha \| \leq c \|f ; L^\infty(0, \infty; W\dot{K}_{p,\infty}^\alpha)\|.$$

Proof. Let $A > 0$ and decompose

$$\begin{aligned} F &:= \int_0^\infty (-\Delta)^{1/2} e^{t\Delta} \mathbb{P}_{j,k} f(t) dt = \int_0^A \cdots dt + \int_A^\infty \cdots dt \\ &= \int_0^A (-\Delta)^{1-\beta} e^{t\Delta} \mathbb{P}_{j,k} (-\Delta)^{-(1/2-\beta)} f(t) dt \\ &\quad + \int_A^\infty (-\Delta)^{1+\beta} e^{t\Delta} \mathbb{P}_{j,k} (-\Delta)^{-(\beta+1/2)} f(t) dt =: G_A + H_A. \end{aligned}$$

Note that $1/p - 1/r_1 = 2(1/2 - \beta)/n$, $1/p - 1/r_2 = 2(1/2 + \beta)/n$ and $-n/r_i < \alpha < n(1 - 1/p)$ for $i = 1, 2$. Therefore, applications of Proposition 2.3 give the following estimates:

$$\begin{aligned} \|G_A ; W\dot{K}_{r_1,q}^\alpha\| &\leq \int_0^A \|(-\Delta)^{1-\beta} e^{t\Delta} \mathbb{P}_{j,k} I_{2(1/2-\beta)} f(t) ; W\dot{K}_{r_1,q}^\alpha\| dt \\ &\leq c \int_0^A t^{\beta-1} \|e^{\frac{t}{2}\Delta} \mathbb{P}_{j,k} I_{2(1/2-\beta)} f(t) ; W\dot{K}_{r_1,q}^\alpha\| dt \\ &\leq c \int_0^A t^{\beta-1} \|I_{2(1/2-\beta)} f(t) ; W\dot{K}_{r_1,q}^\alpha\| dt \\ &\leq c \int_0^A t^{\beta-1} \|f(t) ; W\dot{K}_{p,q}^\alpha\| dt \\ &\leq cA^\beta \|f ; L^\infty(0, \infty; W\dot{K}_{p,q}^\alpha)\|, \end{aligned}$$

where I_s is the fractional integral operator which satisfies the conditions of Proposition 2.3. Here we have used the boundedness of the convolution operator $(\Delta^{1-\beta} G)_{\sqrt{t}} *$ and $e^{t\Delta} \mathbb{P}_{j,k}$ on $W\dot{K}_{r_1,q}^\alpha$. Also, by a similar argument to above, we can obtain

$$\|H_A ; W\dot{K}_{r_2,q}^\alpha\| \leq cA^{-\beta} \|f ; L^\infty(0, \infty; W\dot{K}_{p,q}^\alpha)\|.$$

Then, we reach the desired inequality in the following way:

$$\text{LHS} = \sup_{t>0} t^{-1/2} K(t, F; W\dot{K}_{r_1,q}^\alpha, W\dot{K}_{r_2,q}^\alpha)$$

$$\begin{aligned} &\leq \sup_{t>0} t^{-1/2} (\|G_{t^{1/(2\beta)}} ; W\dot{K}_{r_1,q}^\alpha\| + t \|H_{t^{1/(2\beta)}} ; W\dot{K}_{r_2,q}^\alpha\|) \\ &\leq c \|f ; L^\infty(0, \infty; W\dot{K}_{p,q}^\alpha)\| = \text{RHS}. \end{aligned}$$

It remains to show the continuous inclusion $(W\dot{K}_{r_1,\infty}^\alpha, W\dot{K}_{r_2,\infty}^\alpha)_{1/2, \infty} \hookrightarrow W\dot{K}_{r,\infty}^\alpha$. Once we know the estimate

$$\|g\chi_k ; L^{(r,\infty)}\| \leq c2^{-k\alpha} \|g ; (W\dot{K}_{r_1,\infty}^\alpha, W\dot{K}_{r_2,\infty}^\alpha)_{1/2, \infty}\|,$$

the inclusion is immediately verified. By using the real interpolation theory for Lorentz spaces, we get the estimate as follows:

$$\begin{aligned} \|g\chi_k ; L^{(r,\infty)}\| &\approx \|g\chi_k ; (L^{(r_1,\infty)}, L^{(r_2,\infty)})_{1/2,\infty}\| \\ &= \sup_{\lambda>0} \lambda^{-\frac{1}{2}} \inf_{g\chi_k=g^1+g^2} (\|g^1 ; L^{(r_1,\infty)}\| + \lambda\|g^2 ; L^{(r_2,\infty)}\|) \\ &\leq \sup_{\lambda>0} \lambda^{-\frac{1}{2}} \inf_{\substack{g\chi_k=g^1+g^2 \\ g^1=g^2=0 \text{ on } A_k^c}} (\|g^1 ; L^{(r_1,\infty)}\| + \lambda\|g^2 ; L^{(r_2,\infty)}\|) \\ &= \sup_{\lambda>0} \lambda^{-\frac{1}{2}} \inf_{g=g^1+g^2} (\|g^1\chi_k ; L^{(r_1,\infty)}\| + \lambda\|g^2\chi_k ; L^{(r_2,\infty)}\|) \\ &\leq 2^{-k\alpha} \sup_{\lambda>0} \lambda^{-\frac{1}{2}} \inf_{g=g^1+g^2} (\|g^1 ; W\dot{K}_{r_1,\infty}^\alpha\| + \lambda\|g^2 ; W\dot{K}_{r_2,\infty}^\alpha\|) \\ &= 2^{-k\alpha} \|g ; (W\dot{K}_{r_1,\infty}^\alpha, W\dot{K}_{r_2,\infty}^\alpha)_{1/2,\infty}\|. \quad \square \end{aligned}$$

Remark 2.1. In the proof above, we have used the one-sided inclusion

$$(W\dot{K}_{r_1,\infty}^\alpha, W\dot{K}_{r_2,\infty}^\alpha)_{1/2, \infty} \hookrightarrow W\dot{K}_{r,\infty}^\alpha$$

only. The author does not know whether the reverse inclusion is true or not. On the other hand, for the interpolation with respect to the weight parameter α , we have

$$(W\dot{K}_{p,\infty}^{\alpha_1}, W\dot{K}_{p,\infty}^{\alpha_2})_{\theta, \infty} = W\dot{K}_{p,\infty}^\alpha$$

where $\alpha = (1 - \theta)\alpha_1 + \theta\alpha_2$ and $0 < p \leq \infty$. Then it is natural to try to prove Proposition 2.4 by using interpolation with respect to the parameter α , but we do not have the estimate $W\dot{K}_{p,\infty}^\alpha - W\dot{K}_{\sigma,\infty}^\beta$ of the heat semigroup in the case $\alpha < \beta$.

In order to solve (I.E.), we use the following Picard contraction principle. For example, see [31] for the proof.

Proposition 2.5 (The Picard contraction principle). *Let E be a Banach space and let B be a bounded bilinear transform $E \times E$ to E satisfying*

$$\|B(e, f) ; E\| \leq C_B \|e ; E\| \|f ; E\|.$$

Then, if $0 < \delta < (4C_B)^{-1}$ and $e_0 \in E$ satisfies $\|e_0 ; E\| \leq \delta$, the equation $e = e_0 - B(e, e)$ has a solution with $\|e ; E\| \leq 2\delta$. This solution is unique in the closed ball $\bar{B}(0, 2\delta)$. Moreover, the solution continuously depends on e_0 ; if $\|f_0 ; E\| \leq \delta$, $f = f_0 - B(f, f)$ and $\|f ; E\| \leq 2\delta$, then $\|e - f ; E\| \leq (1 - 4C_B\delta)^{-1} \|e_0 - f_0 ; E\|$.

3. PROOF OF THEOREMS 1.1 AND 1.2

3.1. Proof of Theorem 1.1. The constant $T > 0$ will be chosen later. To prove Theorem 1.1, we divide the proof into four Steps.

Step 1: *The bilinear form B is a map from $X_T \times X_T$ to X_T and has the estimate*

$$\|B(u, v) ; X_T\| \leq C_B T^{(1-n/p-\alpha)/2} \|u ; X_T\| \|v ; X_T\|.$$

The claim is an immediate consequence of Corollary 2.1. Indeed, by using the corollary, we have the following three estimates. Firstly, because $1 - n/p - \alpha > 0$, one has that, for $0 < t \leq T$,

$$\begin{aligned} \|B(u, v)(t) ; \dot{K}_{p,\infty}^\alpha\| &\leq \int_0^t \|\nabla e^{(t-s)\Delta} \mathbb{P}(u \otimes v)(s) ; \dot{K}_{p,\infty}^\alpha\| ds \\ &\leq c \int_0^t (t-s)^{-(1+n/p+\alpha)/2} \|u \otimes v(s) ; W\dot{K}_{p/2,\infty}^{2\alpha}\| ds \\ &\leq c \|u ; X_{T,1}\| \|v ; X_{T,1}\| \int_0^t s^{-(1+n/p+\alpha)/2} ds \\ &\leq (C_B/2) T^{(1-n/p-\alpha)/2} \|u ; X_T\| \|v ; X_T\|, \end{aligned}$$

which implies $\|B(u, v) ; X_{T,1}\| \leq (C_B/2) T^{(1-n/p-\alpha)/2} \|u ; X_T\| \|v ; X_T\|$. Secondly, from the same reason as above, one can show that $\|B(u, v) ; X_{T,2}\| \leq (C_B/2) T^{(1-\alpha-n/p)/2} \|u ; X_T\| \|v ; X_T\|$ in the following way:

$$\begin{aligned} \|\nabla B(u, v)(t) ; \dot{K}_{p,\infty}^\alpha\| &\leq \int_0^t \|\nabla e^{(t-s)\Delta} \mathbb{P}(u \cdot \nabla)v(s) ; \dot{K}_{p,\infty}^\alpha\| ds \\ &\leq c \int_0^t (t-s)^{-(1+n/p+\alpha)/2} \|(u \cdot \nabla)v(s) ; W\dot{K}_{p/2,\infty}^{2\alpha}\| ds \\ &\leq c \|u ; X_{T,1}\| \|v ; X_{T,2}\| \int_0^t (t-s)^{-(1+n/p+\alpha)/2} s^{-1/2} ds \end{aligned}$$

$$\leq (C_B/2)t^{-(n/p+\alpha)/2}\|u ; X_T\| \|v ; X_T\|.$$

As a result, we get the desired estimate.

Step 2: *We verify that*

$$\|e^{\cdot\Delta}u_0 ; X_T\| \leq C_0\|u_0 ; W\dot{K}_{p,\infty}^\alpha\|.$$

Our hypothesis guarantees the following estimates will hold:

$$\begin{aligned} \|e^{t\Delta}u_0 ; W\dot{K}_{p,\infty}^\alpha\| &\leq C_0/2\|u_0 ; W\dot{K}_{p,\infty}^\alpha\|, \\ \|\nabla e^{t\Delta}u_0 ; W\dot{K}_{p,\infty}^\alpha\| &\leq (C_0/2)t^{-1/2}\|u_0 ; W\dot{K}_{p,\infty}^\alpha\|. \end{aligned}$$

Therefore, the above estimate holds.

Now, we define T as

$$\left(\frac{1}{8C_B C_0\|u_0 ; W\dot{K}_{p,\infty}^\alpha\|}\right)^{2/(1-n/p-\alpha)} = \frac{C^*}{\|u_0 ; W\dot{K}_{p,\infty}^\alpha\|^{2/(1-n/p-\alpha)}}.$$

Then, because T satisfies $C_0\|u_0 ; W\dot{K}_{p,\infty}^\alpha\| < (4C_B T^{(1-n/p-\alpha)/2})^{-1}$, we can find a solution $u \in X_T$ to (I.E.) and the solution continuously depends on the initial data u_0 , from the Picard contraction principle. Note that the constant C^* , in the definition of T above, depends on p, n and α only. Then the solution automatically satisfies the condition (1.1). In fact, we have

$$\begin{aligned} \|e^{t\Delta}u_0 ; L^\infty\| &\leq ct^{-(\alpha+n/p)/2}\|u_0 ; W\dot{K}_{p,\infty}^\alpha\| \text{ and} \\ \|B(u, u)(t) ; L^\infty\| &\leq c \int_0^t (t-s)^{-(\alpha+n/p)}\|u(s) ; W\dot{K}_{p,\infty}^\alpha\| \|\nabla u(s) ; W\dot{K}_{p,\infty}^\alpha\| ds \\ &\leq c\|u ; X_{T,1}\|\|u ; X_{T,2}\| \int_0^t (t-s)^{-(\alpha+n/p)} s^{-1/2} ds \\ &\leq ct^{1/2-(\alpha+n/p)}\|u ; X_T\|^2, \end{aligned}$$

which implies

$$\sup_{0 < t \leq T} t^{(\alpha+n/p)/2}\|B(u, u)(t) ; L^\infty\| \leq cT^{(1-\alpha-n/p)/2}\|u ; X_T\|^2 < \infty.$$

Step 3: *In this step, we check the properties (1.2), (1.3), (1.4) and (1.5) for the solution u .*

But, because these assertions can be verified by the same arguments as Steps 3, 4 and 5 in the proof of Theorem 1.3, we omit the details.

Step 4: *Finally, the solution u is the only solution of (I.E.) in the class $L^\infty(0, T; W\dot{K}_{p,\infty}^\alpha)$.*

Let v also be a solution of (I.E.) with the initial data u_0 in the class $L^\infty(0, T; W\dot{K}_{p,\infty}^\alpha)$. Put $w = u - v = -B(u, u) + B(v, v) = -B(u, w) -$

$B(w, v)$. An application of Corollary 2.1 and our condition $-n/p \leq 0 \leq \alpha < 1 - n/p$ yield that, for $0 < t \leq T_0 < T$,

$$\begin{aligned} \|w(t); \dot{W}K_{p,\infty}^\alpha\| &\leq c \int_0^t (t-s)^{-(1+n/p+\alpha)/2} \|u \otimes w(s) + w \otimes v(s); \dot{W}K_{p/2,\infty}^{2\alpha}\| ds \\ &\leq c \left(\sup_{0 < s \leq T_0} \|u(s); \dot{W}K_{p,\infty}^\alpha\| + \sup_{0 < s \leq T_0} \|v(s); \dot{W}K_{p,\infty}^\alpha\| \right) \\ &\quad \times \sup_{0 < s \leq T_0} \|w(s); \dot{W}K_{p,\infty}^\alpha\| T_0^{(1-n/p-\alpha)/2}. \end{aligned}$$

Here, if we take T_0 as

$$c \left(\sup_{0 < s \leq T_0} \|u(s); \dot{W}K_{p,\infty}^\alpha\| + \sup_{0 < s \leq T_0} \|v(s); \dot{W}K_{p,\infty}^\alpha\| \right) T_0^{2/(1-n/p-\alpha)} \leq 1/2,$$

then we can get $w = 0$ on $(0, T_0]$; i.e., $u = v$ on $(0, T_0]$. By a similar method as Proposition 3.1 in [27], we have $u = v$ on $(0, T)$ and the proof of Theorem 1.1 is completed. \square

3.2. Proof of Theorem 1.2. Next we give the proof of Theorem 1.2. In the proof, the following bilinear estimate plays an important role.

Theorem 3.1 ([47]). *Let $1 < p < \infty$, $1 \leq q \leq \infty$, $-n/p < \alpha < \infty$ and $m \in \mathbb{N}$. There exists a constant c such that, for any $f \in \dot{K}_{p,q}^\alpha$ with $\nabla^m f \in \dot{K}_{BMO,\infty}^0$ and $g \in \dot{K}_{p,q}^\alpha$ with $\nabla^m g \in \dot{K}_{BMO,\infty}^0$,*

$$\|f \nabla^m g; \dot{K}_{p,q}^\alpha\| \leq c \left(\|f; \dot{K}_{p,q}^\alpha\| \|\nabla^m g; \dot{K}_{BMO,\infty}^0\| + \|\nabla^m f; \dot{K}_{BMO,\infty}^0\| \|g; \dot{K}_{p,q}^\alpha\| \right).$$

We can conclude that $\sup_{T^*/2 \leq t < T^*} \|u(t); \dot{K}_{p,\infty}^\alpha\|$ is finite by applying Theorem 3.1 to u and ∇u , as follows: for $T^*/2 \leq t < T^*$

$$\begin{aligned} \|u(t); \dot{K}_{p,\infty}^\alpha\| &\leq c \|u_0; \dot{K}_{p,\infty}^\alpha\| + c \int_0^t \|(u \cdot \nabla)u(s); \dot{K}_{p,\infty}^\alpha\| ds \\ &\leq c \|u_0; \dot{K}_{p,\infty}^\alpha\| + c \int_0^t \|u(s); \dot{K}_{p,\infty}^\alpha\| \|\nabla u(s); \dot{K}_{BMO,\infty}^0\| ds, \end{aligned}$$

and from Gronwall's inequality,

$$\|u(t); \dot{K}_{p,\infty}^\alpha\| \leq c \|u_0; \dot{K}_{p,\infty}^\alpha\| \exp \left(c \int_0^{T^*} \|\nabla u(s); \dot{K}_{BMO,\infty}^0\| ds \right) < \infty.$$

Now we suppose that, for every $\tilde{T} > T^*$, u would not be a mild solution of (N-S) with initial data u_0 in the class $L^\infty(0, \tilde{T}; \dot{K}_{p,\infty}^\alpha)$. Then, from Remark

1.1 we know that, for any positive number $\tau < T^*$,

$$\frac{C}{\|u(\tau) ; \dot{K}_{p,\infty}^\alpha\|^{2/(1-n/p-\alpha)}} \leq T^* - \tau$$

with a constant C independent of τ . Therefore, positivity of $1 - n/p - \alpha$ yields

$$\limsup_{\tau \nearrow T^*} \|u(\tau) ; \dot{K}_{p,\infty}^\alpha\| \geq \limsup_{\tau \nearrow T^*} \left(\frac{C}{T^* - \tau} \right)^{(1-n/p-\alpha)/2} = \infty,$$

which contradicts the fact that $u \in L^\infty(0, T^*; \dot{K}_{p,\infty}^\alpha)$. □

4. PROOF OF THEOREMS 1.3, 1.4 AND 1.5

4.1. Proof of Theorem 1.3. We divide the proof into five steps and begin with an estimate of the bilinear form B .

Step 1: *The bilinear form B is the map from $X \times X$ to X and has the estimate*

$$\|B(u, v) ; X\| \leq C_B \|u ; X\| \|v ; X\|.$$

From Corollary 2.1, we know that our hypothesis guarantees the following inequalities to hold: for $1/\sigma = 1/n + 1/p$

$$\begin{aligned} \|B(u, v)(t) ; \dot{K}_{n,\infty}^0\| &\leq c \int_0^t \|e^{(t-s)\Delta}(u \cdot \nabla)v(s) ; \dot{K}_{n,\infty}^0\| ds \\ &\leq c \int_0^t (t-s)^{-n/2p} \|(u \cdot \nabla)v(s) ; W\dot{K}_{\sigma,\infty}^0\| ds \\ &\leq c \|u ; X_2\| \|v ; X_4\| \int_0^t (t-s)^{-n/2p} s^{-(2-n/p)/2} ds \leq C_B/4 \|u ; X\| \|v ; X\|, \end{aligned}$$

which implies $\|B(u, v) ; X_1\| \leq C_B/4 \|u ; X\| \|v ; X\|$. As for the X_2 norm of $B(u, v)$, an application of the same corollary yields

$$\begin{aligned} \|B(u, v)(t) ; \dot{K}_{p,\infty}^0\| &\leq \int_0^t \|\nabla e^{(t-s)\Delta} \mathbb{P}(u \otimes v)(s) ; \dot{K}_{p,\infty}^0\| ds \\ &\leq c \int_0^t (t-s)^{-1/2-n(2/p-1/p)/2} \|u \otimes v(s) ; W\dot{K}_{p/2,\infty}^0\| ds \\ &\leq c \|u ; X_2\| \|v ; X_2\| \int_0^t (t-s)^{-1/2-n/2p} s^{-(1-n/p)} ds \\ &\leq (C_B/4) t^{-(1-n/p)/2} \|u ; X\| \|v ; X\|, \end{aligned}$$

and, as a consequence, we have $\|B(u, v) ; X_2\| \leq C_B/4\|u ; X\| \|v ; X\|$. Furthermore, to estimate the X_3 norm of $B(u, v)$, we use Corollary 2.1 again and obtain

$$\begin{aligned} \|B(u, v)(t); L^\infty\| &\leq \int_0^t \|e^{(t-s)\Delta}\mathbb{P}(u \cdot \nabla)v(s); L^\infty\| ds \\ &\leq c \int_0^t \min\left((t-s)^{-n/2\sigma}\|(u \cdot \nabla)v(s); W\dot{K}_{\sigma,\infty}^0\|, (t-s)^{-1/2}\|u \otimes v(s); L^\infty\|\right) ds \\ &\leq c\|u; X\|\|v; X\| \int_0^t \min\left((t-s)^{-n/2\sigma} s^{-1+n/2p}, (t-s)^{-1/2} s^{-1}\right) ds \\ &\leq (C_B/4)t^{-1/2}\|u; X\|\|v; X\|. \end{aligned}$$

Finally, the fourth term consisting of the X norm of $B(u, v)$ can be bounded by the product of the X norms of u and v as follows:

$$\begin{aligned} \|\nabla B(u, v)(t) \dot{K}_{n,\infty}^0\| &\leq \int_0^t \|\nabla e^{(t-s)\Delta}\mathbb{P}(u \cdot \nabla)v(s) ; \dot{K}_{n,\infty}^0\| ds \\ &\leq c \int_0^t (t-s)^{-1/2-n(1/\sigma-1/n)/2}\|(u \cdot \nabla)v(s) W\dot{K}_{\sigma,\infty}^0\| ds \\ &\leq c\|u X_2\| \|v ; X_4\| \int_0^t (t-s)^{-(1+n/p)/2} s^{-(2-n/p)/2} ds \\ &\leq (C_B/4)t^{-1/2}\|u ; X\| \|v ; X\|, \end{aligned}$$

hence one has $\|B(u, v) ; X_4\| \leq C_B/4\|u ; X\| \|v ; X\|$. By putting the three estimates above together, the desired norm inequality is verified and we finish Step 1.

Step 2: *In this step, we shall check*

$$\|e^{\cdot\Delta}u_0 ; X\| \leq C_0\|u_0 ; W\dot{K}_{n,\infty}^0\|.$$

From our assumption and applications of Corollary 2.1, we conclude the claim in the following way:

$$\begin{aligned} \|e^{t\Delta}u_0 ; W\dot{K}_{n,\infty}^0\| &\leq C_0/4\|u_0 ; W\dot{K}_{n,\infty}^0\|, \\ \|e^{t\Delta}u_0 ; \dot{K}_{p,\infty}^0\| &\leq (C_0/4) t^{-(1-n/p)/2}\|u_0 ; W\dot{K}_{n,\infty}^0\|, \\ \|e^{t\Delta}u_0 ; L^\infty\| &\leq (C_0/4)t^{-1/2}\|u_0 ; W\dot{K}_{n,\infty}^0\| \text{ and} \\ \|\nabla e^{t\Delta}u_0 ; W\dot{K}_{n,\infty}^0\| &\leq (C_0/4) t^{-1/2}\|u_0 ; W\dot{K}_{n,\infty}^0\|. \end{aligned}$$

Therefore, from the Picard contraction principle, we can find a solution $u \in X$ to (I.E.) which continuously depends on u_0 .

Step 3: *In this step, we show the continuity of u .*

Since $e^{(t\pm\tau)\Delta}u_0 - e^{t\Delta}u_0 = u_0 * (G_{\sqrt{t\pm\tau}} - G_{\sqrt{t}})$, by using Proposition 2.3, one has

$$\|e^{(t\pm\tau)\Delta}u_0 - e^{t\Delta}u_0; W\dot{K}_{n,\infty}^0\| \leq c|G_{\sqrt{t\pm\tau}} - G_{\sqrt{t}}|_{\mathcal{S}} \|u_0; W\dot{K}_{n,\infty}^0\|$$

which implies that $e^{t\Delta}u_0 \in C((0, \infty); W\dot{K}_{n,\infty}^0)$, where $|\cdot|_{\mathcal{S}}$ denotes a seminorm of \mathcal{S} . To show the right continuity of B in $\dot{K}_{n,\infty}^0$ on $(0, \infty)$, we write, for $t > 0$ and $\tau > 0$,

$$\begin{aligned} & B(u, u)(t + \tau) - B(u, u)(t) \\ &= \int_0^t (e^{(t+\tau-s)\Delta}\mathbb{P}(u \cdot \nabla)u(s) - e^{(t-s)\Delta}\mathbb{P}(u \cdot \nabla)u(s))ds \\ &+ \int_t^{t+\tau} e^{(t+\tau-s)\Delta}\mathbb{P}(u \cdot \nabla)u(s)ds =: \text{I}_+ + \text{II}_+. \end{aligned}$$

Because we have, for sufficiently small $\tau > 0$,

$$\begin{aligned} & \|e^{(t+\tau-s)\Delta}\mathbb{P}(u \cdot \nabla)u(s); \dot{K}_{n,\infty}^0\| \\ & \leq c(t-s)^{-n/2p} s^{-1+n/2p} \|u; X_2\| \|u; X_4\| \in L^1(0, t), \end{aligned}$$

Lebesgue's convergence theorem yields that $\lim_{\tau \searrow 0} \|\text{I}_+; \dot{K}_{n,\infty}^0\| = 0$. The second term also tends to 0 as $\tau \searrow 0$:

$$\begin{aligned} \|\text{II}_+; \dot{K}_{n,\infty}^0\| & \leq \int_t^{t+\tau} \|e^{(t+\tau-s)\Delta}\mathbb{P}(u \cdot \nabla)u(s); \dot{K}_{n,\infty}^0\| ds \\ & \leq c\|u; X_2\| \|u; X_4\| \int_t^{t+\tau} (t+\tau-s)^{-n/2p} s^{-1+n/2p} ds \rightarrow 0 \text{ as } \tau \searrow 0. \end{aligned}$$

Therefore, (1.9) is proved. Next, we check the left continuity of B in $W\dot{K}_{n,\infty}^0$ on $(0, \infty)$. To this end, we write, for $t > 0$ and $\tau > 0$,

$$\begin{aligned} & B(u, u)(t - \tau) - B(u, u)(t) \\ &= \int_0^{t-\tau} e^{(t-\tau-s)\Delta}\mathbb{P}(u \cdot \nabla)u(s) - e^{(t-s)\Delta}\mathbb{P}(u \cdot \nabla)u(s)ds \\ &+ \int_{t-\tau}^t e^{(t-s)\Delta}\mathbb{P}(u \cdot \nabla)u(s)ds =: \text{I}_- + \text{II}_-. \end{aligned}$$

Since one obtains the estimate

$$\|e^{(t-\tau-s)\Delta}\mathbb{P}(u \cdot \nabla)u(s); W\dot{K}_{n,\infty}^0\|$$

$$\begin{aligned} &\leq c \min\left(\|(u \cdot \nabla)u(s) ; W\dot{K}_{n,\infty}^0\|, (t - \tau - s)^{-1/2}\|u \otimes u(s) ; W\dot{K}_{n,\infty}^0\|\right) \\ &\leq c \min\left(s^{-1}, (t - \tau - s)^{-1/2}s^{-1/2}\right)\|u ; X\|^2, \end{aligned}$$

the convergence $\lim_{\tau \searrow 0} \|\mathbb{I}_- ; W\dot{K}_{n,\infty}^0\| = 0$ is verified from the Lebesgue convergence theorem. It is not hard to show that $\lim_{\tau \searrow 0} \|\mathbb{II}_- ; W\dot{K}_{n,\infty}^0\| = 0$. As a result, we obtain the continuity of the solution u in $W\dot{K}_{n,\infty}^0$ on $(0, \infty)$, and $u(t) \in W\dot{K}_{n,\infty}^0$ for $t > 0$, that is (1.6).

Step 4: Here we check the properties (1.7) and (1.8) for the solution u . Since the property (1.8) is proved in Steps 1 and 3, we have to verify (1.7). To do this, thanks to Theorem 1.2 in [42], it suffices to show

$$\lim_{t \searrow 0} \left| \left\langle \int_0^t e^{(t-s)\Delta} \mathbb{P}(u \cdot \nabla)u(s) ds, \phi \right\rangle \right| = 0$$

for any $\phi \in C_0^\infty$. For $n/2 < r < n$, the equality can be verified by integration by parts:

$$\begin{aligned} \left| \left\langle \int_0^t e^{(t-s)\Delta} \mathbb{P}(u \cdot \nabla)u(s) ds, \phi \right\rangle \right| &\leq \int_0^t \|e^{(t-s)\Delta} \mathbb{P}(u \otimes u)(s) ; W\dot{K}_{r,\infty}^0\| \|\nabla \phi ; Z\| ds \\ &\leq c \int_0^t (t - s)^{-1+n/2r} \|u \otimes u(s) ; W\dot{K}_{n/2,\infty}^0\| \|\nabla \phi ; Z\| ds \\ &\leq ct^{(n/r-1)/2} \|u ; X\|^2 \|\nabla \phi ; Z\| \rightarrow 0, \text{ as } t \searrow 0, \end{aligned}$$

where

$$\|f ; Z\| := \sum_{k \in \mathbb{Z}} \|f ; L^{r',1}(A_k)\|.$$

Step 5: Finally, we consider the case $u_0 \in W\dot{K}_{n,\infty}^0$.

In this case, by the same argument as above, we can find a solution u in

$$Y := \{u \in X : \lim_{t \searrow 0} t^{(1-n/p)/2} \|u(t) ; \dot{K}_{p,\infty}^0\| = \lim_{t \searrow 0} t^{1/2} \|u(t) ; L^\infty\| = 0\}.$$

This means that the required convergences are true. □

4.2. Proof of Theorem 1.4. Next, we shall show Theorem 1.4 by the method due to Meyer [39] and [31]. The main ingredient is Proposition 2.4 which yields the critical estimate of the bilinear form B .

Proof of Theorem 1.4. Let $w(t) := u(t) - v(t)$; then we have that $w \in L^\infty(0, T; W\dot{K}_{n,\infty}^0)$ and

$$w(t) = B(e^{\Delta}u_0 - u, w)(t) + B(w, e^{\Delta}u_0 - v)(t) - B(e^{\Delta}u_0, w)(t) - B(w, e^{\Delta}u_0)(t).$$

Note that

$$B(u, v)(t) = \int_0^t \nabla e^{(t-s)\Delta} \mathbb{P}(u \otimes v)(s) ds = \int_0^\infty \nabla e^{s\Delta} \mathbb{P}(u \otimes v) \chi_{(0,t)}(t-s) ds.$$

By using Proposition 2.4 with $r_1 = 2n/3$ and $r_2 = 2n$, i.e., $\beta = 1/4$, one obtains

$$\begin{aligned} & \|B(e^\Delta u_0 - u, w)(t); W\dot{K}_{n,\infty}^0\| \\ &= \left\| \int_0^\infty \nabla e^{s\Delta} \mathbb{P}((e^\Delta u_0 - u) \otimes w) \chi_{(0,t)}(t-s) ds; W\dot{K}_{n,\infty}^0 \right\| \\ &\leq c \left\| ((e^\Delta u_0 - u) \otimes w) \chi_{(0,t)}(t-\cdot); L^\infty(0, \infty; W\dot{K}_{n/2,\infty}^0) \right\| \\ &\leq c \left\| (e^\Delta u_0 - u) \otimes w; L^\infty(0, t; W\dot{K}_{n/2,\infty}^0) \right\| \\ &\leq c \left(\sup_{0 < s \leq t} \|e^{s\Delta} u_0 - u(s); W\dot{K}_{n,\infty}^0\| \right) \left(\sup_{0 < s \leq t} \|w(s); W\dot{K}_{n,\infty}^0\| \right). \end{aligned}$$

$B(w, e^\Delta u_0 - v)$ also has a similar bound as above. On the other hand, the bilinear form $B(e^\Delta u_0, w)$ is estimated as follows:

$$\begin{aligned} \|B(e^\Delta u_0, w)(t); W\dot{K}_{n,\infty}^0\| &\leq \int_0^t \left\| \nabla e^{\frac{(t-s)}{2}\Delta} e^{\frac{(t-s)}{2}\Delta} \mathbb{P}(e^{s\Delta} u_0 \otimes w)(s); W\dot{K}_{n,\infty}^0 \right\| ds \\ &\leq c \int_0^t (t-s)^{-1/2} \left\| (e^{s\Delta} u_0 \otimes w)(s); W\dot{K}_{n,\infty}^0 \right\| ds \\ &\leq c \left(\sup_{0 < s \leq t} s^{1/2} \|e^{s\Delta} u_0; W\dot{K}_{\infty,\infty}^0\| \right) \left(\sup_{0 < s \leq t} \|w(s); W\dot{K}_{n,\infty}^0\| \right). \end{aligned}$$

We now claim that

$$\sup_{0 < s \leq t} \|e^{s\Delta} u_0 - u(s); W\dot{K}_{n,\infty}^0\| \rightarrow 0, \text{ as } t \searrow 0, \tag{4.1}$$

and

$$\sup_{0 < s \leq t} s^{1/2} \|e^{s\Delta} u_0; W\dot{K}_{\infty,\infty}^0\| \rightarrow 0, \text{ as } t \searrow 0. \tag{4.2}$$

The claim (4.1) follows from the continuity of u at $t = 0$ and the definition of $W\dot{K}_{n,\infty}^0$. To show the claim (4.2), we take a small $\tau > 0$ such that $\|u_0 - \tilde{u}_0; W\dot{K}_{n,\infty}^0\|$ is sufficiently small, where $\tilde{u}_0 = e^\tau \Delta u_0$. Applying Corollary 2.1 to \tilde{u}_0 , one has $\|e^{s\Delta} \tilde{u}_0; W\dot{K}_{\infty,\infty}^0\| \leq cs^{-n/(2(n+1))} \|\tilde{u}_0; W\dot{K}_{n+1,\infty}^0\|$, which implies the claim (4.2). Then, there exists $T_0 > 0$ such that

$$\sup_{0 < s \leq T_0} \|w(s); W\dot{K}_{n,\infty}^0\| \leq \frac{1}{2} \sup_{0 < s \leq T_0} \|w(s); W\dot{K}_{n,\infty}^0\|,$$

which implies $u = v$ on $[0, T_0]$. Next, we put

$$T^* := \sup\{0 < t \leq T : u = v \text{ on } [0, t]\}.$$

Then we assume that $T^* < \infty$, otherwise the proof of Theorem 1.4 is completed. By the continuity of solutions, we have $u = v$ on $[0, T^*]$. Then $\tilde{u}(x, t) := u(x, T^* + t)$ and $\tilde{v}(x, t) := v(x, T^* + t)$ are mild solutions of (N-S) in the class $C([0, T - T^*]; W\dot{K}_{n,\infty}^0)$ with initial data $u(\cdot, T^*) = v(\cdot, T^*) \in W\dot{K}_{n,\infty}^0$. Therefore, from the above argument there exists $\tau > 0$ such that $\tilde{u} = \tilde{v}$ on $[0, \tau]$; i.e., $u = v$ on $[0, T^* + \tau]$ which contradicts the maximal property of T^* . \square

Remark 4.1. In the proof above, we showed that the bilinear operator B is bounded from $\tilde{\mathcal{E}} \times \tilde{\mathcal{E}}$ to $\tilde{\mathcal{E}}$ where

$$\tilde{\mathcal{E}} := \{f \in L^\infty(0, \infty; W\dot{K}_{n,\infty}^0) : \sup_{t>0} \|f(t) ; W\dot{K}_{n,\infty}^0\| < \infty\}.$$

This is similar to the Meyer’s estimate [39]; B is continuous from $\mathcal{E} \times \mathcal{E}$ to \mathcal{E} where $\mathcal{E} := \{f \in L^\infty(0, \infty; L^{n,\infty}) : \sup_{t>0} \|f(t) ; L^{n,\infty}\| < \infty\}$. The discontinuity of B from $C([0, T]; L^n) \times C([0, T]; L^n)$ to $C([0, T]; L^n)$ was showed by Oru [43].

4.3. Proof of Theorem 1.5. We make use of the boundedness of B above to get the stability result. As we mentioned in the first section, the reverse also holds without the smallness condition on initial data. Because the proof is not difficult, we omit the details.

Proof of Theorem 1.5. From the smallness assumption on u_0 and v_0 , we have the following estimate for solutions u and v :

$$\|u ; L^\infty(0, \infty; W\dot{K}_{n,\infty}^0)\|, \|v ; L^\infty(0, \infty; W\dot{K}_{n,\infty}^0)\| \leq c_*\delta.$$

Hence the difference of solutions is dominated by itself with a small constant by applying the boundedness of B mentioned in Remark 4.1 as follows:

$$\begin{aligned} \|u(t) - v(t) ; W\dot{K}_{n,\infty}^0\| &\leq \|e^{t\Delta}(u_0 - v_0) ; W\dot{K}_{n,\infty}^0\| \\ &+ \|B(u, u - v)(t) ; W\dot{K}_{n,\infty}^0\| + \|B(v, u - v)(t) ; W\dot{K}_{n,\infty}^0\| \\ &\leq \|e^{t\Delta}(u_0 - v_0) ; W\dot{K}_{n,\infty}^0\| \\ &+ \|B\| \left(\sup_{0 < s \leq t} \|u(s) ; W\dot{K}_{n,\infty}^0\| + \sup_{0 < s \leq t} \|v(s) ; W\dot{K}_{n,\infty}^0\| \right) \\ &\times \sup_{0 < s \leq t} \|u(s) - v(s) ; W\dot{K}_{n,\infty}^0\| \end{aligned}$$

$$\leq \|e^{t\Delta}(u_0 - v_0) ; W\dot{K}_{n,\infty}^0\| + 2c_*\|B\|\delta \sup_{0 < s \leq t} \|u(s) - v(s) ; W\dot{K}_{n,\infty}^0\|.$$

Therefore, we obtain $\sup_{0 < s \leq t} \|u(s) - v(s) ; W\dot{K}_{n,\infty}^0\| \leq \frac{1}{1 - 2c_*\|B\|\delta} \|e^{t\Delta}(u_0 - v_0) ; W\dot{K}_{n,\infty}^0\|$, which implies $\limsup_{t \nearrow \infty} \|u(t) - v(t) ; W\dot{K}_{n,\infty}^0\| = 0$. \square

5. EMBEDDINGS OF HERZ SPACES INTO BESOV SPACES

5.1. **Embedding results.** We prove Theorem 1.6.

(i): In this case, $-n/\sigma < 0 < \alpha < n(1 - 1/p)$. Then, by using Corollary 2.1, one has

$$\begin{aligned} \|f ; \dot{B}_{\sigma,\infty}^{-(\alpha+n(1/p-1/\sigma))}\| &= \sup_{k \in \mathbb{Z}} 2^{-k(\alpha+n(1/p-1/\sigma))} \|(f_{2^k} * \mathcal{F}^{-1}\phi)_{2^{-k}} ; L^\sigma\| \\ &\approx \sup_{k \in \mathbb{Z}} 2^{-k(\alpha+n(1/p-1))} \|f_{2^k} * \mathcal{F}^{-1}\phi ; \dot{K}_{\sigma,\sigma}^0\| \\ &\leq \sup_{k \in \mathbb{Z}} 2^{-k(\alpha+n(1/p-1))} \|f_{2^k} ; W\dot{K}_{p,\infty}^\alpha\| \approx \|f ; W\dot{K}_{p,\infty}^\alpha\|. \end{aligned}$$

(ii): In this case, $-n/p < 0 < \alpha < n(1 - 1/p)$. The inequality $\|f ; \dot{B}_{p,\infty}^{-\alpha}\| \lesssim \|f ; \dot{K}_{p,\infty}^\alpha\|$ can be verified by the same argument as (i).

(iii): In this case, $-n/\sigma < 0 < n(1 - 1/p)$. By replacing the estimate $\|f_{2^k} * \mathcal{F}^{-1}\phi ; \dot{K}_{\sigma,\sigma}^0\| \lesssim \|f_{2^k} ; W\dot{K}_{p,\infty}^\alpha\|$ in the proof of (i) by the estimate $\|f_{2^k} * \mathcal{F}^{-1}\phi ; \dot{K}_{\sigma,\sigma}^0\| \lesssim \|f_{2^k} ; W\dot{K}_{p,\sigma}^0\|$, we can get the inequality $\|f ; \dot{B}_{\sigma,\infty}^{-n(1/p-1/\sigma)}\| \lesssim \|f ; W\dot{K}_{p,\sigma}^0\|$.

(iv): In this case, $0 \leq \alpha < n(1 - 1/p)$. We omit the details.

(v): In this case, $0 \leq \alpha < n$. We omit the details.

(vi): If $\alpha + n/p = 0$; i.e., $\alpha = 0$ and $p = \infty$, then we have

$$W\dot{K}_{\infty,1}^0 = \dot{K}_{\infty,1}^0 \hookrightarrow \dot{K}_{\infty,\infty}^0 \hookrightarrow \dot{B}_{\infty,\infty}^0.$$

To show the case $\alpha + n/p > 0$, we use the following lemma with $\Psi = G$.

Lemma 5.1. *Let $0 < p, q \leq \infty$. Then, for any $\Psi \in \mathcal{S}$ there exists a constant $C_\Psi > 0$, depending on p, q, α, n and Ψ , such that*

$$\sup_{y \in \mathbb{R}^n} \|\Psi(\cdot + y) ; \dot{K}_{(p,1),q}^\alpha\| \leq C_\Psi, \text{ if } p < \infty$$

and

$$\sup_{y \in \mathbb{R}^n} \|\Psi(\cdot + y) ; \dot{K}_{p,q}^\alpha\| \leq C_\Psi,$$

provided that $q < \infty$ and $-n/p < \alpha \leq 0$ or $q = \infty$ and $-n/p \leq \alpha \leq 0$, where

$$\|f ; \dot{K}_{(p,r),q}^\alpha\| := \left(\sum_{k \in \mathbb{Z}} 2^{k\alpha q} \|f \chi_k ; L^{p,r}\|^q \right)^{1/q}.$$

We verify the first inequality of Lemma 5.1 after the proof of Theorem 1.6 is completed. An application of the lemma yields

$$\begin{aligned} |e^{t\Delta} f(x)| &\leq t^{-n/2} \|f_{1/\sqrt{t}} ; W\dot{K}_{p,1}^\alpha\| \|G(\cdot - \frac{x}{\sqrt{t}}) ; \dot{K}_{(p',1),\infty}^{-\alpha}\| \\ &\leq c t^{-(\alpha+n/p)/2} \|f ; W\dot{K}_{p,1}^\alpha\|, \end{aligned}$$

where $p' = \frac{p}{p-1}$ is the conjugate exponent. Here we have used the fact that $-n/p' \leq -\alpha \leq 0$, i.e., $0 \leq \alpha \leq n(1 - 1/p)$. As a consequence, we have

$$\sup_{t>0} t^{(\alpha+n/p)/2} \|e^{t\Delta} f ; L^\infty\| \leq c \|f ; W\dot{K}_{p,1}^\alpha\|.$$

(vii): Since $L^1 \hookrightarrow \dot{B}_{\infty,\infty}^{-n}$, we have $\|f ; \dot{B}_{\infty,\infty}^{-n}\| \lesssim \|f ; \dot{K}_{1,1}^0\|$.

(viii): From (iv), it readily follows that

$$\begin{aligned} \|f ; bmo^{-1}\| &= \sup_{|Q| \leq 1} \left(\int_Q \int_0^{l(Q)^2} |e^{t\Delta} f(x)|^2 dt dx \right)^{1/2} \\ &\leq c \sup_{|Q| \leq 1} \left(\int_0^{l(Q)^2} t^{-(\alpha+n/p)} \|f ; W\dot{K}_{p,\infty}^\alpha\|^2 dt \right)^{1/2} \leq c \|f ; W\dot{K}_{p,\infty}^\alpha\|. \end{aligned}$$

□

We shall verify Lemma 5.1. To do this, it suffices to prove the following lemma.

Lemma 5.2. *Let $0 < p, q \leq \infty$ and $y \in \mathbb{R}^n$. Then, for any $\Psi \in \mathcal{S}$ the inequalities*

$$\|\Psi(\cdot + y) ; \dot{K}_{(p,1),q}^\alpha\| \leq C_\Psi (1 + \min(|y|^\alpha, |y|^{\alpha+n/p})), \text{ if } p < \infty$$

and

$$\|\Psi(\cdot + y) ; \dot{K}_{p,q}^\alpha\| \leq C_\Psi (1 + \min(|y|^\alpha, |y|^{\alpha+n/p}))$$

hold with a constant C_Ψ depending on p, q, α, n and Ψ , provided that $q < \infty$ and $-n/p < \alpha < \infty$ or $q = \infty$ and $-n/p \leq \alpha < \infty$.

Proof. We show the first inequality in the case $q < \infty$ and $-n/p < \alpha < \infty$ only. The other case can be showed by the same argument. Let k_0 be

an integer satisfying $2^{k_0-1} \leq |y| < 2^{k_0}$; i.e., $y \in A_{k_0}$. We decompose the left-hand side:

$$\begin{aligned} \|\Psi(\cdot + y) ; \dot{K}_{(p,1),q}^\alpha\| &\leq c \left(\sum_{k=-\infty}^{k_0-2} 2^{k\alpha q} \|\Psi(\cdot + y) ; L^{p,1}(A_k)\|^q \right)^{1/q} \\ &\quad + c \left(\sum_{k=k_0-1}^{k_0+1} 2^{k\alpha q} \|\Psi(\cdot + y) ; L^{p,1}(A_k)\|^q \right)^{1/q} \\ &\quad + c \left(\sum_{k=k_0+2}^{\infty} 2^{k\alpha q} \|\Psi(\cdot + y) ; L^{p,1}(A_k)\|^q \right)^{1/q} \\ &=: \text{I} + \text{II} + \text{III}. \end{aligned}$$

Each term is dominated by the right-hand side of the statement in the following way:

$$\text{I} \leq C_\Psi \left(\sum_{k=-\infty}^{k_0-2} 2^{k\alpha q} \|\langle \cdot + y \rangle^{-N} ; L^{p,1}(A_k)\|^q \right)^{1/q} \leq C_\Psi 2^{-k_0 N} |y|^{\alpha+n/p} \leq C_\Psi,$$

where N is a positive integer with $|N - (\alpha + n/p)| \leq 1$,

$$\text{II} \leq c \min(\|\Psi ; L^{p,1}\| |y|^\alpha, \|\Psi ; L^\infty\| |y|^{\alpha+n/p}) \leq C_\Psi \min(|y|^\alpha, |y|^{\alpha+n/p})$$

and

$$\text{III} \leq c \left(\sum_{k=k_0}^{\infty} 2^{k\alpha q} \|\Psi ; L^{p,1}(A_k)\|^q \right)^{1/q} \leq c \|\Psi ; \dot{K}_{(p,1),q}^\alpha\| \leq C_\Psi. \quad \square$$

5.2. Negative results. Next, we shall prove Theorem 1.7.

(i): Let

$$f(x) := \sum_{k \in \mathbb{Z}} \frac{2^{-k\alpha}}{|x - x_k|^{n/p}} \chi_k(x),$$

where $x_k := (\frac{3}{2}2^{k-1}, 0, \dots, 0) \in \mathbb{R}^n$. Since $\|f ; L^{p,\infty}(A_k)\| = \sup_{\lambda>0} \lambda |A_k \cap B(x_k, \lambda^{-p/n})|^{1/p}$, we see $f \in \dot{W}K_{p,\infty}^\alpha$. On the other hand, by using (1.10), we have

$$\begin{aligned} \|f ; \dot{B}_{\sigma,\infty}^{-(\alpha+n(1/p-1/\sigma))}\| &\approx \sup_{t>0} t^{(\alpha+n(1/p-1/\sigma))/2} \|e^{t\Delta} f ; L^\sigma\| \geq \|e^\Delta f ; L^\sigma\| \\ &= \left(\sum_{k \in \mathbb{Z}} \|e^\Delta f ; L^\sigma(A_k)\|^\sigma \right)^{1/\sigma}. \end{aligned}$$

For a sufficiently large k , the term $\|e^{\Delta} f ; L^{\sigma}(A_k)\|$ has the lower bound $2^{-k\alpha}$. In fact,

$$\begin{aligned} \|e^{\Delta} f ; L^{\sigma}(A_k)\| &\geq \frac{1}{(4\pi)^{n/2}} \left(\int_{A_k} \left(\int_{A_k} \frac{2^{-k\alpha}}{|y-x_k|^{n/p}} e^{-|x-y|^2/4} dy \right)^{\sigma} dx \right)^{1/\sigma} \\ &\geq \frac{1}{(4\pi)^{n/2}} \left(\int_{B(x_k,1)} \left(\int_{2 \leq |y-x_k| \leq 3} \frac{2^{-k\alpha}}{|y-x_k|^{n/p}} e^{-|x-y|^2/4} dy \right)^{\sigma} dx \right)^{1/\sigma} \gtrsim 2^{-k\alpha}. \end{aligned}$$

Hence, since $\alpha \leq 0$, we have $\|f ; \dot{B}_{\sigma,\infty}^{-(\alpha+n(1/p-1/\sigma))}\| = \infty$.

(ii): From the same argument as above, one has

$$\|f ; \dot{B}_{\infty,\infty}^{-(\alpha+n/p)}\| \gtrsim \sup_{k \in \mathbb{Z}} \|f * G ; L^{\infty}(B(x_k, 1))\| \geq \sup_{k \gg 0} 2^{-k\alpha} = \infty. \quad \square$$

Furthermore, there is no embedding relation between $\dot{K}_{n,\infty}^{-1}$ and BMO , which includes L^{∞} ; that is, we have the following.

Proposition 5.1. *There is no embedding relation between $\dot{K}_{n,\infty}^{-1}$ and BMO . Moreover, there is no embedding relation between $\dot{K}_{n,\infty}^{-1}$ and BMO^d , where BMO^d stands for dyadic BMO .*

Proof. Because $\log|x| \notin \dot{K}_{n,\infty}^{-1}$, the problem is whether $\dot{K}_{n,\infty}^{-1} \hookrightarrow BMO$ is true or not. Let us consider a function $h(x) := (-\log|x-2|)\chi_{(2,4)}(x)$ on \mathbb{R} . By an easy computation, we obtain $h \in \dot{K}_{n,\infty}^{-1}(\mathbb{R})$ and $h \notin BMO(\mathbb{R}) \cap BMO^d(\mathbb{R})$, which implies the embedding does not hold in general. \square

Acknowledgements. The author would like to express thanks to Professor T. Nishitani for valuable comments. The author is grateful to Professor A. Miyachi for indicating him a proof of the inclusion $W\dot{K}_{n,\infty}^0 \hookrightarrow BMO^{-1}$. The author also would like to thank Professor N. Tomita, Professor H. Miura and Dr. T. Yoneda for valuable conversations. The author would like to thank the referees for many helpful suggestions.

REFERENCES

- [1] A. Baernstein II and E.T. Sawyer, *Embedding and multiplier theorems for $H^p(\mathbb{R}^n)$* , Mem. Amer. Math. Soc., 318 (1985).
- [2] O. Barraza, *Self-similar solutions in weak L^p -spaces of the Navier-Stokes equations*, Rev. Mat. Iberoamericana, 12 (1996), 411–439.
- [3] J. T. Beale, T. Kato and A. Majda, *Remarks on the breakdown of smooth solutions for the 3-D Euler equations*, Commu. Math. Phys., 94 (1984), 61–66.
- [4] A. Beurling, *Construction and analysis of some convolution algebras*, Ann. Inst. Fourier Grenoble, 14 (1964), 1–32.

- [5] J. Bourgain and N. Pavlović, *Ill-posedness of the Navier-Stokes equations in a critical space in 3D*, J. Funct. Anal., 255 (2008), 2233–2247.
- [6] M. Cannone, “Ondelettes, paraproducts et Navier-Stokes, ” Thesis. Université Paris IX-Dauphine, (1994).
- [7] M. Cannone, *A generalization of a theorem by Kato on Navier-Stokes equations*, Rev. Mat. Iberoamericana, 13 (1997), 515–541.
- [8] Y. Z. Chen and K. S. Lau, *On some new classes of Hardy spaces*, J. Funct. Anal., 84 (1989), 255–278.
- [9] E. B. Fabes, J.E. Jones and N. M. Riviere, *The initial value problem for the Navier-Stokes equations with the data in L^p* , Arch. Rational Mech. Anal., 42 (1972), 222–240.
- [10] J. García-Cuerva, *Hardy spaces and Beurling algebras*, J. London Math. Soc., 39 (1989), 499–513.
- [11] J. García-Cuerva and M.-J. L. Herrero, *A theory of Hardy spaces associated to the Herz spaces*, Proc. London Math. Soc., 69 (1994), 605–628.
- [12] Y. Giga, *Solutions for semilinear Parabolic equations in L^p and Regularity of weak solutions of the Navier-Stokes system*, J. Differential Equations, 61 (1986), 186–212.
- [13] Y. Giga, K. Inui and Matsui, *On the Cauchy problem for the Navier-Stokes equations with nondecaying initial data*, Advances in fluid dynamics, 4 (1999), Quad. Mat., Dept. Math., Seconda Univ. Napoli, Caserta, 27–68.
- [14] Y. Giga and T. Miyakawa, *Navier-Stokes flow in \mathbb{R}^3 with measures as initial vorticity and Morrey spaces*, Comm. Partial Differential Equations, 14 (1989), 577–618.
- [15] L. Grafakos, “Classical Fourier Analysis, ” Second Edition Graduate Texts in Mathematics, 249, Springer, New York, 2008.
- [16] H. G. Feichtinger, *An elementary approach to Wiener’s third Tauberian theorem for the Euclidean n -space*, In Symposia Math., XXIX (1988), 267–301.
- [17] H. Fujita and T. Kato, *On the Navier-Stokes initial value problem, I*, Arch. Rat. Mech. Anal., 16 (1964), 269–315.
- [18] G. Furioli, P. G. Lemarié-Rieusset and E. Terraneo, *Sur l’unicité dans $L^3(\mathbb{R}^3)$ des solutions « mild » des équations de Navier-Stokes*, C. R. Acad. Sci. Paris Ser. I Math., 325 (1997), 1253–1256.
- [19] G. Furioli, P. G. Lemarié-Rieusset and E. Terraneo, *Unicité dans $L^3(\mathbb{R}^3)$ et d’autres espaces fonctionnels limites pour Navier-Stokes*, Rev. Mat. Iberoamericana, 16 (2000), 605–667.
- [20] C. Herz, *Lipschitz spaces and Bernstein’s theorem on absolutely convergent Fourier transforms*, J. Math. Mech., 205 (1968), 283–324.
- [21] T. Iwabuchi, *Navier-Stokes equations and nonlinear heat equations in modulation spaces with negative derivative indices*, J. Differential Equations, 248 (2010), 1972–2002.
- [22] G. Karch, *Scaling in nonlinear parabolic equations*, J. Math. Anal. Appl., 234 (1999), 534–558.
- [23] T. Kato, *Strong L^p -solutions of the Navier-Stokes equation in \mathbb{R}^n , with applications to weak solutions*, Math. Z., 187 (1984), 471–480.
- [24] H. Koch and D. Tataru, *Well-posedness for the Navier-Stokes equations*, Adv. Math., 157 (2001), 22–35.
- [25] Y. Komori, *Weak type estimates for Calderón-Zygmund operators on Herz spaces at critical indexes*, Math. Nachr., 259 (2003), 42–50.

- [26] P. Konieczny and T. Yoneda, *On dispersive effect of the Coriolis force for the stationary Navier-Stokes equations*, arXiv:1007.1181v1.
- [27] H. Kozono, T. Ogawa and Y. Taniuchi, *Navier-Stokes equations in the Besov space near L^∞ and BMO*, Kyushu J. Math., 57 (2003), 303–324.
- [28] H. Kozono and Y. Shimada, *Bilinear estimates in homogeneous Triebel-Lizorkin spaces and the Navier-Stokes equations*, Math. Nachr., 276 (2004), 63–74.
- [29] H. Kozono and Y. Taniuchi, *Bilinear estimates in BMO and the Navier-Stokes equations*, Math. Z., 235 (2000), 173–194.
- [30] H. Kozono and M. Yamazaki, *Semilinear heat equations and the Navier-Stokes equation with distributions in new function spaces as initial data*, Comm. Partial Differential Equations, 19 (1994), 959–1014.
- [31] P. G. Lemarié-Rieusset, “Recent Development in the Navier-Stokes Problem,” Chapman & Hall/CRC Research Notes in Mathematics, 431, Chapman & Hall/CRC, Boca Raton, FL. (2002).
- [32] P. G. Lemarié-Rieusset, *The Navier-Stokes equations in the critical Morrey-Campanato space*, Rev. Mat. Iberoamericana, 23 (2007), 897–930.
- [33] J. Leray, *Essai sur le mouvement d’un fluide visqueux emplissant l’espace*, Acta Math., 63 (1934), 193–248.
- [34] X. Li and D. Yang, *Boundedness of some sublinear operators on Herz spaces*, Illinois J. Math., 40 (1996), 484–501.
- [35] S. Lu and D. Yang, *The decomposition of weighted Herz space on \mathbf{R}^n and its applications*, Sci. China Ser. A, 38 (1995), 147–158.
- [36] S. Lu and D. Yang, *The weighted Herz-type Hardy spaces and its applications*, Sci. China Ser. A., 38 (1995), 662–673.
- [37] S. Lu, K. Yabuta and D. Yang, *Boundedness of some sublinear operators in weighted Herz-type spaces*, Kodai Math. J., 23 (2000), 391–410.
- [38] Y. Maekawa and Y. Terasawa, *The Navier-Stokes equations with initial data in uniformly local L^p spaces*, Differential Integral Equations, 19 (2006), 369–400.
- [39] Y. Meyer, *Wavelet, Paraproduct and Navier-Stokes Equations*, Current developments in mathematics. International Press, (1996), 105–212.
- [40] C. Miao and B. Yuan, *Weak Morrey spaces and strong solutions to the Navier-Stokes equations*, Sci. China Ser. A., 50 (2007), 1401–1417.
- [41] A. Miyachi, *Remarks on Herz-type Hardy spaces*, Acta Math. Sinica. English Series, 17 (2001), 339–360.
- [42] E. Nakai, N. Tomita and K. Yabuta, *Density of the set of all infinitely differentiable functions with compact support in weighted sobolev spaces*, Sci. Math. Jpn., 60 (2004), 121–127.
- [43] F. Oru, *Rôle des oscillations dans quelques problèmes d’analyse non linéaire*, Thèse, École Normale Supérieure de Cachan (1998).
- [44] F. Planchon, *Asymptotic behavior of global solutions to the Navier-Stokes equations in \mathbb{R}^n* , Rev. Mat. Iberoamericana, 14 (1998), 71–93.
- [45] O. Sawada, *On time-local solvability of the Navier-Stokes equations in Besov spaces*, Adv. Differential Equations, 8 (2003), 385–412.
- [46] H. Triebel, *Characterizations of Besov-Hardy-Sobolev Spaces; A Unified Approach*, J. Approx. Theory, 52 (1988), 162–203.

- [47] Y. Tsutsui, *Sharp maximal inequalities and its application to some bilinear estimates*, J. Fourier Anal. Appl., 17 (2011), 265–289.
- [48] F. Weisz, *Wiener amalgams, Hardy spaces and summability of Fourier series*, Math. Proc. Cambridge Philos. Soc., 145 (2008), 419–442.
- [49] T. Yoneda, *Ill-posedness of the 3D-Navier-Stokes equations in a generalized Besov space near BMO^{-1}* , J. Funct. Anal., 258 (2010), 3376–3387.