

SIMPLICIAL INTERSECTION CHAINS FOR AN ABSTRACT COMPLEX

W. W. FLEXNER

Abstract complexes have been defined by J. W. Alexander, W. Mayer, A. W. Tucker and S. Lefschetz.¹ The definition here adopted is that of Lefschetz which derives directly from that of Tucker. To conform to Lefschetz's present usage the notation here will differ from that in the article cited by having $K = \{E_i^p\}$ and $K^* = \{E_p^i\}$. It should be remarked that when K is infinite $FF=0$ implies that for given E_i^p and E_j^{p-2} ,

$$[E_i^p : E_k^{p-1}] [E_k^{p-1} : E_j^{p-2}] \neq 0$$

for at most a finite number of E_k^{p-1} . The dual K^* here is to have the property

$$[E_i^p : E_j^{p-1}] = (-1)^{p+1} [E_{p-1}^j : E_p^i] = \mu_{ij}^p$$

where μ_{ij}^p is an abbreviated notation (compare loc. cit., p. 346 (c)). This is in order that a fourth postulate

$$\text{IV. } \text{Ki} (E_i^p \cdot E_p^i) = 1$$

may be added to I-III (loc. cit., p. 350), where Ki means Kronecker index, the sum of the coefficients of the elements of the zero-chain in question. Here C is a p -chain of K , D a q -chain of K^* , $p > q$.

It is well known that when K is a simplicial or polyhedral manifold, for every pair E_i^p, E_q^j , $p \geq q$, an intersection can be defined which is a chain of the simplicial regular subdivision of K . And when K is an arbitrary simplicial complex, the intersection is on K and hence simplicial. Here it is shown that *for any complex the intersections $E_i^p \cdot E_q^j$ can be regarded as integral chains of a simplicial complex K^π .*

This complex will be defined abstractly by means of its vertices σ_i^p . The first definition is for $p = q$:

$$(1) \quad E_i^p \cdot E_p^i = \delta_i^i \sigma_i^p$$

(where \cdot replaces the \odot of the article cited). Then for $p > q$, if $r = p - q - 1$,

$$(2) \quad E_i^p \cdot E_q^j = \sum_{k_1, k_2, \dots, k_r} \mu_{i k_1}^p \mu_{k_1 k_2}^{p-1} \cdots \mu_{k_r j}^{q+1} \sigma_i^p \sigma_{k_1}^{p-1} \cdots \sigma_{k_r}^{q+1} \sigma_j^q$$

¹ S. Lefschetz, this Bulletin, vol. 43 (1937), pp. 345-359. (References to the other authors will be found on page 345.)

where $\sigma_i^p \sigma_{k_1}^{p-1} \cdots \sigma_{k_r}^{q+1} \sigma_j^q$ denotes the simplex of K^π with the given vertices; so $E_i^p \cdot E_q^j$ is an integral chain of K^π .

To verify that $E_i^p \cdot E_q^j$ is an intersection, it is sufficient, in view of (1), (2), to prove the formula III (p. 350, loc. cit.):

$$\text{III. } F(E_i^p \cdot E_q^j) = (FE_i^p) \cdot E_q^j + (-1)^p E_i^p \cdot (FE_q^j).$$

When $q = p - 1, r = 0$, (2) gives $E_i^p \cdot E_{p-1}^j = \mu_{ij}^p \sigma_i^p \sigma_j^{p-1}$. Using the rule for the boundary of a simplex given by its vertices, we obtain $F(E_i^p \cdot E_{p-1}^j) = \mu_{ij}^p (\sigma_j^{p-1} - \sigma_i^p)$. By formula (2) (loc. cit.), its dual, and (1),

$$\begin{aligned} (FE_i^p) \cdot E_{p-1}^j &= \sum_k \mu_{ik}^p E_k^{p-1} \cdot E_{p-1}^j = \mu_{ij}^p \sigma_j^{p-1}, \\ E_i^p \cdot (FE_{p-1}^j) &= (-1)^{p+1} \mu_{ij}^p \sigma_i^p \end{aligned}$$

which verifies III in this case. When $q < p - 1$,

$$\sum_k \mu_{ik_1}^p \cdots \mu_{k_{s-1} k_s}^{p-s+1} \mu_{k_s k_{s+1}}^{p-s} \cdots \mu_{k_r j}^{q+1} \sigma_i^p \sigma_{k_1}^{p-1} \cdots \sigma_{k_{s-1}}^{p-s+1} \sigma_{k_{s+1}}^{p-s-1} \cdots \sigma_{k_r}^{q+1} \sigma_j^q = 0, \quad 1 \leq s \leq r,$$

since $FF = 0$ implies $\sum_k \mu_{k_s k_{s-1} k_s}^{p-s+1} \mu_{k_s k_{s+1}}^{p-s} = 0$. Hence

$$\begin{aligned} F(E_i^p \cdot E_q^j) &= \sum_k \mu_{ik_1}^p \cdots \mu_{k_r j}^{q+1} (\sigma_{k_1}^{p-1} \cdots \sigma_{k_r}^{q+1} \sigma_j^q \\ &\quad + (-1)^{p-q} \sigma_i^p \sigma_{k_1}^{p-1} \cdots \sigma_{k_r}^{q+1}) \\ &= \sum_{k_1} \mu_{ik_1}^p (E_{k_1}^{p-1} \cdot E_q^j) + (-1)^{p-q} \sum_{k_r} \mu_{k_r j}^{q+1} (E_i^p \cdot E_{q+1}^{k_r}) \end{aligned}$$

which by Lefschetz's article, formulas (2) and (1), is the right side of III.

CORNELL UNIVERSITY AND
INSTITUTE FOR ADVANCED STUDY