

THE PERMANENT ANALOGUE OF THE HADAMARD DETERMINANT THEOREM

BY MARVIN MARCUS

Communicated by A. M. Gleason, April 11, 1963

1. **Statement of results.** In [2; 3] it was conjectured that if $A = (a_{ij})$ is an n -square positive semi-definite hermitian matrix then

$$(1) \quad \text{per } A \geq \prod_{i=1}^n a_{ii}.$$

Here $\text{per } A$ denotes the permanent of A : $\text{per } A = \sum_{\sigma} \prod_{i=1}^n a_{i\sigma(i)}$ where the summation is over the whole symmetric group of degree n . It was announced [1] and later proved [2] that $\text{per } (A) \geq \det A$ and the Hadamard determinant theorem suggests that the product of the main diagonal entries of A in fact separates the permanent and the determinant of A . In this note we sketch a proof of an inequality that is substantially stronger than (1). Let $A(i)$ denote the principal submatrix of A obtained by deleting row and column i .

THEOREM. *If A is an $(r+1)$ -square positive semi-definite hermitian matrix then*

$$(2) \quad (r+1)a_{11} \text{per } A(1) \geq \text{per } A \geq a_{11} \text{per } A(1).$$

If A has a zero row then (2) is equality throughout. If A has no zero row then the lower equality holds if and only if $A = a_{11}I + A(1)$; the upper equality holds if and only if A is of rank 1.

We remark that what is true for $A(1)$ is true for any $A(i)$ because the permanent is unaltered by permutation of the rows and columns.

By an obvious induction on r we have the

COROLLARY. *If A is an n -square positive semi-definite hermitian matrix then*

$$(3) \quad \text{per } A \geq \prod_{i=1}^n a_{ii}$$

with equality if and only if A has a zero row or A is a diagonal matrix.

2. **Proof of theorem.** We outline the proof of the theorem. Let U be an n -dimensional unitary space with inner product (x, y) . For $1 \leq r \leq n$ define $U^{(r)}$ to be the space of r -tensors on U ; that is, $U^{(r)}$ is the dual space of the space of all multilinear complex valued func-

tions of r -tuples of vectors from U . If x_1, \dots, x_r are in U then $x_1 \otimes \dots \otimes x_r \in U^{(r)}$ is defined by $x_1 \otimes \dots \otimes x_r(\phi) = \phi(x_1, \dots, x_r)$ for any multilinear functional ϕ . Define the completely symmetric operator $S: U^{(r)} \rightarrow U^{(r)}$ by

$$S(x_1 \otimes \dots \otimes x_r) = \frac{1}{r!} \sum_{\sigma} x_{\sigma(1)} \otimes \dots \otimes x_{\sigma(r)}$$

where σ runs over the symmetric group of degree r . Then define the symmetric product of x_1, \dots, x_r by $x_1 \cdots x_r = S(x_1 \otimes \dots \otimes x_r)$. The range of S is the symmetry class of completely symmetric tensors denoted by $U_{(r)}$. It is well known that an inner product in $U_{(r)}$ is given by

$$(4) \quad (x_1 \cdots x_r, y_1 \cdots y_r) = \frac{1}{r!} \text{per}((x_i, y_j)), \quad i, j = 1, \dots, r.$$

Now let $G_{r,n}$ denote the totality of nondecreasing sequences of length r chosen from $1, \dots, n$ and for $\omega \in G_{r,n}$ define $\mu(\omega)$ to be the product of the factorials of the multiplicities of the distinct integers in ω ; e.g. $\mu(1, 1, 4, 4, 4, 5) = 2!3!$. From (4) it follows that if e_1, \dots, e_n is an orthonormal (o.n.) basis in U then $\sqrt{(r!/\mu(\omega))}e_{\omega}$, $\omega \in G_{r,n}$, is an o.n. basis in $U_{(r)}$: here $e_{\omega} = e_{\omega_1} \cdots e_{\omega_r}$.

Now let e_1 be a unit vector in U and complete to an o.n. basis of U with e_2, \dots, e_n . Next define a linear map $T: U_{(r)} \rightarrow U_{(r+1)}$ by $T(x_1 \cdots x_r) = e_1 \cdot x_1 \cdots x_r$. From (4) it follows that if no $x_i = 0$, $i = 1, \dots, r$, then the Rayleigh quotient

$$(5) \quad \frac{(Tx_1 \cdots x_r, Tx_1 \cdots x_r)}{(x_1 \cdots x_r, x_1 \cdots x_r)}$$

is given by

$$\frac{1}{r+1} \text{per } A / \text{per } A(1)$$

where $A = a_{ij}$ is the Gram matrix based on the ordered set e_1, x_1, \dots, x_r : that is,

$$\begin{aligned} a_{i+1, j+1} &= (x_i, x_j), & 1 \leq i, j \leq r, \\ a_{11} &= (e_1, e_1), \\ a_{1 j+1} &= \bar{a}_{j+1 1} = (e_1, x_j), & j = 1, \dots, r. \end{aligned}$$

Let $H = T^*T$, then from (5) the problem becomes equivalent to determining the maximum and minimum eigenvalues of H and the cor-

responding eigenvectors. It turns out that a matrix representation of H with respect to the lexicographically ordered basis $\sqrt{(r!/\mu(\omega))}e_\omega$, $\omega \in G_{r,n}$, is diagonal and the eigenvalues of H are

$$(6) \quad \frac{\mu(1, \omega)}{\mu(\omega)} \cdot \frac{1}{r+1}$$

with corresponding eigenvectors e_ω : here $(1, \omega)$ designates the sequence $(1, \omega_1, \dots, \omega_r) \in G_{r+1,n}$. The expression (6) is bounded below by $1/(r+1)$ and takes on this value if and only if $\omega_1 > 1$. This means that the lower inequality can hold in (2) if and only if $x_1 \cdots x_r$ is in the space spanned by e_ω , $\omega \in G_{r,n}$, $\omega_1 > 1$. We then prove that this happens if and only if $(x_i, e_1) = 0, i = 1, \dots, r$, i.e., that $A = a_{11}I + A(1)$. Similarly, the expression (6) is bounded above by 1 and takes on this value only for the sequence $\omega_i = 1, i = 1, \dots, r$. This means that the upper equality can hold in (2) if and only if $x_1 \cdots x_r$ is a multiple of $e_1 \cdots e_1$. According to [2, Theorem 3] this happens only in case $x_i = d_i e_1, i = 1, \dots, r$, and this leads to the condition that rank of A is 1.

REFERENCES

1. Marvin Marcus and Morris Newman, *The permanent function as an inner product*, Bull. Amer. Math. Soc. **67** (1961), 223-224.
2. ———, *Inequalities for the permanent function*, Ann. of Math. (2) **75** (1962), 47-62.
3. Marvin Marcus and Henryk Minc, *The Pythagorean theorem in certain symmetry classes of tensors*, Trans. Amer. Math. Soc. **104** (1962), 510-515.

UNIVERSITY OF CALIFORNIA, SANTA BARBARA