

Theorem 3 and the Lemma give precisely the amount of uniformity required to obtain two sequences of pushes  $\phi^1, \phi^2, \dots$  and  $\psi^1, \psi^2, \dots$  of  $(Q, f(M))$  (converging to  $\epsilon/2$ -pushes  $\phi$  and  $\psi$  of  $(Q, f(M))$ ), respectively, such that  $\phi f = \psi g$ . The desired  $\epsilon$ -push of  $(Q, f(M))$  is the composition  $h = \psi^{-1}\phi$ .

Complete proofs of these results will appear in full elsewhere.

*Added in proof.* Since this announcement was submitted, some errors in the proof of Homma's approximation theorem have been discovered. Should these discrepancies not be rectified, then the restriction  $q - m \geq 3$  must be replaced by  $m \leq (2/3)q - 1$  in the statements of Theorem 1 and the corollaries.

#### REFERENCES

1. J. L. Bryant and C. L. Seebeck, III, *Locally nice embeddings of polyhedra*, Quart. J. Math. Oxford Ser., (to appear).
2. H. Gluck, *Embeddings in the trivial range*, Ann. of Math. (2) **81** (1965), 195–210.
3. J. P. Hempel and D. R. McMillan, Jr., *Locally nice embeddings of manifolds*, Amer. J. Math. **88** (1966), 1–19.
4. T. Homma, *On the imbedding of polyhedra in manifolds*, Yokohama Math. J. **10** (1962), 5–10.
5. ———, *Piecewise linear approximations of embeddings of manifolds*, Mineographed notes, Florida State University, 1965.
6. J. F. P. Hudson, *Concordance and isotopy of PL embeddings*, Bull. Amer. Math. Soc. **72** (1966), 534–535.
7. D. R. McMillan, Jr., *A criterion for cellularity in a manifold*, Ann. of Math. (2) **79** (1964), 327–337.

FLORIDA STATE UNIVERSITY, AND  
MICHIGAN STATE UNIVERSITY

#### ERRATUM, VOLUME 73

Index to Volume 73, p. 1004.

Line 29 should read:

Kahn, Peter J. *Chern numbers and oriented homotopy type*, 932