

Since the factors in the first two sets of brackets are finite Blaschke products and the zero in the third is a convex combination of such, and since the coefficients are nonnegative and sum to 1, the proof is complete.

REFERENCES

1. C. Caratheodory, *Theory of functions*. Vol. 2, Chelsea, New York, 1954.
2. R. R. Phelps, *Extreme points in functions algebras*, Duke Math. J. **32** (1965), 267-278.

MASSACHUSETTS INSTITUTE OF TECHNOLOGY

EXACTNESS OF INVERSE LIMITS

BY ULRICH OBERST

Communicated by Saunders Mac Lane, May 20, 1968

I. The problem of this investigation is to characterize those small categories X for which the inverse limit

$$\text{proj lim}_X: AB^X \rightarrow AB$$

is exact. Here AB is the category of abelian groups, and AB^X is the category of functors from X to AB . In this context I conjecture the following

THEOREM I. *Let X be a small category. Then the following assertions are equivalent:*

- (1) *The inverse limit $\text{proj lim}_X: AB^X \rightarrow AB$ is exact.*
- (2) *For every abelian category \mathfrak{A} with exact direct products, the inverse limit $\text{proj lim}_X: \mathfrak{A}^X \rightarrow \mathfrak{A}$ is exact.*
- (3) *Every connected component Y of X contains an object y together with an endomorphism $e \in Y(y, y)$ such that the following properties are satisfied:*
 - (i) *y is a smallest object of Y , i.e., for any object $z \in Y$ there is a morphism $y \rightarrow z$.*
 - (ii) *e equalizes any two morphisms with the same codomain and domain y , i.e., any diagram $y \xrightarrow{\alpha} y \xrightarrow{\beta} z$ is commutative.*

At present, I can prove the equivalence of (1) and (2) and the implication (3) \Rightarrow (1) in general, i.e., without any additional condition on X . The implication (1) \Rightarrow (3) holds at least if one of the following conditions on X is satisfied:

(a) X is a monoid, i.e., has exactly one object. In this case (3) means that there is a right zero element $e \in X$ with $\alpha e = e$ for all $\alpha \in X$.

(b) X satisfies the condition

(F2')^o: Every commutative diagram $x_1 \rightrightarrows x_2 \rightarrow x_3$ can be extended to a commutative diagram $x_0 \rightarrow x_1 \rightrightarrows x_2 \rightarrow x_3$.

This condition is satisfied in particular if the connected components Y of X are filtered from below (see also [1, p. 7]), i.e., if X satisfies the condition

(F2)^o: Any diagram $x_1 \rightrightarrows x_2$ in X can be completed to a commutative diagram $x_0 \rightarrow x_1 \rightrightarrows x_2$.

II. The following steps are taken in the proof of Theorem I.

Step 1. It is clear that for any abelian category \mathfrak{A} with exact direct products the inverse limit $\text{proj lim}_X: \mathfrak{A}^X \rightarrow \mathfrak{A}$ is exact if and only if for every connected component Y of X the inverse limit $\text{proj lim}_Y: \mathfrak{A}^Y \rightarrow \mathfrak{A}$ is exact. Hence one may assume without loss of generality that X itself is connected.

Step 2. Let \mathbf{Z} denote the group of integers, and let $\mathfrak{Z}: X \rightarrow AB$ be the constant functor with values \mathbf{Z} . Let

$$\text{Hom}_X: (AB^X)^0 \times \mathfrak{A}^X \rightarrow \mathfrak{A}$$

be the formal Hom-functor, defined for any abelian category \mathfrak{A} with direct products (see e.g. [4, p. 145ff]). In the same fashion let

$$\otimes_X: AB^X \times \mathfrak{A}^{X^0} \rightarrow \mathfrak{A}$$

be the tensor product, defined for any abelian category \mathfrak{A} with direct sums (loc. cit.). It is easily seen that

$$\text{proj lim}_X = \text{Hom}_X(\mathfrak{Z}, ?): \mathfrak{A}^X \rightarrow \mathfrak{A},$$

respectively

$$\text{inj lim}_{X^0} = \mathfrak{Z} \otimes_{X^0} ? : \mathfrak{A}^{X^0} \rightarrow \mathfrak{A}$$

where X^0 is the dual category of X .

Using these notions we prove the

PROPOSITION. *Let X be a small category. Then the following properties are equivalent.*

- (1) $\text{proj lim}_X: AB^X \rightarrow AB$ is exact.
- (2) For any abelian category \mathfrak{A} with exact direct products the functor $\text{proj lim}_X: \mathfrak{A}^X \rightarrow \mathfrak{A}$ is exact.
- (3) The object $\mathfrak{Z} \in AB^X$ is projective.

Moreover, if (1)–(3) are satisfied then for any abelian category \mathfrak{A} with exact direct sums the direct limit

$$\text{inj lim}_{X^0} \mathfrak{A}^{X^0} \rightarrow \mathfrak{A}$$

is exact.

The basic observation for this proof is that \mathfrak{Z} is projective if and only if it is X -representable in the sense of Eilenberg-Mac Lane [2], i.e., if and only if the canonical epimorphism

$$\begin{aligned} \phi: \prod_{x \in X} \mathfrak{Z}X(x, ?) &\rightarrow \mathfrak{Z}, \\ id_x \rightsquigarrow 1 \in \mathfrak{Z}x &= \mathfrak{Z} \end{aligned}$$

splits. Here $\mathfrak{Z}X(x, y)$ is the free abelian group with basis $X(x, y)$.

The preceding proposition can be generalized to the

PROPOSITION. For a small category X and a nonnegative integer n the following assertions are equivalent:

- (1) The $(n+1)$ st right derived functor $(\text{proj lim}_X)_{n+1}$ of $\text{proj lim}_X: AB^X \rightarrow AB$ is zero.
- (2) The $(n+1)$ st right derived functor $(\text{proj lim}_X)_{n+1}$ of $\text{proj lim}_X: \mathfrak{A}^X \rightarrow \mathfrak{A}$ is zero for any abelian category \mathfrak{A} with exact direct products.
- (3) The projective dimension of \mathfrak{Z} is smaller than or equal to n .

Step 3. The most important step in proving Theorem I is the following proposition which is a special case of the implication (1) \Rightarrow (3) of Theorem I.

PROPOSITION. Let X be a small, connected category. Assume that $\text{proj lim}_X: AB^X \rightarrow AB$ is exact, and that X is filtered from below, i.e., satisfies (F2)^o. Then there are an object $x \in X$ and an endomorphism $e \in X(x, x)$ satisfying the conditions (i) and (ii) of (3), Theorem I.

The proof of this proposition starts with the fact that the canonical isomorphism

$$\phi: \prod_{x \in X} \mathfrak{Z}X(x, ?) \rightarrow \mathfrak{Z}$$

splits.

Step 4. Under the hypotheses (a), respectively (b), on X the implication (1) \Rightarrow (3) of Theorem I (assume X connected without loss of generality) is proven in the following way: assume that $\text{proj lim}_X: AB^X \rightarrow AB$ is exact. From the proposition of the second step we conclude that $\text{inj lim}_{X^0}: AB^{X^0} \rightarrow AB$ is exact. Moreover X^0 satisfies the dual conditions (a)^o respectively (b)^o of (a) respectively (b). Under these

hypotheses it was shown in [3] respectively [5] that X^0 is filtered from above. Hence X is filtered from below. Using this fact and the proposition of the third step we obtain that X satisfies the condition (3) of Theorem I.

Step 5. In order to prove Theorem I in general it is sufficient to prove the following

CONJECTURE: Let X be a connected, small category and assume that $\text{inj } \lim_X: AB^{\mathbf{x}} \rightarrow AB$ is exact. Then X is filtered from above.

I conjecture this result because it is true if X is a monoid [3] or satisfies (F2') [5], and no counterexamples are known.

REFERENCES

1. M. Artin, *Grothendieck topologies*, Seminar Notes, Harvard University 1962.
2. S. Eilenberg and S. Mac Lane, *Acyclic models*, Amer. J. Math. **75** (1953), 189–199.
3. J. R. Isbell, (to appear).
4. B. Mitchell, *Theory of categories*, Academic Press, New York, 1965.
5. U. Oberst, *Homology of categories and exactness of direct limits*, Math. Z. (to appear.)

UNIVERSITY OF CHICAGO