

## LOCAL SPECTRAL MAPPING THEOREMS

BY ROBERT G. BARTLE AND CONSTANTINE A. KARIOTIS<sup>1</sup>

Communicated September 26, 1972

This note is concerned with various “localized versions” of the spectral mapping theorem (cf. [3, VII.3.11]) for bounded linear operators in a complex Banach space. The details of the proof and some additional results will be published elsewhere.

Let  $X$  be a complex Banach space and let  $T \in B(X)$ , the Banach algebra of all bounded linear operators on  $X$ . We recall (cf. [2, p. 1], [3, p. 1931]) that if  $T$  has the single-valued extension property then there exist a maximal open set  $\rho_T(x)$  containing  $\rho(T)$  and a unique holomorphic function  $\tilde{x}_T: \rho_T(x) \rightarrow X$  such that  $(\lambda I - T)\tilde{x}_T(\lambda) = x$  for all  $\lambda \in \rho_T(x)$ . The complementary set  $\sigma_T(x) = \mathbf{C} - \rho_T(x)$ , which we call the *local spectrum of  $x$*  (with respect to  $T$ ), is compact and is contained in  $\sigma(T)$ , the spectrum of  $T$ . If  $F \subseteq \mathbf{C}$  is closed we introduce the spectral manifold  $X_T(F) = \{x \in X: \sigma_T(x) \subseteq F\}$ .

**THEOREM 1.** *Let  $f$  be holomorphic on a neighborhood of  $\sigma(T)$  and suppose that  $T$  and  $f(T)$  have the single-valued extension property. Then  $f(\sigma_T(x)) = \sigma_{f(T)}(x)$  for all  $x \in X$ .*

This result<sup>2</sup> was proved by the second-named author in his dissertation [4]. The proof given there is similar to the proof of a theorem due to Colojoară and Foiaş ([1], [2, p. 71]). In fact, the conclusion of Theorem 1 is equivalent to the condition that if  $F$  is a closed subset of  $\sigma(f(T)) = f(\sigma(T))$ , then

$$X_{f(T)}(F) = X_T(f^{-1}(F)).$$

If  $T \in B(X)$  and  $Y$  is a (closed) subspace of  $X$  which is invariant under  $T$ , then the spectrum of the restriction  $T|_Y$  may be either smaller or larger than  $\sigma(T)$ . It is often desirable to limit attention to subspaces which do not increase the spectrum under restriction (e.g., ultra-invariant subspaces).

**THEOREM 2.** *Let  $Y$  be a subspace invariant under  $T$  such that  $\sigma(T|_Y) \subseteq \sigma(T)$  and let  $f$  be holomorphic on a neighborhood of  $\sigma(T)$ . Then  $Y$  is invariant*

---

AMS(MOS) subject classifications (1970). Primary 47A10, 47A60; Secondary 47A15.

<sup>1</sup> The authors were partially supported by a grant from the National Science Foundation.

<sup>2</sup> ADDED IN PROOF. After this note was communicated we discovered this theorem with a similar proof in C. Apostol, *Teorie spectrală și calcul funcțional*, Stud. Cerc. Mat. **20**, no. 5 (1968), 635–668.

under  $f(T)$ ,  $f(T)|Y = f(T|Y)$ , and

$$f(\sigma(T|Y)) = \sigma(f(T|Y)) = \sigma(f(T)|Y) \subseteq f(\sigma(T)).$$

If  $Y$  is an invariant subspace for  $T$  such that  $\sigma(T|Y) \subseteq \sigma(T)$  and if  $y \in Y$ , then  $\sigma_T(y) \subseteq \sigma_{T|Y}(y)$ , but this inclusion can be proper. The next result holds for subspaces for which equality holds (e.g., spectral maximal spaces).

**THEOREM 3.** *Suppose that  $Y$  is a subspace invariant under  $T$  such that  $\sigma_{T|Y}(y) = \sigma_T(y)$  for all  $y \in Y$ . Let  $f$  be holomorphic on a neighborhood of  $\sigma(T)$  and let  $T$  and  $f(T)$  have the single-valued extension property. If  $y \in Y$ , we have*

$$\sigma_{f(T)|Y}(y) = \sigma_{f(T|Y)}(y) = f(\sigma_{T|Y}(y)) = f(\sigma_T(y)) = \sigma_{f(T)}(y).$$

If  $T \in B(X)$  and  $Y$  is a (closed) subspace of  $X$  which is invariant under  $T$  and if  $T|Y$  is the operator in the quotient space  $X/Y$  defined by  $(T|Y)[x] = [Tx]$ , then the spectrum of  $T|Y$  may be either smaller or larger than  $\sigma(T)$ . It is often desirable to limit attention to subspaces which do not increase the spectrum under quotients (e.g., ultra-invariant subspaces).

**THEOREM 4.** *Let  $Y$  be a subspace invariant under  $T$  such that  $\sigma(T|Y) \subseteq \sigma(T)$  and let  $f$  be holomorphic on a neighborhood of  $\sigma(T)$ . Then  $Y$  is invariant under  $f(T)$ ,  $f(T)|Y = f(T|Y)$ , and*

$$f(\sigma(T|Y)) = \sigma(f(T|Y)) = \sigma(f(T)|Y) \subseteq f(\sigma(T)).$$

*In addition, if  $T$ ,  $f(T)$ ,  $T|Y$ , and  $f(T)|Y$  have the single-valued extension property and  $x \in X$ , then we have*

$$\sigma_{f(T)|Y}([x]) = \sigma_{f(T|Y)}([x]) = f(\sigma_{T|Y}([x])) \subseteq f(\sigma_T(x)) = \sigma_{f(T)}(x).$$

#### REFERENCES

1. I. Colojoară and C. Foiaş, *The Riesz-Dunford functional calculus with decomposable operators*, Rev. Roumaine Math. Pures Appl. **12** (1967), 627–641. MR **39** #3334.
2. ———, *Theory of generalized spectral operators*, Gordon and Breach, New York, 1968.
- 3a. N. Dunford and J. T. Schwartz, *Linear operators. I: General theory*, Pure and Appl. Math., vol. 7, Interscience, New York, 1958. MR **22** #8302.
- 3b. ———, *Linear operators. III: Spectral operators*, Interscience, New York, 1971.
4. C. A. Kariotis, *Spectral properties of certain classes of operators*, Dissertation, University of Illinois, Urbana-Champaign, Ill., 1966.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ILLINOIS, URBANA-CHAMPAIGN, ILLINOIS 61801.

DEPARTMENT OF MATHEMATICS, GEORGIA SOUTHERN COLLEGE, STATESBORO, GEORGIA 30458.