QUASICONFORMAL EXTENSION OF HOLOMORPHIC MAPPINGS OF A BALL IN C^n

BY J. A. PFALTZGRAFF¹

Communicated by F. W. Gehring, October 16, 1973

Let C^n denote the space of n complex variables $z=(z_1, \dots, z_n)$ with Euclidean norm ||z||. The open unit ball $\{z \in C^n : ||z|| < 1\}$ is denoted by B^n . We consider holomorphic functions $f(z)=(f_1(z), \dots, f_n(z)), z \in B^n$, from B^n into C^n . The second derivative of such a function is a symmetric bilinear operator, $D^2f(z)(\cdot, \cdot)$ on $C^n \times C^n$, and $D^2f(z)(z, \cdot)$ is the linear operator obtained by restricting $D^2f(z)$ to $z \times C^n$, with matrix representation

$$D^2 f(z)(z,\cdot) = \left(\sum_{m=1}^n \frac{\partial^2 f_k(z)}{\partial z_i \partial z_m} z_m\right), \qquad 1 \leq j, k \leq n.$$

A locally biholomorphic mapping f(z) from a domain $G \subset C^n$ into C^n is said to be K-quasiconformal in G if $||Df(z)||^n \le K|\det Df(z)|$, $z \in G$, where || || denotes the standard operator norm $||A|| = \sup\{||Aw|| : ||w|| \le 1\}$, $A \in \mathcal{L}(C^n)$.

The purpose of this note is to announce the following n-dimensional $(n \ge 1)$ generalizations of one-dimensional results due to J. Becker [1].

THEOREM. Let f(z) with Df(0)=I be locally biholomorphic in B^n and satisfy

(1)
$$(1 - ||z||^2) ||(Df(z))^{-1}D^2f(z)(z, \cdot)|| \le c, \quad z \in B^n.$$

If $c \leq 1$ then f is univalent in B^n and

$$||z||/(1+c||z||)^2 \le ||f(z)|| \le ||z||/(1-c||z||)^2, \quad z \in B^n.$$

If f is K-quasiconformal in B^n and c<1 then f is univalent and continuous in the closed ball, \bar{B}^n , and f can be extended to a quasiconformal homeomorphism of R^{2n} onto R^{2n} .

For n=1, (1) is $|zf''(z)|f'(z)| \le c/(1-|z|^2)$, the local biholomorphy implies f is 1-quasiconformal in B^1 , and our theorem coincides with

AMS (MOS) subject classifications (1970). Primary 32A10; Secondary 30A36, 30A60

¹ Research supported in part by the U.S. Army Research Office—Durham Grant 31—124-72G182E.

Becker's results. The quasiconformal extension of f is not holomorphic on all of C^n , but viewed as a mapping of R^{2n} to R^{2n} , it is ACL, differentiable a.e., and the dilatation is uniformly bounded a.e. (cf. [5, p. 115]). By arguments similar to Becker's, we derive these results from our n-dimensional generalization [3] of Pommerenke's theory of subordination chains [4], and from the following lemma.

LEMMA. Let f(z) be locally biholomorphic and K-quasiconformal in B^n . If f satisfies (1) with c < 2 then

$$||Df(z)|| = O(1/(1 - ||z||)^c), \quad z \in B^n,$$

and f has a continuous extension to \bar{B}^n that satisfies a Lipschitz condition

(3)
$$||f(z) - f(w)|| \le M ||z - w||^{1-c}, \quad z, w \in \overline{B}^n.$$

The proof of (2) is fairly elementary, and does not require the use of subordination chains. The proof that (2) implies (3) depends upon n-dimensional versions of classical theorems of Hardy and Littlewood [2, pp. 361–363].

Complete proofs of our results and details of the theory of n-dimensional subordination chains will be submitted for publication elsewhere [3].

REFERENCES

- 1. J. Becker, Löwnersche Differentialgleichung und quasikonform fortsetzbare schlichte Funktionen, J. Reine Angew. Math. 255 (1972), 23-43. MR 45 #8828.
- 2. G. M. Goluzin, Geometric theory of functions of a complex variable, GITTL, Moscow, 1952; German transl., VEB Deutscher Verlag, Berlin, 1957; English transl., Transl. Math. Monographs, vol. 26, Amer. Math. Soc., Providence, R.I., 1969. MR 15, 112; 19, 735; 40 #308.
- 3. J. A. Pfaltzgraff, Subordination chains, univalence, and quasi-conformal extension of holomorphic mappings in C^n (manuscript).
- 4. Ch. Pommerenke, Über die Subordination analytischer Funktionen, J. Reine Angew. Math. 218 (1965), 159-173. MR 31 #4900.
- 5. J. Väisälä, Lectures on n-dimensional quasiconformal mappings, Lecture Notes in Math., vol. 229, Springer-Verlag, Berlin and New York, 1971.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NORTH CAROLINA AT CHAPEL HILL, NORTH CAROLINA 27514