

## EIGENVALUES ASSOCIATED WITH A CLOSED GEODESIC<sup>1</sup>

BY VICTOR GUILLEMIN AND ALAN WEINSTEIN

Communicated by I. M. Singer, September 15, 1974

1. **Background.** Intuitive arguments drawn from quantum mechanics and optics suggest that there should be some relation between the closed geodesics (periodic “particles”) on a compact Riemannian manifold  $X$  and the eigenvalues (periodic “waves”) of the Laplace-Beltrami operator  $\Delta_X$ . Indeed, in 1959, Huber [8] proved, for  $X$  a surface of constant negative curvature, that the set of lengths of closed geodesics on  $X$  and the spectrum of  $\Delta_X$  determine one another. The relation given by Huber between these two sequences of numbers is sufficiently complicated to make it extremely difficult to find one sequence explicitly, given the other.

Recently, Colin de Verdière [2], then Chazarain [1] and Duistermaat and Guillemin [3] have shown that, for most Riemannian metrics on any differentiable manifold, the spectrum of the Laplacian determines the lengths of the closed geodesics and their Morse indices modulo 4. Here, the lengths of the closed geodesics appear as the singular points of the distribution

$$\hat{o}(t) = \text{Trace}(e^{it\sqrt{\Delta_X}}) = \sum_{\lambda_j \in \text{Spec } \Delta_X} e^{i\sqrt{\lambda_j}t}$$

on the real line. This result, although very striking, leaves open some important questions. To apply it in any particular case, one would need to know a lot about  $\text{Spec } \Delta_X$  to get any information about the closed geodesics; even then, a formidable calculation would be involved in all but the simplest cases. In fact, when one is “handed” a Riemannian manifold, it is more likely that one knows something about the closed geodesics than about the spectrum.

In quantum mechanics, the eigenvalues are energy levels, and any available information about them is of interest. In a series of four papers culminating in [6], Gutzwiller used the method of Feynman integrals to find a contribution to the spectral density distribution

$$\sigma(\lambda) = \sum_{\lambda_j \in \text{Spec } \Delta_X} \delta(\lambda - \lambda_j)$$

corresponding to a single closed geodesic  $\gamma$ . For stable  $\gamma$ , this contribution is a series of  $\delta$ -functions at locations which are presumably approximate eigenvalues;

---

AMS (MOS) subject classifications (1970). Primary 35P20.

<sup>1</sup>The research described here was supported by NSF grants P22927 and GP34785X3.

for  $\gamma$  unstable, the contribution is a series of smooth peaks. Gutzwiller's arguments are quite physical and make no pretense of rigor, but his results have been very suggestive.

Recently, Voros [12] has clarified Gutzwiller's work in the case of stable closed geodesics and has provided simpler, though still heuristic, arguments based on the Keller-Maslov correction of the Bohr-Sommerfeld quantization condition. Essentially, Voros replaces the given classical system by its completely integrable linearization at the closed geodesic (in the cotangent bundle  $T^*X$ ) and sees which of the invariant tori satisfy the quantization condition.

The purpose of the present note is to announce a proof of Voros' result, based upon an extension of Hörmander's calculus of Fourier integral operators [5], [11].

**2. Statement of Theorem.** Before stating our theorem, we introduce some terminology and notation. Let  $\gamma$  be a closed geodesic on the  $k + 1$  dimensional manifold  $X$ . Denote by  $P_\gamma$  the linearized Poincaré map of  $\gamma$ ; it is a symplectic automorphism of the tangent space  $V_\gamma$  of a  $2k$ -dimensional manifold transverse to  $\gamma$  in the unit sphere bundle in  $T^*X$ .  $\gamma$  is called *nondegenerate elliptic* if the eigenvalues of  $P_\gamma$  are all distinct and have modulus one. The map  $P_\gamma$  then splits into a direct sum of two-dimensional rotations through angles  $\theta_1, \dots, \theta_k$  between 0 and  $2\pi$ . The *Morse index* of  $\gamma$  is the index of  $\gamma$  as a critical point of the length functional on the loop space of  $\gamma$ . Finally, our proof requires the assumption that  $T^*X$  admits a metaplectic structure [5]. This is a very weak requirement, satisfied for example if  $X$  is orientable, and probably not really necessary for the result to be true.

**THEOREM.** *If  $\gamma$  is a nondegenerate elliptic closed geodesic on  $X$  with length  $L$ , rotation angles  $\theta_1, \dots, \theta_k$ , and Morse index  $\mu$ , then for every  $k$ -tuple  $(n_1, \dots, n_k)$  of nonnegative integers there exists a sequence  $\lambda_1, \lambda_2, \dots$  of eigenvalues of  $\Delta_X$  satisfying*

$$\sqrt{\lambda_n} = L^{-1}(n_1\theta_1 + \dots + n_k\theta_k + 2\pi n + \mu) + O(n^{-1/2}).$$

**REMARK.** If one drops the term  $O(n^{-1/2})$  in (\*) and computes the resulting contribution from all these eigenvalues to  $\hat{\sigma}(t)$ , one obtains a pole at  $t = L$  whose residue is exactly that calculated in [3]. It remains to be seen whether the singularities of  $\hat{\sigma}(t)$  corresponding to unstable closed geodesics can also be traced to the spectrum itself.

**3. Comments on the proof.** The idea for the proof comes from a technique used by one of us [13] to demonstrate a theorem of Moslov [10] relating a sequence of eigenvalues of  $\Delta_X$  to a Lagrangian submanifold  $L$  contained in the unit sphere bundle of  $T^*X$ . (It is this theorem which Voros applies to the invariant tori.) The technique involves constructing out of  $L$  a *conic* Lagrangian submanifold  $\Lambda \subset T^*X \times T^*S^1$  and considering an associated Fourier integral

operator as an approximate intertwining operator between  $\Delta_{S^1}$  and  $\Delta_X$ . It was clear that an extension of the Fourier integral operator calculus from Lagrangian to isotropic submanifolds would permit an extension of Maslov's result to the case where  $L$  is isotropic and invariant under the geodesic flow. One such extension is provided by the complex Fourier integral operators of Melin and Sjöstrand [11], which leads to the part of our Theorem with  $(n_1, \dots, n_k) = 0$ . Another is given by the calculus developed in Guillemin [5]. The symbols of the intertwining operators in this calculus turn out to be the symplectic spinors [9] on  $V_\gamma$  which are invariant under  $P_\gamma$ . This accounts for the presence of the rotation angles in (\*) and the restriction of the Theorem to metilinear manifolds.

Further details of our proof may be found in the last section of [5].

Our collaboration was made possible by invitations to the Colloquia on Fourier Integral Operators (Nice, May 1974) and Symplectic Geometry and Mathematical Physics (Aix-en-Provence, June 1974), for which we would like to thank Professors J. Chazarain and J. M. Sourian.

#### REFERENCES

1. J. Chazarain, *Formule de Poisson pour les variétés riemanniennes*, Invent. Math. **24** (1974), 65–82. MR 49 #8062.
2. V. Colin de Verdière, *Spectre du Laplacien et longueurs des géodésiques périodiques*. II, Compositio Math. **27** (1973), 150–184.
3. J. J. Duistermaat and V. Guillemin, *The spectrum of positive elliptic operators and periodic geodesics*, Invent. Math. (to appear).
4. J. J. Duistermaat and L. Hörmander, *Fourier integral operators*. II, Acta Math. **128** (1972), 183–269.
5. V. Guillemin, *Symplectic spinors and partial differential equations*, C.N.R.S. Sympos. on Symplectic Geometry and Mathematical Physics, Aix-en-Provence, 1974 (to appear).
6. M. Gutzwiller, *Periodic orbits and classical quantization conditions*, J. Mathematical Phys. **12** (1971), 343–358.
7. L. Hörmander, *Fourier integral operators*. I, Acta Math. **127** (1971), 79–183.
8. H. Huber, *Zur analytischen Theorie hyperbolischer Raumformen und Bewegungsgruppen*, Math. Ann. **138** (1959), 1–26. MR 22 #99.
9. B. Kostant, *Symplectic spinors*, Symposia Mathematica (to appear).
10. V. P. Moslov, *Théorie des perturbations et méthodes asymptotiques*, Izdat. Moskov. Gos. Univ., Moscow, 1965; French transl., Dunod, Paris, 1972.
11. A. Melin and J. Sjöstrand, *Fourier integral operators with complex-valued phase functions*, Sympos. on Fourier Integral Operators (Nice, 1974), Lecture Notes in Math., Springer-Verlag, Berlin and New York (to appear).
12. A. Voros, *The WKB-Moslov method for nonseparable systems*, C.N.R.S. Sympos. on Symplectic Geometry and Mathematical Physics, Aix-en-Provence, 1974 (to appear).
13. A. Weinstein, *On Moslov's quantization condition*, Sympos. on Fourier Integral Operators (Nice, 1974), Lecture Notes in Math., Springer-Verlag, Berlin and New York (to appear).

DEPARTMENT OF MATHEMATICS, MASSACHUSETTS INSTITUTE OF TECHNOLOGY, CAMBRIDGE, MASSACHUSETTS 02139

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, BERKELEY, CALIFORNIA 94720