

this the reader will have to consult other works (such as Stein-Weiss [S-W], Stein [S], and the more recent Garcia-Cuerva–Rubio de Francia [G-R]).

This is a book for mature readers. The author does not hesitate to use ideas and results from diverse branches of mathematics—special functions, functional analysis, partial differential equations, differential geometry. But for the reader with a strong background, or a willingness to accept a non-self-contained presentation, this book offers many pleasures. In addition to the concrete computational results already mentioned, the book contains concise and insightful presentations of a number of important abstract topics, including induced representations, representations of compact groups, conformal transformations, Clifford algebras and spinors. Taylor has an original point of view, and is able to bring new insights to familiar topics.

#### REFERENCES

[F-S] G. Folland and E. M. Stein, *Estimates for the  $\bar{\partial}_b$  complex and analysis on the Heisenberg group*, *Comm. Pure Appl. Math.* **27** (1974), 429–522.

[G-R] J. Garcia-Cuerva and J. L. Rubio de Francia, *Weighted norm inequalities and related topics*, North-Holland, 1985.

[H] R. Howe, *On the role of the Heisenberg group in harmonic analysis*, *Bull. Amer. Math. Soc. (N.S.)* **3** (1980), 821–843.

[S] E. M. Stein, *Singular integrals and differentiability properties of functions*, Princeton Univ. Press, Princeton, N. J., 1970.

[S-W] E. M. Stein and G. Weiss, *Introduction to Fourier analysis on Euclidean spaces*, Princeton Univ. Press, Princeton, N. J., 1971.

ROBERT S. STRICHARTZ

BULLETIN (New Series) OF THE  
AMERICAN MATHEMATICAL SOCIETY  
Volume 17, Number 1, July 1987  
©1987 American Mathematical Society  
0273-0979/87 \$1.00 + \$.25 per page

*Near-rings and their links with groups*, by J. D. P. Meldrum, Pitman Publishing, Boston, London, Melbourne, 275 pp., \$44.95. ISBN 0-470-20648-9.

Many mathematicians do not highly appreciate theories which have prefixes like near-, semi-, hemi-, para-, quasi-, and so on. This certainly should not apply in the case of near-rings. As the name suggests, a *near-ring* is a “generalized ring”; more precisely, the commutativity of addition is not required and just one of the distributive laws is postulated.

Near-rings arise very naturally in the study of mappings on groups. If  $(G, +)$  is a group (not necessarily abelian) then the set  $M(G)$  of all mappings from  $G$  to  $G$  is a near-ring with respect to pointwise addition and composition of mappings. If  $G$  is abelian and if one only takes “linear” maps

(= endomorphisms of  $G$ ), one gets the well-known endomorphism ring  $\text{End } G$ . From that, one already sees that the generalization from rings to near-rings basically is the transition from linearity to nonlinearity. And this is a matter truly worth being considered.

At the beginning of this century, L. E. Dickson [6] constructed the first examples of “proper” near-rings. Actually, these near-rings were *near-fields* (= near-rings in which the nonzero elements form a multiplicative group). Dickson started with a field  $F = (F, +, \cdot)$ , took a “coupling map”  $\Phi: F - \{0\} \rightarrow \text{Aut}(F, +)$ ,  $f \rightarrow \phi_f$  with  $\phi_f \circ \phi_g = \phi_{(f\phi_g) \cdot g}$  (we write  $xh$  for the image of  $x$  under the function  $h$ ). Defining  $f \cdot_{\Phi} g := (g\phi_f) \cdot g$  and  $0 \cdot_{\Phi} f := 0$ ,

$$F^{\Phi} := (F, +, \cdot, \cdot_{\Phi}),$$

the “ $\Phi$ -derivation” of  $F$ , is a near-field, and in general not a field. Near-fields which arise as  $F^{\Phi}$  from some field  $F$  are now called *Dickson near-fields*. Hence Dickson showed that the second distributive law does not follow from the remaining axioms of a skew-field. But the commutativity of addition does, and this is certainly not trivial.

**THEOREM 1** (B. H. NEUMANN [12], KARZEL [10], ZEMMER [19], ET AL.) *The additive group of a near-field is abelian.*

In 1936, H. Zassenhaus succeeded in finding all finite near-fields:

**THEOREM 2** (ZASSENHAUS [18]). *A finite near-field is either a Dickson near-field, or it belongs to 8 exceptional near-fields or orders 2, 5<sup>2</sup>, 7<sup>2</sup>, 11<sup>2</sup> (two near-fields), 23<sup>2</sup>, 29<sup>2</sup> or 59<sup>2</sup>.*

The 7 exceptional near-fields of order  $p^2$  in Theorem 2 can easily be described by means of  $2 \times 2$ -matrices. The exceptional near-field of order 2 is  $(\mathbf{Z}_2, +, *)$  with  $x * y = y$ . For more details see, e.g., [3].

From 1907 on, Veblen, and Wedderburn [15] and many other authors used near-fields  $N$  in order to coordinatize geometric planes  $\mathcal{G}$ , so the points of  $\mathcal{G}$  are just the elements of  $N \times N$  and the lines of  $\mathcal{G}$  are given by all  $\{(x, xa + b) \mid x \in N\}$  and  $\{(c, x) \mid x \in N\}$  for  $a, b, c \in N$ ,  $a \neq 0$ . For more information on this topic see e.g. [8] or [5].

Another application of near-fields is in the description of all *sharply 2-transitive* permutation groups  $G$  on some set  $X$ . This means that for all distinct  $x_1, x_2 \in X$  and all distinct  $y_1, y_2 \in X$  there is precisely one  $g \in G$  with  $x_i g = y_i$  ( $i = 1, 2$ ).

**THEOREM 3** (ZASSENHAUS [18], BLACKBURN AND HUPPERT [3]). *For every finite sharply 2-transitive permutation group  $G$  there is a finite near-field  $F$  such that  $G$  is isomorphic to all transformations  $x \rightarrow ax + b$  ( $a, b \in F$ ,  $a \neq 0$ ) from  $F$  to  $F$ .*

Thus, by Theorem 3 in conjunction with Theorem 2, all finite sharply 2-transitive groups can be considered to be “known.” More can be found in [3]

and [11]. Near-fields have numerous other applications, mostly to geometry, which are described in the new book [16].

J. D. P. Meldrum's book does give some information on these near-fields, but it concentrates much more on the exciting (and mostly quite recent) theories and applications of "proper" near-rings (which are neither near-fields nor rings). It consists of two parts. Part 1 gives a very readable and thoughtful introduction to the subject, mentions many examples and some applications, and then proceeds quickly to the structure theory of near-rings (radicals, semisimplicity, primitive and simple near-rings). The main difference from the existing book [13] is that Meldrum's book is much easier to read and concentrates on the core of the theory, while [13] was in some way intended to be encyclopedic.

At a first glance, the structure theory (initiated in [17] and in the dissertation [4], supervised by E. Artin) of near-rings seems to run along very similar lines as the one of ring theory. It does, to a large extent, in the results; but most of the deeper results need completely different proofs and approaches. As an example, let me present Theorem 3.35 in the book, which many people consider to be the basic structure result on near-rings.

If  $N$  is a near-ring and if  $(G, +)$  is a group such that for  $n \in N$  and  $g \in G$  a "product"  $gn$  is defined such that  $g(n_1 + n_2) = gn_1 + gn_2$ ,  $(gn_1)n_2 = g(n_1n_2)$  hold for all  $g \in G$  and  $n_1, n_2 \in G$ , then  $G$  is called an  $N$ -group (or  $N$ -near-module). Every group  $G$  is, for example, an  $M(G)$ -group and also a  $\mathbf{Z}$ -group in the obvious ways. For any near-ring  $N$ ,  $N_0 := \{n \in N \mid 0n = 0\}$  is called its *zero-symmetric part*.  $N_0$  is a subnear-ring of  $N$ . If  $N = M(G)$ ,  $N_0 := M_0(G)$  consists of the maps  $G \rightarrow G$  which map 0 into 0. If  $G$  is an  $N$ -group with  $GN \neq 0$  and without proper subgroups  $H$  such that  $HN_0 \subseteq H$ ,  $G$  is called *irreducible*. If the *annihilator*  $A(G_N) := \{n \in N \mid Gn = 0\}$  is 0,  $G$  is said to be a *faithful*  $N$ -group. A near-ring  $N$  with a faithful, irreducible  $N$ -group  $G$  is called *primitive* (on  $G$ ). Since for near-rings it makes sense to study different concepts of primitivity, the described concept is more precisely called "2-primitivity" in the book. An endomorphism  $h \in \text{End } G$  is called an  $N$ -homomorphism if  $h(gn) = h(g)n$  for all  $g \in G$ ,  $n \in N$ .  $\text{Hom}_N(G, G)$  is the collection of all  $N$ -homomorphisms of the  $N$ -group  $G$ .

If  $S$  is a set of endomorphisms of a group  $(G, +)$ ,

$$M_S(G) := \{f \in M_0(G) \mid f \circ s = s \circ f \text{ for all } s \in S\}$$

is an important subnear-ring of  $M(G)$ . If  $N$  and  $M$  are subnear-rings of  $M(G)$  with  $N \subseteq M$ , we say that  $N$  is *dense* in  $M$  if for all  $k \in \mathbf{N}$ , all distinct  $g_1, \dots, g_k \in G$  and all  $m \in M$  there is some  $n \in N$  with  $g_i n = g_i m$  ( $i = 1, \dots, k$ ). If we take, for  $g \in G$  and  $m \in M$ , all  $\{m' \in M \mid gm' = gm\}$  as a base of a topology, the resulting concept of density is the one just defined.

We are now ready for the following result; parts of it were first obtained by H. Wielandt (unpublished manuscripts). The form as given now is due to G. Betsch [1].

**DENSITY THEOREM.** *Let  $N = N_0$  be a near-ring with identity which is primitive on  $G$ .*

(i) *If  $N$  is a ring then Jacobson's density theorem [9] applies:  $N$  is a dense subring of  $\text{Hom}_F(G, G)$ , where  $F$  is the skew-field  $\text{Hom}_N(G, G)$ .*

(ii) *If  $N$  is not a ring then  $N$  is a dense subnear-ring of  $M_S(G)$ , where  $S := \text{Aut}(G, +) \cap \text{Hom}_N(G, G)$  is a fixed-point-free automorphism group on  $G$ .*

The proof is certainly nontrivial. Part (ii) has to be set apart from case (i) by a careful investigation of the lattice of "ideals" of the  $N$ -group  $G$ . Many variants, generalizations, specializations, and applications of this theorem are known. A convenient special case is

**COROLLARY.** *Let the "non-ring"  $N = N_0$  with identity be primitive on  $G$ , such that  $S$  (as in the Density Theorem) =  $\{\text{id}\}$ . Then  $N$  is dense in  $M_0(G)$ .*

Of course, for finite  $G$  (more generally for near-rings  $N$  with a suitable descending chain condition), "density" is the same as "equality." Some applications of the Density Theorem are mentioned in [14].

Part 2 of Meldrum's book considerably differs from the first part. It starts from the observation that the sum of two endomorphisms of a nonabelian group  $(G, +)$  is, in general, not an endomorphism any more. Hence it makes sense to study the collection of all sums/differences of endomorphisms of  $G$ , i.e., the additive closure  $E(G)$  of  $\text{End } G$  in  $M(G)$ . It quickly turns out that  $E(G)$  is a near-ring (subnear-ring of  $M(G)$ );  $E(G)$  is the most prominent example of a "distributively generated" near-ring (that is, a near-ring which is additively generated by "distributive elements"). These distributively generated near-rings are thoroughly investigated in part 2 of J. D. P. Meldrum's book. After presenting the basic facts of distributively generated near-rings (in particular: of *endomorphism near-rings*  $E(G)$ ), the author studies their structure, considers free distributively generated near-rings and "group distributively generated near-rings", thus initiating a "nonlinear representation theory" for groups. A similar sounding, but different, topic is how to represent an arbitrary distributively generated near-ring as a subnear-ring of a suitable  $E(G)$ . Since the relation between  $G$  and  $E(G)$  is, of course, much stronger than that between  $G$  and  $M(G)$ , a lot of group theory enters part 2 of the book. The information flow is by no means just one-sided from groups to endomorphism near-rings, but goes the other way, too: near-ring theory faithfully pays back to groups (part of) the amount groups have invested into near-rings. For example, the following result can be derived from the Density Theorem mentioned above (J. D. P. Meldrum gives a very nice new proof of it):

**THEOREM 4.** *For a group  $G$ , every map  $G \rightarrow G$  fixing 0 is the sum/difference of inner automorphisms iff  $G$  is a finite, simple, nonabelian group.*

There are, of course, many interesting open problems in this area. A "nonabelian homological algebra," initiated by A. Fröhlich in [7], is still to be developed. A striking problem, withstanding all attacks up to now, is the following.

**PROBLEM.** Can every zero-symmetric near-ring be embedded in an endomorphism near-ring?

In other words: given a collection of 0-fixing mappings from a group  $G$  into itself, can we enlarge  $G$  to some bigger group  $H$  such that all of the given maps become “partly linear” (i.e., the sums/differences of endomorphisms of  $H$ )?

Summarizing: Meldrum’s book is an excellent piece of work. Part 1 gives an easy introduction into the subject, most useful both for beginners as well as other mathematicians who just want to see the main ideas. Part 2 goes considerably deeper and reflects the author’s mastery of group theory; it also contains research results of the author which appear in a book for the first time.

The book is written very carefully and thoughtfully. This applies to the mathematical precision as well as to the style and also to the typing; I know that the author put the manuscript on a word processor himself; this guaranteed that everything is so arranged that it provides maximal clarity. The reader might miss more detailed descriptions of applications of near-rings, like the ones to block designs, but this can be done in further reading ([13], the new proceedings volume [2] or, e.g., [14]). Altogether, this book should *not* stand on a shelf: it should be read and used.

#### REFERENCES

1. G. Betsch, *Primitive near-rings*, Math. Z. **130** (1973), 351–361.
2. G. Betsch, (ed.), *Near-rings and near-fields*, North-Holland, Amsterdam, 1987.
3. N. Blackburn and B. Huppert, *Finite groups*. III, Springer, Berlin-Heidelberg-New York, 1982.
4. D. W. Blackett, *Simple and semi-simple near-rings*, doctoral dissertation, Princeton University, 1950.
5. P. Dembrowski, *Finite geometries*, Springer, Berlin, 1968.
6. L. E. Dickson, *Definitions of a group and a field by independent postulates*, Trans. Amer. Math. Soc. **6** (1905), 198–204.
7. A. Fröhlich, *Non-abelian homological algebra*. I, *Derived functors and satellites*, Proc. London Math. Soc. **11** (1961), 239–275.
8. M. Hall, *The theory of groups*, McMillan, New York, 1959.
9. N. Jacobson, *Structure of rings*, Amer. Math. Soc. Colloq. Publ., vol. 37, Providence, R. I., 1968.
10. H. Karzel, *Bericht über projektive Inzidenzgruppen*, Jahresber. Dt. Math. Ver. **67** (1965), 58–92.
11. W. E. Kerby, *On infinite sharply multiply transitive groups*, Vandenhoeck and Ruprecht, Göttingen, 1974.
12. B. H. Neumann, *On the commutativity of addition*, J. London Math. Soc. **15** (1940), 203–208.
13. G. Pilz, *Near-rings*, North-Holland, Amsterdam, second edition, 1983.
14. ———, *What near-rings can do for you*, Contrib. to General Algebra 4, Teubner, Stuttgart, 1987.
15. O. Veblen and J. H. Wedderburn, *Non-desarguesian and non-pascalian geometries*, Trans. Amer. Math. Soc. **8** (1907), 379–388.
16. H. Wähling, *Theorie der Fastkörper*, Thales Verlag, Essen, 1987.
17. H. Wielandt, *Über Bereiche aus Gruppenabbildungen*, Deutsche Mathematik **3** (1938), 9–10.
18. H. Zassenhaus, *Über endliche Fastkörper*, Abh. Math. Sem. Univ. Hamburg **11** (1935/36), 187–220.
19. J. L. Zemmer, *The additive group of an infinite near-field is abelian*, J. London Math. Soc. **44** (1969), 65–67.