

Existence of Solitary Waves in Higher Dimensions*

Walter A. Strauss

Department of Mathematics, Brown University, Providence, Rhode Island 02912, USA

Abstract. The elliptic equation $\Delta u = F(u)$ possesses non-trivial solutions in \mathbb{R}^n which are exponentially small at infinity, for a large class of functions F . Each of them provides a solitary wave of the nonlinear Klein-Gordon equation.

1. Introduction

We define a solitary wave as a solution $\phi(x, t)$ of a wave equation whose maximum amplitude at time t , $\sup_x |\phi(x, t)|$, does not tend to zero as $t \rightarrow \infty$, but which tends to zero in some convenient sense as $|x| \rightarrow \infty$ for each t . The convergence should have the property that physical quantities, such as the energy and charge, are finite. Particular types of solitary waves are (1) traveling waves $\phi = u(x - ct)$ where c is a constant vector and (2) standing waves $\phi = \exp(i\omega t)u(x)$ where ω is a real constant. Traditionally, solitary waves have been traveling waves, but in recent years oscillatory factors have been allowed. The above definition includes all uses of the term. Solitary waves have also been called “solitons” but, properly speaking, the latter word should be reserved for those special solitary waves which exactly preserve their shapes after interaction. Many examples of these special solitons have been discovered in recent years in the case of two space-time dimensions. In higher dimensions, however, even the existence of solitary waves seems to be elusive.

We consider the scalar NLKG equation

$$\phi_{tt} - \Delta \phi + m^2 \phi + f(\phi) = 0, \tag{3}$$

where $x = (x_1, \dots, x_n) \in \mathbb{R}^n$, Δ is the Laplacian in x and $m > 0$. We assume $f(0) = 0$ and $f(re^{i\theta}) = f(r)e^{i\theta}$. If ϕ is a standing wave (2), Equation (3) reduces to

$$-\Delta u + (m^2 - \omega^2)u + f(u) = 0. \tag{4}$$

We shall show that (4) possesses non-trivial solutions exponentially small at infinity provided $|\omega| < m$ and f satisfies certain conditions. In particular, if $\omega = 0$, we have $\phi(x) = u(x)$. Since we may change to a different Lorentz frame, it follows that there exist traveling solitary waves (1) for any $|c| < 1$. Alternatively we may proceed by

* This work was supported in part by NSF Grant MCS 75-08827

putting (1) directly into (3) to obtain the equation

$$-\sum a_{ij} \frac{\partial^2 u}{\partial x_i \partial x_j} + m^2 u + f(u) = 0, \tag{5}$$

where $a_{ij} = \delta_{ij} + c_i c_j$. Note that (a_{ij}) is a positive-definite matrix because

$$\sum a_{ij} \xi_i \xi_j = |\xi|^2 - (c \cdot \xi)^2 \geq (1 - |c|^2) |\xi|^2$$

for any $\xi = (\xi_1, \dots, \xi_n) \in \mathbb{R}^n$. Thus Equation (5) is elliptic and it can be converted into an equation with leading term $-\Delta v$ by means of a rotation and stretching of axes. The same sort of equation comes from standing wave solutions (2) of the NLS equation $i\phi_t - \Delta\phi + f(\phi) = 0$.

In each of these cases we get an equation of the form

$$-\Delta u + F(u) = 0, \quad x \in \mathbb{R}^n, \tag{6}$$

where $F(u) = f(u) + (\text{const.})u$.

We assume that $F(0) = 0$, which means that (6) always possesses the trivial solution $u = 0$. Let F be a real continuous function and let $G' = F$, $G(0) = 0$. Consider real-valued solutions of (6); for the complex case, see Section 2.

Some necessary conditions on F , found in part by Pohozaev [7], are the following.

Theorem 1. *If u is a solution of (6) which is sufficiently small at infinity, then*

$$(n-2) \int |\nabla u|^2 dx = -(n-2) \int uF(u) dx = -2n \int G(u) dx. \tag{7}$$

Hence if $sF(s)$ or $G(s)$ (for $n \neq 1$) or $H(s) = (n-2)sF(s) - 2nG(s)$ or $-H(s)$ is positive (for $s \neq 0$), then the only solution is the trivial one. For any non-trivial solution, the energy is positive:

$$E = \int [\frac{1}{2} |\nabla u|^2 + G(u)] dx = \frac{1}{n} \int |\nabla u|^2 dx > 0.$$

The main result of this paper is the existence of non-trivial solutions of (6) for a large class of functions F . Let

$$L = -\sum a_{ij} \frac{\partial^2}{\partial x_i \partial x_j} + a_0,$$

where the constant matrix (a_{ij}) is positive-definite and a_0 is a positive constant. Let $F_1(s)$ and $F_2(s)$ be real continuous functions defined for $0 \leq s < \infty$. Denote by G_1 and G_2 their indefinite integrals. Assume the following.

$$F_1(s) \geq 0, \quad F_2(s) > 0 \quad \text{for } s > 0. \tag{8}$$

$$\text{As } s \rightarrow 0, \quad F_1(s) = O(s) \quad \text{and} \quad F_2(s) = o(s). \tag{9}$$

$$F_2(s) = o(s^l + F_1(s)) \quad \text{as } s \rightarrow \infty. \tag{10}$$

$$F_2(s) = O(s^l + G_1(s)/s) \quad \text{as } s \rightarrow \infty, \tag{11}$$

where $l = (n+2)/(n-2)$ and $n \geq 3$. (For the cases $n = 1$ or 2 , see Section 3.)

Theorem 2. *Under these assumptions, there exists $\lambda > 0$ and a solution u of*

$$Lu + F_1(u) = \lambda F_2(u) \tag{12}$$

which is non-negative, belongs to the Sobolev space H^1 , decays exponentially as $|x| \rightarrow \infty$, and $\int G_1(u(x)) dx < \infty$.

Examples are presented in Section 2.

A number of studies have been made of Equation (6) by restricting attention to the radial solutions ($u = u(r)$, $r = |x|$) and analyzing the resulting ordinary differential equation; see Synge [14], Nehari [6], Anderson and Derrick [1], and Berger [2]. One-dimensional problems have also been studied by Stuart [13] and Dancer [4]. In addition, there are a large number of papers on the existence of a solution of an equation with an inhomogeneous term $g(x)$; in the present case, when $g = 0$, there is always the trivial solution $u = 0$.

Nonlinear eigenvalue problems of this type have also been much studied for the case of a bounded domain $\Omega \subset \mathbf{R}^n$ with Dirichlet boundary conditions. Methods of the calculus of variations, bifurcation theory and topological degree have been used; see Pohozaev [7], Rabinowitz [9, 10], for example. These methods fail to extend to unbounded Ω because H^1 is no longer compact in L^2 . To be compact in L^2 on an unbounded domain, a class of functions has to be uniformly small at infinity. We use the variational method with the extra constraint that the functions be radial, which ensures the compactness.

We allow rapid growth of the nonlinearities in Theorem 2. Almost all authors (for Ω bounded or not) require $F_i(s) = o(s^i)$ as $s \rightarrow \infty$, $i = 1, 2$. Clément [3], for the case of (12) in a bounded domain, relaxes this condition for F_1 but requires it for F_2 . Rabinowitz [11] exhibits an example where F_2 does not satisfy the growth condition and $F_1 = 0$.

In Section 3 we prove Theorem 2 in two stages. We apply the direct variational method to truncated versions of F_1 and F_2 . Then we remove the truncations by a limiting process, using techniques of Strauss [12] and Clément [3].

Classical eigenvalue problems possess an infinite sequence of solutions. By the minimax technique, the same should be true in our case. We have not carried this out; instead, in Section 4, we apply abstract theorems of Rabinowitz [10]. These theorems unfortunately require the growth conditions on F_1 and F_2 mentioned above.

I wish to thank L. Vazquez for many stimulating discussions on this topic and P. Rabinowitz and C. Dafermos for some helpful suggestions.

2. Examples and Necessary Conditions

We use the notation $L^p = L^p(\mathbf{R}^n)$, $\| \cdot \|_p = L^p$ -norm, $H^1 = H^1(\mathbf{R}^n) = \{u \in L^2 \mid \partial u / \partial x_i \in L^2, i = 1, \dots, n\}$. Integrals are taken over all of \mathbf{R}^n unless otherwise specified.

Example 1 (Pohozaev [7]).

$$-\Delta u + u - |u|^{q-1}u = 0,$$

$x \in \mathbf{R}^n$, $n \geq 3$, $q > 1$. We have scaled out the coefficients to be 1. To apply Theorem 1, $F(s) = s - |s|^{q-1}s$, $G(s) = s^2/2 - |s|^{q+1}/(q+1)$. Let $\alpha^{-1} = 2^{-1} - (q+1)^{-1}$. So there is no non-trivial solution if $((n-2)/2)sF(s) - nG(s) = -s^2 + (1 - \alpha^{-1}n)|s|^{q+1}$ is of one sign; that is, $\alpha \leq n$ or $q \geq (n+2)/(n-2)$. So assume $1 < q < (n+2)/(n-2)$. Any solution must satisfy identity (7), which reduces to:

$$\alpha(n-2) \int |\nabla u|^2 dx = \frac{n\alpha}{\alpha-n} \int |u|^2 dx = n \int |u|^{q+1} dx.$$

Theorem 2 asserts the existence of a solution, where $F_1(s)=0$, $F_2(s)=|s|^{q-1}s$, $L=-\Delta+I$ and a scale change is used to make $\lambda=1$. In Section 4 we show that this solution is the first one of an infinite sequence of distinct solutions.

Example 2. $-\Delta u+(m^2-\omega^2)u+|u|^{p-1}u-\lambda|u|^{q-1}u=0$ where $x\in\mathbf{R}^n$, $m^2-\omega^2>0$ and p and q are distinct numbers larger than 1. We distinguish four cases.

Case A: $q<p$. Theorem 2 asserts the existence of a non-trivial solution for some $\lambda>0$. Note that

$$G(s)=\frac{1}{2}(m^2-\omega^2)s^2+\frac{1}{p+1}|s|^{p+1}-\frac{\lambda}{q+1}|s|^{q+1}$$

is bounded below. There is a number λ_* so that, for $\lambda\leq\lambda_*$, $G(s)$ is non-negative and the only solution is the trivial one, according to Theorem 1.

According to Theorem 1, the (integrated) energy is necessarily positive, even though the function $G(s)$ is not allowed to be positive if a non-trivial solution is to exist. On the other hand, for a standing wave solution (2) of the NLKG Equation (3), the energy density

$$\frac{1}{2}|\phi_t|^2+\frac{1}{2}|\nabla\phi|^2+G(\phi)=\frac{1}{2}|\nabla u|^2+\omega^2u^2+G(u)$$

may be positive. This is the case if $\omega>0$ and λ is slightly larger than λ_* .

Anderson [1] has computed these solutions in the case $n=3$, $p=5$, $q=3$. His most interesting result is that the positive solution appears to be stable with respect to perturbations of the initial data of Equation (3) in the cases when the energy density is positive. For this choice of p , q , and n , the inequalities

$$\lambda s^4=(2s)(\frac{1}{2}\lambda s^3)\leq\frac{1}{2}(2s)^2+\frac{1}{2}(\frac{1}{2}\lambda s^3)^2=2s^2+\frac{1}{8}\lambda^2s^6,$$

$$G(s)=\frac{1}{2}s^2+\frac{1}{6}s^6-\frac{\lambda}{4}s^4\geq(\frac{1}{6}-\frac{1}{32}\lambda^2)s^6$$

show that $\lambda_*=4\cdot 3^{-1/2}$, if we normalize to make $m^2-\omega^2=1$. This explains the rather mysterious fraction $3/16=\lambda_*^{-2}$ appearing in [1].

Case B: $p<q<(n+2)/(n-2)$. Theorem 2 is again applicable. In Section 4 we prove the existence of an infinite sequence of non-trivial solutions for each $\lambda>0$.

Case C: $p\leq(n+2)/(n-2)\leq q$. Let $\alpha^{-1}=2^{-1}+(q+1)^{-1}$ and $\beta^{-1}=2^{-1}+(p+1)^{-1}$. Then $\alpha\leq n\leq\beta$ and

$$\frac{n-2}{2}sF(s)-nG(s)=-s^2-\left(1-\frac{n}{\beta}\right)|s|^{p+1}+\left(1-\frac{n}{\alpha}\right)\lambda|s|^{q+1}.$$

By Theorem 1, there can be no non-trivial solution.

Case D: $(n+2)/(n-2)<p<q$. This case remains open: we do not know whether or not there exists a non-trivial solution.

Remark. For the equation $-\Delta u+F(u)=0$, existence essentially requires $F'(0)\geq 0$. Indeed, suppose $-\alpha=F'(0)<0$ and let $f(s)=F(s)/s+\alpha$. Let $u(x)$ be a non-trivial

solution small at infinity and let $q(x) = f(u(x))$. Then the equation can be rewritten as $-\Delta u + qu = \alpha u$. Assuming $u(x)$ is small enough at infinity that $q(x) = o(|x|^{-1})$, it is well-known that the operator $-\Delta + q$ has no positive eigenvalues (Kato [5]), so that we reach a contradiction.

Remark. A word about real vs. complex solutions. If we restrict our consideration to real solutions $u(x)$, then we consider real functions $F(s)$ and $G(s) = \int_0^s F(\sigma) d\sigma$. If we allow $u(x)$ to be complex-valued, then we assume the following.

$$g : [0, \infty] \rightarrow \mathbf{R} ,$$

$$G(z) = g(|z|) \quad \text{for complex } z ,$$

$$F(z) = g'(z)z/|z| .$$

It follows that

$$\frac{\partial}{\partial \varepsilon} G(z + \varepsilon \zeta)|_{\varepsilon=0} = \operatorname{Re} F(z) \bar{\zeta}$$

and

$$\frac{\partial}{\partial x_j} G(u(x)) = \operatorname{Re} F(u(x)) \frac{\partial \bar{u}}{\partial x_j} .$$

In this case, the middle term in Equation (7) should be $(2 - n) \operatorname{Re} \int \bar{u} F(u) dx$.

Proof of Theorem 1. We have $-(\Delta u)\bar{u} = -\nabla(\nabla u \bar{u}) + |\nabla u|^2$. So if we multiply Equation (6) by \bar{u} , integrate, and assume that u and its derivatives are small enough at infinity, we obtain

$$\int \{ |\nabla u|^2 + \operatorname{Re} \bar{u} F(u) \} dx = 0 .$$

On the other hand, the multiplier $r\partial\bar{u}/\partial r = \sum_i x_i \bar{u}_i$, where we denote derivatives by subscripts, gives the identities:

$$-\operatorname{Re} u_{jj} x_i \bar{u}_i = -\operatorname{Re} (u_j x_i \bar{u}_i)_j + (\frac{1}{2} x_i |u_j|^2)_i + \left(1 - \frac{n}{2} \right) |u_j|^2 ,$$

$$\operatorname{Re} F(u) x_i \bar{u}_i = (x_i G(u))_i - nG(u) ,$$

$$\int \{ (n-2) |\nabla u|^2 + 2nG(u) \} dx = 0 .$$

This proves (7). It may be illuminating to give a different proof of the second equality in (7). We regard u as a solution of the variational problem $\delta \mathcal{L}(u) = 0$, where

$$\mathcal{L}(u) = \int \{ \frac{1}{2} |\nabla u|^2 + G(u) \} dx .$$

Motivated by the fact that $r\partial/\partial r$ is the infinitesimal generator of the scale change $u(x) \rightarrow u(\lambda x)$, we define $u_\lambda(x) = \lambda^{(n-2)/2} u(\lambda x)$ for $\lambda > 0$. Then

$$\mathcal{L}(u_\lambda) = \int \{ \frac{1}{2} \lambda^n |(\nabla u)(\lambda x)|^2 + G(\lambda^{(n-2)/2} u(\lambda x)) \} dx$$

Hence

$$0 = \frac{d}{d\lambda} \mathcal{L}(u_\lambda)|_{\lambda=1} = \int \left\{ -nG(u) + \frac{n-2}{2} \operatorname{Re} \bar{u}F(u) \right\} dy .$$

Remark. The last proof is based on the transformation $u \rightarrow u_\lambda$, which leaves the Dirichlet integral invariant. For which functions G does it also leave the integral $\int G(u)dx$ invariant? Exactly those G for which $2nG(z) = (n-2) \operatorname{Re} \bar{z}F(z)$. That is, $2ng(s) = (n-2)sg'(s)$, whence

$$G(z) = -C|z|^{q+1}, \quad q = \frac{n+2}{n-2}, \quad \text{if } n > 2 .$$

Now a critical point of the functional $\mathcal{L}(u)$ is the same as a critical point of the Dirichlet integral subject to the constraint $\int |u|^{q+1} dx = 1$. Let

$$m = \inf \{ \int |\nabla u|^2 dx \}^{1/2} \quad \text{subject to} \quad \int |u|^{q+1} dx = 1$$

(u a test function), then $m \|u\|_{q+1} \leq \|\nabla u\|_2$. This is the classical Sobolev inequality.

We are going to prove Theorem 2 by means of the variational problem

$$\min \int \left\{ \frac{1}{2} |\nabla u|^2 + \frac{1}{2} u^2 + G_1(u) \right\} dx$$

subject to the constraint $\int G_2(u) dx = 1, u \in H^1$. The following theorem serves as motivation for our proof. We have made some simplifications; in particular, we consider the real case only.

Theorem 3. *Let F_1 and F_2 be continuous functions satisfying (8) and (9) and $F_i(s) = O(s)$ as $s \rightarrow \infty$. Extend $F_i(s)$ to be zero for $s < 0$. If the above problem has a solution, then it has a solution which is non-negative, radial and exponentially decaying.*

Proof. Let u be a solution. Let $u^+(x) = \max(u(x), 0)$. Then $G_i(u^+) = G_i(u), \nabla(u^+) = (\nabla u)^+$ and $u^+ \in H^1$. So u^+ is a solution of the same minimum problem and of course u^+ is non-negative. In the rest of the proof we assume $u \geq 0$. To find a radial solution we use Steiner symmetrization. In the classical use of this technique (Polya and Szegö [8], Weinberger [15]), the integrands are quadratic, the equation is linear and the domain is bounded. However, the proof easily extends to the present case. Let $D = \{(x, t) \in \mathbf{R}^{n+1} : 0 \leq t \leq u(x)\}$. Let D^* be the symmetrization of D in the hyperplane $x_1 = 0$. This means that each line L in \mathbf{R}^{n+1} perpendicular to $x_1 = 0$ intersects D^* in a single line segment symmetric about $x_1 = 0$ such that the length of $L \cap D^*$ equals the length of $L \cap D$. D^* is of the form $D^* = \{(x, t) \in \mathbf{R}^{n+1} : 0 \leq t \leq u^*(x)\}$. Then (for $i = 1, 2$)

$$\begin{aligned} \int G_i(u(x)) dx &= \int_0^{u(x)} \int F_i(t) dt dx = \int_D F_i(t) dx dt \\ &= \int_{D^*} \int F_i(t) dx dt = \int_0^{u^*(x)} \int F_i(t) dt dx = \int G_i(u^*(x)) dx , \end{aligned}$$

where the middle equality follows from the definition of D^* . On the other hand, symmetrization does not increase surface area. We apply this fact to the boundary of

$$D_{R,\varepsilon} = \{(x, t) \in \mathbf{R}^{n+1} : |x| < R, 0 \leq t \leq \varepsilon u(x)\}$$

where R and ε are positive numbers. Thus

$$\int_{|x|<R} \{1 + \varepsilon^2 |\nabla u|^2\}^{1/2} dx + \int_{|x|=R} \varepsilon u(x) dS \geq \int_{|x|<R} \left\{ 1 + \varepsilon^2 |\nabla u^*|^2 \right\}^{1/2} dx + \int_{|x|=R} \varepsilon u^*(x) dS .$$

Since $u - u^* \in L^2$, we may choose a sequence $R_n \rightarrow \infty$ so that the integral of $|u - u^*|$ over $|x| = R_n$ tends to zero. Noting that $0 \leq \{1 + \varepsilon^2 |\nabla u|^2\}^{1/2} - 1 \leq \frac{1}{2} \varepsilon^2 |\nabla u|^2$, and taking $R = R_n \rightarrow \infty$, we obtain

$$\int [\{1 + \varepsilon^2 |\nabla u|^2\}^{1/2} - 1] dx \geq \int \{1 + \varepsilon^2 |\nabla u^*|^2\}^{1/2} - 1] dx .$$

Dividing both sides by ε^2 and letting $\varepsilon \rightarrow 0$, we get

$$\int |\nabla u|^2 dx \geq \int |\nabla u^*|^2 dx .$$

This shows that u^* is a solution of the same minimum problem as u . If we repeat this procedure successively over an appropriate sequence of hyperplanes, we get a non-negative radial solution.

Call the radial solution u . It is continuous for $r = |x| \neq 0$, by the Radial Lemma of Section 3. It satisfies an equation

$$u_{rr} + \frac{n-1}{r} u_r - F(u) = 0, \quad 0 < r < \infty ,$$

where $F(u) = u + F_1(u) - \lambda F_2(u)$. Thus $r^{1-n}(r^{n-1}u_r)_r = u_{rr} + ((n-1)/r)u_r$ is continuous and hence u is a C^2 function for $r \neq 0$. Let

$$q(r) = \frac{F(u(r))}{u(r)} = 1 + F_1'(0) + p(r) .$$

By (8) and (9), $p(r) \rightarrow 0$ as $r \rightarrow \infty$. Hence $q(r) \geq \frac{1}{2}$ for large enough r . Now $v = r^{(n-1)/2}u$ satisfies the equation

$$v_{rr} - \left[q(r) + \frac{(n-1)(n-3)}{4r^2} \right] v = 0 ,$$

whence

$$\left(\frac{1}{2}v^2\right)_{rr} = v_r^2 + \left[q(r) + \frac{(n-1)(n-3)}{4r^2} \right] v^2 .$$

Thus $w = v^2$ satisfies the inequality $w_{rr} \geq w$ for large enough r . From this inequality follows the exponential decay of w , hence of u .

In fact, it implies that $Q = e^{-r}(w_r + w)$ in non-decreasing for large r . If Q remains non-positive for large r , then $(e^r w)_r = e^{2r}Q \leq 0$, which implies that $w = O(e^{-r})$ as $r \rightarrow \infty$. If, on the other hand, $Q \geq 2\delta > 0$, then $w_r + w$ is certainly not integrable near ∞ . But the functions v^2, v_r^2, w and w_r are all integrable on an interval $R < r < \infty$ because $u \in H^1$. This contradiction proves the exponential decay.

3. Existence

Radial Lemma 1. *Let $n \geq 2$. Every radial function $u \in H^1$ is almost everywhere equal to a function $U(x)$, continuous for $x \neq 0$, such that*

$$|U(x)| \leq c|x|^{(1-n)/2} \|u\|_{H^1} \quad \text{for } |x| \geq 1 ,$$

where c depends only on n .

Proof. Let $m=(n-1)/2$. For $v \in C^\infty \cap H^1$, we have

$$\begin{aligned} (r^{2m}v^2)_r &= ((r^m v)^2)_r = 2(r^m v)_r r^m v \leq (r^m v)_r^2 + (r^m v)^2 \\ &= r^{n-1}(v_r^2 + v^2) + m(r^{n-2}v^2)_r - \frac{(n-1)(n-3)}{4} r^{n-3}v^2 . \end{aligned}$$

If $n \geq 3$, we integrate over an interval $[0, r]$ to obtain

$$\begin{aligned} r^{n-1}v^2(r) &\leq \int_0^r (v_r^2 + v^2) \varrho^{n-1} d\varrho \\ &= |S^{n-1}|^{-1} \|v\|_{H^1}^2 , \end{aligned}$$

where $|S^{n-1}|$ is the area of the unit sphere in \mathbf{R}^n . If $u \in H^1$, we let

$$v_\varepsilon(r) = \int_{-\varepsilon}^0 \zeta_\varepsilon(\varrho) u(r-\varrho) d\varrho ,$$

where ζ_ε is the usual approximate delta function, and pass to the limit to obtain the result. In case $n=2$, the differential inequality takes the form

$$-(rv^2)_r \leq r(v_r^2 + v^2) + (\frac{1}{2}v^2)_r + \frac{1}{4r}v^2 .$$

We integrate over an interval $[r, \infty]$ to obtain

$$(r + \frac{1}{2})v^2(r) \leq \int_r^\infty (v_r^2 + v^2) \varrho d\varrho + \int_r^\infty \frac{1}{4\varrho} v^2 d\varrho .$$

For $r \geq 1$, we may replace $1/\varrho$ by ϱ to obtain

$$2\pi(r + \frac{1}{2})v^2(r) \leq \frac{5}{4} \|v\|_{H^1}^2 .$$

The proof is completed as before.

Compactness Lemma 2. Let $\{P_N\}$ and $\{Q_N\}$ be two sequences of continuous functions: $\mathbf{R} \rightarrow \mathbf{R}$. For $c > 0$, let $\gamma(c) = \sup \{ |t| : t = P_N(s) \text{ for some } N \text{ and } s \text{ such that } |Q_N(s)| \leq c |P_N(s)| \}$.

Assume i) $\gamma(c) < \infty$ for all $c > 0$. (In other words, $P_N/Q_N \rightarrow 0$ uniformly as $P_N \rightarrow \infty$.)

ii) $\{u_N\}$ is a sequence of measurable functions: $\mathbf{R}^n \rightarrow \mathbf{R}$ such that

$$q = \sup_N \int |Q_N(u_N(x))| dx < \infty .$$

iii) $P_N(u_N(x)) \rightarrow v(x)$ for a.e. $x \in \mathbf{R}^n$.

a) Then

$$\int_B |P_N(u_N) - v| dx \rightarrow 0$$

for all bounded sets B .

b) Assume in addition that

iv) $P_N(s) = o(Q_N(s))$ as $s \rightarrow 0$ uniformly in N .

v) $u_N(x) \rightarrow 0$ as $|x| \rightarrow \infty$, uniformly in x and N .

Then $\int |P_N(u_N) - v| dx \rightarrow 0$.

Proof. We use the method of [12]. Fix N and $\varepsilon > 0$. Let $c > 4q/\varepsilon$. Let B be a bounded set. By iii) and Egorov's lemma, choose a set $Z \subset B$ of measure less than $\varepsilon/(4\gamma(c))$ such that $P_N(u_N) \rightarrow v$ uniformly on B/Z . Let

$$Z' = \{x \in Z : |Q_N(u_N(x))| \leq c|P_N(u_N(x))|\}, \quad Z'' = Z \setminus Z'.$$

Thus $|P_N(u_N)| \leq \gamma(c)$ on Z' and $|P_N(u_N)| \leq c^{-1}|Q_N(u_N)|$ on Z'' . Hence

$$\int_Z |P_N(u_N)| dx \leq q/c + \gamma(c)|Z''| < \varepsilon/2,$$

$$\overline{\lim}_B \int |P_N(u_N) - v| dx \leq \varepsilon + \lim_{B \setminus Z} \int |P_N(u_N) - v| dx = \varepsilon.$$

This proves a). To prove b), let $\varepsilon > 0$. By iv) choose $\delta > 0$ so that $\sup_N |P_N(s)/Q_N(s)| \leq \varepsilon/2q$ for $|s| \leq \delta$. By v) choose $R > 0$ so that $\sup_N |u_N(x)| < \delta$ for $|x| > R$. Then

$$\int_{|x| > R} |P_N(u_N(x))| dx \leq \frac{\varepsilon}{2q} \int |Q_N(u_N(x))| dx \leq \frac{\varepsilon}{2}.$$

Applying Part a) of this lemma to $B = \{|x| \leq R\}$, we conclude that

$$\overline{\lim} \int |P_N(u_N) - v| dx \leq \varepsilon.$$

This completes the proof.

We are now ready to begin the proof of Theorem 2. Let F_1 and F_2 satisfy conditions (8)–(11), where

$$G_i(s) = \int_0^s F_i(t) dt.$$

We extend them by zero for negative s : $F_i(s) = 0$ for $s < 0$. We may also assume $L = -\Delta + I$.

Lemma 3. *There exists a non-trivial solution of (12) in case $F_i(s) \leq c_i s$ for all s ($i = 1, 2$).*

Proof. In this case $0 \leq G_i(s) \leq \frac{1}{2}c_i s^2$. Let

$$M = \inf \int \left\{ \frac{1}{2}(|\nabla u|^2 + u^2) + G_1(u) \right\} dx = \inf \mathcal{L}(u)$$

subject to the conditions

$$u \in H^1, \quad u \text{ radial}, \quad \mathcal{M}(u) = \int G_2(u) dx = 1.$$

By (8), $G_2(s) > 0$ for $s > 0$. So for any $\gamma > 0$, there exists $u_0 \in C_c^\infty$, u_0 radial, with $\mathcal{M}(u_0) = \gamma$. Thus M is finite. Choose any minimizing sequence: $u_\nu \in H^1$, u_ν radial, $\mathcal{M}(u_\nu) = 1$, $\mathcal{L}(u_\nu) \rightarrow M$. Then $\{u_\nu\}$ is bounded in H^1 and $\{G_1(u_\nu)\}$ is bounded in L^1 . By Sobolev's inequality, $\{u_\nu\}$ is also bounded in L^{l+1} , where $l+1 = 2n/(n-2)$ and $n \geq 3$. By compactness on bounded subsets of \mathbf{R}^n , there is a subsequence, which we still denote by $\{u_\nu\}$, converging to a limit u weakly in H^1 and almost everywhere. Hence $G_i(u_\nu) \rightarrow G_i(u)$ a.e. ($i = 1, 2$). We shall apply the Compactness Lemma with $u_N = u_\nu$, $P_N = G_2$, and $Q_N(s) = |s|^{l+1} + G_1(s) + s^2$. Hypothesis i) is satisfied because $|s|^{l+1} + G_1(s) \leq cG_2(s) \Rightarrow |s|$ bounded $\Rightarrow G_2(s)$ bounded.

Hypothesis v) is satisfied because of the Radial Lemma. Thus $\mathcal{M}(u_v) \rightarrow \mathcal{M}(u)$. Hence $\mathcal{M}(u) = 1$ and also u is radial.

By Fatou's Lemma and weak convergence, we have $\mathcal{L}(u) \leq \liminf \mathcal{L}(u_v) \leq M$. By definition of M , $\mathcal{L}(u) = M$. Thus u attains the minimum. Now, because F_1 and F_2 are sublinear and continuous, it is easy to check that \mathcal{L} and \mathcal{M} are C^1 functionals on $H_r^1 = \{w \in H^1 : w \text{ is radial}\}$. Their Frechet derivatives are $\mathcal{M}'(w)v = \int F_2(w)v dx$ and $\mathcal{L}'(w)v = \int \{Vw \cdot Vv + wv + F_1(w)v\} dx$. Thus there exists $\lambda \in \mathbf{R}$, $u \in H^1$ such that $\mathcal{L}'(u)v = \lambda \mathcal{M}'(u)v$ for all $v \in H_r^1$. Let $h = -\Delta u + u + F_1(u) - \lambda F_2(u)$. Then h is radial and is orthogonal to all radial test functions. Hence $h = 0$.

Remark. If the space $H_r^1 \cap L^p$ were used, the preceding method could be made to work for functions F_i of polynomial growth at infinity; the F_i could have faster growth if Orlicz spaces were used. Instead, in the following argument we allow more general conditions on F_i by using an extra passage to the limit.

Lemma 4. *Let $F_i(s)$ be as in Theorem 2 (zero for $s < 0$). Then*

$$G_2(s) = o(s^{l+1} + G_1(s)) \quad \text{as } s \rightarrow \infty. \tag{30}$$

For $N > 0$ define the truncated functions

$$F_{iN}(s) = \begin{cases} F_i(s) & \text{for } s \leq N \\ F_i(N) & \text{for } s > N \end{cases}$$

and let $G_{iN}(s)$ be their indefinite integrals. Then

$$sF_{2N}(s) = O(s^{l+1} + G_{1N}(s)) \quad \text{as } s \rightarrow \infty \text{ uniformly in } N. \tag{31}$$

For any $c > 0$, there exists $\gamma(c) > 0$ such that

$$s^{l+1} + G_{1N}(s) \leq cG_{2N}(s) \Rightarrow G_{2N}(s) \leq \gamma(c). \tag{32}$$

For any $\sigma > 0$, let $c_\sigma = \sup_{s \leq \sigma} F_1(s)$; then

$$F_{1N}(s) \leq c_\sigma + \frac{1}{\sigma} sF_{1N}(s) \quad \text{for } N > \sigma, \quad s \in \mathbf{R}. \tag{33}$$

Proof. To prove (33), we let $\sigma < N$ and consider 3 cases. For $s \leq \sigma$, we have $F_{1N}(s) = F_1(s) \leq c_\sigma$. For $\sigma \leq s \leq N$, we have $F_{1N}(s) \leq sF_1(s)/\sigma = sF_{1N}(s)/\sigma$. For $N \leq s$, we have $F_{1N}(s) = F_1(N) \leq sF_1(N)/\sigma = sF_{1N}(s)/\sigma$.

We also prove (31) in 3 cases. For $s \leq N$, (31) is identical with (11). For $s > N$, we have

$$G_{iN}(s) = \int_0^s F_{iN} = G_i(N) + (s - N)F_i(N). \tag{34}$$

For $N < s < 2N$, we have from (11) and (34)

$$F_2(N) = O(N^l + G_1(N)/N) = O(s^l + 2G_1(N)/s) = O(s^l + G_{1N}(s)/s).$$

For $s \geq 2N$, we have from (10) and (34),

$$F_2(N) = o(N^l + F_1(N)) = o\left(s^l + 2 \frac{s - N}{s} F_1(N)\right) = O(s^l + G_{1N}(s)/s).$$

To prove (30), let $\varepsilon > 0$. By (10) there exists $T = T(\varepsilon)$ so large that

$$F_2(t) < \varepsilon(t^l + F_1(t)) \quad \text{for } t \geq T. \tag{35}$$

For $s \geq T$, we integrate (35) over the interval $[T, s]$ to get

$$G_2(s) \leq \varepsilon \left(\frac{s^{l+1}}{l+1} + G_1(s) \right) + G_2(T) - \varepsilon G_1(T) - \varepsilon \frac{T^{l+1}}{l+1}.$$

Thus (30) holds. That is, there exists $T' = T'(\varepsilon) \geq T$ so that

$$G_2(t) < \varepsilon(t^{l+1} + G_1(t)) \quad \text{for } t \geq T'. \tag{36}$$

To prove (32), let $c > 0$. Let $\varepsilon = 1/(2c)$. Let s satisfy the inequality

$$s^{l+1} + G_{1N}(s) \leq cG_{2N}(s). \tag{37}$$

Case 1: $s \leq N$. Then $G_{iN}(s) = G_i(s)$, so that (36) and (37) imply that $s \leq T'$. Hence $G_{2N}(s) \leq G_2(T') = \gamma(c)$.

Case 2: $s > N$. By (34), (37) takes the form

$$s^{l+1} + G_1(N) + (s - N)F_1(N) \leq c\{G_2(N) + (s - N)F_2(N)\}. \tag{37'}$$

In this case, we will show that $N < T' = T'(1/(2c))$. Thus N runs through a finite number of integers. By (37), s is also bounded by a function of c . Hence so is $G_{2N}(s)$. It remains to prove a contradiction if we suppose that $N \geq T'$. In that case, we may choose $t = N$ in both (35) and (36). Then we add $(s - N)$ times (35) to (36) to obtain

$$G_2(N) + (s - N)F_2(N) < \frac{1}{2c} \{N^{l+1} + (s - N)N^l + G_1(N) + (s - N)F_1(N)\}.$$

This indeed contradicts (37').

Proof of Theorem 2. For fixed N , we apply Lemma 3. There exists $u_N \in H^1$, u_N radial, u_n minimizes

$$\mathcal{L}_N(v) = \int \frac{1}{2}(|\nabla v|^2 + v^2) + G_{1N}(v) dx$$

subject to the constraints that $v \in H^1$, v is radial and $G_{2N}(v) dx = 1$. Furthermore,

$$-\Delta u_N + u_N + F_{1N}(u_N) = \lambda_N F_{2N}(u_N). \tag{38}$$

Multiplying (38) by u_N and integrating, we find

$$\int \{|\nabla u_N|^2 + u_N^2 + u_N F_{1N}(u_N)\} dx = \lambda_N \int u_N F_{2N}(u_N) dx. \tag{39}$$

Now choose u_o as in the proof of Lemma 3. Let $N > \sup|u_o(x)|$. Then $G_{iN}(u_o) = G_i(u_o)$. By the minimization property of u_N , we have

$$\mathcal{L}_N(u_N) \leq \mathcal{L}_N(u_o) = \mathcal{L}(u_o) < \infty.$$

Thus $\{u_N\}$ is bounded in H^1 and $\int G_{1N}(u_N) dx$ is bounded. By Sobolev's inequality, $\{u_N\}$ is bounded in L^{l+1} . By (31), $\int u_N F_{2N}(u_N) dx$ is also bounded.

By compactness, there is a subsequence, still denoted by $\{u_N\}$, such that u_N converges to some limit u weakly in H^1 and a.e. Hence $G_{iN}(u_N) \rightarrow G_i(u)$ and $F_{iN}(u_N) \rightarrow F_i(u)$ a.e. We now apply the Compactness Lemma 2b) with

$$P_N(s) = G_{2N}(s) \quad \text{and} \quad Q_N(s) = |s|^{l+1} + G_{1N}(s) + s^2.$$

Assumption i) of Lemma 2 is identical with (32), while Assumption v) follows from the Radial Lemma 1. Therefore $\int G_2(u)dx = 1$.

Lemma 5. $\{\lambda_N\}$ is a bounded sequence of positive numbers.

Assuming the validity of Lemma 5, the right side of (39) is bounded, whence $\int u_N F_{1N}(u_N)dx$ is also bounded. By the Compactness Lemma 2a) with $P_N = F_{iN}$ and $Q_N(s) = sF_{iN}(s)$, we deduce that $F_{iN}(u_N) \rightarrow F_i(u)$ in L^1_{loc} for $i=1, 2$. Taking a subsequence $\lambda_N \rightarrow \lambda$, we see that each term in (38) converges in the sense of distributions. We conclude that

$$-\Delta u + u + F_1(u) = \lambda F_2(u) .$$

To show u is non-negative, let $u^-(x) = \min(u(x), 0)$ and let $A = \{x : u(x) < 0\}$. Since $u \in H^1$, it follows that $u^- \in H^1$ and $\partial(u^-)/\partial x_j = (\partial u/\partial x_j)^-$. Multiplying the differential equation by u^- , we obtain

$$\begin{aligned} 0 &= \int \{ \nabla u \cdot \nabla u^- + uu^- + F(u)u^- \} dx \\ &= \int_A \{ |\nabla u|^2 + |u|^2 \} dx . \end{aligned}$$

Hence $u=0$ on A so that $u \geq 0$. (Another proof could be based on the method of Theorem 3.) Finally, by the proof of Theorem 3, u is a C^2 function for $x \neq 0$ which decays exponentially as $r \rightarrow \infty$. We also note that from the differential equation we have

$$\int \{ |\nabla u|^2 + u^2 + uF_1(u) \} dx = \lambda \int uF_2(u)dx ,$$

from which it follows that $\lambda \geq 0$. If $\lambda = 0$, then $u = 0$ and $\int G_2(u)dx = 0$, which is a contradiction.

Proof of Lemma 5. It follows from (39) that $\lambda_N > 0$. Now fix $v \in C_c^\infty$. By (38),

$$\lambda_N (F_{2N}(u_N), v) = (\nabla u_N, \nabla v) + (u_N, v) + (F_{1N}(u_N), v) .$$

By (33),

$$\begin{aligned} \lambda_N |(F_{2N}(u_N), v)| &\leq \|u_N\|_{H^1} \|v\|_{H^1} + c_\sigma \|v\|_{L^1} + \sigma^{-1} \|v\|_{L^\infty} (u_N, F_{1N}(u_N)) \\ &\leq k(1 + c_\sigma + \sigma^{-1} \lambda_N) \end{aligned}$$

with k independent of σ and N , where we have used (39) and the known bounds. Given $\varepsilon > 0$, choose $N > \sigma = k/\varepsilon$. Then

$$|(F_{2N}(u_N), v)| \leq \lambda_N^{-1} k(1 + c_\sigma) + \varepsilon .$$

If $\{\lambda_N\}$ were unbounded, then a subsequence would satisfy $\lambda_N \rightarrow \infty$ and so $(F_{2N}(u_N), v) \rightarrow 0$. On the other hand, by the Compactness Lemma 2a) just as above, $(F_{2N}(u_N), v) \rightarrow (F_2(u), v)$. Thus $(F_2(u), v) = 0$ for all test functions $v \in C_c^\infty$. Hence $F_2(u) = 0$. By (8), $G_2(u) = 0$. This contradicts $\int G_2(u)dx = 1$ and completes the proof of Lemma 5 and Theorem 2.

Remark. If the dimension $n = 1$ or 2 , Theorem 2 is valid under weaker conditions. If $n = 2$, we replace $(n+2)/(n-2)$ by an arbitrary positive number. Since H^1 is contained in a certain exponential Orlicz space L^ϕ , we actually need only replace

s^{l+1} by $\Phi(s)$. The proof is unchanged. If $n = 1$, strongly convergent sequences in H^1 converge uniformly; hence the direct method of Lemma 3 works. Theorem 2 is valid without any growth conditions (10), (11).

Remark. Our proof shows that, for each $\gamma > 0$, there is a solution with $\int G_2(u)dx = \gamma$.

Remark. There exists a non-positive as well as a non-negative solution if F_i are continuous real functions satisfying

$$sF_1(s) \geq 0, \quad sF_2(s) > 0 \tag{8*}$$

and (9), (10), (11) with s^l replaced by $|s|^l$, and ∞ by $\pm \infty$.

4. Existence of Many Solutions

Theorem 4. Let $L = I - \Delta$. Let $F_1 = 0$. Let F_2 be a continuous real function such that

$$sF_2(s) > 0 \quad \text{for } s \neq 0, \tag{41}$$

$$F_2(s) = O(|s|^p) \quad \text{as } |s| \rightarrow \infty \quad \text{where } p < (n+2)/(n-2) = l, \tag{42}$$

$$F_2 \text{ is odd and } F_2(0) = 0. \tag{43}$$

For any $\gamma > 0$, there exists an infinite sequence of distinct pairs of radial solutions of (12),

$$(\lambda_k, \pm u_k) \quad k = 0, 1, 2, \dots,$$

with $(Lu_k, u_k) = \gamma$.

Proof. We apply Theorem 2.10 of Rabinowitz [10], which is based on the minimax characterization of the higher eigenvalues. We use the space $E = H_r^1$ (the radial functions in H^1) and the functional $f(u) = \int G_2(u)dx$. We need only verify the hypothesis of Lemma 2.11 of [10]. Let $u_N \rightarrow u$ weakly in H_r^1 , hence in L^{l+1} . By the Compactness and Radial Lemmas and (42), we have $F_2(u_N) \rightarrow F_2(u)$ strongly in $L^{p'}$, where $(p')^{-1} + p^{-1} = 1$, and hence strongly in $(H_r^1)^*$.

A special case of Theorem 4 can be found in Berger [2]. The following theorem is based on a variational method without constraints.

Theorem 5. Let F be a real continuous function which satisfies (42), as well as

$$F(s)/s \rightarrow -\infty \quad \text{as } s \rightarrow +\infty, \tag{44}$$

$$sF(s) \geq \alpha G(s) \quad \text{for some } \alpha > 2, \tag{45}$$

$$F(s) = o(s) \quad \text{as } s \rightarrow 0. \tag{46}$$

Then the equation $Lu + F(u) = 0$ possesses at least one non-trivial solution. If F is odd, there is an infinite number of distinct pairs of solutions, $\pm u_k$ ($k = 0, 1, 2, \dots$).

Proof. We apply Theorems 3.9 and 3.37 of [10]. We use $L = I - \Delta$, $E = H_r^1$ and

$$f(u) = \int \left\{ \frac{1}{2}(|\nabla u|^2 + u^2) + G(u) \right\} dx.$$

The only novel feature is the verification of condition (PS)⁺ of [10]. Let $\{u_N\}$ be a sequence satisfying

$$f(u_N) \leq d,$$

$$h_N = -\Delta u_N + u_N + F(u_N) \rightarrow 0 \quad \text{in } H^{-1}.$$

Just as in [10], p. 172, these conditions imply that $\{u_N\}$ is bounded in H^1 . As in the proof of Theorem 4, $F(u_N) \rightarrow F(u)$ strongly in L^p and $h_N \rightarrow 0$ strongly in H^{-1} , for a subsequence which converges weakly to some u in H^1 and a.e. Hence

$$\|u_N - u\|_{H^1}^2 = -(F(u_N) - F(u), u_N - u) + (h_N, u_N - u) \rightarrow 0.$$

This proves (PS)⁺ and Theorem 5.

Example 1. $-\Delta u + u - |u|^{q-1}u = 0$, where $1 < q < (n+2)/(n-2)$. Both Theorems 4 and 5 imply the existence of an infinite sequence of radial solutions u_0, u_1, u_2, \dots . (If we apply Theorem 4, the eigenvalue λ_k has to be scaled out.) The solution u_k has exactly k nodes as a function of r . For some computer-generated solutions, see [1].

Example 2. $-\Delta u + u + |u|^{p-1}u - \lambda|u|^{q-1}u = 0$, where $1 < p < q < (n+2)/(n-2)$. Theorem 5 is applicable for every $\lambda > 0$. The case of larger p and q is open.

References

1. Anderson, D., Derrick, G.: Stability of time-dependent particlelike solutions in nonlinear field theories. *J. Math. Phys.* **11**, 1336—1346 (1970); **12**, 945—952 (1971)
2. Berger, M.S.: On the existence and structure of stationary states for a nonlinear Klein-Gordon equation. *J. Funct. Anal.* **9**, 249—261 (1972)
3. Clément, P.: Positive eigenfunctions for a class of second-order elliptic equations with strong nonlinearity (preprint)
4. Dancer, E.N.: Boundary value problems for ordinary differential equations in infinite intervals. *Proc. London Math. Soc.* **30**, 76—94 (1975)
5. Kato, T.: Growth properties of solutions of the reduced wave equation with a variable coefficient. *Comm. Pure Appl. Math.* **12**, 403—425 (1959)
6. Nehari, Z.: On a nonlinear differential equation arising in nuclear physics. *Proc. Roy. Irish Acad.* **62**, 117—135 (1963)
7. Pohozaev, S.I.: Eigenfunctions of the equation $\Delta u + \lambda f(u) = 0$. *Sov. Math. Doklady* **5**, 1408—1411 (1965)
8. Polya, G., Szegő, G.: *Isoperimetric inequalities in mathematical physics*. Princeton: University Press 1951
9. Rabinowitz, P.H.: Some aspects of nonlinear eigenvalue problems. *Rocky Mtn. J. Math.* **3**, 161—202 (1973)
10. Rabinowitz, P.H.: Variational methods for nonlinear eigenvalue problems. In: *Eigenvalues of nonlinear problems*. Rome: C.I.M.E., Edizioni Cremonese 1974
11. Rabinowitz, P.H.: Variational methods for nonlinear elliptic eigenvalue problems. *Indiana U. Math. J.* **23**, 729—754 (1974)
12. Strauss, W.A.: On weak solutions of semi-linear hyperbolic equations. *Anais Acad. Brasil. Ciencias* **42**, 645—651 (1970)
13. Stuart, C.A.: Battelle Inst. math. report No. 75, 1973
14. Syngé, J.L.: On a certain non-linear differential equation. *Proc. Roy. Irish Acad.* **62**, 17—41 (1961)
15. Weinberger, H.: *Variational methods for eigenvalue approximation*. Philadelphia: SIAM 1974

Communicated by A. Jaffe

Received February 2, 1977; in revised form April 4, 1977