

MOVIE COMMUNICATION

NUMERICAL SOLUTION OF THE FOKKER-PLANCK-LANDAU EQUATION BY SPECTRAL METHODS*

FRANCIS FILBET[†] AND LORENZO PARESCHI[‡]

Abstract. A new approach for the accurate numerical solution of the Fokker-Planck-Landau (FPL) equation has been presented recently in [1, 2]. The method is based on a fast spectral solver for the efficient solution of the collision operator. The use of a suitable explicit Runge-Kutta solver for the time integration of the collision phase avoids excessive small time steps induced by the stiffness of the diffusive collision operator. Here we present the details of a numerical simulation of the relaxation process in a three-dimensional Coulomb gas.

Key words. Landau-Fokker-Planck equation, spectral methods, Runge-Kutta methods

1. The FPL Equation

In the homogeneous case, the Landau or Fokker-Planck-Landau (FPL) model is described by

$$\begin{aligned} \frac{\partial f}{\partial t} &= \nabla_v \cdot \int_{R^3} \Phi(v - v^*) [\nabla_v f(v) f(v^*) - \nabla_{v^*} f(v^*) f(v)] dv^* \\ &= Q(f, f), \quad v \in R^3, \end{aligned} \tag{1.1}$$

where the non-negative function $f(t, v)$ depends on time t and velocity v of particles. In the collision operator $Q(f, f)$, Φ is a 3×3 non-negative and symmetric matrix of the form $\Phi(v) = |v|^{\gamma+2} S(v)$, with $S(v) = Id - \frac{v \otimes v}{|v|^2}$. Different values of $\gamma \in R$ lead to the usual classification in hard potentials $\gamma > 0$, Maxwellian molecules $\gamma = 0$, or soft potentials $\gamma < 0$ (which involves the Coulombian case $\gamma = -3$, of primary importance for applications). The structure of the FPL operator is similar to the Boltzmann one; this gives the physical properties of mass, momentum, and energy conservation and the decay of the entropy $H(t) = \int f \ln f dv$.

2. The Relaxation Process in 3D

We have considered the relaxation process for a three-dimensional Coulomb gas ($\gamma = -3$). The initial data is chosen as the sum of two symmetric Maxwellian functions

$$f_0(v) = \frac{1}{2} \frac{5}{(2\pi v_{th}^2)^{3/2}} \left[\exp\left(-\frac{|v - v_1|^2}{2 v_{th}^2}\right) + \exp\left(-\frac{|v - v_2|^2}{2 v_{th}^2}\right) \right],$$

with $v_1 = (1.25, 1.25, 0)$, $v_2 = (-1.25, -1.25, 0)$ and the thermal velocity is $v_{th} = 0.4$. The final time of the simulation is $T_{end} = 80$. The movie reports the time evolution of the level set of the distribution function $f(t, v_x, v_y, v_z) = 0.02$ obtained with $n = 32$

* Received: July 23, 2002; Accepted (in revised version): July 29, 2002.

Movie link: <http://www.unife.it/~prl/movies/lfp.html>

[†]IRMA, Université Louis Pasteur, 7 rue René Descartes, 67084 Strasbourg, France (filbet@math.u-strasbg.fr).

[‡]Department of Mathematics, University of Ferrara, via Machiavelli 35, I-44100 Ferrara, Italy (pareschi@dm.unife.it).

modes. Initially the level set of the initial data corresponds to two “spheres” in the velocity space. Then, the two distributions start to merge until the stationary state, characterized by a Maxwellian distribution with zero mean velocity, is reached. This is represented by a single centered sphere. We refer to [2, 1] for the details of the numerical method.

REFERENCES

- [1] F. Filbet and L. Pareschi, *A numerical method for the accurate solution of the Fokker-Planck-Landau equation in the nonhomogeneous case*. J. Comput. Phys., 179:1–26, 2002.
- [2] L. Pareschi, G. Russo, and G. Toscani, *Fast spectral methods for the Fokker-Planck-Landau collision operator*. J. Comput. Phys., 165:1–21, 2000.