

NON-EXISTENCE THEOREMS FOR SYSTEMS OF QUASILINEAR PARTIAL DIFFERENTIAL EQUATIONS

ENZO MITIDIERI¹

Dipartimento di Scienze Matematiche, Università degli Studi di Trieste
Piazzale Europa 1, 34100 Trieste, Italy

GUIDO SWEERS

Department of Pure Mathematics, Delft University of Technology, 2600 GA Delft, The Netherlands

ROB VAN DER VORST

Mathematisch Instituut, Leiden University, P.O. Box 9512, 2300 RA Leiden, The Netherlands

(Submitted by: L.A. Peletier)

1. Introduction. In this paper we shall investigate the non-existence of ground states of quasilinear systems of the form

$$\begin{cases} -\operatorname{div}(A(|Du|)Du) = f(|x|, u, v), \\ -\operatorname{div}(B(|Dv|)Dv) = g(|x|, u, v), \end{cases} \quad (1)$$

in \mathbb{R}^N , $N \geq 3$. Here D denotes the gradient operator, A, B are positive scalar functions and f, g are given nonlinearities that will be specified later.

We recall that a ground state of (1) is a positive radially symmetric solution of (1) such that

$$\lim_{|x| \rightarrow \infty} u(|x|) = \lim_{|x| \rightarrow \infty} v(|x|) = 0. \quad (2)$$

The corresponding scalar problem, i.e.,

$$-\operatorname{div}(A(|Du|)Du) = f(|x|, u) \quad \text{in } \mathbb{R}^N \quad (3)$$

has been studied extensively in a series of pioneering papers by Ni and Serrin ([15], [16] and [17]). In this paper we shall consider some extensions of their results to the system in (1). As a consequence we shall obtain some results which to our knowledge are new even in the scalar case.

Received for publication May 1994.

This work was reported on at the International Conference on Differential Equations in August, 1993.

¹Work supported in part by MURST 40%.

AMS Subject Classifications: 35J60, 35J70, 35R45.

Besides their intrinsic interest, non-existence results are a useful tool for proving related existence theorems for the corresponding Dirichlet problem

$$\begin{cases} -\operatorname{div}(A(|Du|)Du) = f(|x|, u, v) & \text{in } B_{r_0}(0) \subset \mathbb{R}^N, \\ -\operatorname{div}(B(|Dv|)Dv) = g(|x|, u, v) & \text{in } B_{r_0}(0) \subset \mathbb{R}^N, \\ u(x) = v(x) = 0 & \text{for } |x| = r_0. \end{cases} \quad (4)$$

This aspect is by now well understood in the case when the operator on the left hand side of (4) are linear, that is, in the semilinear case. See for example the paper by Gidas and Spruck, [9], or [22]. In view of new applications it is natural to investigate the possibility of extending the blow-up method (see [9] and [22]) to the quasilinear situation. Other strategies are available for the study of (4) (see for example [1], [2], [5], [6] [10], [18], [23], [24] and the references therein). However, it seems that for not necessarily variational systems containing strong nonlinearities (i.e., ‘superlinear’) methods based on non existence results are more powerful. Some results concerning this last aspect are contained in [3]. Our study in this paper is based on a property shared by positive radial supersolutions of the scalar problem

$$\begin{cases} -\operatorname{div}(A(|Du|)Du) \geq 0 & \text{in } \mathbb{R}^N \setminus B_{r_0}(0), \\ \frac{\partial}{\partial n} u(x) \leq 0 & \text{for } |x| = r_0, \end{cases} \quad (5)$$

where n denotes the normal at $x \in \partial B_{r_0}$ ($n = x/|x|$).

This paper is organized as follows. In Section 2, we state the lemma that allows us to control the behavior at $+\infty$ of positive radial supersolutions of (5). In Section 3, we will use this control to obtain non-existence results for quasilinear elliptic systems. Whenever the system has a variational structure it is possible to obtain a refined result by using a variational identity from [20], [25]. This is done in the fourth section.

The sections end with some examples. In the examples quasilinear elliptic operators such as the p -Laplacian and the mean curvature operator in non-parametric form will appear.

2. A fundamental Lemma. In this section we shall prove a lemma concerning positive radial supersolutions of (5). This result is our fundamental tool in the rest of the paper.

Let us specify the assumption.

(A1) Let $A : [0, \infty) \rightarrow [0, \infty)$ be such that

$$\exists A_M \in \mathbb{R} : \quad 0 < A(t) \leq A_M \quad \text{for all } t \in [0, \infty). \quad (6)$$

Lemma 2.1. *Let A satisfy (A1) and let $r_0 > 0$. If $N \geq 3$ and $u \in C^1(r_0, \infty)$, with $r \rightarrow A(|u'(r)|)u'(r) \in C^1(r_0, \infty)$, is a positive radial supersolution satisfying (5), then*

$$(1) \quad u'(r) \leq 0 \quad \text{for } r \geq r_0;$$

(2) the function $M(\cdot)$ defined by

$$M(r) = A(|u'(r)|)ru'(r) + A_M (N - 2)(u(r) - u_\infty), \tag{7}$$

where $u_\infty = \inf_{r \geq r_0} u(r) = \lim_{r \rightarrow \infty} u(r)$, is non-negative and non-increasing on (r_0, ∞) .

Proof. We may suppose that u is not constant. First observe that the positivity of u and $u'(r_0) \leq 0$ imply that $u'(r) < 0$ for $r > r_0$. Let us put

$$-\operatorname{div}(A(|Du|) Du) = f,$$

which can be written in radial coordinates as

$$-(r^{N-1}A(|u'(r)|)u'(r))' = r^{N-1}f(r) \quad \text{for } r > r_0,$$

i.e.,

$$-(r^{N-2}r A(|u'(r)|)u'(r))' = r^{N-1}f(r)$$

or

$$-(N - 2)A(|u'(r)|)u'(r) - \frac{d}{dr}[rA(|u'(r)|)u'(r)] = rf(r). \tag{8}$$

Hence, by (8), (6) and $u'(r) < 0$ it follows that

$$-M'(r) = -\frac{d}{dr}[rA(|u'(r)|)u'(r) + A_M(N - 2)(u(r) - u_\infty)] \geq rf(r) \tag{9}$$

which shows that M is non-increasing. It remains to prove that M is non-negative.

By contradiction, let us suppose that there exists $r_1 > r_0$ such that $M(r_1) < 0$. Integrating (9) on (r_1, r) we obtain

$$rA(|u'(r)|)u'(r) + A_M(N - 2)(u(r) - u_\infty) \leq M(r_1). \tag{10}$$

The positivity of u and the fact that u is decreasing gives

$$A_M u'(r) \leq A(|u'(r)|)u'(r) + \frac{1}{r}A_M(N - 2)(u(r) - u_\infty) \leq \frac{1}{r}M(r_1). \tag{11}$$

Integrating (11) on (s, t) we obtain for $r_1 < s < t$ that

$$u(t) - u(s) \leq \frac{M(r_1)}{A_M} \log\left(\frac{t}{s}\right). \tag{12}$$

By letting $t \rightarrow \infty$ we obtain a contradiction. \square

Remark. In the above lemma one of the crucial assumption is that $A(\cdot)$ is uniformly bounded on $(0, \infty)$. This assumption can be weakened somewhat if we assume that $A \in C^1(\mathbb{R}^+)$ holds and that A is bounded near 0. This is due to the fact that, if u is a positive supersolution of (5), then $u'(r) < 0$ for large r . Hence, the function $\tilde{u}(r) = u(r) - u_\infty$, where $u_\infty = \lim_{r \rightarrow \infty} u(r)$, is a positive supersolution of (5) such that $\lim_{r \rightarrow \infty} \tilde{u}(r) = 0$. By using an argument of Ni and Serrin (see page 10 of [16]) for $\tilde{u}(r)$, it follows that $\tilde{u}'(r) = u'(r) \rightarrow 0$ as r goes to infinity. Then, if $A(t) \leq M$ for small t , we can apply the same kind of proof as in Lemma 2.1. For easy reference we state this result.

Lemma 2.2. *Let $A \in C([0, \infty)) \cap C^1((0, \infty))$ satisfy*

$$0 < A(t) \leq A_M \quad \text{for sufficiently small } t > 0. \tag{13}$$

If $N \geq 3$ and $u \in C^1(r_0, \infty)$, with $r \rightarrow A(|u'(r)|)u'(r) \in C^1(r_0, \infty)$, is a positive radial supersolution satisfying (5), then

- (1) $u'(r) \leq 0$ for $r \geq r_0$,
- (2) there is $\tilde{r}_0 \geq r_0$ such that the function $M(\cdot)$ defined by (7) is non-negative and non-increasing on (\tilde{r}_0, ∞) .

The following lemma is essentially contained in [15].

Lemma 2.3. *Let A satisfy (A1) and let $r_0 > 0$. If $N \geq 3$ and $u \in C^1(r_0, \infty)$, with $r \rightarrow A(|u'(r)|)u'(r) \in C^1(r_0, \infty)$ is a non-constant positive radial supersolution satisfying*

$$-(r^{N-1}A(|u'|)u')' \geq 0 \quad \text{for } r > r_0, \quad u'(r_0) \leq 0, \tag{14}$$

then there exists $c > 0$ such that for all r sufficiently large, we have

$$u(r) - u_\infty \geq c r^{2-N}. \tag{15}$$

Proof. We may assume that $u'(r_0) < 0$. By integrating (14) on (r_0, r) we obtain

$$-u'(r)A(|u'(r)|)r^{N-1} \geq -u'(r_0)A(|u'(r_0)|)r_0^{N-1},$$

and hence for some $c_0 > 0$

$$-u'(r) \geq \frac{c_0}{A(|u'(r)|)}r^{1-N} \geq \frac{c_0}{A_M} r^{1-N}.$$

A further integration on (r, t) gives

$$u(r) - u_\infty \geq -u(t) + u(r) \geq \frac{1}{2-N} \frac{c_0}{A_M} (t^{2-N} - r^{2-N}).$$

By letting $t \rightarrow \infty$ we obtain

$$u(r) - u_\infty \geq \frac{c_0}{(N-2)A_M} r^{2-N}. \quad \square$$

Now we shall illustrate with some examples how we can apply Lemma 2.1.

Example 1. The mean curvature operator in non-parametric form. Let $u \in C^2(r_0, \infty)$ be radial and positive, and such that

$$-\operatorname{div}\left(\frac{Du}{\sqrt{1+|Du|^2}}\right) \geq 0 \quad \text{for } r \geq r_0, \quad u'(r_0) \leq 0.$$

Therefore, Lemma 2.1 applies with $A(t) = \frac{1}{\sqrt{1+t^2}} \leq 1$. The conclusion is that the function

$$M(r) = \frac{r u'(r)}{\sqrt{1 + (u'(r))^2}} + (N - 2)(u(r) - u_\infty)$$

is non-negative and non-increasing for $r \geq r_0$. Hence,

$$\frac{-r u'(r)}{\sqrt{1 + (u'(r))^2}} \leq (N - 2)(u(r) - u_\infty), \tag{16}$$

and thus $\lim_{r \rightarrow \infty} r u'(r) = 0$. This implies that there exists $\varepsilon > 0$ such that for $r \geq r_1$ sufficiently large we have

$$(u'(r))^2 < \varepsilon. \tag{17}$$

From (16) and (17) it follows that

$$-r u'(r) \leq (1 + \varepsilon)(N - 2)(u(r) - u_\infty) \quad \text{for } r > r_1,$$

and hence

$$(r^{(N-2)(1+\varepsilon)}u(r))' \geq (r^{(N-2)(1+\varepsilon)}(u(r) - u_\infty))' \geq 0 \quad \text{for } r > r_1. \tag{18}$$

If we assume that

$$0 < A_{\inf} = \inf_{t \in [0, \infty)} A(t) \quad \text{and} \quad \sup_{t \in [0, \infty)} A(t) = A_{\sup} < \infty \tag{19}$$

holds, then we have the same conclusion as in (18). That is, if u satisfies

$$-(r^{N-1}A(|u'|)u')' \geq 0 \quad \text{for } r > r_0, \quad u'(r_0) \leq 0,$$

then the function $M(\cdot)$, defined on $[r_0, \infty)$ by

$$M(r) = r u'(r) + (N - 2) \frac{A_{\sup}}{A_{\inf}} (u(r) - u_\infty)$$

satisfies

$$M(r) \geq 0 \quad \text{for } r \in [r_0, \infty).$$

Putting $\theta = (N - 2) \frac{A_{\sup}}{A_{\inf}}$ we find

$$(r^\theta u(r))' \geq (r^\theta (u(r) - u_\infty))' \geq 0. \tag{20}$$

An example of such a situation is the following.

Example 2. Let $\varepsilon \in (0, 1]$ and consider

$$A_\varepsilon(t) = \varepsilon + \frac{1 - \varepsilon}{\sqrt{1 + t^2}}, \quad t \geq 0.$$

The corresponding differential inequality associated with A_ε is

$$-(\varepsilon \Delta u + (1 - \varepsilon) \operatorname{div}(\frac{Du}{\sqrt{1 + |Du|^2}})) \geq 0 \quad \text{in } \mathbb{R}^N.$$

This regularizing operator is used, for example, in [11]. It follows from the preceding analysis that (20) holds with $\theta = (N - 2)\varepsilon^{-1}$. The particular case when $\varepsilon = 1$ is already known, see for example [12].

Another example is given by the following generalized mean curvature operator. (See [15], [16]).

Example 3. Let $u \in C^2(r_0, \infty)$ be radial, positive and satisfying

$$-\operatorname{div}(\frac{Du}{(1 + |Du|^2)^{\frac{2-m}{2}}}) \geq 0 \quad \text{for } r \geq r_0, \quad u'(r_0) \leq 0,$$

where $m \in (1, 2]$. The case $m = 1$ is treated above. Lemma 2.1 applies with

$$A(t) = \frac{1}{(1 + t^2)^{\frac{2-m}{2}}}, \quad t \geq 0.$$

We find for r sufficiently large that

$$\frac{r u'(r)}{(1 + (u'(r))^2)^{\frac{2-m}{2}}} + (N - 2)(u(r) - u_\infty) \geq 0.$$

Since $A(t) \leq 1$ we find for r sufficiently large that

$$(r^{(N-2)(1+\varepsilon)} u(r))' \geq (r^{(N-2)(1+\varepsilon)} (u(r) - u_\infty))' \geq 0. \quad (21)$$

One may also proceed using Lemma 2.1 of [3]. If A is continuously differentiable and

$$\begin{aligned} 1) \quad & (tA(t))' > 0 \quad \forall t > 0, \\ 2) \quad & 0 < \frac{A(t)}{(tA(t))'} \leq C \quad \forall t > 0, \end{aligned}$$

then the function $M(r) = ru'(r) + \theta(u(r) - u_\infty)$, with $\theta = (N - 1)C - 1$, satisfies $M(r) \geq 0$ for $r \geq r_0$. We have $C = \frac{1}{m-1}$ and with $\theta = \frac{N-m}{m-1}$, we obtain

$$(r^{\frac{N-m}{m-1}} u(r))' \geq (r^{\frac{N-m}{m-1}} (u(r) - u_\infty))' \geq 0. \quad (22)$$

For small ε the exponent in (21) is larger than in (22) and hence gives a better estimate. However, since the estimate depends on ε and is valid only for large r , it is not uniform.

3. Non-existence results.

3.1. A system with mean curvature operators. As a first example of a non-existence result we consider the following system

$$\begin{cases} -(r^{N-1} \frac{u'(r)}{\sqrt{1 + (u'(r))^2}})' = r^{N-1} f(v(r)) & \text{for } r > 0, \\ -(r^{N-1} \frac{v'(r)}{\sqrt{1 + (v'(r))^2}})' = r^{N-1} g(u(r)) & \text{for } r > 0. \end{cases} \quad (23)$$

Proposition 3.1. *Let $f, g \in C(\mathbb{R})$ with $f(0) = g(0) = 0$. Suppose that there exists $c > 0$ such that for small u and v we have*

$$f(v) \geq c|v|^{p-1}v, \quad (24)$$

$$g(u) \geq c|u|^{q-1}u, \quad (25)$$

where $p, q > 1$. If $N \geq 3$ and

$$\frac{(N-2)(pq-1)}{2} \leq \max\{p+1, q+1\}, \quad (26)$$

then problem (23) has no non-trivial ground states.

Proof. Let (u, v) be a non-trivial ground state of (23). By integrating the first equation in (23) on (r_0, r) for r large, and using $u'(r) < 0$ respectively $v'(r) < 0$ jointly with (24) and (25), we obtain

$$\frac{-ru'(r)}{\sqrt{1+(u'(r))^2}} \geq cr^2 v(r)^p, \quad (27)$$

$$\frac{-rv'(r)}{\sqrt{1+(v'(r))^2}} \geq cr^2 u(r)^q, \quad (28)$$

and hence by Lemma 2.1,

$$(N-2)u(r) \geq cr^2 v(r)^p, \quad (29)$$

$$(N-2)v(r) \geq cr^2 u(r)^q. \quad (30)$$

This implies that there exist $c_1, c_2 > 0$ such that for large r

$$u(r) \leq c_1 r^{-2\frac{p+1}{pq-1}}, \quad (31)$$

and

$$v(r) \leq c_2 r^{-2\frac{q+1}{pq-1}}. \quad (32)$$

By Lemma 2.3 we also deduce that

$$c_3 r^{2-N} \leq u(r), \quad (33)$$

and

$$c_4 r^{2-N} \leq v(r), \quad (34)$$

for some $c_3, c_4 > 0$ and r sufficiently large. Combining (31)–(33) and (32)–(34) we obtain a contradiction for $r \rightarrow \infty$ when the strict inequality holds in (26). Next, consider the case that

$$\frac{(N-2)(pq-1)}{2} = \max\{p+1, q+1\}.$$

Without loss of generality we may assume that $p \geq q$. In this case we have $pq = \frac{2p+N}{N-2}$. From the first equation of (23) and the assumption that $f(v) \geq cv^p$ for small v , we find that for large r

$$-(r^{N-1} \frac{u'(r)}{\sqrt{1+(u'(r))^2}})' \geq cr^{N-1} u^{pq} r^{2p}.$$

Integrating on (s, r) and using (33), we get for some $c^* > 0$,

$$\begin{aligned} \frac{-r^{N-1}u'(r)}{\sqrt{1+(u'(r))^2}} &\geq c \int_s^r \xi^{N-1}u(\xi)^{pq}\xi^{2p}d\xi \\ &\geq c^* \int_s^r \xi^{N-1}\xi^{(2-N)pq}\xi^{2p}d\xi = c^* \int_s^r \xi^{-1}d\xi = c^* \log \frac{r}{s}. \end{aligned}$$

Hence, by Lemma 2.1

$$(N-2)r^{N-2}u(r) \geq \frac{-r^{N-1}u'}{\sqrt{1+(u')^2}} \geq c \log r,$$

which is a contradiction to

$$r^{N-2}u(r) \leq cr^{-2\frac{p+1}{pq-1}}r^{N-2} = c.$$

Remark. The same result holds if we replace the mean curvature operators by

$$T_A = -\operatorname{div}(A(|Du|)Du), \quad T_B = -\operatorname{div}(B(|Dv|)Dv),$$

with A and B satisfying the assumption (A1).

3.2. Uniformly bounded elliptic operators. In this section we shall consider some non-existence results for ground states of systems of the form

$$\begin{cases} -\operatorname{div}(A(|Du|)Du) = f(|x|, u, v), \\ -\operatorname{div}(B(|Dv|)Dv) = g(|x|, u, v), \end{cases} \quad \text{in } \mathbb{R}^N. \tag{35}$$

Let $N \geq 3$ and suppose that

(A2) $A, B : [0, \infty) \rightarrow (0, \infty)$ are C^1 functions that satisfy

$$0 < A_{\inf} = \inf_{t \in [0, \infty)} A(t) \leq A(t) \leq \sup_{t \in [0, \infty)} A(t) = A_{\sup} < \infty,$$

$$0 < B_{\inf} = \inf_{t \in [0, \infty)} B(t) \leq B(t) \leq \sup_{t \in [0, \infty)} B(t) = B_{\sup} < \infty.$$

Define

$$\alpha_A = (N-2) \frac{A_{\sup}}{A_{\inf}}, \quad \alpha_B = (N-2) \frac{B_{\sup}}{B_{\inf}}.$$

The following result generalizes Proposition 3.1.

Theorem 3.2. *Suppose that condition (A2) is satisfied. Let $f, g \in C(\mathbb{R}^+ \times \mathbb{R}^2)$ with $f(r, 0, 0) = g(r, 0, 0) = 0 \forall r \geq 0$. Moreover, suppose that for large r and small u, v we have*

$$f(r, u, v) \geq a(r)|v|^{q-1}v, \quad (36)$$

$$g(r, u, v) \geq b(r)|u|^{p-1}u, \quad (37)$$

for some $p, q > 1$, where $a, b : [0, \infty) \rightarrow [0, \infty)$ are continuous. If, either

$$(i) \quad \int_{r_0}^{\infty} \vartheta^{1+\alpha_A} a(\vartheta) \left(\int_{\vartheta}^{\infty} b(s) s^{1-q\alpha_A} ds \right)^p d\vartheta = +\infty, \quad (38)$$

or

$$(ii) \quad \int_{r_0}^{\infty} \vartheta^{1+\alpha_B} b(\vartheta) \left(\int_{\vartheta}^{\infty} a(s) s^{1-p\alpha_B} ds \right)^q d\vartheta = +\infty, \quad (39)$$

then system (35) has no non-trivial ground state.

Proof. The proof is by contradiction. Let us prove the assertion in the case when (38) holds. The other case is similar. Let (u, v) be a non-trivial positive and radial ground state. Then (u, v) satisfies, for $r > 0$,

$$\begin{cases} -(r^{N-1}A(|u'(r)|)u'(r))' = r^{N-1}f(r, u, v), \\ -(r^{N-1}B(|v'(r)|)v'(r))' = r^{N-1}g(r, u, v). \end{cases} \quad (40)$$

Since (u, v) is positive and $u(r), v(r) \rightarrow 0$ as $r \rightarrow \infty$, it follows that there exists r_0 sufficiently large such that $u'(r_0) \leq 0$, and hence $u'(r) \leq 0$ for $r \geq r_0$. A similar result holds for v . We may assume that for some large r_0 we have

$$u'(r) \leq 0, \quad v'(r) \leq 0 \quad \text{for } r \geq r_0. \quad (41)$$

Let us define

$$\begin{cases} \tilde{M}_A(r) = rA(|u'(r)|)u'(r) + A_{\sup}(N-2)u(r), \\ \tilde{M}_B(r) = rB(|v'(r)|)v'(r) + B_{\sup}(N-2)v(r). \end{cases} \quad (42)$$

Similar to (9), one shows that

$$-\frac{d}{dr}\tilde{M}_A(r) \geq ra(r)v^p, \quad -\frac{d}{dr}\tilde{M}_B(r) \geq rb(r)u^q, \quad (43)$$

on (r_0, ∞) . By applying Lemma 2.1, it also follows that the two functions $\tilde{M}_A(\cdot)$ and $\tilde{M}_B(\cdot)$ are non-negative on (r_0, ∞) . Integrating both inequalities in (43) on (r, t) with $r > r_0$ it follows that

$$\begin{cases} \tilde{M}_A(r) \geq -\tilde{M}_A(t) + \tilde{M}_A(r) \geq \int_r^t sa(s)v(s)^p ds, \\ \tilde{M}_B(r) \geq -\tilde{M}_B(t) + \tilde{M}_B(r) \geq \int_r^t sb(s)u(s)^q ds, \end{cases} \quad (44)$$

and hence, using the fact that $A_{\inf}, B_{\inf} > 0$ and $u', v' \leq 0$ for $r \geq r_0$ we obtain, with $C_A = A_{\sup}(N - 2)u(r_0) > 0$, that

$$C_A \geq A_{\sup}(N - 2)u(r) \geq \tilde{M}_A(r) \geq \int_r^t sa(s)v(s)^p ds \geq (r^{\alpha_B}v(r))^p \int_r^t sa(s)s^{1-p\alpha_B} ds, \tag{45}$$

and similarly, with $C_B = B_{\sup}(N - 2)v(r_0) > 0$, that

$$C_B \geq B_{\sup}(N - 2)v(r) \geq (r^{\alpha_A}u(r))^q \int_r^t sb(s)s^{1-q\alpha_A} ds. \tag{46}$$

In the last two estimates we have used the fact that on (r_0, ∞)

$$(r^{\alpha_A}u(r))' \geq 0, \quad (r^{\alpha_B}v(r))' \geq 0. \tag{47}$$

Indeed, the first estimate of (47) follows from (44) and

$$(r^{\alpha_A}u(r))' = \frac{1}{A_{\inf}}r^{\alpha_A-1}(A_{\inf}r u'(r) + (N - 2)A_{\sup}u(r)) \geq \frac{1}{A_{\inf}}r^{\alpha_A-1}\tilde{M}_A(r) \tag{48}$$

for $r > r_0$. By (45) and (46) we find that for $r > r_0$

$$\int_r^\infty a(s)s^{1-p\alpha_B} ds < \infty, \quad \int_r^\infty b(s)s^{1-q\alpha_A} ds < \infty. \tag{49}$$

If one of the integrals in (49) equals $+\infty$, the proof is complete. If not, we proceed as follows. For some $c_a, c_b > 0$ it follows from (45)–(46) that

$$\begin{aligned} (u(r))^q &\geq c_a(r^{\alpha_B}v(r))^{pq} \left(\int_r^\infty a(s)s^{1-p\alpha_B} ds \right)^q, \\ (v(r))^p &\geq c_b(r^{\alpha_A}u(r))^{pq} \left(\int_r^\infty b(s)s^{1-q\alpha_A} ds \right)^p. \end{aligned} \tag{50}$$

From the first inequality of (43) and (50) it follows that

$$-\frac{d}{dr}\tilde{M}_A(r) \geq c_b r a(r)(r^{\alpha_A} u(r))^{pq} \left(\int_r^\infty b(s)s^{1-q\alpha_A} ds \right)^p. \tag{51}$$

Integrating (51) on (r, t) yields

$$\begin{aligned} \tilde{M}_A(r) &\geq -\tilde{M}_A(t) + \tilde{M}_A(r) \geq c_b \int_r^t sa(s)(s^{\alpha_A}u(s))^{pq} \left(\int_s^\infty b(\xi)\xi^{1-q\alpha_A} d\xi \right)^p ds \\ &\geq c_b(r^{\alpha_A}u(r))^{pq} \int_r^t sa(s) \left(\int_s^\infty b(\xi)\xi^{1-q\alpha_A} d\xi \right)^p ds, \end{aligned} \tag{52}$$

and then, using (48) again,

$$\begin{aligned} (r^{\alpha_A} u(r))' &\geq \frac{1}{A_{\inf}} r^{\alpha_A-1} \tilde{M}_A(r) \\ &\geq \frac{c_b}{A_{\inf}} r^{\alpha_A-1} (r^{\alpha_A} u(r))^{pq} \int_r^t sa(s) \left(\int_s^\infty b(\xi) \xi^{1-q\alpha_A} d\xi \right)^p ds. \end{aligned} \tag{53}$$

Setting $\phi(r) = r^{\alpha_A} u(r)$ we find from (53) that

$$\frac{1}{1-pq} \frac{d}{dr} (\phi(r))^{1-pq} \geq c'_a r^{\alpha_A-1} \int_r^t sa(s) \left(\int_s^\infty b(\xi) \xi^{1-q\alpha_A} d\xi \right)^p ds, \tag{54}$$

with $c'_a = \frac{c_b}{A_{\inf}}$. An integration of (54) on (r, \tilde{r}) gives

$$\begin{aligned} \frac{-1}{1-pq} (\phi(r))^{1-pq} &\geq \frac{1}{1-pq} (\phi(\tilde{r}))^{1-pq} - \frac{1}{1-pq} (\phi(r))^{1-pq} \\ &\geq c'_b \int_r^{\tilde{r}} \zeta^{\alpha_A-1} \int_\zeta^t s a(s) \left(\int_s^\infty b(\xi) \xi^{1-q\alpha_A} d\xi \right)^p ds d\zeta \\ &= c''_b \left[\zeta^{\alpha_A} \int_\zeta^t s a(s) \left(\int_s^\infty b(\xi) \xi^{1-q\alpha_A} d\xi \right)^p ds \right] \Big|_{\zeta=r}^{\zeta=\tilde{r}} \\ &\quad + c''_b \int_r^{\tilde{r}} \zeta^{\alpha_A+1} a(\zeta) \left(\int_\zeta^\infty b(\xi) \xi^{1-q\alpha_A} d\xi \right)^p d\zeta \\ &= c''_b \tilde{r}^{\alpha_A} \int_{\tilde{r}}^t sa(s) \left(\int_s^\infty b(\xi) \xi^{1-q\alpha_A} d\xi \right)^p ds \\ &\quad - c''_b r^{\alpha_A} \int_r^t sa(s) \left(\int_s^\infty b(\xi) \xi^{1-q\alpha_A} d\xi \right)^p ds \\ &\quad + c''_b \int_r^{\tilde{r}} \zeta^{\alpha_A+1} a(\zeta) \left(\int_\zeta^\infty b(\xi) \xi^{1-q\alpha_A} d\xi \right)^p d\zeta, \end{aligned} \tag{55}$$

with $c''_b = c'_b/\alpha_A$. Now we first let t go to infinity and then \tilde{r} . Since the second term is bounded by (45) and (52), and the third term goes to infinity by assumption, one finds that

$$\frac{-1}{1-pq} (\phi(r))^{1-pq} \rightarrow \infty \quad \text{when } r \rightarrow \infty$$

which shows that $\lim_{r \rightarrow \infty} \phi(r) = 0$. But this contradicts the fact that $\phi'(r) \geq 0$ for $r > r_0$.

3.3. Possibly unbounded elliptic operators. Next we shall consider a general non-existence result for a different class of quasilinear operators. Our theorem can be applied for example to the case when the left hand side of (1) involves two generalized Laplacians. The explicit statement can be found in the last theorem of this section.

(A1') Let $A, B : \mathbb{R}^+ \rightarrow (0, \infty)$ be C^1 and suppose that for some $\delta_A, \delta_B \geq 0, c_A, c_B > 0$ and $m_A, m_B > \frac{1}{N-1}$ we have

$$\begin{aligned} (i)_A \quad A(t) &\leq m_A \frac{\partial}{\partial t}(tA(t)) \leq c_A t^{\delta_A}, \quad t > 0, \\ (i)_B \quad B(t) &\leq m_B \frac{\partial}{\partial t}(tB(t)) \leq c_B t^{\delta_B}, \quad t > 0. \end{aligned} \tag{56}$$

(A2') Let $f, g : \mathbb{R}^+ \times \mathbb{R}^2 \rightarrow \mathbb{R}$ be C^0 with $f(x, 0, 0) = g(x, 0, 0) = 0$ and suppose that for small u, v and large $x \in \mathbb{R}^N$ there are continuous $a_{ij} : [0, \infty) \rightarrow [0, \infty)$, with $i, j \in \{1, 2\}$, such that we have

$$\begin{aligned} (ii)_A \quad f(x, u, v) &\geq a_{11}(|x|)|u|^{p_{11}-1}u + a_{12}(|x|)|v|^{p_{12}-1}v, \\ (ii)_B \quad g(x, u, v) &\geq a_{21}(|x|)|u|^{p_{21}-1}u + a_{22}(|x|)|v|^{p_{22}-1}v. \end{aligned} \tag{57}$$

Theorem 3.3. *Suppose that conditions (56) and (57) are satisfied. Moreover, assume that*

(1) $p_{12}p_{21} > (1 + \delta_A)(1 + \delta_B)$ and

$$\int_{r_0}^\infty \xi^{(1+\delta_B)(1+\theta_B)} a_{21}(\xi) \left(\int_\xi^\infty s^{1+\delta_A-\theta_B p_{12}} a_{12}(s) ds \right)^{\frac{p_{21}}{1+\delta_A}} d\xi = +\infty, \tag{58}$$

or

(2) $p_{12}p_{21} > (1 + \delta_A)(1 + \delta_B)$ and

$$\int_{r_0}^\infty \xi^{(1+\delta_A)(1+\theta_A)} a_{12}(\xi) \left(\int_\xi^\infty s^{1+\delta_B-\theta_A p_{21}} a_{21}(s) ds \right)^{\frac{p_{12}}{1+\delta_B}} d\xi = +\infty, \tag{59}$$

or

(3) $p_{11} > 1 + \delta_A$ and

$$\int_{r_0}^\infty \xi^{(1+\delta_A)(1+\theta_A)-\theta_A p_{11}} a_{11}(\xi) d\xi = +\infty, \tag{60}$$

or

(4) $p_{22} > 1 + \delta_B$ and

$$\int_{r_0}^\infty \xi^{(1+\delta_B)(1+\theta_B)-\theta_B p_{22}} a_{22}(\xi) d\xi = +\infty, \tag{61}$$

with $\theta_A = m_A(N - 1) - 1$ and $\theta_B = m_B(N - 1) - 1$.

Then, problem

$$\begin{cases} -\operatorname{div}(A(|Du|)Du) = f(|x|, u, v), \\ -\operatorname{div}(B(|Dv|)Dv) = g(|x|, u, v), \end{cases} \quad \text{in } \mathbb{R}^N. \tag{62}$$

has no non-trivial ground state.

Proof. We start by showing that there is no non-trivial ground state when (58) holds. First we set

$$E_A(t) = \frac{\partial}{\partial t}(tA(t)), \quad E_B(t) = \frac{\partial}{\partial t}(tB(t)). \quad (63)$$

System (62) can be rewritten as

$$\begin{cases} -E_A(|u'|)u'' + \frac{1-N}{r}u'A(|u'|) = f(r, u, v) & \text{for } r > 0, \\ -E_B(|v'|)v'' + \frac{1-N}{r}v'B(|v'|) = g(r, u, v) & \text{for } r > 0, \end{cases} \quad (64)$$

and by the estimates in (57) we obtain for large r

$$\begin{cases} -rE_A(|u'|)u'' + (1-N)u'A(|u'|) \geq a_{12}(r)rv^{p_{12}}, \\ -rE_B(|v'|)v'' + (1-N)v'B(|v'|) \geq a_{21}(r)ru^{p_{21}}, \end{cases} \quad (65)$$

and hence

$$\begin{cases} -ru'' + (1-N)u' \frac{A(|u'|)}{E_A(|u'|)} \geq \frac{a_{12}(r)r}{E_A(|u'|)}v^{p_{12}}, \\ -rv'' + (1-N)v' \frac{B(|v'|)}{E_B(|v'|)} \geq \frac{a_{21}(r)r}{E_B(|v'|)}u^{p_{21}}. \end{cases} \quad (66)$$

By (56) and $u' < 0, v' < 0$ we find

$$-ru'' + (1-N)m_A u' \geq \frac{a_{12}(r)r}{E_A(|u'|)}v^{p_{12}} \geq \frac{c_A}{m_A}|u'|^{-\delta_A}a_{12}(r)rv^{p_{12}}, \quad (67)$$

and

$$-rv'' + (1-N)m_B v' \geq \frac{c_B}{m_B}|v'|^{-\delta_B}a_{21}(r)ru^{p_{21}}. \quad (68)$$

We put

$$\tilde{M}_A(r) = ru' + (m_A(N-1) - 1)u, \quad \tilde{M}_B(r) = rv' + (m_B(N-1) - 1)v. \quad (69)$$

From (67) and (68) it follows that for r large enough, we have

$$\frac{d}{dr}\tilde{M}_A(r) \leq 0, \quad \frac{d}{dr}\tilde{M}_B(r) \leq 0. \quad (70)$$

With θ_A, θ_B as above, this is equivalent to

$$\begin{cases} (r^{\theta_A}u(r))' \geq 0 & \text{for } r > r_1, \\ (r^{\theta_B}v(r))' \geq 0 & \text{for } r > r_1, \end{cases} \quad (71)$$

for some $r_1 > 0$. Hence, by Lemma 2.1 of [3] we find that

$$\tilde{M}_A(r) \geq 0, \quad \tilde{M}_B(r) \geq 0. \tag{72}$$

Since $u' < 0$ and $v' < 0$, we deduce from (72) that

$$\begin{cases} \theta_A u(r) \geq -ru'(r) = r|u'(r)| & \text{for } r > r_1, \\ \theta_B v(r) \geq -rv'(r) = r|v'(r)| & \text{for } r > r_1. \end{cases} \tag{73}$$

By integrating (68) on (s, r) with $s \geq r_1$ and using (73) we find that, for some positive c_i ,

$$\begin{aligned} c_0 v(s) &\geq \tilde{M}_B(s) \geq -\tilde{M}_B(r) + \tilde{M}_B(s) \geq c_1 \int_s^r |v'(\xi)|^{-\delta_B} a_{21}(\xi) \xi u(\xi)^{p_{21}} d\xi \\ &\geq c_2 \int_s^r \xi^{\delta_B} v(\xi)^{-\delta_B} a_{21}(\xi) \xi u(\xi)^{p_{21}} d\xi \geq c_2 v(s)^{-\delta_B} \int_s^r \xi^{1+\delta_B-\theta_A p_{21}} a_{21}(\xi) (\xi^{\theta_A} u(\xi))^{p_{21}} d\xi \\ &\geq c_2 v(s)^{-\delta_B} (s^{\theta_A} u(s))^{p_{21}} \int_s^r \xi^{1+\delta_B-\theta_A p_{21}} a_{21}(\xi) d\xi. \end{aligned} \tag{74}$$

Hence, there exists $c > 0$ such that

$$v(s)^{1+\delta_B} \geq c (s^{\theta_A} u(s))^{p_{21}} \int_s^\infty \xi^{1+\delta_B-\theta_A p_{21}} a_{21}(\xi) d\xi. \tag{75}$$

Using (67) and (73) it follows that there exist positive constants c_i such that for all $r > r_1$

$$\begin{aligned} -\frac{d}{dr} \tilde{M}_A(r) &\geq c_1 r^{\delta_A} u(r)^{-\delta_A} a_{12}(r) r v(r)^{p_{12}} \\ &\geq c_2 r^{\delta_A} u(r)^{-\delta_A} a_{12}(r) r \left((r^{\theta_A} u(r))^{p_{21}} \int_r^\infty \xi^{1+\delta_B-\theta_A p_{21}} a_{21}(\xi) d\xi \right)^{\frac{p_{12}}{1+\delta_B}}. \end{aligned} \tag{76}$$

We continue as in the uniform case. Integrating (76) on (s, t) for $s \geq r_1$ yields

$$\begin{aligned} c_0 u(s) &\geq \tilde{M}_A(s) \geq -\tilde{M}_A(t) + \tilde{M}_A(s) \\ &\geq c_2 \int_s^t r^{1+\delta_A(1+\theta_A)} a_{12}(r) (r^{\theta_A} u(r))^{\frac{p_{12}p_{21}}{1+\delta_B}-\delta_A} \left(\int_r^\infty \xi^{1+\delta_B-\theta_A p_{21}} a_{21}(\xi) d\xi \right)^{\frac{p_{12}}{1+\delta_B}} dr \\ &\geq c_2 (s^{\theta_A} u(s))^{\frac{p_{12}p_{21}}{1+\delta_B}-\delta_A} \int_s^t r^{1+\delta_A(1+\theta_A)} a_{12}(r) \left(\int_r^\infty \xi^{1+\delta_B-\theta_A p_{21}} a_{21}(\xi) d\xi \right)^{\frac{p_{12}}{1+\delta_B}} dr. \end{aligned} \tag{77}$$

And again for some $c > 0$ we have

$$\begin{aligned} (r^{\theta_A} u(r))' &= r^{\theta_A-1} \tilde{M}_A(r) \\ &\geq c r^{\theta_A-1} (r^{\theta_A} u(r))^{\frac{p_{12}p_{21}}{1+\delta_B}-\delta_A} \int_r^t \eta^{1+\delta_A(1+\theta_A)} a_{12}(\eta) \left(\int_\eta^\infty \xi^{1+\delta_B-\theta_A p_{21}} a_{21}(\xi) d\xi \right)^{\frac{p_{12}}{1+\delta_B}} d\eta. \end{aligned} \tag{78}$$

Hence,

$$\begin{aligned} & ((r^{\theta_A} u(r))^{1+\delta_A - \frac{p_{12}p_{21}}{1+\delta_B}})' \\ & \geq cr^{\theta_A - 1} \int_r^t \eta^{1+\delta_A(1+\theta_A)} a_{12}(\eta) \left(\int_\eta^\infty \xi^{1+\delta_B - \theta_A p_{21}} a_{21}(\xi) d\xi \right)^{\frac{p_{12}}{1+\delta_B}} d\eta. \end{aligned} \tag{79}$$

Integrating (79) on (s_1, s_2) gives

$$\begin{aligned} & (r^{\theta_A} u(s_2))^{1+\delta_A - \frac{p_{12}p_{21}}{1+\delta_B}} \\ & \geq c \int_{s_1}^{s_2} r^{\theta_A - 1} \int_r^t \eta^{1+\delta_A(1+\theta_A)} a_{12}(\eta) \left(\int_\eta^\infty \xi^{1+\delta_B - \theta_A p_{21}} a_{21}(\xi) d\xi \right)^{\frac{p_{12}}{1+\delta_B}} d\eta dr \\ & = c' \left[r^{\theta_A} \int_r^t \eta^{1+\delta_A(1+\theta_A)} a_{12}(\eta) \left(\int_\eta^\infty \xi^{1+\delta_B - \theta_A p_{21}} a_{21}(\xi) d\xi \right)^{\frac{p_{12}}{1+\delta_B}} d\eta \right]_{r=s_1}^{r=s_2} \\ & \quad + c' \int_{s_1}^{s_2} r^{\theta_A} r^{1+\delta_A(1+\theta_A)} a_{12}(r) \left(\int_r^\infty \xi^{1+\delta_B - \theta_A p_{21}} a_{21}(\xi) d\xi \right)^{\frac{p_{12}}{1+\delta_B}} dr \\ & = c' s_2^{\theta_A} \int_{s_2}^t \eta^{1+\delta_A(1+\theta_A)} a_{12}(\eta) \left(\int_\eta^\infty \xi^{1+\delta_B - \theta_A p_{21}} a_{21}(\xi) d\xi \right)^{\frac{p_{12}}{1+\delta_B}} d\eta \\ & \quad - c' s_1^{\theta_A} \int_{s_1}^t \eta^{1+\delta_A(1+\theta_A)} a_{12}(\eta) \left(\int_\eta^\infty \xi^{1+\delta_B - \theta_A p_{21}} a_{21}(\xi) d\xi \right)^{\frac{p_{12}}{1+\delta_B}} d\eta \\ & \quad + c' \int_{s_1}^{s_2} r^{(1+\delta_A)(1+\theta_A)} a_{12}(r) \left(\int_r^\infty \xi^{1+\delta_B - \theta_A p_{21}} a_{21}(\xi) d\xi \right)^{\frac{p_{12}}{1+\delta_B}} dr. \end{aligned} \tag{80}$$

Now we let $t \rightarrow \infty$. The second term is bounded independently of s_2 . Next, we let $s_2 \rightarrow \infty$ and with (58) we obtain a contradiction. Similarly, a contradiction is obtained in the case in which (59) holds. This concludes the proof in the cases when (58) or (59) hold.

In order to obtain a contradiction in the cases when (60) or (61) hold, we proceed as before until (67). This estimate is replaced by

$$\begin{aligned} -\tilde{M}'_A(r) &= -ru''(r) + (1 - N)m_A u'(r) \geq \frac{a_{11}(r)r}{E_A(|u'|)} u(r)^{p_{11}} \\ &\geq \frac{c_A}{m_A} |u'(r)|^{-\delta_A} a_{11}(r) ru(r)^{p_{11}}. \end{aligned} \tag{81}$$

By (73), we find that there exists $c > 0$ such that

$$\frac{c_A}{m_A} |u'(r)|^{-\delta_A} a_{11}(r) ru(r)^{p_{11}} \geq cr^{\delta_A} u(r)^{-\delta_A} a_{11}(r) ru(r)^{p_{11}},$$

and as in (74), since $p_{11} > \delta_A + 1 > \delta_A$, we obtain

$$\begin{aligned} \tilde{M}_A(s) &\geq -\tilde{M}_A(t) + \tilde{M}_A(s) \geq c \int_s^t r^{1+\delta_A} u(r)^{p_{11} - \delta_A} a_{11}(r) dr \\ &\geq c(s^{\theta_A} u(s))^{p_{11} - \delta_A} \int_s^t r^{1+\delta_A - \theta_A(p_{11} - \delta_A)} a_{11}(r) dr. \end{aligned} \tag{82}$$

The last inequality is equivalent to

$$(s^{\theta_A} u(s))' \geq c(s^{\theta_A} u(s))^{p_{11}-\delta_A} s^{\theta_A-1} \int_s^t r^{1+\delta_A-\theta_A(p_{11}-\delta_A)} a_{11}(r) dr$$

or

$$\frac{1}{1 + \delta_A - p_{11}} ((s^{\theta_A} u(s))^{1+\delta_A-p_{11}})' \geq c s^{\theta_A-1} \int_s^t r^{1+\delta_A-\theta_A(p_{11}-\delta_A)} a_{11}(r) dr. \tag{83}$$

From (83) and $p_{11} > 1 + \delta_A$ we find for some $c' > 0$ and for $s_2 > s_1$ that

$$\begin{aligned} \frac{1}{p_{11} - (1 + \delta_A)} (s_1^{\theta_A} u(s_1))^{1+\delta_A-p_{11}} &\geq c \int_{s_1}^{s_2} s^{\theta_A-1} \int_s^t \xi^{1+\delta_A-\theta_A(p_{11}-\delta_A)} a_{11}(\xi) d\xi ds \\ &= c' \left[s^{\theta_A} \int_s^t \xi^{1+\delta_A-\theta_A(p_{11}-\delta_A)} a_{11}(\xi) d\xi ds \right] \Big|_{s_1}^{s_2} + c' \int_{s_1}^{s_2} s^{(1+\delta_A)(1+\theta_A)-\theta_A p_{11}} a_{11}(s) ds. \end{aligned} \tag{84}$$

A contradiction follows if (60) holds. A similar argument is used when (61) holds. This concludes the proof of the theorem.

Example 4. Consider (1) with

$$f(x, u, v) = |x|^a |u|^{p_{11}-1} u + |x|^b |v|^{p_{12}-1} v, \quad g(x, u, v) = |x|^c |u|^{p_{21}-1} u + |x|^d |v|^{p_{22}-1} v.$$

If $p_{12}p_{21} > (1 + \delta_A)(1 + \delta_B)$ and

$$N - 1 \leq (m_B)^{-1} \frac{(1 + c + p_{21})(1 + \delta_A) + (1 + b + p_{12})p_{21}}{p_{12}p_{21} - (1 + \delta_A)(1 + \delta_B)},$$

or $p_{12}p_{21} > (1 + \delta_A)(1 + \delta_B)$ and

$$N - 1 \leq (m_A)^{-1} \frac{(1 + b + p_{12})(1 + \delta_B) + (1 + c + p_{21})p_{12}}{p_{12}p_{21} - (1 + \delta_A)(1 + \delta_B)},$$

or $p_{11} > 1 + \delta_A$ and

$$N - 1 \leq (m_A)^{-1} \frac{1 + a + p_{11}}{p_{11} - (1 + \delta_A)},$$

or $p_{22} > 1 + \delta_B$ and

$$N - 1 \leq (m_B)^{-1} \frac{1 + d + p_{22}}{p_{22} - (1 + \delta_B)},$$

then (1) has no non-trivial ground state.

Example 5. (A generalization of Corollary 2.2 of [12]). Consider

$$\begin{cases} -\operatorname{div}\left(\frac{Du}{(1 + |Du|^2)^{\frac{2-m}{2}}}\right) = |x|^\alpha |v|^{q-1} v \\ -\operatorname{div}\left(\frac{Dv}{(1 + |Dv|^2)^{\frac{2-n}{2}}}\right) = |x|^\beta |u|^{p-1} u \end{cases} \quad \text{in } \mathbb{R}^N \tag{85}$$

with $m, n \in (1, 2]$. If $N > m, n$ and $p, q > 1$ and

$$N - 1 \leq (n - 1) \frac{1 + \beta + (1 + \alpha + q)p}{pq - 1} \quad \text{or} \quad N - 1 \leq (m - 1) \frac{1 + \alpha + (1 + \beta + p)q}{pq - 1},$$

then (85) has no non-trivial ground state.

Remark. The technique used above allows us to obtain non-existence theorems for ground states of systems containing an arbitrary (finite) number of equations. However, for systems containing more than two equations the conditions will become rather involved. Some examples in this direction are studied in [14].

An application of Theorem 3.3 is the following. Consider a system that involves two degenerate generalized Laplacians, namely

$$\begin{cases} -\operatorname{div}(|Du|^{p-2}Du) = f(|x|, u, v) \\ -\operatorname{div}(|Dv|^{q-2}Dv) = g(|x|, u, v) \end{cases} \quad \text{in } \mathbb{R}^N, \tag{86}$$

where p, q are such that $2 < p, q < N$ and $f, g : \mathbb{R}^+ \times \mathbb{R}^2 \rightarrow \mathbb{R}$ are given functions specified next.

The result is an analogy of Theorem 3.2 in the degenerate case (i.e., $p, q > 2$). It generalizes Theorem 3.1 of [3].

Theorem 3.4. *Suppose that $N > p, q > 2$. Let $f, g \in C(\mathbb{R}^+ \times \mathbb{R}^2)$ with $f(r, 0, 0) = g(r, 0, 0) = 0 \ \forall r \geq 0$. Further, assume that there exist continuous $a, b, c, d : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ with $a(r), b(r), c(r), d(r) > 0$ for $r > 0$ and such that for large r and small u, v we have*

$$\begin{cases} f(r, u, v) \geq a(r)|u|^{\alpha-1}u + b(r)|v|^{\beta-1}v, \\ g(r, u, v) \geq c(r)|u|^{\gamma-1}u + d(r)|v|^{\delta-1}v, \end{cases} \tag{87}$$

where $\alpha, \beta, \gamma, \delta$ are positive constants. If one of the following conditions is satisfied,

(1) $\beta\gamma > pq$ and

$$\int_{r_0}^{\infty} s^{N-1}b(s) \left(\int_s^{\infty} \vartheta^{p-1-\beta\frac{N-q}{q-1}} c(\vartheta) d\vartheta \right)^{\frac{\gamma}{p-1}} ds = +\infty, \tag{88}$$

(2) $\beta\gamma > pq$ and

$$\int_{r_0}^{\infty} s^{N-1}c(s) \left(\int_s^{\infty} \vartheta^{q-1-\gamma\frac{N-p}{p-1}} b(\vartheta) d\vartheta \right)^{\frac{\beta}{p-1}} ds = +\infty, \tag{89}$$

(3) $\alpha > p$ and

$$\int_{r_0}^{\infty} s^{N-1-\alpha\frac{N-p}{p-1}} a(s) ds = +\infty, \tag{90}$$

(4) $\delta > q$ and

$$\int_{r_0}^{\infty} s^{N-1-\delta\frac{N-q}{q-1}} d(s) ds = +\infty, \tag{91}$$

then (86) has no non-trivial ground states.

Proof. The demonstration follows straightforwardly from Theorem 3.3 by choosing

$$\begin{aligned} \delta_A &= p - 1, & \delta_B &= q - 1, & \theta_A &= \frac{N - p}{p - 1}, \\ \theta_B &= \frac{N - q}{q - 1}, & m_A &= \frac{1}{p - 1}, & m_B &= \frac{1}{q - 1}, \\ p_{11} &= \alpha, & p_{21} &= \gamma, & p_{12} &= \beta, & p_{22} &= \delta. \end{aligned}$$

Remark. The above theorem can be applied to the scalar case

$$-\operatorname{div}(|Du|^{q-2}Du) = f(x, u) \quad \text{in } \mathbb{R}^N,$$

for some continuous function $a : [0, \infty) \rightarrow [0, \infty)$

$$f(x, u) \geq a(|x|)|u|^{\alpha-1}u \quad \text{for } r \text{ large and } u \text{ small.}$$

In this case we have non-existence of ground states if

$$\int_{r_0}^{\infty} s^{N-1-\alpha\frac{N-q}{q-1}} a(s) ds = +\infty.$$

For $a(r) \equiv a > 0$ condition (92) is equivalent to $\alpha \leq \frac{N}{N-q}(q - 1)$ (see [16], [17]).

4. Variational systems. In the case that the system under consideration has a variational structure it is possible to obtain refined versions of the preceding results by using the variational identities that were proved in [19], [4], [20] and [25]. In this section we shall consider a general result of this type and discuss some particular case.

We will start by recalling some facts about the scalar case. Consider the Lagrangian density

$$J(u) = \frac{1}{p}|Du|^p - \frac{1}{\alpha + 1}|x|^\delta|u|^{\alpha+1},$$

with $1 < p < N$. From [20] one may derive the following identity:

$$\frac{1-p}{p} \oint_{\partial\Omega} \left| \frac{\partial u}{\partial n} \right|^p (x \cdot n) d\sigma = \left(\frac{N-p}{p} - \frac{N+\delta}{\alpha+1} \right) \int_{\Omega} |x|^\delta |u|^{\alpha+1} dx. \tag{93}$$

Using identity (93) one finds that the boundary value problem

$$\begin{cases} -\operatorname{div}(|Du|^{p-2}Du) = |x|^\delta |u|^{\alpha-1} & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases} \tag{94}$$

has no positive solution when Ω is a bounded starshaped domain in \mathbb{R}^N and

$$\alpha + 1 \geq \frac{p(N + \delta)}{N - p}. \tag{95}$$

See for instance [4]. By variational techniques it can be shown that (94) has a non-trivial solution if

$$\alpha + 1 < \frac{p(N + \delta)}{N - p}, \tag{96}$$

which is equivalent to

$$N + \delta < \left(\frac{\alpha + 1}{p}(p + \delta)\right) / \left(-1 + \frac{\alpha + 1}{p}\right). \tag{97}$$

Taking $\Omega = \mathbb{R}^N$ and restricting it to radially symmetric solutions, the situation reverses, and (94) does not have positive radially symmetric solutions when (96) holds. However, when (95) holds, radially symmetric positive solutions are known to exist (see [16]). In the case of systems and more general nonlinearities we can derive a similar non-existence result by imposing natural growth conditions on H .

Our next result deals with the case when the system is the Euler-Lagrange equation $\delta\{J(u)\} = 0$ of the functional

$$J(u) = \sum_{i=1}^k \frac{1}{p_i} \int_{\mathbb{R}^N} |Du_i|^{p_i} dx - \int_{\mathbb{R}^N} H(|x|, u) dx, \tag{98}$$

that is, with a system of the form

$$\begin{cases} -\operatorname{div}(|Du_1|^{p_1-2} Du_1) = \frac{\partial}{\partial u_1} H(|x|, u) \\ \vdots \\ -\operatorname{div}(|Du_k|^{p_k-2} Du_k) = \frac{\partial}{\partial u_k} H(|x|, u) \end{cases} \quad \text{in } \mathbb{R}^N. \tag{99}$$

Here $k \geq 1$, $N > p_i > 1$ for all $i \in \{1, \dots, k\}$ and H is a given potential that satisfies the following assumptions.

(H) Let $H : \mathbb{R}^+ \times \mathbb{R}^k \rightarrow \mathbb{R}$ be a C^1 -function such that $\frac{\partial}{\partial u_i} H(r, 0) = 0 \ \forall r \geq 0$, $i \in \{1, \dots, k\}$ and

i) for all $(r, u) \in \mathbb{R}^+ \times \mathbb{R}^k$:

$$NH(r, u) + r \frac{\partial}{\partial r} H(r, u) > \sum_{i=1}^k \frac{N - p_i}{p_i} u_i \frac{\partial}{\partial u_i} H(r, u), \tag{100}$$

ii) there exist δ_i, α_i and $c_i > 0$ such that for all large r and sufficiently small $u \in \mathbb{R}^k$ we have

$$u_i \frac{\partial}{\partial u_i} H(r, u) \geq c_i r^{\delta_i} \prod_{i=1}^k |u_i|^{\alpha_i+1},$$

iii) there exists $\delta \in \mathbb{R}, c > 0$ such that for all large r and sufficiently small $u \in \mathbb{R}^k$ we have

$$H(r, u) \leq cr^\delta \prod_{i=1}^k |u_i|^{\alpha_i+1},$$

- iv) $\sum_{i=1}^k \frac{\alpha_i + 1}{p_i} > 1,$
- v) subcriticality

$$N + \delta < \left(\sum_{i=1}^k \frac{\alpha_i + 1}{p_i} (p_i + \delta_i) \right) / \left(-1 + \sum_{i=1}^k \frac{\alpha_i + 1}{p_i} \right). \tag{101}$$

Remark 1. For $k = 1$ (i.e., the scalar case), condition (101) reduces to the usual subcriticality condition (96). For the pure power case, that is, $H(r, u) = cr^\delta \prod_{i=1}^k |u_i|^{\alpha_i+1}$, (100) coincides with (101).

Remark 2. The strict inequality in (100) can be replaced by \geq if one assumes that there is a sequence $\{r_n\}$ with $r_n \rightarrow \infty$ for which the strict inequality holds.

Theorem 4.1. *Assume that the above condition (H) is satisfied. Then system (99) has no non-trivial ground states in $C^2(\mathbb{R}^N - \{0\})^k$.*

The proof of the theorem is based on the following version of the identity proved in [20].

Proposition 4.2. *Let $u \in C^2(\mathbb{R}^N - \{0\})^k$ be a radial solution of (99). Then the following identity holds. For any $R > 0$, we have*

$$\begin{aligned} & \int_0^R (NH(r, u) + r \frac{\partial}{\partial r} H(r, u)) r^{N-1} dr + \sum_{i=1}^k \left(\frac{p_i - N}{p_i} \int_0^R r^{N-1} u_i \frac{\partial}{\partial u_i} H(r, u) dr \right) \\ &= \sum_{i=1}^k \left(\frac{p_i - 1}{p_i} R^N |u'_i(R)|^p + \frac{N - p_i}{p_i} R^{N-1} |u'_i(R)|^{p-2} u'_i(R) u_i(R) \right) + R^N H(R, u(R)). \end{aligned} \tag{102}$$

In order to prove the theorem we shall also need some asymptotic estimates.

Lemma 4.3. *Let $u \in C^2(\mathbb{R}^N - \{0\})^k$ be a ground state of (99). If (H) ii), iii) and iv) hold, then there exists $c > 0$ such that for all sufficiently large r we have*

$$H(r, u(r)) \leq cr^{\delta-\mu}$$

with

$$\mu = \left(\sum_{i=1}^k \frac{\alpha_i + 1}{p_i} (p_i + \delta_i) \right) / \left(-1 + \sum_{i=1}^k \frac{\alpha_i + 1}{p_i} \right).$$

Proof. Let $u \in C^2(\mathbb{R}^N - \{0\})^k$ be a radial ground state of (99). By ii) it follows that for r sufficiently large, we have $u'_i(r) < 0$ for all i , and, for some $c > 0$ and all i ,

$$-(r^{N-1} |u'_i(r)|^{p_i-2} u'_i(r))' \geq cr^{\delta_i+N-1} u_1(r)^{\alpha_1+1} \dots u_i(r)^{\alpha_i} \dots u_k(r)^{\alpha_k+1}. \tag{103}$$

Integrating on (r, t) we obtain for all i

$$\begin{aligned} & t^{N-1}|u'_i(t)|^{p_i-1} + r^{N-1}|u'_i(r)|^{p_i-2}u'_i(r) \\ & \geq c \int_r^t s^{\delta_i+N-1}u_1(s)^{\alpha_1+1} \dots u_i(s)^{\alpha_i} \dots u_k(s)^{\alpha_k+1} ds, \end{aligned} \tag{104}$$

and hence, by using Lemma 2.2, that is (see (71)),

$$(r^{\theta_i}u_i(r))' \geq 0 \quad \text{for large } r,$$

with $\theta_i = \frac{N-p_i}{p_i-1}$, we obtain

$$t^{N-1}|u'_i(t)|^{p_i-1} \geq c^*(r^{\theta_1}u_1(r))^{\alpha_1+1} \dots (r^{\theta_i}u_i(r))^{\alpha_i} \dots (r^{\theta_k}u_k(r))^{\alpha_k+1} \int_r^t s^{\beta_i} ds, \tag{105}$$

where $\beta_i = \delta_i + N - 1 + \theta_i - \sum_{j=1}^k \theta_j(\alpha_j + 1)$. Using again the inequality

$$ru'_i(r) + \theta_i u_i(r) \geq 0 \quad \text{for } r \geq r_0, \tag{106}$$

and $u'_i(r) \leq 0$, we deduce from (105) that there exist $c > 0$ such that

$$t^{N-p_i}u_i(t)^{p_i-1} \geq c(r^{\theta_1}u_1(r))^{\alpha_1+1} \dots (r^{\theta_i}u_i(r))^{\alpha_i} \dots (r^{\theta_k}u_k(r))^{\alpha_k+1} \int_r^t s^{\beta_i} ds. \tag{107}$$

By choosing $t = 2r$ in (107), we obtain

$$2^{N-p_i}r^{N-p_i}u_i(r)^{p_i-1} \geq c(r^{\theta_1}u_1(r))^{\alpha_1+1} \dots (r^{\theta_i}u_i(r))^{\alpha_i} \dots (r^{\theta_k}u_k(r))^{\alpha_k+1} \gamma_i r^{\beta_i+1}, \tag{108}$$

with $\gamma_i = \frac{2^{\beta_i+1}-1}{\beta_i+1}$ if $\beta_i + 1 \neq 0$ and $\gamma_i = \log 2$ if $\beta_i + 1 = 0$. Thus, for some C and for every i and sufficiently large r , we have

$$Cu_i(r)^{\alpha_i+1} \geq \left(r^{p_i+\delta_i} \prod_{j=1}^k u_j(r)^{\alpha_j+1} \right)^{\frac{\alpha_i+1}{p_i}}. \tag{109}$$

After combining the k inequalities in (109), we obtain

$$\begin{aligned} C^k \prod_{i=1}^k u_i(r)^{\alpha_i+1} & \geq \prod_{i=1}^k \left(r^{p_i+\delta_i} \prod_{j=1}^k u_j(r)^{\alpha_j+1} \right)^{\frac{\alpha_i+1}{p_i}} \\ & = r^{(\sum_{i=1}^k (p_i+\delta_i) \frac{\alpha_i+1}{p_i})} \left(\prod_{j=1}^k u_j(r)^{\alpha_j+1} \right)^{\sum_{i=1}^k \frac{\alpha_i+1}{p_i}}, \end{aligned} \tag{110}$$

or in other words

$$C^k \geq r^{(\sum_{i=1}^k \frac{\alpha_i+1}{p_i}(p_i+\delta_i))} \left(\prod_{j=1}^k u_j(r)^{\alpha_j+1} \right)^{(\sum_{i=1}^k \frac{\alpha_i+1}{p_i})-1}. \tag{111}$$

Now, by using condition **(H)**(iv) in (111) it follows that

$$C^* \geq r^\mu \prod_{j=1}^k u_j(r)^{\alpha_j+1}, \tag{112}$$

with

$$\mu = \left(\sum_{i=1}^k \frac{\alpha_i+1}{p_i}(p_i+\delta_i) \right) / \left(-1 + \sum_{i=1}^k \frac{\alpha_i+1}{p_i} \right). \tag{113}$$

From (112) and condition **(H)**(iii), we find for large r that

$$H(r, u(r)) \leq cr^\delta \prod_{j=1}^k u_j(r)^{\alpha_j+1} \leq c_1 r^{\delta-\mu}.$$

This concludes the proof of Lemma 4.3.

Proof of Theorem 4.1. We argue by contradiction. Let u be a non-trivial ground state of (99). Applying identity (102) to this specific situation it follows that the left hand side of (102) is bounded away from zero for R large. On the other hand, by using Lemma 2.1 of [3] and the fact that $u'_i(R) \leq 0 \forall i$ we know that the first term of the right hand side of (102) is non-positive. We conclude the proof by using Lemma 4.3 in order to show that the second term of the right hand side of (102), namely $H(R, u(R)) R^N$, converges to zero as $R \rightarrow \infty$. As a result we obtain a contradiction.

Remark. The same technique used in the proof of the last theorem allows us to study more general systems than (99). It is possible to show that non-existence results can be obtained for systems of the form

$$\begin{cases} -\operatorname{div}(A_1(|Du_1|)Du_1) = \frac{\partial}{\partial u_1} H(|x|, u) \\ \vdots \\ -\operatorname{div}(A_k(|Du_k|)Du_k) = \frac{\partial}{\partial u_k} H(|x|, u), \end{cases}$$

where each operator A_i satisfies (i)_{A_i} of (56) and where the potential $H(|x|, u)$ is controlled by a suitable growth condition. This problem will be considered in a forthcoming paper ([14]).

Example 6. See [24]. As an example of a system with a variational structure we consider

$$\begin{cases} -\operatorname{div}(|Du|^{p-2}Du) = |x|^\delta |u|^{\alpha-1} |v|^{\beta+1} u, \\ -\operatorname{div}(|Dv|^{q-2}Dv) = |x|^\delta |u|^{\alpha+1} |v|^{\beta-1} v. \end{cases} \tag{114}$$

We have, after a possible rescaling, that

$$H(r, u) = r^\delta |u|^{\alpha+1} |v|^{\beta+1}.$$

Suppose $\frac{(\alpha+1)}{p} + \frac{(\beta+1)}{q} > 1$ and $N > p, q > 2$. If

$$N + \delta < \frac{\frac{(\alpha+1)}{p}(p + \delta) + \frac{(\beta+1)}{q}(q + \delta)}{\frac{(\alpha+1)}{p} + \frac{(\beta+1)}{q} - 1},$$

then there is no non-trivial ground state of (114). The proof follows from a straightforward calculation using Theorem 4.1.

Acknowledgment. This paper was completed during a visit of Enzo Mitidieri at the Stieltjes Institute/Delft University in October 1993. He wishes to thank this institution for its support and for the kind atmosphere during his visit. He also thanks Professor Pohozaev for bringing the paper [19] to his attention.

REFERENCES

- [1] Ph. Clément, D.G. de Figueiredo, and E. Mitidieri, *Positive solutions of semilinear elliptic systems*, Comm. in P.D.E., 17 (1992), 923–940.
- [2] Ph. Clément, D.G. de Figueiredo, and E. Mitidieri, *A priori estimates for positive solutions of semilinear elliptic stems via Hardy-Sobolev estimates*, Relatorio de Pesquisa, Julho RP 20/92 Universidade Estadual de Campinas.
- [3] Ph. Clément, R. Manásevich, and E. Mitidieri, *Positive solutions for a quasilinear system via blow-up*, Comm. in P.D.E., 18 (1993), 2071–2106.
- [4] H. Egnell, *Existence and non-existence results for the m -Laplacian equation involving critical Sobolev exponents*, Ph.D. Thesis Uppsala University 1987.
- [5] D.G. de Figueiredo and P.L. Felmer, *On superquadratic elliptic systems*, Trans. Amer. Math. Soc., 343 (1994), 99–116.
- [6] D.G. de Figueiredo and P.L. Felmer, *A Liouville-type theorem for elliptic systems*, preprint.
- [7] B. Gidas, *Symmetry properties and isolated singularities of positive solutions of nonlinear elliptic equations*, in “Nonlinear Differential Equations in Engineering and Applied Sciences,” ed. R.L. Sternberg, Marcel Dekker, New York 1979, 255–273.
- [8] B. Gidas and J. Spruck, *Global and local behavior of positive solutions on nonlinear elliptic equations*, Comm. Pure Appl. Math., 34 (1981), 525–598.
- [9] B. Gidas and J. Spruck, *A priori bounds for positive solutions of nonlinear elliptic equations*, Comm. in P.D.E., 6 (1981), 883–901.
- [10] J. Hulshof and R.C.A.M. van der Vorst, *Differential systems with strongly indefinite variational structure*, J. Functional Analysis, 114 (1993), 32–58.
- [11] P. Marcellini and K. Miller, *Asymptotic growth for evolutionary surfaces of prescribed mean curvature*, in “Nonlinear Functional Analysis and its Applications,” Proceedings of Symposia in Pure Mathematics of the Amer. Math. Soc., 45 Part 2 (1986), 175–178.
- [12] E. Mitidieri, *Nonexistence of positive solutions of semilinear elliptic systems in \mathbb{R}^N* , Quaderno Matematici 285, Report Università Trieste 1992.
- [13] E. Mitidieri, *A Rellich type identity and applications*, Comm. in P.D.E., 18 (1993), 125–151.
- [14] E. Mitidieri, G. Sweers, and R.C.A.M. van der Vorst, *Existence and non existence of ground states for quasilinear systems*, in preparation.
- [15] W.M. Ni and J. Serrin, *Non-existence theorems for quasilinear partial differential equations*, Rend. Circ. Mat. Palermo, suppl., 8 (1985), 171–185.

- [16] W.M. Ni and J. Serrin, *Existence and nonexistence for ground states of quasilinear partial differential equations, the anomalous case*, Atti Convegni Lincei, 77 (1985), 231–257.
- [17] W.M. Ni and J. Serrin, *Non-existence theorems for singular solutions of quasilinear partial differential equations*, Comm. Pure Appl. Math., 39 (1986), 379–399.
- [18] L.A. Peletier and R.C.A.M. van der Vorst, *Existence and non-existence of positive solutions of non-linear elliptic systems and the biharmonic equation*, Differential and Integral Equations, 6 (1992), 747–767.
- [19] S.I. Pohozaev, *On the eigenfunction of quasilinear elliptic problems*, Mat. Sbornik, 82 (1-4) 2 (1970), 171–188.
- [20] P. Pucci and J. Serrin, *A general variational identity*, Indiana Univ. Math. J., 35 (1986), 681–703.
- [21] P. Pucci and J. Serrin, *Continuation and limit properties for solutions of strongly nonlinear second order differential equations*, Asymptotic Analysis, 4 (1991), 97–160.
- [22] M.A.S. Souto, *Sobre a existencia de solucoes positivas para sistemas cooperativos nao lineare*, PhD thesis, Unicamp Brasil 1992.
- [23] R. Soranzo, *A priori estimates and existence of positive solutions of a superlinear polyharmonic equation*, Dynamic System and Applications, to appear.
- [24] F. de Thelin and J. Velin, *Existence and nonexistence of nontrivial solutions for some nonlinear elliptic systems*, preprint.
- [25] R.C.A.M. van der Vorst, *Variational identities and applications to differential systems*, Arch. Rational Mech. Anal., 116 (1991), 375–398.