

Kashaev's Conjecture and the Chern-Simons Invariants of Knots and Links

Hitoshi Murakami, Jun Murakami, Miyuki Okamoto, Toshie Takata, and Yoshiyuki Yokota

CONTENTS

1. Introduction
 2. Preliminaries
 3. Knot 6_3
 4. Knot 8_9
 5. Knot 8_{20}
 6. Whitehead Link
 7. Topological Chern-Simons Invariant and Some Examples
 8. Conclusion
- Acknowledgments
References

R. M. Kashaev conjectured that the asymptotic behavior of the link invariant he introduced [Kashaev 95], which equals the colored Jones polynomial evaluated at a root of unity, determines the hyperbolic volume of any hyperbolic link complement. We observe numerically that for knots 6_3 , 8_9 and 8_{20} and for the Whitehead link, the colored Jones polynomials are related to the hyperbolic volumes and the Chern–Simons invariants and propose a complexification of Kashaev's conjecture.

1. INTRODUCTION

In [Kashaev 95], R.M. Kashaev defined a link invariant associated with the quantum dilogarithm, depending on a positive integer N , which is denoted by $\langle L \rangle_N$ for a link L . Moreover, in [Kashaev 97], he conjectured that for any hyperbolic link L , the asymptotics at $N \rightarrow \infty$ of $|\langle L \rangle_N|$ gives its volume, that is

$$\text{vol}(L) = 2\pi \lim_{N \rightarrow \infty} \frac{\log |\langle L \rangle_N|}{N}$$

with $\text{vol}(L)$ the hyperbolic volume of the complement of L . He showed that this conjecture is true for three doubled knots 4_1 , 5_2 , and 6_1 . Unfortunately, his proof is not mathematically rigorous.

Afterwards, in [Murakami and Murakami 01], the first two authors proved that for any link L , Kashaev's invariant $\langle L \rangle_N$ is equal to the colored Jones polynomial evaluated at $\exp(2\pi\sqrt{-1}/N)$, which is written by $J_N(L)$, and extended Kashaev's conjecture as follows.

Conjecture 1.1. (Volume conjecture.)

$$\|L\| = \frac{2\pi}{v_3} \lim_{N \rightarrow \infty} \frac{\log |J_N(L)|}{N},$$

where $\|L\|$ is the simplicial volume of the complement of L and v_3 is the volume of the ideal regular tetrahedron.

2000 AMS Subject Classification: Primary 57M27, 57M25, 57M50; Secondary 41A60, 17B37, 81R50

Keywords: Volume conjecture, Kashaev's conjecture, colored Jones polynomial, Chern-Simons invariant, volume

Note that the hyperbolic volume $\text{vol}(L)$ of a hyperbolic link L is equal to $\|L\|$ multiplied by v_3 . This conjecture is not true for links in general, as $J_N(L)$ vanishes for a split link L . It is shown by Kashaev and O. Tirkkonen in [Kashaev and Tirkkonen 00] that the volume conjecture holds for torus knots. See [Thurston 99] and [Yokota 00, Yokota 02] for discussions about Kashaev’s conjecture for hyperbolic knots from the viewpoint of tetrahedron decomposition.

In this paper, following Kashaev’s way to analyze the asymptotic behavior of the invariant, we observe numerically, by using MAPLE V (a product of Waterloo Maple Inc.) and SnapPea [Weeks 02], that for the hyperbolic knots $6_3, 8_9, 8_{20}$, and for the Whitehead link, the colored Jones polynomials are related to the hyperbolic volumes and the Chern–Simons invariants. Note that the knots 6_3 and 8_9 are not doubles of the unknot.

We also discuss a relation between the asymptotic behavior of $J_N(L)$ and the Chern–Simons invariant of the complement of the above-mentioned links L , and propose the following conjecture.

Conjecture 1.2. (Complexification of Kashaev’s conjecture.) *Let L be a hyperbolic link. Then the following formula holds.*

$$J_N(L) \sim \exp \frac{N}{2\pi} (\text{vol}(L) + \sqrt{-1} \text{CS}(L)) \quad (N \rightarrow \infty)$$

where $\text{CS}(L)$ is the Chern–Simons invariant of L [Chern and Simons 74, Meyerhoff 86]. Note that the complement of L is a hyperbolic manifold with cusps.

The statement of this conjecture will be given more properly in the last section.

2. PRELIMINARIES

First, we will briefly review the colored Jones polynomials of links following [Kirby and Melvin 91]. It is obtained from the quantum group $U_q(\mathfrak{sl}(2, \mathbb{C}))$ and its N -dimensional irreducible representation.

Let L be an oriented link. We consider a $(1, 1)$ -tangle presentation of L , obtained by cutting a component of the link. We assume that all crossing and local extreme points are as in Figure 1. We can calculate the N -colored Jones polynomial $J_L(N)$ evaluated at the N -th root of unity for L in the following way. We start with a labeling of the edges of the $(1, 1)$ -tangle presentation with labels $\{0, 1, \dots, N - 1\}$. Here we label the two edges containing the end points of the tangle by 0. Following the labeling,




FIGURE 1. Crossings and local extrema in a link diagram.

we associate a positive (respectively, negative) crossing with the element R_{kl}^{ij} (respectively, \bar{R}_{kl}^{ij}), a maximal point \cap labeled by i with the element $-s^{-2i-1}$, and a minimal point \cup labeled by i with the element $-s^{2i+1}$ with $s = \exp \left(\frac{\pi\sqrt{-1}}{N} \right)$ as in Figure 1.

Here R_{kl}^{ij} and \bar{R}_{kl}^{ij} are given by

$$R_{kl}^{ij} = \sum_{n=0}^{\min(N-1-i, j)} \delta_{l, i+n} \delta_{k, j-n} \frac{(i+n)!(N-1+n-j)!}{(i)!(N-1-j)!(n)!} \times s^{2(i-\frac{N-1}{2})(j-\frac{N-1}{2})-n(i-j)-\frac{n(n+1)}{2}},$$

$$\bar{R}_{kl}^{ij} = \sum_{n=0}^{\min(N-1-j, i)} \delta_{l, i-n} \delta_{k, j+n} \frac{(j+n)!(N-1+n-i)!}{(j)!(N-1-i)!(n)!} (-1)^n \times s^{-2(i-\frac{N-1}{2})(j-\frac{N-1}{2})-n(i-j)+\frac{n(n+1)}{2}}$$

with $(n)! = (s - s^{-1})(s^2 - s^{-2}) \dots (s^n - s^{-n})$.

After multiplying all elements associated with the critical points, we sum over all indices, ignoring framings of links.

We calculate the colored Jones polynomial of the Whitehead link as an example. We can label each edge in the following way, noting Kronecker’s deltas in R_{kl}^{ij} and \bar{R}_{kl}^{ij} .

We have to rotate a crossing where edges go up. In that case, we use \cup and/or \cap to calculate the invariant.




FIGURE 2. Labeling of the Whitehead link.

$$J_N(6_3) = \sum_{\substack{0 \leq k, l, m \\ k+l+m \leq N-1}} (-1)^{k+l} s^{\frac{(l+k)(l+k+1)}{2} - \frac{(m+k)(m+k+1)}{2} + \frac{k(k+1)}{2} + 2(m-l)(k+1) + N(m-l+k)} \\ \times \frac{(N-1-l)!(N-1-m)!(l+m+k)!(N-1)!(1-s^{-2N-2}) \cdots (1-s^{-2N-2k})}{(N-1-l-m-k)!(N-1-l-k)!(N-1-m-k)!(l)!(m)!(k)!}.$$

FIGURE 3. Calculation of $J_N(6_3)$.

Then we calculate the formula

$$J_N(L) = \sum_{\substack{0 \leq i, j, k \leq N-1 \\ i, j \geq k}} \frac{(q)_i (q)_j \{(q)_{N-1-k}\}^2}{\{(q)_k\}^2 (q)_{N-1-i} (q)_{N-1-j} (q)_{i-k} (q)_{j-k}} q^{-k(i+j+1)}, \tag{2-1}$$

where $q = s^2 = \exp\left(\frac{2\pi\sqrt{-1}}{N}\right)$. Here $(x)_k = (1-x)(1-x^2) \cdots (1-x^k)$.

Next, the Chern–Simons invariant of a link is defined. Let \mathcal{A} be the set of all $SO(3)$ -connections of the trivial $SO(3)$ -bundle of a closed three-manifold M and $cs: \mathcal{A} \rightarrow \mathbb{R}$ the Chern–Simons functional defined by

$$cs(A) = \frac{1}{8\pi^2} \text{Tr} \left(A \wedge dA + \frac{2}{3} A \wedge A \wedge A \right).$$

The Chern–Simons invariant of the connection A is then defined to be the integral

$$cs_M(A) = \int_{s(M)} cs(A) \in \mathbb{R}/\mathbb{Z},$$

where the integral is over a section s of the $SO(3)$ -bundle (i.e., an orthonormal frame field on M) [Chern and Simons 74]. If M is hyperbolic, we define $cs(M)$ to be the Chern–Simons invariant of the connection defined by the hyperbolic metric.

The definition of the Chern–Simons invariant for hyperbolic three-manifolds with cusps is due to R. Meyerhoff [Meyerhoff 86]. It is defined modulo $1/2$ by using a special singular frame field which is linear near the cusps. See [Meyerhoff 86] for details. See [Coulson et al. 00] to examine how it is computed by SnapPea [Weeks 02]. Throughout this paper, we use another normalization $CS(M) = -2\pi^2 cs(M)$ so that $\text{vol}(M) + \sqrt{-1} CS(M)$ is a natural complexification of the hyperbolic volume $\text{vol}(M)$ (see [Neumann and Zagier 85, Yoshida 85]).

3. KNOT 6_3

Let us calculate the colored Jones polynomial of the knot 6_3 using the labeling as in Figure 4.

Putting $k = n_1 + n_2$ and using the formula in [Murakami and Murakami 01]

$$\sum_{i=0}^{N-1} (-1)^i s^{\beta i} \begin{bmatrix} \alpha \\ i \end{bmatrix} = \prod_{j=1}^{\alpha} (1 - s^{\beta+\alpha+1-2j}) \tag{3-1}$$

with $\alpha = k$, $i = n_1$, and $\beta = -k - 1 - 2N$, we calculate $J_N(6_3)$ as shown in Figure 3.

The colored Jones polynomial of the knot 6_3 is given by

$$J_N(6_3) = \sum_{\substack{k, l, m \geq 0 \\ k+l+m \leq N-1}} \left| \frac{(q)_{k+l+m}}{(q)_l (q)_m} \right|^2 (q)_{k+l} (\bar{q})_{m+k} q^{(m-l)(k+1)}. \tag{3-2}$$

We review the technique in [Kashaev 97]. For a complex number p and a positive real number γ with $|\text{Re } p| < \pi + \gamma$, we define

$$S_\gamma(p) = \exp \frac{1}{4} \int_{-\infty}^{\infty} \frac{e^{px}}{\sinh(\pi x) \sinh(\gamma x)} \frac{dx}{x}.$$

Here Re denotes the real part. This function has two properties:

- (a) $(1 + \exp(\sqrt{-1}p)) S_\gamma(p + \gamma) = S_\gamma(p - \gamma)$,
- (b) $S_\gamma(p) \sim \exp\left(\frac{1}{2\gamma\sqrt{-1}} \text{Li}_2(-\exp(\sqrt{-1}p))\right)$ ($\gamma \rightarrow 0$),




FIGURE 4. Labeling of 6_3 .

where

$$\text{Li}_2(z) = - \int_0^z \frac{\log(1-u)}{u} du.$$

We put

$$f_\gamma(p) = \frac{S_\gamma(\gamma - \pi)}{S_\gamma(p)}, \quad \bar{f}_\gamma(p) = \frac{S_\gamma(-p)}{S_\gamma(\pi - \gamma)},$$

so that

$$(q)_k = f_\gamma(-\pi + (2k + 1)\gamma), \quad (\bar{q})_k = \bar{f}_\gamma(-\pi + (2k + 1)\gamma).$$

Following Kashaev's analysis, we rewrite the formula (3-2) as a multiple integral with appropriately chosen contours. (Note that there is considerable doubt as to the contours.) By using the property (b), it can be asymptotically approximated by

$$\iiint \exp \frac{\sqrt{-1}}{2\gamma} V_{6_3}(z, u, v) dz du dv$$

with $\gamma = \pi/N$. Here z , u , and v correspond to q^k , q^m , and q^l , respectively, and

$$\begin{aligned} V_{6_3}(z, u, v) = & \text{Li}_2(zuv) - \text{Li}_2\left(\frac{1}{zuv}\right) + \text{Li}_2(zv) \\ & - \text{Li}_2\left(\frac{1}{zu}\right) - \text{Li}_2(u) + \text{Li}_2\left(\frac{1}{u}\right) \\ & - \text{Li}_2(v) + \text{Li}_2\left(\frac{1}{v}\right) - \log z \log \frac{u}{v}. \end{aligned}$$

Then there exists a stationary point

$$(z_0, u_0, v_0) = (0.204323 - 0.978904\sqrt{-1}, 1.60838 + 0.558752\sqrt{-1}, 0.554788 + 0.192734\sqrt{-1})$$

$$\text{Im } V_{6_3}(z_0, u_0, v_0) < 0, \quad \arg z_0 + \arg u_0 + \arg v_0 \leq 2\pi,$$

and we have

$$- \text{Im } V_{6_3}(z_0, u_0, v_0) = 5.693021 \dots,$$

$$\text{Re } V_{6_3}(z_0, u_0, v_0) = 0.$$

From values of $\text{vol}(6_3)$ and $\text{CS}(6_3)$ given by SnapPea, we see that the equation

$$\exp \frac{\sqrt{-1}}{2\gamma} V_{6_3}(z_0, u_0, v_0) = \exp \frac{\text{vol}(6_3) + \sqrt{-1} \text{CS}(6_3)}{2\gamma}$$

holds up to digits shown above.

4. KNOT 8_9

We label the edges of the $(1, 1)$ -tangle presentation of the knot 8_9 as in Figure 5.




FIGURE 5. Labeling of 8_9 .

We obtain the following formula of the colored Jones polynomial of the knot 8_9 , where we put $l = m_1 + m_2 + k_1 + k_2$ and use the formula (3-1):

$$\begin{aligned} J_N(8_9) = & \sum_{\substack{0 \leq l, m_1, m_2, n_1, n_2 \leq N-1 \\ m_1+n_1, m_2+n_2 \leq l \\ m_1+m_2 \leq l}} \left| \frac{(q)_{l-m_1}(q)_l(q)_{l-m_2}}{(q)_{m_1}(q)_{m_2}(q)_{n_1}(q)_{n_2}} \right|^2 \\ & \times \frac{(\bar{q})_{l-n_1}(q)_{l-n_2}}{(q)_{l-m_1-n_1}(\bar{q})_{l-m_2-n_2}} \\ & \times q^{(m_2-m_1)(l-m_1-m_2)+(n_2-n_1)(l-n_1-n_2)+m_2-m_1+n_2-n_1}, \end{aligned}$$

which can be asymptotically approximated by

$$\int \dots \int \exp \frac{\sqrt{-1}}{2\gamma} V_{8_9}(x, y, z, u, v) dx dy dz du dv,$$

where x , y , z , u , and v correspond to q^{-l} , q^{m_1} , q^{m_2} , q^{n_1} , and q^{n_2} respectively, and

$$\begin{aligned} V_{8_9}(x, y, z, u, v) = & -\text{Li}_2(xy) + \text{Li}_2\left(\frac{1}{xy}\right) - \text{Li}_2(xz) + \text{Li}_2\left(\frac{1}{xz}\right) \\ & - \text{Li}_2(xu) + \text{Li}_2\left(\frac{1}{xv}\right) - \text{Li}_2(x) + \text{Li}_2\left(\frac{1}{x}\right) - \text{Li}_2(y) \\ & + \text{Li}_2\left(\frac{1}{y}\right) - \text{Li}_2(z) + \text{Li}_2\left(\frac{1}{z}\right) - \text{Li}_2(u) + \text{Li}_2\left(\frac{1}{u}\right) \\ & - \text{Li}_2(v) + \text{Li}_2\left(\frac{1}{v}\right) + \text{Li}_2(xzv) - \text{Li}_2\left(\frac{1}{xyu}\right) \\ & - \log \frac{y}{z} \log(xzv) - \log \frac{u}{v} \log(xyu). \end{aligned}$$

Thus, we have

$$- \text{Im } V_{8_9}(x_0, y_0, z_0, u_0, v_0) = 7.5881802 \dots,$$

$$\text{Re } V_{8_9}(x_0, y_0, z_0, u_0, v_0) = 0$$

for

$$\begin{aligned} x_0 &= 0.7366011609 - 0.6763273835\sqrt{-1}, \\ y_0 &= 0.4472176075 - 0.1647027124\sqrt{-1}, \\ z_0 &= 1.968989044 - 0.7251455025\sqrt{-1}, \\ u_0 &= 0.3859112582 - 0.0202712198\sqrt{-1}, \\ v_0 &= 2.584139126 - 0.1357401508\sqrt{-1} \end{aligned}$$

satisfying

$$\begin{aligned} \operatorname{Im} V_{8_9}(x_0, y_0, z_0, u_0, v_0) &< 0, \\ \arg x_0 + \arg y_0 + \arg u_0 &\leq 2\pi, \\ \arg x_0 + \arg z_0 + \arg v_0 &\leq 2\pi, \\ \arg x_0 + \arg u_0 + \arg v_0 &\leq 2\pi. \end{aligned}$$

It follows from the calculation by SnapPea that

$$\begin{aligned} &\exp \frac{\sqrt{-1}}{2\gamma} V_{8_9}(x_0, y_0, z_0, u_0, v_0) \\ &= \exp \frac{\operatorname{vol}(8_9) + \sqrt{-1} \operatorname{CS}(8_9)}{2\gamma}, \end{aligned}$$

up to digits shown above.

5. KNOT 8₂₀

In this section, we discuss a relation between the asymptotic behavior of the colored Jones polynomial and the Chern–Simons invariant for the knot 8₂₀. We label each edge in the diagram of the knot in Figure 6.




FIGURE 6. Labeling of 8₂₀.

The N -colored Jones polynomial of the knot 8₂₀ is given by

$$\sum_{\substack{j,l \leq k \leq i+l \leq j+m \\ j \leq i \\ 0 \leq i,j,k,l,m \leq N-1}} \frac{\{(\bar{q})_i(q)_k(\bar{q})_m\}^2}{\{(\bar{q})_j(q)_l\}^2 (q)_{k-l}(\bar{q})_{i-k+l}(\bar{q})_{j+m-i-l}(q)_{i-j}(q)_{k-j}} \times q^{k+m+im+km-il}}, \quad (5-1)$$

which can be rewritten in the integral

$$\int \cdots \int \exp \frac{\sqrt{-1}}{2\gamma} V_{8_{20}}(x, y, z, u, v) dx dy dz du dv$$

with

$$\begin{aligned} V_{8_{20}}(x, y, z, u, v) &= -2\operatorname{Li}_2(x) + 2\operatorname{Li}_2\left(\frac{1}{y}\right) + 2\operatorname{Li}_2(z) \\ &\quad - 2\operatorname{Li}_2\left(\frac{1}{u}\right) - 2\operatorname{Li}_2\left(\frac{1}{x}\right) - \operatorname{Li}_2\left(\frac{1}{xy}\right) \\ &\quad - \operatorname{Li}_2\left(\frac{z}{y}\right) - \operatorname{Li}_2(zu) + \operatorname{Li}_2(xzu) \\ &\quad + \operatorname{Li}_2\left(\frac{1}{xyuv}\right) + \log x \log u \\ &\quad + \log x \log v - \log z \log v + \frac{\pi^2}{2}. \end{aligned}$$

Here $x, y, z, u,$ and v correspond to $q^{-i}, q^j, q^k, q^{-l},$ and $q^m,$ respectively.

Stationary points are solutions to partial differential equations,

$$\frac{\partial V_{8_{20}}}{\partial x} = \frac{\partial V_{8_{20}}}{\partial y} = \frac{\partial V_{8_{20}}}{\partial z} = \frac{\partial V_{8_{20}}}{\partial u} = \frac{\partial V_{8_{20}}}{\partial v} = 0.$$

From these equations, we have the following system of algebraic equations:

$$\begin{aligned} (1-x)^2 \left(1 - \frac{1}{xyuv}\right) uv &= \left(1 - \frac{1}{xy}\right) (1-xzu), \\ \left(1 - \frac{1}{xy}\right) \left(1 - \frac{z}{y}\right) &= \left(1 - \frac{1}{y}\right)^2 \left(1 - \frac{1}{xyuv}\right), \\ (1-z)^2 (1-xzu) v &= (1-zu) \left(1 - \frac{z}{y}\right), \\ (1-zu) \left(1 - \frac{1}{xyuv}\right) x &= \left(1 - \frac{1}{u}\right)^2 (1-xzu), \\ \left(1 - \frac{1}{v}\right)^2 z &= \left(1 - \frac{1}{xyuv}\right) x. \end{aligned}$$

Using MAPLE V, we get a stationary point $(x_0, y_0, z_0, u_0, v_0)$ which satisfies the conditions

$$\arg \frac{1}{u_0} \leq \arg z_0, \quad \arg z_0 \leq \arg \frac{1}{x_0} + \arg \frac{1}{u_0}$$

from the range in the summation in (5-1), and

$$\text{Im } V_{8_{20}}(x_0, y_0, z_0, u_0, v_0) < 0,$$

where Im denotes the imaginary part. Note that the range of (5-1) can be read as

$$\begin{aligned} \arg \frac{1}{u} &\leq \arg z \leq \arg \frac{1}{x} + \arg \frac{1}{u}, \\ \arg \frac{1}{x} + \arg \frac{1}{y} + \arg \frac{1}{u} &\leq \arg v, \\ 0 &\leq \arg \frac{1}{x} + \arg \frac{1}{y}, \\ 0 &\leq \arg \frac{1}{x}, \arg z, \arg \frac{1}{u}, \arg v \leq 2\pi. \end{aligned}$$

To put it concretely,

$$\begin{aligned} x_0 &= 2.878599677 + 2.657408013\sqrt{-1}, \\ y_0 &= \infty, \\ z_0 &= -0.4425377456 - 0.4544788919\sqrt{-1}, \\ u_0 &= 0.3542198353 - 0.02180673815\sqrt{-1}, \\ v_0 &= 0.1458832937 - 0.3399257634\sqrt{-1}. \end{aligned}$$

Then we obtain

$$\begin{aligned} -\text{Im } V_{8_{20}}(x_0, y_0, z_0, u_0, v_0) &= 4.1249032\dots, \\ -\frac{\text{Re } V_{8_{20}}(x_0, y_0, z_0, u_0, v_0) + \pi^2}{2\pi^2} &= 0.1033634\dots \end{aligned}$$

Applying values of $\text{vol}(8_{20})$ and $\text{CS}(8_{20})$ given by SnapPea [Weeks 02], we see that the following equation holds up to digits shown above.

$$\begin{aligned} \exp \frac{\sqrt{-1}}{2\gamma} V_{8_{20}}(x_0, y_0, z_0, u_0, v_0) \\ = \exp \frac{\text{vol}(8_{20}) + \sqrt{-1} \text{CS}(8_{20})}{2\gamma}. \end{aligned}$$

Note that $\text{CS}(8_{20})$ is defined modulo π^2 .

6. WHITEHEAD LINK

For the final example, we calculate the limit of the colored Jones polynomial of the Whitehead link given by (2-1), which can be changed to the formula

$$J_N(L) = \sum_{\substack{0 \leq i, j, k \leq N-1 \\ k \leq i, j}} \frac{\{(\bar{q})_i(\bar{q})_j\}^2}{(q)_k^4(\bar{q})_{i-k}(\bar{q})_{j-k}} q^{-(N-1)N/2}.$$

This can be asymptotically approximated by

$$\iiint \exp \frac{\sqrt{-1}}{2\gamma} V_L(x, y, z) dx dy dz,$$

where

$$\begin{aligned} V_L(x, y, z) &= -2\text{Li}_2\left(\frac{1}{x}\right) - 2\text{Li}_2\left(\frac{1}{y}\right) - 4\text{Li}_2(z) \\ &\quad + \text{Li}_2\left(\frac{z}{x}\right) + \text{Li}_2\left(\frac{z}{y}\right) + \pi^2, \end{aligned}$$

and $x, y,$ and z correspond to $q^i, q^j,$ and q^k respectively. For a stationary point $(x_0, y_0, z_0) = (\infty, \infty, 1 + \sqrt{-1})$, we obtain

$$\begin{aligned} -\text{Im } V_L(x_0, y_0, z_0) &= 3.663862\dots, \\ -\frac{\text{Re } V_L(x_0, y_0, z_0)}{2\pi^2} &= -0.1250000\dots \end{aligned}$$

Since these values agree with SnapPea, the equation

$$\exp \frac{\sqrt{-1}}{2\gamma} V_L(x_0, y_0, z_0) = \exp \frac{\text{vol}(L) + \sqrt{-1} \text{CS}(L)}{2\gamma}$$

holds up to digits shown above.

7. TOPOLOGICAL CHERN-SIMONS INVARIANT AND SOME EXAMPLES

We propose a topological definition of the Chern-Simons invariant for links.

For a link L , if there exists the limit

$$2\pi \text{Im} \lim_{N \rightarrow \infty} \log \frac{J_{N+1}(L)}{J_N(L)} \pmod{\pi^2},$$

then we denote it by $\text{CS}_{\text{TOP}}(L)$ and call it the *topological Chern-Simons invariant* of L .

Let us give some numerical examples.

For the knot 5_2 , we list some values of $(N, 2\pi \log(J_{N+1}(5_2)/J_N(5_2)))$ by Pari-Gp in Table 1.

By fitting the above data to quadratic functions on $1/N$, we can obtain the limit value

$$2.82813 - 3.02414\sqrt{-1}$$

of $2\pi \log(J_{N+1}(5_2)/J_N(5_2))$ as $N \rightarrow \infty$ numerically, which agrees with the value

$$2.8281220 - 3.02412837\sqrt{-1}$$

by SnapPea. We display our data graphically in Figures 7 and 8, which help us to see the limit.

Similarly, for the Whitehead link L , we illustrate our numerical check in Table 2, Figure 9, and Figure 10.

Fitting, we get the numerical limit value $3.66386 + 2.46742\sqrt{-1}$ of $2\pi \log(J_{N+1}(L)/J_N(L))$ as $N \rightarrow \infty$, which agrees with our result in Section 6.

(40, 3.058223721261842722613885956 – 3.022924613281720287391974968√–1)
(50, 3.013081508530188353573854822 – 3.023340368517507069134855780√–1)
(60, 2.982744318753580696821772299 – 3.023574042878935429645720640√–1)
(70, 2.960955404961739170749114151 – 3.023717381786374852930574631√–1)
(80, 2.944548269170450112446966301 – 3.023811574968472287718611711√–1)
(100, 2.921483906108228993018469212 – 3.023923719027833555669502480√–1)
(120, 2.906046421388666000282542398 – 3.023985374930307234443986632√–1)
(150, 2.890559881907537128372001511 – 3.024036295143969179028770901√–1)
(200, 2.875024234226941620327156350 – 3.024076266558545340852410631√–1)
(250, 2.865679250969538531562099056 – 3.024094905811349375139149331√–1)

TABLE 1.

(40, 3.892920359101811097809525583 + 2.457483997330866045812504703√–1)
(50, 3.848161466402914225154530180 + 2.461039474018016569869745301√–1)
(60, 3.818029013349499312708236153 + 2.462976748675980254703390855√–1)
(70, 3.796362501209537691078944556 + 2.464147191795881614582476451√–1)
(80, 3.780034327560022195082015385 + 2.464907923404764622274395868√–1)
(100, 3.757062258985477857247991239 + 2.465803785962819679236327339√–1)
(120, 3.741674608179023673159144258 + 2.466291085896660260688606142√–1)
(150, 3.726228649726558590507828429 + 2.466690204011030007962113880√–1)

TABLE 2. $(N, 2\pi \log(J_{N+1}(L)/J_N(L)))$ for the Whitehead link L




FIGURE 7.
Dots indicate $(1/N, 2\pi \operatorname{Re} \log(J_{N+1}(5_2)/J_N(5_2)))$ for $N = 40, 50, 60, 70, 80, 100, 120, 150, 200, 250$. The origin corresponds to $(0, 2.82)$.




FIGURE 8.
Dots indicate $(1/N, 2\pi \operatorname{Im} \log(J_{N+1}(5_2)/J_N(5_2)))$ for $N = 40, 50, 60, 70, 80, 100, 120, 150, 200, 250$. The origin corresponds to $(0, -3.0242)$.




FIGURE 9. Dots indicate $(1/N, 2\pi \operatorname{Re} \log(J_{N+1}(L)/J_N(L)))$ for $N = 40, 50, 60, 70, 80, 100, 120, 150$. The origin corresponds to $(0, 3.66)$.




FIGURE 10. Dots indicate $(1/N, 2\pi \operatorname{Im} \log(J_{N+1}(L)/J_N(L)))$ for $N = 40, 50, 60, 70, 80, 100, 120, 150$. The origin corresponds to $(0, 2.4674)$.

8. CONCLUSION

We have shown the following by direct calculation.

Observation 8.1. Let L be one of the hyperbolic knots 6_3 , 8_9 , and 8_{20} , or the Whitehead link. Following Kashaev’s way, we approximate the colored Jones polynomial $J_N(L)$ of L asymptotically by

$$\int \cdots \int \exp \frac{N\sqrt{-1}}{2\pi} V_L(\mathbf{x}) d\mathbf{x}.$$

Then there exists a stationary point \mathbf{x}_0 of V_L such that the formula

$$\exp \frac{N\sqrt{-1}}{2\pi} V_L(\mathbf{x}_0) = \exp \frac{N}{2\pi} (\operatorname{vol}(L) + \sqrt{-1} \operatorname{CS}(L))$$

holds up to 6 digits.

Conjecture 8.2. (Complexification of Kashaev’s conjecture.) Let L be a hyperbolic link. Then, it holds that

$$\operatorname{vol}(L) = 2\pi \lim_{N \rightarrow \infty} \frac{\log |\langle L \rangle_N|}{N}$$

with $\operatorname{vol}(L)$ the hyperbolic volume of the complement of L . Moreover, there exists the topological Chern–Simons invariant $\operatorname{CS}_{TOP}(L)$ of L

$$\operatorname{CS}_{TOP}(L) = 2\pi \operatorname{Im} \lim_{N \rightarrow \infty} \log \frac{J_{N+1}(L)}{J_N(L)} \pmod{\pi^2},$$

and $\operatorname{CS}_{TOP}(L)$ equals to $\operatorname{CS}(L)$ modulo π^2 . Here $\operatorname{CS}(L)$ is the Chern–Simons invariant of L [Chern and Simons 74, Meyerhoff 86]. Note that the complement of L is a hyperbolic manifold with cusps.

We note that Observation 8.1 also holds for the knots 4_1 , 5_2 and 6_1 by calculating Kashaev’s examples in [Kashaev 97] using MAPLE V and SnapPea.

Therefore we conclude that the complexified Kashaev conjecture is true, up to several digits, up to choices of contours when we change summations into integrals, and up to choices of saddle (stationary) points when we approximate integrals by the saddle point method, for the six hyperbolic knots above and for the Whitehead link.

Note that if the complexified Kashaev conjecture is true then the topological Chern–Simons invariant of a hyperbolic link coincides with its Chern–Simons invariant associated with the hyperbolic metric. Moreover if the volume conjecture is true then the colored Jones polynomial would give both the simplicial volume and the topological Chern–Simons invariant for any knot.

ACKNOWLEDGMENTS

We thank the participants in the meeting “Volume conjecture,” October 1999 and those in the workshop “Recent Progress toward the Volume Conjecture,” March 2000, both of which were held at the International Institute for Advanced Study. The latter was financially supported by the Research Institute for Mathematical Sciences, Kyoto University. We are grateful to both of the institutes.

H. M., J. M. and M. O. express their gratitude to the Graduate School of Mathematics, Kyushu University, where the essential part of this work was carried out in December 1999.

Thanks are also due to S. Kojima for his suggestion of the Chern–Simons invariant, to K. Mimachi for valuable discussions, and to K. Hikami for information on Pari-Gp [Cohen 02] and fitting.

This research is partially supported by Grant-in-Aid for Scientific Research, The Ministry of Education, Science, Sports and Culture.

REFERENCES

- [Chern and Simons 74] S.-S. Chern and J. Simons. "Characteristic forms and geometric invariants." *Ann. of Math.* (2) **99** (1974) 48–69.
- [Cohen 02] H. Cohen. *Pari-Gp: a computer program for number theory*. available at <http://www.parigp-home.de/>.
- [Coulson et al. 00] D. Coulson, O. A. Goodman, C. D Hodgson, and W. D. Neumann. "Computing arithmetic invariants of 3-manifolds." *Experiment. Math.* **9**:1 (2000) 127–152.
- [Kashaev and Tirkkonen 00] R. M. Kashaev and O. Tirkkonen. "A proof of the volume conjecture on torus knots." *Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI)*, 269(Vopr. Kvant. Teor. Polya i Stat. Fiz. 16) (2000) 262–268, 370.
- [Kashaev 95] R. M. Kashaev. "A link invariant from quantum dilogarithm." *Modern Phys. Lett. A* **10**:19 (1995), 1409–1418.
- [Kashaev 97] R. M. Kashaev. "The hyperbolic volume of knots from the quantum dilogarithm." *Lett. Math. Phys.* **39**:3 (1997) 269–275.
- [Kirby and Melvin 91] R. Kirby and P. Melvin. "The 3-manifold invariants of Witten and Reshetikhin–Turaev for $sl(2, \mathbb{C})$." 105: (1991) 473–545.
- [Meyerhoff 86] R. Meyerhoff. "Density of the Chern–Simons invariant for hyperbolic 3-manifolds." In *Low-dimensional topology and Kleinian groups (Coventry/Durham, 1984)*, pp. 217–239, Cambridge Univ. Press, Cambridge, 1986.
- [Murakami and Murakami 01] H. Murakami and J. Murakami. "The colored Jones polynomials and the simplicial volume of a knot." *Acta Math.* **186**:1 (2001) 85–104.
- [Neumann and Zagier 85] W. D. Neumann and D. Zagier. "Volumes of hyperbolic three-manifolds." *Topology* **24**:3 (1985) 307–332.
- [Thurston 99] D. Thurston. *Hyperbolic volume and the Jones polynomial*. Lecture notes, École d'été de Mathématiques 'Invariants de nœuds et de variétés de dimension 3', Institut Fourier - UMR 5582 du CNRS et de l'UJF Grenoble (France) du 21 juin au 9 juillet 1999.
- [Weeks 02] J. Weeks. *SnapPea: a computer program for creating and studying hyperbolic 3-manifolds*. available at <http://www.northnet.org/weeks/index/SnapPea.html>.
- [Yokota 00] Y. Yokota. "On the volume conjecture of hyperbolic knots." In *Knot Theory – dedicated to Professor Kunio Murasugi for his 70th birthday*, M. Sakuma, editor, pp 362–367, March 2000.
- [Yokota 02] Y. Yokota. "On the volume conjecture for hyperbolic knots." preprint, available at www.comp.metro-u.ac.jp/~jojo/volume_conjecture.ps.
- [Yoshida 85] T. Yoshida. "The η -invariant of hyperbolic 3-manifolds." *Invent. Math.* **81**:3 (1985) 473–514.

Hitoshi Murakami, Department of Mathematics, Tokyo Institute of Technology, Oh-okayama, Meguro, Tokyo 152-8551, Japan (starshea@tky3.3web.ne.jp)

Jun Murakami, Department of Mathematical Sciences, School of Science and Engineering, Waseda University, 3-4-1, Ohkubo, Shinjuku-ku, Tokyo, 169-8555 Japan (murakami@waseda.jp)

Miyuki Okamoto, University of the Sacred Heart, Hiroo, Shibuya, Tokyo 150-8938, Japan (miyuki3@hh.ij4u.or.jp)

Toshie Takata, Department of Mathematics, Faculty of Science, Niigata University, Niigata 950-2181, Japan (takata@math.sc.niigata-u.ac.jp)

Yoshiyuki Yokota, Department of Mathematics, Tokyo Metropolitan University, Tokyo 192-0397, Japan (jojo@math.metro-u.ac.jp)

Received April 30, 2001; accepted in revised form October 24, 2002.