

Isometric Composition Operators Between Two Weighted Hardy Spaces

Takahiko Nakazi *

Abstract

We study isometric composition operators C_ϕ between two weighted Hardy spaces $H^2(\nu)$ and $H^2(\mu)$ when ν is a radial measure. The isometric C_ϕ is related to a moment sequence and such a ϕ is studied by the Nevanlinna counting function of ϕ when μ is the normalized Lebesgue measure on the unit circle.

§1. Introduction

Let D be the open unit disc in the complex plane \mathbb{C} . We denote by \mathcal{P} the set of all analytic polynomials and H the set of all analytic functions on D . Let μ be a positive Borel measure on \overline{D} with $\mu(\overline{D}) = 1$. $H^p(\mu)$ denotes the closure of all analytic polynomials in $L^p(d\mu)$ for $0 < p < \infty$. If $d\mu = d\theta/2\pi$, then $H^p(\mu) = H^p$ is the classical Hardy space. If $d\mu = 2rdrd\theta/2\pi$, then $H^p(\mu) = L^p_a$ is the classical Bergman space. H^p and L^p_a can be embedded in H . In this paper, we assume that $H^p(\mu)$ is embedded in H for a general μ . H^∞ denotes the set of all bounded analytic functions on D . We also assume that $H^\infty = H \cap L^\infty(d\mu)$.

For an analytic self map ϕ of D , the composition operator C_ϕ is defined by $(C_\phi f)(z) = f(\phi(z))$ ($z \in D$) for f in H . Throughout this paper, we assume that ν and μ are positive Borel measures on \overline{D} with $\nu(\overline{D}) = \mu(\overline{D}) = 1$. ν is called a radial measure if $d\nu = d\nu_0(r)d\theta/2\pi$ for a positive Borel measure ν_0 on $[0, 1]$. Since $d\theta/2\pi = d\delta_{r=1}d\theta/2\pi$, $d\theta/2\pi$ is a radial measure.

In this paper, we studied isometric composition operators from $H^2(\nu)$ into $H^2(\mu)$ when ν is a radial measure. As we show in the final section, our isometric composition operator C_ϕ is related to an isometric operator T from $H^p(\nu)$ into $H^p(\mu)$ with

*This research was partially supported by Grant-in-Aid for Scientific Research, Japan Society for the Promotion of Science.

$T1 = 1$ when $p \neq 2$. We have a long history for such isometric operators (see [8]). The onto isometries on H^p or L_a^p for $p \neq 2$ were described completely. Unfortunately into isometries have been known very little.

Problem 1. For given measures ν and μ , does there exist an isometric composition operator C_ϕ from $H^2(\nu)$ into $H^2(\mu)$? If there exists such a C_ϕ , describe ϕ .

A function F in $H^2(\mu)$ is called an inner function in $H^2(\mu)$ if

$$\int_{\overline{D}} f|F|^2 d\mu = \int_{\overline{D}} f d\mu \int_{\overline{D}} |F|^2 d\mu \quad (f \in \mathcal{P}).$$

If ϕ^n is an inner function in $H^2(\mu)$ with $\int_{\overline{D}} \phi^n d\mu = 0$ for any $n \geq 0$ then there exists a unique radial measure ν such that C_ϕ is isometric from $H^2(\nu)$ into $H^2(\mu)$ where $d\nu = d\nu_0(r)d\theta/2\pi$ and $1 \in \text{supp } \nu_0$. This is not difficult to prove. However we don't know whether the converse is true.

Problem 2. If a composition operator C_ϕ is isometric from $H^2(\nu)$ into $H^2(\mu)$ then is ϕ^n an inner function in $H^2(\mu)$ with $\int_{\overline{D}} \phi^n d\mu = 0$ for any $n \geq 0$?

A function ϕ in H^∞ with $\|\phi\|_\infty = 1$ is called a Rudin's orthogonal function in $H^2(\mu)$ if $\{\phi^n; n = 0, 1, 2, \dots\}$ is a set of orthogonal functions in $H^2(\mu)$. If ϕ^n is an inner function in $H^2(\mu)$ with $\int_{\overline{D}} \phi^n d\mu = 0$ for any $n \geq 0$ and $\|\phi\|_\infty = 1$ then ϕ is a Rudin's orthogonal function in $H^2(\mu)$ because \mathcal{P} is dense in $H^2(\mu)$ by its definition. We can ask whether the converse is true or not.

Problem 3. If ϕ is a Rudin's orthogonal function in $H^2(\mu)$ then is ϕ^n an inner function in $H^2(\mu)$ with $\int_{\overline{D}} \phi^n d\mu = 0$ for any $n \geq 0$?

When $d\mu = d\theta/2\pi$, Problem 3 was studied by several people, for example, [2],[3],[5],[6] and [10]. C. Bishop [2] and C. Sundberg [10] gave counter examples. Hence there exists a Rudin's orthogonal function which is not an inner function in $H^2(d\theta/2\pi)$.

Problem 3 has a strong connection with Problem 2. In fact, if C_ϕ is an isometric operator from $H^2(\nu)$ into $H^2(\mu)$ then by Theorem 1 ϕ is a Rudin's orthogonal function in $H^2(\mu)$. Conversely if ϕ is a Rudin's orthogonal function in $H^2(\mu)$ then

by Proposition 8 there exists a unique radial measure ν such that C_ϕ is an isometric operator from $H^2(\nu)$ into $H^2(\mu)$.

For each ϕ , we will use two Borel measures μ_ϕ on \bar{D} and $\mu_{|\phi|}$ on $[0,1]$. For a Borel set E in \bar{D} $\mu_\phi(E) = \mu(\{z \in \bar{D}; \phi(z) \in E\})$ and for a Borel set G in $[0,1]$ $\mu_{|\phi|}(G) = \mu(\{z \in \bar{D}; |\phi(z)| \in G\})$.

§2. General case

In this section we assume that ν is a radial measure, μ is an arbitrary measure and ϕ is an analytic selfmap with $\|\phi\|_\infty = 1$. We say that $\{a_n\}$ is a moment sequence of ν_0 , a positive Borel measure on $[0,1]$, if $a_n = \int_0^1 r^n d\nu_0$ ($n = 0, 1, 2, \dots$).

Theorem 1. *Suppose $d\nu = d\nu_0(r)d\theta/2\pi$. Then C_ϕ is an isometric operator from $H^2(\nu)$ into $H^2(\mu)$ if and only if $\int_{\bar{D}} \phi^n \bar{\phi}^m d\mu = 0$ ($n \neq m$) and $\left\{ \int_{\bar{D}} |\phi|^n d\mu \right\}$ is a moment sequence of ν_0 .*

Proof. If C_ϕ is isometric, by the polarization formula

$$\delta_{nm} \int_0^1 r^n r^m d\nu_0(r) = \int_{\bar{D}} z^n \bar{z}^m d\nu = \int_{\bar{D}} \phi^n \bar{\phi}^m d\mu$$

because ν is a radial measure. Hence

$$\int_{\bar{D}} |\phi|^{2n} d\mu = \int_0^1 r^{2n} d\nu_0 \quad (n = 0, 1, 2, \dots).$$

It is elementary to see that $x = \sqrt{1 - (1 - x^2)} = \sum_{n=0}^{\infty} a_n (1 - x^2)^n$ and $\sum_{n=0}^{\infty} |a_n| (1 - x^2)^n < \infty$ ($0 \leq x \leq 1$). Hence by Lebesgue's dominated convergence theorem

$$\begin{aligned} \int_{\bar{D}} |\phi| d\mu &= \int_{\bar{D}} \sum_{n=0}^{\infty} a_n (1 - |\phi|^2)^n d\mu = \sum_{n=0}^{\infty} a_n \int_{\bar{D}} (1 - |\phi|^2)^n d\mu \\ &= \sum_{n=0}^{\infty} a_n \int_0^1 (1 - r^2)^n d\nu_0 = \int_0^1 \sum_{n=0}^{\infty} a_n (1 - r^2)^n d\nu_0 = \int_0^1 r d\nu_0 \end{aligned}$$

because $\left| \sum_{n=0}^k a_n (1 - |\phi|^2)^n \right| \leq \sum_{n=0}^{\infty} |a_n|$ and $\left| \sum_{n=0}^k a_n (1 - r^2)^n \right| \leq \sum_{n=0}^{\infty} |a_n| < \infty$. Similarly, as $x^{2\ell+1} = \sqrt{1 - (1 - x^{4\ell+2})}$ we can show that $\int_{\bar{D}} |\phi|^{2n+1} d\mu = \int_0^1 r^{2n+1} d\nu_0$ ($n =$

$0, 1, 2, \dots$). Thus $\left\{ \int_{\overline{D}} |\phi|^n d\mu \right\}$ is a moment sequence of ν_0 .

Conversely if $\int_{\overline{D}} \phi^n \bar{\phi}^m d\mu = 0$ ($n \neq m$) and $\left\{ \int_{\overline{D}} |\phi|^n d\mu \right\}$ is a moment sequence of ν_0 , then

$$\begin{aligned} \int_{\overline{D}} \left| \sum_{n=0}^k a_n \phi^n \right|^2 d\mu &= \sum_{n=0}^k |a_n|^2 \int_{\overline{D}} |\phi|^{2n} d\mu \\ &= \sum_{n=0}^k |a_n|^2 \int_0^1 r^{2n} d\nu_0 = \int_{\overline{D}} \left| \sum_{n=0}^k a_n z^n \right|^2 d\nu. \end{aligned}$$

Hence C_ϕ is isometric. \square

Theorem 2. *If $d\nu = d\nu_0(r)d\theta/2\pi$ then the following conditions are equivalent.*

(1) C_ϕ is an isometric operator from $H^2(\nu)$ into $H^2(\mu)$.

(2) $\nu_0 = \mu|_\phi$

(3) $\int_0^1 F(r) d\nu_0 = \int_{\overline{D}} F(|\phi|) d\mu$ for any Borel nonnegative function F on $[0, 1]$.

Proof. (1) \Rightarrow (2) If G is a Borel set in $[0, 1]$, then $\nu_0(G) = \inf\{\nu_0(V) ; G \subset V, V \text{ is open in } [0, 1]\}$ because ν_0 is a Borel measure. Hence there exists a sequence of continuous functions $\{f_m\}$ such that $f_m \rightarrow \chi_G$ a.e. ν_0 on $[0, 1]$ and $\|f_m\|_\infty \leq \gamma < \infty$ ($m = 1, 2, \dots$). By the Stone-Weierstrass theorem,

$$\int_0^1 f_m(r) d\nu_0 = \int_{\overline{D}} f_m(|\phi|) d\mu \quad (m = 1, 2, \dots)$$

because $\int_0^1 r^n d\nu_0 = \int_{\overline{D}} |\phi|^n d\mu$ ($n = 0, 1, 2, \dots$). Thus $\nu_0(G) = \mu(\{z \in \overline{D} ; |\phi(z)| \in G\})$. (2) \Rightarrow (3) is clear. (3) \Rightarrow (1) is a result of Theorem 1. \square

The following theorem shows that we can solve Problem 2 in the Introduction when C_ϕ is onto.

Theorem 3. *Suppose $d\nu = d\nu_0(r)d\theta/2\pi$. If C_ϕ is an isometric operator from $H^2(\nu)$ onto $H^2(\mu)$ then ϕ^n is an inner function in $H^2(\mu)$ for any $n \geq 0$.*

Proof. Let $F \in \mathcal{P}$ then there exists $f \in H^2(\nu)$ such that $F = f \circ \phi$. Let $f =$

$\sum_{j=0}^{\infty} a_j z^j$, since $\sum_{j=0}^{\infty} |a_j|^2 \int_0^1 r^{2j} d\nu_0(r) < \infty$, $F = \sum_{j=0}^{\infty} a_j \phi^j$ and $\sum_{j=0}^{\infty} |a_j|^2 \int_{\bar{D}} |\phi|^{2j} d\mu < \infty$.

By Theorem 1, for any $\ell \geq 0$

$$\int_{\bar{D}} F |\phi|^{2\ell} d\mu = a_0 \int_{\bar{D}} |\phi|^{2\ell} d\mu = \int_{\bar{D}} F d\mu \int_{\bar{D}} |\phi|^{2\ell} d\mu$$

because $\int_{\bar{D}} \phi d\mu = 0$. This implies that ϕ^ℓ is an inner function in $H^2(\mu)$ for any $\ell \geq 0$.

When $d\nu = d\nu_0(r)d\theta/2\pi$, if C_ϕ is an isometric operator from $H^2(\nu)$ into $H^2(\mu)$ then C_z is isometric from $H^2(\nu)$ onto $H^2(\mu_\phi)$.

Corollary 1. *Suppose $d\nu = d\nu_0(r)d\theta/2\pi$. If C_z is an isometric operator then z^n is an inner function in $H^2(\mu)$ for any $n \geq 0$. Moreover $d\mu = d\nu_1(r)d\theta/2\pi + d\delta_{r=0}d\mu_1(\theta)$, where ν_1 is a Borel measure on $[0,1]$ and μ_1 is a Borel measure on ∂D . If ν_0 does not have point mass on $\{r=0\}$ then $\nu = \mu$.*

Proof. By the remark above, C_z is isometric from $H^2(\nu)$ onto $H^2(\mu)$ because $\mu_z = \mu$. By Theorem 3, z^n is inner in $H^2(\mu)$ for any $n \geq 0$. Put $C_0[0,1] = \{u; u \text{ is continuous on } [0,1] \text{ and } u(0) = 0\}$ and $C_0(\partial D) = \{f; f \text{ is continuous on } \partial D \text{ and } f(1) = 0\}$. Since $r^n d\mu$ annihilates $z\mathcal{P} + \bar{z}\bar{\mathcal{P}}$ for any $n \geq 0$, for any $j \neq 0$, $d\mu \perp \{r^{2n+|j|}e^{ij\theta}; n = 0, 1, 2, \dots\}$. By the Müntz-Szasz theorem [6], $d\mu \perp C_0[0,1]e^{ij\theta}$ for any $j \neq 0$ and so $d\mu \perp C_0[0,1] \otimes C_0(\partial D)$. This implies that $d\mu = d\nu_1(r)d\theta/2\pi + d\delta_{r=0}d\mu_1(\theta)$ where ν_1 is a Borel measure on $[0,1]$ and μ_1 is a Borel measure on T . If ν_0 does not have point mass on $\{r=0\}$ then we may assume that $\mu_1 = 0$ and so $d\mu = d\nu_1(r)d\theta/2\pi$. By Theorem 2 $\nu_0 = \mu_{|z|}$ and $\mu_{|z|} = \nu_1$ because $d\mu = d\nu_1(r)d\theta/2\pi$. \square

§3. Radial measure

In this section we assume that ν and μ are radial measures, that is, $d\nu = d\nu_0(r)d\theta/2\pi$ and $d\mu = d\mu_0(r)d\theta/2\pi$. Proposition 1 solves Problem 2 when $\nu = \mu$. By Theorem 2, if C_ϕ is isometric from $H^2(\nu)$ into $H^2(\mu)$, then for some positive integer k

$$\int_0^1 \log r d\nu_0 \leq k \int_0^1 \log r d\mu_0 + \int_0^{2\pi} \log |\phi(e^{i\theta})| d\theta$$

as $F(t) = \log t$, using the inner outer factorization of ϕ . Proposition 2 gives an exact formula for this.

Proposition 1. Suppose ν is a radial measure. If C_ϕ is an isometric operator from $H^2(\nu)$ into $H^2(\nu)$, then ϕ^n is an inner function in $H^2(\nu)$ for any $n \geq 0$.

Proof. By Theorem 1, $\phi(0) = 0$ because ν is a radial measure and so by Schwarz's lemma, $|\phi(z)| \leq |z|$ ($z \in D$). Since $\int_{\overline{D}} |\phi(z)|^2 d\nu = \int_{\overline{D}} |z|^2 d\nu$, $|\phi(z)| = |z|$ a.e. ν . For $f \in \mathcal{P}$,

$$\int_{\overline{D}} f |\phi|^{2n} d\nu = \int_{\overline{D}} f |z|^{2n} d\nu = f(0) \int_0^1 r^{2n} d\nu_0 = \int_{\overline{D}} f d\nu \int_{\overline{D}} |\phi|^{2n} d\nu. \quad \square$$

Proposition 2. Suppose ν and μ are radial measures, that is, $d\nu = d\nu_0(r)d\theta/2\pi$ and $d\mu = d\mu_0(r)d\theta/2\pi$. Let $\phi = z^k BQh$ where k is a positive integer, B is a Blaschke product with $B(0) \neq 0$, $Q(z) = \exp - \int_0^{2\pi} \frac{e^{it} + z}{e^{it} - z} d\lambda(t)$ is a singular inner function and h is an outer function. If C_ϕ is an isometric operator from $H^2(\nu)$ into $H^2(\mu)$, then

$$\begin{aligned} \int_0^1 \log r d\nu_0 &= k \int_0^1 \log r d\mu_0 + \int_0^1 d\mu_0 \int_0^r n(s, B) \frac{ds}{s} + \\ &\quad \log |B(0)| - \mu_0([0, 1))\lambda([0, 2\pi]) + \int_0^{2\pi} \log |\phi(e^{i\theta})| d\theta/2\pi \end{aligned}$$

where $n(s, B)$ is the number of zeros of B on the closed disc $\{z \in \mathbb{C}; |z| \leq r\}$.

Proof. Let $n(s, B) = n(s, BQh)$ is the number of zeros of BQh on the closed disc $\{z \in \mathbb{C}; |z| \leq r\}$. Then, by Theorem 2 and [1, §2 of Chapter 5]

$$\begin{aligned} &\int_0^1 \log r d\nu_0 \\ &= \int_0^{1-} d\mu_0 \int_0^{2\pi} \log |\phi(re^{i\theta})| d\theta/2\pi + \mu_0(\{1\}) \int_0^{2\pi} \log |\phi(e^{i\theta})| d\theta/2\pi \\ &= \int_0^{1-} d\mu_0 \left\{ \log r^k + \int_0^r n(s, B) \frac{ds}{s} \right\} + \mu_0([0, 1)) \log |B(0)Q(0)h(0)| \\ &\quad + \mu_0(\{1\}) \int_0^{2\pi} \log |\phi(e^{i\theta})| d\theta/2\pi \\ &= k \int_0^1 \log r d\mu_0 + \int_0^1 d\mu_0 \int_0^r n(s, B) \frac{ds}{s} + \log |B(0)| \\ &\quad - \mu_0([0, 1))\lambda([0, 2\pi]) + \int_0^{2\pi} \log |\phi(e^{i\theta})| d\theta/2\pi \end{aligned}$$

because $\mu_0(\{1\}) \int_0^1 n(s, B) \frac{ds}{s} = -\mu_0(\{1\}) \log |B(0)|$. \square

§4. Special cases

In this section we assume that ν or μ is the normalized Lebesgue measure or the normalized area measure. Proposition 3 solves Problems 1 and 2 when ν is the normalized Lebesgue measure on the circle and μ is a radial measure. Proposition 5 solves Problem 2 when ν is a radial measure or the Lebesgue measure on the circle. Corollary 3 solves Problem 2 negatively when $d\nu = 2rdrd\theta/2\pi$ and $d\mu = d\theta/2\pi$.

Proposition 3. *Let μ be a radial measure. C_ϕ is an isometric operator from H^2 into $H^2(\mu)$ if and only if ϕ^n is an inner function with $\int_{\overline{D}} \phi d\mu = 0$ in $H^2(\mu)$ for any $n \geq 1$ and $H^2(d\mu) = H^2$.*

Proof. If C_ϕ is isometric, by Theorem 1 $\int_{\overline{D}} \phi^n \bar{\phi}^m d\mu = 0$ ($n \neq m$) and we have

$$1 = \int_0^{2\pi} |z|^2 d\theta/2\pi = \int_{\overline{D}} |\phi|^2 d\mu \leq 1.$$

Hence $|\phi(z)| = 1$ a.e. μ and so $\text{supp } \mu \subset \partial D$. This implies that $d\mu = d\delta_{r=1}d\theta/2\pi$ because μ is a radial measure. The converse is clear. \square

Proposition 4. *Suppose $d\nu = d\nu_0(r)d\theta/2\pi$ and C_ϕ is an isometric operator from $H^2(\nu)$ into H^2 .*

(1) $\nu_0(\{a\}) > 0$ for $0 \leq a \leq 1$ if and only if $d\theta/2\pi(\{e^{i\theta}; |\phi(e^{i\theta})| = a\}) > 0$.

(2) $d\nu_0 = d\delta_{r=1}$ if and only if ϕ is an inner function in H^2 .

(3) ν_0 is a discrete measure if and only if $|\phi| = \sum_{n=1}^{\infty} a_n \chi_{E_n}$ where $0 \leq a_n \leq 1$, and $d\theta/2\pi(E_n) = \nu_0(\{a_n\})$ ($n = 1, 2, \dots$).

Proof. Since $\nu_0(G) = d\theta/2\pi(\{e^{i\theta}; |\phi(e^{i\theta})| \in G\})$ for a Borel set G in $[0,1]$ by Theorem 2, it is easy to see. \square

Proof. This is just (2) of Proposition 4.

Now we consider when $d\nu = r dr d\theta/\pi$ or $d\mu = r dr d\theta/\pi$.

Proposition 5. *If C_ϕ is an isometric operator from L_a^2 into $H^2(\mu)$, then $\mu(\{z \in \overline{D}; |\phi| = b\}) = 0$ and $\int_{\overline{D}} (b - |\phi|)^{-1} d\mu = \infty$ for any $0 \leq b \leq 1$.*

Proof. It is clear by Theorem 2. \square

Corollary 2. *If C_ϕ is an isometric operator from L_a^2 into H^2 , then ϕ is not inner in H^2 .*

Proposition 6. *Suppose $d\nu = d\nu_0(r)d\theta/2\pi$. If C_ϕ is an isometric operator from $H^2(\nu)$ into L_a^2 , then $\int_0^1 \log r d\nu_0 = -\frac{k}{4} + \int_0^1 2r dr \int_0^r n(s, B) \frac{ds}{s} + \log |B(0)| - \lambda([0, 2\pi]) + \int_0^{2\pi} \log |\phi(e^{i\theta})| d\theta/2\pi$, where the inner part of ϕ is $z^k BQ$, B is a Blaschke product with $B(0) \neq 0$, $Q(z) = \exp - \int_0^{2\pi} \frac{e^{it} + z}{e^{it} - z} d\lambda$ is a singular inner function. Hence if ϕ is a strictly function, then $\int_0^1 \log r d\nu_0 = -\frac{1}{4} + \int_0^{2\pi} \log |\phi(e^{i\theta})| d\theta/2\pi$.*

Proof. It is clear by Proposition 2. \square

§5. Nevanlinna counting function

Suppose ν or μ is the normalized Lebesgue measure or the normalized area measure. We assume that ϕ is a non-constant function in H^∞ with $\|\phi\|_\infty = 1$. The Nevanlinna counting function of ϕ , N_ϕ , is defined on $D \setminus \{\phi(0)\}$ by

$$N_\phi(w) = \sum_{\phi(z)=w} \log \frac{1}{|z|},$$

where multiplicities are counted and $N_\phi(w)$ is taken to be zero if w is not in the range of ϕ . Corollary 4 seems to be interesting in spite of Corollary 3.

Theorem 4. *Suppose $d\nu = d\nu_0(r)d\theta/2\pi$. Then, C_ϕ is an isometric operator*

from $H^2(\nu)$ into H^2 if and only if

$$N_\phi(z) = \int_{|z|}^1 \log \frac{r}{|z|} d\nu_0(r)$$

for nearly all z in D .

Proof. The 'only if' part was proved in [6, Lemma 3]. If $N_\phi(z) = \int_{|z|}^1 \log \frac{r}{|z|} d\nu_0(r)$ for nearly all z in D , by the Littlewood-Paley theorem (see [3]),

$$\begin{aligned} & \int_0^{2\pi} \phi^n(e^{i\theta}) \bar{\phi}^m(e^{i\theta}) d\theta / 2\pi \\ &= 2nm \int_D z^{n-1} \bar{z}^{m-1} N_\phi(|z|) dA(z) \\ &= 4nm \delta_{nm} \int_0^1 r^{n+m-1} \left(\int_r^1 \log \frac{s}{r} d\nu_0(s) \right) dr \\ &= 4nm \delta_{nm} \int_0^1 d\nu_0(s) \int_0^s r^{m+n-1} \left(\log \frac{s}{r} \right) dr \\ &= \frac{4nm}{(n+m)^2} \delta_{nm} \int_0^1 s^{n+m} d\nu_0(s). \end{aligned}$$

When $n = m$, $\int_0^{2\pi} |\phi(e^{i\theta})|^{2n} d\theta / 2\pi = \int_0^1 s^{2n} d\nu_0(s)$ for $n = 0, 1, 2, \dots$. Hence by Theorem 1 and its proof, C_ϕ is an isometric operator from $H^2(\nu)$ into H^2 .

Lemma. $D \setminus \{z \in D ; \phi'(z) = 0\}$ can be decomposed into an at most countable disjoint collection $\{R_n\}$ of "semi-closed" polar rectangles, on each of which ϕ is schricht.

Proof. It is known in [9, p186]. \square

Corollary 3. Suppose ϕ is a finite-to-one map. Then C_ϕ is not an isometric operator from L_a^2 into H^2 .

Proof. By Lemma, there exists the inverse ψ_n of the restriction of ϕ to R_n . Let $w \in \phi(R_{j_1})$. If ϕ is an ℓ to 1 map, then there exist j_2, \dots, j_ℓ such that $\psi_{j_1}(z) = \psi_{j_2}(z) = \dots = \psi_{j_\ell}(z) = w$. Hence there exists a small disc Δ in $\phi(R_{j_1})$ such that

$$N_\phi(z) = \sum_{z=\phi(w)} \log \frac{1}{|w|} = \sum_{i=1}^{\ell} \log \frac{1}{|\psi_{j_i}(z)|}$$

for all $w \in \Delta$. Therefore there exists a subdisc Δ_0 in Δ such that $N_\phi(z)$ is harmonic on Δ_0 . On the other hand, by Proposition 5

$$N_\phi(z) = 2 \int_{|z|}^1 \left(\log \frac{r}{|z|} \right) r dr = \frac{|z|^2 - 1}{2} + \log \frac{1}{|z|}.$$

This contradicts that $N_\phi(z)$ is harmonic on Δ_0 . \square

Theorem 5. *Suppose ϕ is a contractive function in H^∞ such that ϕ is a finite-to-one map and $|\phi| = \sum_{j=1}^{\ell} a_j \chi_{E_j}$ where $0 < a_j < a_{j+1}$, $\sum_{j=1}^{\ell} \chi_{E_j} = 1$ and E_j is a measurable set in ∂D where $1 \leq \ell \leq \infty$. If the inner part of $z - \phi$ is a Blaschke product for each $z \in D$, then C_ϕ is not an isometric operator from $H^2(\nu)$ into H^2 for any $d\nu = d\nu_0(r)d\theta/2\pi$ if $\ell \neq 1$.*

Proof. Suppose C_ϕ is an isometric operator from $H^2(\nu)$ into H^2 for some $d\nu = d\nu_0(r)d\theta/2\pi$. By Proposition 4, ν_0 is a discrete measure and $d\theta/2\pi(E_j) = \nu_0(\{a_j\})$ ($j = 1, 2, \dots$). Since $\phi(0) = 0$, by Lemma 2 in [6] and Proposition 7

$$\begin{aligned} N_\phi(z) &= \int_0^{2\pi} \log |z - \phi(e^{i\theta})| d\theta/2\pi + \log \frac{1}{|z|} \\ &= \int_{|z|}^1 \log \frac{r}{|z|} d\nu_0(r) \end{aligned}$$

for $z \in D \setminus \{0\}$. If $|z| \leq a_1$, then

$$\begin{aligned} \int_{|z|}^1 \log \frac{r}{|z|} d\nu_0(r) &= \sum_{j=1}^{\infty} \left(\log \frac{a_j}{|z|} \right) \nu_0(\{a_j\}) \\ &= \sum_{j=1}^{\infty} \nu_0(\{a_j\}) \log \frac{1}{|z|} + \sum_{j=1}^{\infty} \nu_0(\{a_j\}) \log a_j \\ &= \log \frac{1}{|z|} + \sum_{j=1}^{\infty} \nu_0(\{a_j\}) \log a_j. \end{aligned}$$

Hence if $|z| \leq a_1$ then

$$\int_0^{2\pi} \log |z - \phi(e^{i\theta})| d\theta/2\pi = \sum_{j=1}^{\infty} \nu_0(\{a_j\}) \log a_j = \alpha.$$

If $a_1 < |z| \leq a_2$, then

$$\begin{aligned} \int_{|z|}^1 \log \frac{r}{|z|} d\nu_0(r) &= \sum_{j=2}^{\infty} \left(\log \frac{a_j}{|z|} \right) \nu_0(\{a_j\}) \\ &= \sum_{j=2}^{\infty} \nu_0(\{a_j\}) \log \frac{1}{|z|} + \sum_{j=2}^{\infty} \nu_0(\{a_j\}) \log a_j \end{aligned}$$

$$\begin{aligned} \int_0^{2\pi} \log |z - \phi(e^{i\theta})| d\theta / 2\pi &= -\nu_0(\{a_1\}) \log \frac{1}{|z|} + \sum_{j=2}^{\infty} \nu_0(\{a_j\}) \log a_j \\ &= \beta \log \frac{1}{|z|} + \gamma. \end{aligned}$$

where $\beta \neq 0$

For each $z \in D$, put

$$z - \phi(\zeta) = q_z(\zeta) h_z(\zeta) \quad (\zeta \in D)$$

where $q_z(\zeta)$ is inner and $h_z(\zeta)$ is outer. Since ϕ is a finite-to-one map, q_z is a finite Blaschke product by hypothesis and so

$$q_{\phi(t)}(\zeta) = \prod_{j=1}^n \frac{\zeta - b_j(t)}{1 - \overline{b_j(t)}\zeta} \quad (t \in D).$$

Then, since $\phi(0) = 0$,

$$\phi(t) = (-1)^n \left(\prod_{j=1}^n b_j(t) \right) h_{\phi(t)}(0) \quad (t \in D).$$

Put $D_r = \{t \in \mathbb{C}; |t| \leq r\}$ for $0 < r < 1$. If both ϕ and ϕ' have no zeros on ∂D_r then there is a division $\{D_r^j\}_{1 \leq j \leq n}$ of D_r such that ϕ is one-to-one on D_r^j for $1 \leq j \leq n$. For, ϕ is conformal in a neighborhood of each point on ∂D_r and so $\arg \phi$ is increasing on, ∂D_r . Put $\phi_j = \phi | D_r^j$ and $b_j(t) = \phi_j^{-1}(\phi(t))$ for $1 \leq j \leq n$. Then $b_j(t)$ is analytic except $\phi'(t) = 0$ when $\phi(t)$ in $\phi(D_r)$. Hence $h_{\phi(t)}(0)$ is analytic except $\phi'(t) = 0$ and $\bigcup_{j=1}^n \{t \in D; b_j(t) = 0\}$ when $\phi(t)$ in $\phi(D_r)$. Since $\phi(0) = 0$, $\{t \in D; |\phi(t)| < a_1\}$ is a nonempty open set. We can choose r such that $\{t \in D; |\phi(t)| < a_1\} \cap \phi(D_r) \neq \emptyset$. If $|\phi(t)| \leq a_1$, by what was proved above,

$$\begin{aligned} \alpha &= \int_0^{2\pi} \log |\phi(t) - \phi(e^{i\theta})| d\theta / 2\pi \\ &= \int_0^{2\pi} \log |h_{\phi(t)}(e^{i\theta})| d\theta / 2\pi = \log |h_{\phi(t)}(0)|. \end{aligned}$$

Hence $|h_{\phi(t)}(0)| = e^\alpha$. and so $h_{\phi(t)}(0)$ is constant on D_r . If $a_1 < |\phi(t)| \leq a_2$, by what was proved above,

$$\begin{aligned} \beta \log \frac{1}{|\phi(t)|} + \gamma &= \int_0^{2\pi} \log |\phi(t) - \phi(e^{i\theta})| d\theta / 2\pi \\ &= \int_0^{2\pi} \log |h_{\phi(t)}(e^{i\theta})| d\theta / 2\pi = \log |h_{\phi(t)}(0)| \end{aligned}$$

and so $|h_{\phi(t)}(0)| = e^\gamma |\phi(t)|^\beta$. Since there exists $0 < r < 1$ such that $\{t \in D; a_1 < |\phi(t)| < a_2\} \cap \phi(D_r) \neq \emptyset$, this implies that $|\phi(t)|$ is constant there and so ϕ is constant on D . This contradicts that ϕ is a finite-to-one map. Therefore C_ϕ is not isometric. \square

If ϕ is a one-to-one map then it is known [4, Theorem 3.17] that the inner part of $z - \phi$ is a Blaschke product for each $z \in D$. Hence we need not such a hypothesis in Theorem 5. Unfortunately we could not prove it in general, that is, for a finite-to-one map.

§6. Rudin's orthogonal function

In this section, we study Rudin's orthogonal functions. By Theorem 1, if C_ϕ is an isometric operator from $H^2(\nu)$ into $H^2(\mu)$ then ϕ is a Rudin's orthogonal function. Proposition 7 implies the converse. This was proved by the author [6] when $d\mu = d\theta/2\pi$. The proof is valid for an arbitrary μ . However we give a new proof due to K. Izuchi.

Proposition 7. *If ϕ is a Rudin's orthogonal function in $H^2(\mu)$ then there exists a unique radial measure ν such that C_ϕ is an isometric operator from $H^2(\nu)$ into $H^2(\mu)$ where $d\nu = d\nu_0(r)d\theta/2\pi$ and $1 \in \text{supp } \nu_0$.*

Proof. Put $\nu_0 = \mu_{|\phi|}$ and $d\nu = d\nu_0 d\theta/2\pi$, then Theorems 1 and 2 imply the proposition. \square

Corollary 4. *Suppose ϕ is a finite-to-one map and ϕ is a Rudin's orthogonal function. If the inner part of $z - \phi$ is a Blaschke product for each $z \in D$ and $|\phi| = \sum_{j=1}^{\ell} a_j \chi_{E_j}$ where $0 \leq a_j < a_{j+1}$, $\sum_{j=1}^{\ell} \chi_{E_j} = 1$ and E_j is a measurable set in ∂D where $1 \leq \ell \leq \infty$, then $|\phi| = 1$ and so ϕ is a finite Blaschke product.*

Proof. If ϕ is a Rudin's orthogonal function, then by Proposition 7 and Theorem 5, $\ell = 1$ and so ϕ is a finite Blaschke product. \square

In Corollary 4, if ϕ is one-to-one map then the inner part of $z - \phi$ is a Blaschke product (see [4.Theorem 3.17]). Hence we can take off such a condition. However in such a case Corollary 4 is not new. In fact, P. S. Bourdon [3] showed that if ϕ is univalent and a Rudin's orthogonal function then ϕ is just the coordinate function z .

§7. Final remark

The research in this paper gives more general one. Suppose $0 < p < \infty$ and $p \neq 2$. T is an isometric operator from $H^p(\nu)$ into $H^p(\mu)$ with $T1 = 1$ if and only if $T = C_\phi$ for some ϕ in H^∞ with $\|\phi\|_\infty = 1$ and C_ϕ is an isometric operator from $H^p(\nu)$ into $H^p(\mu)$. For the 'if' part is trivial. For the 'only if' part, if T is isometric and $T1 = 1$, then by [5, Theorem 7.5.3] $T(fg) = Tf \cdot Tg$ a.e. μ and $\|Tf\|_\infty = \|f\|_\infty$ for all $f \in \mathcal{P}$, $g \in \mathcal{P}$. Hence if $\phi = Tz$ then ϕ belongs to H^∞ and $\|\phi\|_\infty = 1$. Therefore $Tf = C_\phi f$ ($f \in \mathcal{P}$) and so $Tf = C_\phi f$ ($f \in H^p(\nu)$). When $p \neq 2$, if C_ϕ is an isometric operator from $H^p(\nu)$ into $H^p(\mu)$, then C_ϕ is an isometric operator from $H^2(\nu)$ into $H^2(\mu)$. For by [5, Theorem 8.5.3], for all $f \in \mathcal{P}$ and $g \in \mathcal{P}$

$$\int_{\overline{D}} C_\phi f \cdot \overline{C_\phi g} d\mu = \int_{\overline{D}} f \overline{g} d\mu$$

and $\|C_\phi f\|_\infty = \|f\|_\infty$. This implies that C_ϕ is an isometric operator from $H^2(\nu)$ into $H^2(\mu)$.

We give two open problems :

(1) Are there any isometric C_ϕ from L_a^2 into H^2 ?

(2) When ν_0 is a discrete measure and not a dirac measure, are there any isometric C_ϕ from $H^2(\nu)$ to H^2 where $d\nu = d\nu_0(r)d\theta/2\pi$?

Acknowledgement. The author wishes to express his sincere gratitude to the referee for his many helpful suggestions and advices. In particular, he formulates and improves the original Theorem 2 to the present one.

References

1. M. Anderson, Topics in Complex Analysis, Universetext : Tracts in Mathematics, Springer.
2. C. Bishop, Orthogonal functions in H^p , preprint.
3. P. S. Bourdon, Rudin's orthogonality problem and the Nevanlinna counting function, Proc. Amer. Math. Soc. 125(1997), 1187-1192.
4. P. L. Duren, Theory of H^p Spaces, Academic Press, New York, 1970.
5. T. Nakazi and T. Watanabe, Properties of a Rudin's orthogonal function which is a linear combination of two inner functions, Scientia Mathematica Japonica Online, 7(2002), 347-352.
6. T. Nakazi, The Nevanlinna counting functions for Rudin's orthogonal functions, Proc. Amer. Math. Soc. 131(2003), 1267-1271.
7. W. Rudin, Real Complex Analysis, Third Edition, McGraw-Hill Series in Higher Mathematics
8. W. Rudin, Function Theory in the Unit Ball of \mathbb{C}^n , Springer-Verlag, 1980
9. J. H. Shapiro, Composition Operators and Classical Function Theory, Springer-Verlag, 1993
10. C. Sundberg, Measures induced by analytic functions and a problem of Walter Rudin, preprint.

Department of Mathematics
Hokkaido University
Sapporo 060-0810, Japan

nakazi@math.sci.hokudai.ac.jp

Received July 28, 2005

Revised July 20, 2006