

## A DETERMINANT FORMULA FOR A CLASS OF RATIONAL SOLUTIONS OF PAINLEVÉ V EQUATION

TETSU MASUDA, YASUHIRO OHTA AND KENJI KAJIWARA

**Abstract.** We give an explicit determinant formula for a class of rational solutions of the Painlevé V equation in terms of the universal characters.

### §1. Introduction and main result

It is known that six Painlevé equations are in general irreducible, namely, their solutions cannot be expressed by “classical functions” in the sense of Umemura [18]. However, it is also known that they admit classical solutions for special values of parameters except for  $P_I$ . Much effort have been made for the investigation of classical solutions. As a result, it has been recognized that there are two classes of classical solutions. One is transcendental classical solutions expressible in terms of functions of hypergeometric type. Another one is algebraic or rational solutions. It is also known that the Painlevé equations (except for  $P_I$ ) admit action of the affine Weyl groups as groups of the Bäcklund transformations. It is remarkable that such classical solutions are located on special places from a point of view of symmetry in the parameter spaces [12], [13], [14], [15]. For example,  $P_{II}$ ,  $P_{III}$  and  $P_{IV}$ , whose symmetry is described by the affine Weyl group of type  $A_1^{(1)}$ ,  $A_1^{(1)} \oplus A_1^{(1)}$  and  $A_2^{(1)}$ , respectively, admit transcendental classical solutions on the reflection hyperplanes, and rational solutions on the barycenters of Weyl chambers of the corresponding affine Weyl group.

Umemura et al have investigated the class of solutions on the barycenters of Weyl chambers and found that (1) these solutions are expressed by some characteristic polynomials generated by the Toda type bilinear equations, (2) the coefficients of such polynomials admit mysterious combinatorial properties [17], [7], [16]. These special polynomials are sometimes referred as *Yablonskii-Vorob'ev polynomials* for  $P_{II}$  [19], *Okamoto polynomials* for  $P_{IV}$  [14], *Umemura polynomials* for  $P_{III}$ ,  $P_V$  and  $P_{VI}$ .

---

Received February 2, 2001.

2000 Mathematics Subject Classification: 33E17; 34M55.

One important aspect among such polynomials is that they are expressed as special cases of the Schur functions. As is well known, the Schur functions are characters of the irreducible polynomial representations of  $GL(n)$  and arise as  $\tau$ -functions of the KP hierarchy [1]. For example, it is known that the special polynomials for  $P_{II}$  and  $P_{III}$  are expressible by 2-reduced Schur functions, and those for  $P_{IV}$  by 3-reduced Schur functions [2], [3], [4], [8].

In this paper, we consider  $P_V$ ,

$$(1.1) \quad \frac{d^2y}{dt^2} = \left( \frac{1}{2y} + \frac{1}{y-1} \right) \left( \frac{dy}{dt} \right)^2 - \frac{1}{t} \frac{dy}{dt} + \frac{(y-1)^2}{2t^2} \left( \kappa_\infty^2 y - \frac{\kappa_0^2}{y} \right) - (\theta+1) \frac{y}{t} - \frac{y(y+1)}{2(y-1)},$$

with parameters  $\kappa_\infty$ ,  $\kappa_0$  and  $\theta$ , whose symmetry is described by the affine Weyl group  $W(A_3^{(1)})$ . The aim of this paper is to investigate a class of rational solutions on the barycenters of Weyl chambers and to present an explicit formula for them.

By the analogy from the known cases, it is naively expected that they are expressed in terms of 4-reduced Schur functions. However, our formula is expressed by a generalization of Schur functions, which is called the *universal characters* and defined as follows [6].

DEFINITION 1. Let  $p_k = p_k(t^{(1)})$  and  $q_k = q_k(t^{(2)})$ ,  $k \in \mathbb{Z}$ , be two families of polynomials defined by

$$(1.2) \quad \begin{aligned} \sum_{k=0}^{\infty} p_k \eta^k &= \exp \left( \sum_{j=1}^{\infty} t_j^{(1)} \eta^j \right), & p_k &= 0 \text{ for } k < 0, \\ \sum_{k=0}^{\infty} q_k \eta^k &= \exp \left( \sum_{j=1}^{\infty} t_j^{(2)} \eta^j \right), & q_k &= 0 \text{ for } k < 0, \end{aligned}$$

where  $t^{(1)} = (t_1^{(1)}, t_2^{(1)}, \dots)$  and  $t^{(2)} = (t_1^{(2)}, t_2^{(2)}, \dots)$  are the sets of infinite numbers of variables. For any partitions  $\lambda^{(1)} = (\lambda_1^{(1)}, \lambda_2^{(1)}, \dots, \lambda_n^{(1)})$  and  $\lambda^{(2)} = (\lambda_1^{(2)}, \lambda_2^{(2)}, \dots, \lambda_m^{(2)})$ , the universal character  $S_{\lambda^{(1)}, \lambda^{(2)}}(t^{(1)}, t^{(2)})$  is defined as

$$(1.3) \quad S_{\lambda^{(1)}, \lambda^{(2)}}(t^{(1)}, t^{(2)}) = \det^t \left( q_{\lambda_m^{(2)}}^-, q_{\lambda_{m-1}^{(2)}+1}^-, \dots, q_{\lambda_1^{(2)}+m-1}^-, \right. \\ \left. p_{\lambda_1^{(1)}-m}^+, p_{\lambda_2^{(1)}-m-1}^+, \dots, p_{\lambda_n^{(1)}-m-n+1}^+ \right),$$

where

$$(1.4) \quad p_j^+ = {}^t(p_j, p_{j+1}, \dots, p_{j+m+n-1}), \quad q_j^- = {}^t(q_j, q_{j-1}, \dots, q_{j-m-n+1}).$$

Our main result is stated as follows.

**THEOREM 1.1.** *For  $m, n \in \mathbb{Z}_{\geq 0}$ , we define a family of polynomials  $S_{m,n} = S_{m,n}(t, s)$  by specializing  $S_{\lambda^{(1)}, \lambda^{(2)}}(t^{(1)}, t^{(2)})$  as*

$$(1.5) \quad \lambda^{(1)} = (n, n-1, \dots, 2, 1), \quad \lambda^{(2)} = (m, m-1, \dots, 2, 1),$$

$$(1.6) \quad t_j^{(1)} = -\frac{t}{2} + \frac{2s-m+n}{j}, \quad t_j^{(2)} = \frac{t}{2} + \frac{2s-m+n}{j},$$

where  $s$  is a parameter. For  $m, n \in \mathbb{Z}_{< 0}$ , we define  $S_{m,n}$  through

$$(1.7) \quad \begin{aligned} S_{m,n}(t, s) &= (-1)^{m(m+1)/2} S_{-m-1, n}(t, s-m-1/2), \\ S_{m,n}(t, s) &= (-1)^{n(n+1)/2} S_{m, -n-1}(t, s-n-1/2). \end{aligned}$$

Then,

$$(1.8) \quad y = -\frac{S_{m,n-1}(t, s)S_{m-1,n}(t, s)}{S_{m-1,n}(t, s-1)S_{m,n-1}(t, s+1)},$$

gives the rational solutions of  $P_V$  (1.1) with the parameters

$$(1.9) \quad \kappa_\infty = s, \quad \kappa_0 = s-m+n, \quad \theta = m+n-1,$$

and

$$(1.10) \quad \kappa_\infty = -s, \quad \kappa_0 = s-m+n, \quad \theta = m+n-1.$$

Similarly,

$$(1.11) \quad y = \frac{2n+1}{2m+1} \frac{S_{m,n-1}(t, s+1/2)S_{m,n+1}(t, s-1/2)}{S_{m-1,n}(t, s-1/2)S_{m+1,n}(t, s+1/2)},$$

gives the rational solutions of  $P_V$  (1.1) with the parameters

$$(1.12) \quad \kappa_\infty = m+1/2, \quad \kappa_0 = n+1/2, \quad \theta = 2s-m-n-1.$$

This result covers all the rational solutions obtained by applying the Bäcklund transformations on the particular solution of  $P_V$  (1.1),

$$(1.13) \quad y = -1, \quad \kappa_\infty = s, \quad \kappa_0 = s, \quad \theta = -1.$$

*Remark.* In ref. [5], Kitaev et al gave a complete classification of rational solutions for  $P_V$ . Our result covers all the rational solutions of the cases (III) and (IV) in their classification. The first half corresponds to the case (III) and the other does to (IV). Also, Noumi and Yamada presented a determinant formula for a class of rational solutions of  $P_V$  in terms of 2-reduced Schur functions [9]. Our result includes their formula as a special case, which is explained in Appendix A.

This paper is organized as follows. In Section 2, we give a brief review for the theory of symmetric form of  $P_V$  [8], [10], [11], which provides us with a clear description of symmetry structure and  $\tau$ -functions for  $P_V$ . In Section 3, we construct the rational solutions of  $P_V$  by using the theory of symmetric form. Proof of our result is given in Section 4. We mention the relationship between our result and Yamada's general determinant formula [21] of Jacobi-Trudi type in Section 5.

## §2. Symmetric form of Painlevé V equation

By using the theory of symmetric form for  $P_V$ , it is possible to describe the structure of Bäcklund transformations in a unified manner and to construct particular solutions systematically. In this section, we summarize the symmetric form of  $P_V$  following refs. [10], [11], and derive bilinear equations satisfied by  $\tau$ -functions.

### 2.1. Symmetric form of $P_V$

$P_V$  (1.1) is equivalent to the Hamilton system [20]

$$(2.1) \quad q' = \frac{\partial H}{\partial p}, \quad p' = -\frac{\partial H}{\partial q}, \quad ' = t \frac{d}{dt},$$

with the Hamiltonian

$$(2.2) \quad H = p(p+t)q(q-1) + \alpha_2 qt - \alpha_3 pq - \alpha_1 p(q-1).$$

In fact, putting

$$(2.3) \quad \kappa_\infty = \alpha_1, \quad \kappa_0 = \alpha_3, \quad \theta = \alpha_2 - \alpha_0 - 1,$$

with

$$(2.4) \quad \alpha_0 = 1 - \alpha_1 - \alpha_2 - \alpha_3,$$

we see that equation for  $y = 1 - 1/q$  is nothing but  $P_V$  (1.1). Setting

$$(2.5) \quad f_0 = \frac{1}{\sqrt{t}}(t + p), \quad f_1 = \sqrt{t}q, \quad f_2 = -\frac{1}{\sqrt{t}}p, \quad f_3 = \sqrt{t}(1 - q),$$

we obtain the symmetric form of  $P_V$

$$(2.6) \quad \begin{aligned} f_0' &= f_0 f_2 (f_1 - f_3) + \left(\frac{1}{2} - \alpha_2\right) f_0 + \alpha_0 f_2, \\ f_1' &= f_1 f_3 (f_2 - f_0) + \left(\frac{1}{2} - \alpha_3\right) f_1 + \alpha_1 f_3, \\ f_2' &= f_2 f_0 (f_3 - f_1) + \left(\frac{1}{2} - \alpha_0\right) f_2 + \alpha_2 f_0, \\ f_3' &= f_3 f_1 (f_0 - f_2) + \left(\frac{1}{2} - \alpha_1\right) f_3 + \alpha_3 f_1. \end{aligned}$$

We note that the original dependent variable  $y$  of  $P_V$  is expressed as

$$(2.7) \quad y = -\frac{f_3}{f_1}.$$

In terms of variables  $f_i$  and  $\alpha_i$ , the Bäcklund transformations of  $P_V$  are described by a simple form,

$$(2.8) \quad \begin{aligned} s_i(\alpha_i) &= -\alpha_i, \quad s_i(\alpha_j) = \alpha_j + \alpha_i \quad (j = i \pm 1), \quad s_i(\alpha_j) = \alpha_j \quad (j \neq i, i \pm 1), \\ s_i(f_i) &= f_i, \quad s_i(f_j) = f_j \pm \frac{\alpha_i}{f_i} \quad (j = i \pm 1), \quad s_i(f_j) = f_j \quad (j \neq i, i \pm 1), \\ \pi(\alpha_j) &= \alpha_{j+1}, \quad \pi(f_j) = f_{j+1}, \end{aligned}$$

where the subscripts  $i = 0, 1, 2, 3$  are understood as elements of  $\mathbb{Z}/4\mathbb{Z}$ . These transformations commute with derivation  $'$  and satisfy the fundamental relations

$$(2.9) \quad \begin{aligned} s_i^2 &= 1, \quad s_i s_j = s_j s_i \quad (j \neq i, i \pm 1), \quad s_i s_j s_i = s_j s_i s_j \quad (j = i \pm 1), \\ \pi^4 &= 1, \quad \pi s_j = s_{j+1} \pi, \end{aligned}$$

which means that transformations  $s_i$  ( $i = 0, 1, 2, 3$ ) generate the affine Weyl group  $W(A_3^{(1)})$ , and  $s_i$  and  $\pi$  generate its extension including the Dynkin diagram automorphisms.

## 2.2. $\tau$ -functions and bilinear equations

In order to obtain simpler transformation properties, we add a correction term which depends only on  $t$  to the Hamiltonian (2.2). The corrected

Hamiltonian  $h_0$  is introduced as

$$(2.10) \quad h_0 = f_0 f_1 f_2 f_3 + \frac{\alpha_1 + 2\alpha_2 - \alpha_3}{4} f_0 f_1 + \frac{\alpha_1 + 2\alpha_2 + 3\alpha_3}{4} f_1 f_2 - \frac{3\alpha_1 + 2\alpha_2 + \alpha_3}{4} f_2 f_3 + \frac{\alpha_1 - 2\alpha_2 - \alpha_3}{4} f_3 f_0 + \frac{(\alpha_1 + \alpha_3)^2}{4},$$

and we put  $h_j = \pi^j(h_0)$ . Then, we have

$$(2.11) \quad s_i(h_j) = h_j \quad (i \neq j), \quad s_i(h_i) = h_i + \sqrt{t} \frac{\alpha_i}{f_i}, \quad \pi(h_i) = h_{i+1}.$$

We also introduce  $\tau$ -functions  $\tau_i$  ( $i = 0, 1, 2, 3$ ) by

$$(2.12) \quad h_i = \frac{\tau_i'}{\tau_i}.$$

Then, defining the action of  $s_i$  ( $i = 0, 1, 2, 3$ ) and  $\pi$  on the  $\tau$ -functions by

$$(2.13) \quad s_i(\tau_j) = \tau_j \quad (i \neq j), \quad s_i(\tau_i) = f_i \frac{\tau_{i-1} \tau_{i+1}}{\tau_i}, \quad \pi(\tau_i) = \tau_{i+1},$$

we see that the fundamental relations (2.9) are preserved, which implies that the Bäcklund transformations can be lifted to the level of  $\tau$ -functions. It should be remarked that we have from (2.7) and (2.13)

$$(2.14) \quad y = -\frac{\tau_3 s_3(\tau_3)}{\tau_1 s_1(\tau_1)}.$$

By (2.13), the Bäcklund transformations (2.8) lead to a set of bilinear equations for  $\tau$ -functions,

$$(2.15) \quad \begin{aligned} \tau_0 s_0 s_1(\tau_1) &= s_0(\tau_0) s_1(\tau_1) + \alpha_0 \tau_2 \tau_3, \\ \tau_1 s_1 s_0(\tau_0) &= s_0(\tau_0) s_1(\tau_1) - \alpha_1 \tau_2 \tau_3, \\ \tau_1 s_1 s_2(\tau_2) &= s_1(\tau_1) s_2(\tau_2) + \alpha_1 \tau_3 \tau_0, \\ \tau_2 s_2 s_1(\tau_1) &= s_1(\tau_1) s_2(\tau_2) - \alpha_2 \tau_3 \tau_0, \\ \tau_2 s_2 s_3(\tau_3) &= s_2(\tau_2) s_3(\tau_3) + \alpha_2 \tau_0 \tau_1, \\ \tau_3 s_3 s_2(\tau_2) &= s_2(\tau_2) s_3(\tau_3) - \alpha_3 \tau_0 \tau_1, \\ \tau_3 s_3 s_0(\tau_0) &= s_3(\tau_3) s_0(\tau_0) + \alpha_3 \tau_1 \tau_2, \\ \tau_0 s_0 s_3(\tau_3) &= s_3(\tau_3) s_0(\tau_0) - \alpha_0 \tau_1 \tau_2. \end{aligned}$$

Let us define the translation operators  $T_i$  ( $i = 0, 1, 2, 3$ ) by

$$(2.16) \quad T_1 = \pi s_3 s_2 s_1, \quad T_2 = s_1 \pi s_3 s_2, \quad T_3 = s_2 s_1 \pi s_3, \quad T_0 = s_3 s_2 s_1 \pi,$$

which commute with each other and satisfy  $T_1 T_2 T_3 T_0 = 1$ . These operators act on parameters  $\alpha_i$  as

$$(2.17) \quad T_i(\alpha_{i-1}) = \alpha_{i-1} + 1, \quad T_i(\alpha_i) = \alpha_i - 1, \quad T_i(\alpha_j) = \alpha_j \quad (j \neq i - 1, i),$$

and generate the weight lattice of  $A_3^{(1)}$ . In terms of  $T_i$ ,  $\tau$ -functions in (2.15) are expressed as

$$(2.18) \quad \begin{aligned} \tau_1 &= T_1(\tau_0), & \tau_2 &= T_1 T_2(\tau_0), & \tau_3 &= T_0^{-1}(\tau_0), \\ s_0(\tau_0) &= T_0^{-1} T_1(\tau_0), & s_1(\tau_1) &= T_2(\tau_0), \\ s_2(\tau_2) &= T_1 T_3(\tau_0), & s_3(\tau_3) &= T_3^{-1}(\tau_0), \\ s_0 s_1(\tau_1) &= T_1 T_2 T_0^{-1}(\tau_0), & s_1 s_0(\tau_0) &= T_2 T_0^{-1}(\tau_0), \\ s_1 s_2(\tau_2) &= T_2 T_3(\tau_0), & s_2 s_1(\tau_1) &= T_3(\tau_0), \\ s_2 s_3(\tau_3) &= T_2^{-1}(\tau_0), & s_3 s_2(\tau_2) &= T_1 T_0(\tau_0), \\ s_3 s_0(\tau_0) &= T_1 T_3^{-1}(\tau_0), & s_0 s_3(\tau_3) &= T_1 T_3^{-1} T_0^{-1}(\tau_0). \end{aligned}$$

Furthermore, we can derive bilinear equations of Toda type.

PROPOSITION 2.1. *We have*

$$(2.19) \quad \begin{aligned} T_1(\tau_0) T_1^{-1}(\tau_0) &= \frac{1}{\sqrt{t}} \left( \frac{1}{2} D_T^2 + \frac{3\alpha_1 + 2\alpha_2 + \alpha_3}{4} t \right) \tau_0 \cdot \tau_0, \\ T_2(\tau_0) T_2^{-1}(\tau_0) &= \frac{1}{\sqrt{t}} \left( \frac{1}{2} D_T^2 - \frac{\alpha_1 - 2\alpha_2 - \alpha_3}{4} t \right) \tau_0 \cdot \tau_0, \\ T_3(\tau_0) T_3^{-1}(\tau_0) &= \frac{1}{\sqrt{t}} \left( \frac{1}{2} D_T^2 - \frac{\alpha_1 + 2\alpha_2 - \alpha_3}{4} t \right) \tau_0 \cdot \tau_0, \\ T_0(\tau_0) T_0^{-1}(\tau_0) &= \frac{1}{\sqrt{t}} \left( \frac{1}{2} D_T^2 - \frac{\alpha_1 + 2\alpha_2 + 3\alpha_3}{4} t \right) \tau_0 \cdot \tau_0, \end{aligned}$$

where  $D_T$  is the Hirota's differential operator defined by

$$(2.20) \quad D_T^m f \cdot g = \left( \frac{d}{dT} - \frac{d}{dT'} \right)^m f(T) g(T') \Big|_{T=T'},$$

and  $\frac{d}{dT} = t \frac{d}{dt}$ .

*Proof.* Using (2.8), (2.13) and (2.16), we have

$$(2.21) \quad T_1(\tau_0)T_1^{-1}(\tau_0) = [f_1f_2f_3 + (\alpha_1 + \alpha_2)f_1 + \alpha_1f_3]\tau_0^2.$$

Noticing that

$$(2.22) \quad h'_0 = \sqrt{t} \left( f_1f_2f_3 + \frac{\alpha_1 + 2\alpha_2 - \alpha_3}{4}f_1 + \frac{\alpha_1 - 2\alpha_2 - \alpha_1}{4}f_3 \right),$$

and  $f_1 + f_3 = \sqrt{t}$ , we get the first equation in (2.19). The other equations are obtained in similar way.  $\square$

*Remark.* Bilinear equations (2.15) and (2.19) are overdetermined systems when they are regarded as equations to determine the  $\tau$ -functions. However, by construction, the consistency of these equations is guaranteed.

### 2.3. $\tau$ -cocycles

For simplicity, we introduce a notation,

$$(2.23) \quad \tau_{k,l,m,n} = T_1^k T_2^l T_3^m T_0^n(\tau_0).$$

For small  $k, l, m, n$ , we observe that  $\tau_{k,l,m,n}$  are factorized as

$$(2.24) \quad \tau_{k,l,m,n} = \phi_{k,l,m,n} \tau_0 \left( \frac{\tau_1}{\tau_0} \right)^k \left( \frac{\tau_2}{\tau_1} \right)^l \left( \frac{\tau_3}{\tau_2} \right)^m \left( \frac{\tau_0}{\tau_3} \right)^n,$$

where  $\phi_{k,l,m,n}$  are some functions of  $f_i$  and  $\alpha_i$ . Conversely, if we define  $\phi_{k,l,m,n}$  by (2.24), it is shown that  $\phi_{k,l,m,n}$ 's are polynomials in  $f_i$  and  $\alpha_i$  for any  $k, l, m, n \in \mathbb{Z}$  [21]. The functions  $\phi_{k,l,m,n}$  are called the  $\tau$ -cocycles.

It is easy to see from (2.14), (2.18), (2.23) and (2.24) that we have

$$(2.25) \quad T_1^k T_2^l T_3^m T_0^n(y) = -\frac{\phi_{k,l,m,n-1}\phi_{k,l,m-1,n}}{\phi_{k+1,l,m,n}\phi_{k,l+1,m,n}}, \quad \text{for } k, l, m, n \in \mathbb{Z}.$$

Moreover, it follows from (2.13), (2.16), (2.23) and (2.24) that  $\phi_{k,l,m,n}$  are



determined by the recurrence relations,

(2.26)

$$\begin{aligned}
& \phi_{k+1,l,m,n} \\
&= T_1(\phi_{k,l,m,n})[f_2 f_3 f_0 - (\alpha_3 + \alpha_0) f_0 - \alpha_0 f_2]^{k-l} (f_3 f_0 - \alpha_0)^{l-m} f_0^{m-n}, \\
& \phi_{k,l+1,m,n} \\
&= T_2(\phi_{k,l,m,n})[f_3 f_0 f_1 - (\alpha_0 + \alpha_1) f_1 - \alpha_1 f_3]^{l-m} (f_0 f_1 - \alpha_1)^{m-n} f_1^{1+n-k}, \\
& \phi_{k,l,m+1,n} \\
&= T_3(\phi_{k,l,m,n})[f_0 f_1 f_2 - (\alpha_1 + \alpha_2) f_2 - \alpha_2 f_0]^{m-n} (f_1 f_2 - \alpha_2)^{1+n-k} f_2^{k-l}, \\
& \phi_{k,l,m,n+1} \\
&= T_0(\phi_{k,l,m,n})[f_1 f_2 f_3 - (\alpha_2 + \alpha_3) f_3 - \alpha_3 f_1]^{1+n-k} (f_2 f_3 - \alpha_3)^{k-l} f_3^{l-m},
\end{aligned}$$

with  $\phi_{0,0,0,0} = 1$ .

It is possible to write down the bilinear equations for  $\phi_{k,l,m,n}$ . From (2.18), (2.23) and (2.24), bilinear Bäcklund transformations (2.15) yield

$$\begin{aligned}
& \phi_{k,l,m,n} \phi_{k+1,l+1,m,n-1} \\
&= \phi_{k+1,l,m,n-1} \phi_{k,l+1,m,n} + (\alpha_0 - n + k) \phi_{k+1,l+1,m,n} \phi_{k,l,m,n-1}, \\
& \phi_{k+1,l,m,n} \phi_{k,l+1,m,n-1} \\
&= \phi_{k+1,l,m,n-1} \phi_{k,l+1,m,n} - (\alpha_1 - k + l) \phi_{k+1,l+1,m,n} \phi_{k,l,m,n-1}, \\
& \phi_{k+1,l,m,n} \phi_{k,l+1,m+1,n} \\
&= \phi_{k,l+1,m,n} \phi_{k+1,l,m+1,n} + (\alpha_1 - k + l) \phi_{k,l,m,n-1} \phi_{k,l,m,n}, \\
& \phi_{k+1,l+1,m,n} \phi_{k,l,m+1,n} \\
&= \phi_{k,l+1,m,n} \phi_{k+1,l,m+1,n} - (\alpha_2 - l + m) \phi_{k,l,m,n-1} \phi_{k,l,m,n}, \\
& \phi_{k+1,l+1,m,n} \phi_{k,l-1,m,n} \\
&= \phi_{k+1,l,m+1,n} \phi_{k,l,m-1,n} + (\alpha_2 - l + m) \phi_{k,l,m,n} \phi_{k+1,l,m,n}, \\
& \phi_{k,l,m,n-1} \phi_{k+1,l,m,n+1} \\
&= \phi_{k+1,l,m+1,n} \phi_{k,l,m-1,n} - (\alpha_3 - m + n) \phi_{k,l,m,n} \phi_{k+1,l,m,n}, \\
& \phi_{k,l,m,n-1} \phi_{k+1,l,m-1,n} \\
&= \phi_{k,l,m-1,n} \phi_{k+1,l,m,n-1} + (\alpha_3 - m + n) \phi_{k+1,l,m,n} \phi_{k+1,l+1,m,n}, \\
& \phi_{k,l,m,n} \phi_{k+1,l,m-1,n-1} \\
&= \phi_{k,l,m-1,n} \phi_{k+1,l,m,n-1} - (\alpha_0 - n + k) \phi_{k+1,l,m,n} \phi_{k+1,l+1,m,n}.
\end{aligned}
\tag{2.27}$$

Similarly, the last two equations in (2.19) are lead to

(2.28)

$$\begin{aligned} & \phi_{k,l,m+1,n} \phi_{k,l,m-1,n} \\ &= \frac{1}{\sqrt{t}} \left( \frac{1}{2} D_T^2 + \omega_{k,l,m,n} - \frac{\alpha_1 + 2\alpha_2 - \alpha_3 - k - l + 3m - n}{4} t \right) \phi_{k,l,m,n} \cdot \phi_{k,l,m,n}, \\ & \phi_{k,l,m,n+1} \phi_{k,l,m,n-1} \\ &= \frac{1}{\sqrt{t}} \left( \frac{1}{2} D_T^2 + \omega_{k,l,m,n} - \frac{\alpha_1 + 2\alpha_2 + 3\alpha_3 - k - l - m + 3n}{4} t \right) \phi_{k,l,m,n} \cdot \phi_{k,l,m,n}, \end{aligned}$$

with

$$(2.29) \quad \begin{aligned} \omega_{k,l,m,n} &= (\log \tau_0)'' + k \left( \log \frac{\tau_1}{\tau_0} \right)'' + l \left( \log \frac{\tau_2}{\tau_1} \right)'' \\ &\quad + m \left( \log \frac{\tau_3}{\tau_2} \right)'' + n \left( \log \frac{\tau_0}{\tau_3} \right)'' . \end{aligned}$$

### §3. Construction of rational solutions

In this section, we construct the rational solutions of  $P_V$  by using the results in the previous section.

It is obvious that the symmetric form of  $P_V$  (2.6) with (2.4) has a solution,

$$(3.1) \quad (\alpha_0, \alpha_1, \alpha_2, \alpha_3) = \left( \frac{1}{2} - s, s, \frac{1}{2} - s, s \right), \quad f_i = \frac{\sqrt{t}}{2} \text{ for } i = 0, 1, 2, 3,$$

which is on the fixed points with respect to the transformation  $\pi^2$  and is equivalent to the following solution of  $P_V$ ,

$$(3.2) \quad y = -1, \quad \kappa_\infty = s, \quad \kappa_0 = s, \quad \theta = -1.$$

This is the unique rational solution in the fundamental region of the affine Weyl group  $W(A_3^{(1)})$  in the parameter space, except for the special cases of transcendental classical solutions [20]. Applying Bäcklund transformations to the seed solution (3.1), we obtain the family of rational solutions of  $P_V$ . Note that we have

$$(3.3) \quad T_2^l T_0^l(\alpha_0, \alpha_1, \alpha_2, \alpha_3) = \left( \frac{1}{2} - \tilde{s}, \tilde{s}, \frac{1}{2} - \tilde{s}, \tilde{s} \right), \quad \tilde{s} = s + l, \quad l \in \mathbb{Z},$$

under the specialization (3.1). Comparing (3.3) with (3.1), we see that the effect of  $T_2$  is absorbed by that of  $T_0^{-1}$  and shift of the parameter  $s$ . Then,

we do not need to consider the Bäcklund transformation  $T_2$  for constructing the family of rational solutions of  $P_V$ . Taking the initial condition (3.1) into account, we consider the bilinear Bäcklund transformations in terms of the  $\tau$ -cocycles (2.27). Denoting  $\phi_{0,l,m,n} = \phi_{l,m,n}$  in view of the relation  $T_1 T_2 T_3 T_0 = 1$ , it is easy to derive the following.

LEMMA 3.1. *Under the specialization (3.1), bilinear Bäcklund transformations (2.27) are reduced to*

$$\begin{aligned}
& \phi_{0,m,n} \phi_{0,m-1,n-2} \\
&= \phi_{-1,m-1,n-2} \phi_{1,m,n} + (1/2 - s - n) \phi_{0,m-1,n-1} \phi_{0,m,n-1}, \\
& \phi_{-1,m-1,n-1} \phi_{1,m,n-1} \\
&= \phi_{-1,m-1,n-2} \phi_{1,m,n} - s \phi_{0,m-1,n-1} \phi_{0,m,n-1}, \\
& \phi_{-1,m-1,n-1} \phi_{1,m+1,n} \\
&= \phi_{1,m,n} \phi_{-1,m,n-1} + s \phi_{0,m,n-1} \phi_{0,m,n}, \\
& \phi_{0,m-1,n-1} \phi_{0,m+1,n} \\
&= \phi_{1,m,n} \phi_{-1,m,n-1} - (1/2 - s + m) \phi_{0,m,n-1} \phi_{0,m,n}, \\
(3.4) \quad & \phi_{0,m-1,n-1} \phi_{-1,m,n} \\
&= \phi_{-1,m,n-1} \phi_{0,m-1,n} + (1/2 - s + m) \phi_{0,m,n} \phi_{-1,m-1,n-1}, \\
& \phi_{0,m,n-1} \phi_{-1,m-1,n} \\
&= \phi_{-1,m,n-1} \phi_{0,m-1,n} - (s - m + n) \phi_{0,m,n} \phi_{-1,m-1,n-1}, \\
& \phi_{0,m,n-1} \phi_{-1,m-2,n-1} \\
&= \phi_{0,m-1,n} \phi_{-1,m-1,n-2} + (s - m + n) \phi_{-1,m-1,n-1} \phi_{0,m-1,n-1}, \\
& \phi_{0,m,n} \phi_{-1,m-2,n-2} \\
&= \phi_{0,m-1,n} \phi_{-1,m-1,n-2} - (1/2 - s - n) \phi_{-1,m-1,n-1} \phi_{0,m-1,n-1}.
\end{aligned}$$

Moreover, from (2.25), the function

$$(3.5) \quad y = -\frac{\phi_{0,m,n-1} \phi_{0,m-1,n}}{\phi_{-1,m-1,n-1} \phi_{1,m,n}},$$

solves  $P_V$  (1.1) with parameters

$$(3.6) \quad \kappa_\infty = s, \quad \kappa_0 = s - m + n, \quad \theta = m + n - 1.$$

From the recurrence relations (2.26), we observe that  $\phi_{l,m,n}$  for small  $l, m, n$  are expressed as

$$(3.7) \quad \phi_{l,m,n} = \left(\frac{\sqrt{t}}{2}\right)^{(m-n-l-1)(m-n-l)/2} U_{l,m,n},$$

where  $U_{l,m,n}$  are some polynomials in  $t$  and  $s$ . Therefore, we next rewrite (3.4) in terms of  $U$ . The polynomials  $U_{l,m,n}$  have symmetry described by the following lemma.

LEMMA 3.2. *The polynomials  $U_{l,m,n}$  defined by (3.7) satisfy*

$$(3.8) \quad U_{1,m,n}(t,s) = U_{0,m,n-1}(t,s+1), \quad U_{-1,m,n}(t,s) = U_{0,m,n+1}(t,s-1).$$

*Proof.* Lemma 3.2 is proved by considering the Toda type equation for  $\tau$ -cocycles. Under the specialization (3.1), the Hamiltonians and  $\tau$ -functions are calculated as

$$(3.9) \quad h_0 = h_2 = \frac{t^2}{16} + s^2, \quad h_1 = h_3 = \frac{t^2}{16} + \left(\frac{1}{2} - s\right)^2,$$

and

$$(3.10) \quad \tau_0 = \tau_2 = t^{s^2} \exp\left(\frac{t^2}{32}\right), \quad \tau_1 = \tau_3 = t^{(1/2-s)^2} \exp\left(\frac{t^2}{32}\right),$$

up to the multiplication by some constants, respectively. Then, Toda type bilinear equations (2.28) yield

$$(3.11) \quad \begin{aligned} & \phi_{l,m+1,n} \phi_{l,m-1,n} \\ &= \frac{1}{\sqrt{t}} \left( \frac{1}{2} D_T^2 + \frac{t^2}{8} - \frac{-2s+1-l+3m-n}{4} t \right) \phi_{l,m,n} \cdot \phi_{l,m,n}, \\ & \phi_{l,m,n+1} \phi_{l,m,n-1} \\ &= \frac{1}{\sqrt{t}} \left( \frac{1}{2} D_T^2 + \frac{t^2}{8} - \frac{2s+1-l-m+3n}{4} t \right) \phi_{l,m,n} \cdot \phi_{l,m,n}. \end{aligned}$$

Substituting (3.7) into (3.11), we obtain Toda type bilinear equations to be satisfied by  $U_{m,n} = U_{0,m,n}(t,s)$

$$(3.12) \quad \begin{aligned} U_{m+1,n} U_{m-1,n} &= 2t \left[ \left( \frac{d^2 U_{m,n}}{dt^2} \right) U_{m,n} - \left( \frac{dU_{m,n}}{dt} \right)^2 \right] \\ &\quad + 2 \frac{dU_{m,n}}{dt} U_{m,n} + \left( \frac{t}{4} - \frac{-2s+1+3m-n}{2} \right) U_{m,n}^2, \\ U_{m,n+1} U_{m,n-1} &= 2t \left[ \left( \frac{d^2 U_{m,n}}{dt^2} \right) U_{m,n} - \left( \frac{dU_{m,n}}{dt} \right)^2 \right] \\ &\quad + 2 \frac{dU_{m,n}}{dt} U_{m,n} + \left( \frac{t}{4} - \frac{2s+1-m+3n}{2} \right) U_{m,n}^2, \end{aligned}$$

with initial conditions

$$(3.13) \quad U_{-1,-1} = U_{-1,0} = U_{0,-1} = U_{0,0} = 1.$$

The functions  $U_{m,n} = U_{m,n}(t, s)$  are uniquely determined by Toda equations (3.12) from the initial conditions (3.13) for any  $m, n \in \mathbb{Z}$ . Moreover, we see that  $U_{\pm 1, m, n}(t, s)$  satisfy the same Toda equations as  $U_{0, m, n \mp 1}(t, s \pm 1)$ , respectively, by the similar calculation. Since the initial conditions for  $U_{1, m, n}$  and  $U_{-1, m, n}$  are given by

$$(3.14) \quad \begin{aligned} U_{1,-1,0} &= U_{1,0,0} = U_{1,-1,1} = U_{1,0,1} = 1, \\ U_{-1,-1,-2} &= U_{-1,-1,-1} = U_{-1,0,-2} = U_{-1,0,-1} = 1, \end{aligned}$$

the lemma is proved.  $\square$

From Lemma 3.2, bilinear Bäcklund transformations (3.4) are rewritten in terms of  $U$ .

**PROPOSITION 3.3.** *Let  $U_{m,n} = U_{m,n}(t, s)$  ( $m, n \in \mathbb{Z}$ ) be polynomials which satisfy the bilinear equations,*

$$(3.15) \quad \begin{aligned} 4U_{m,n+1}U_{m-1,n-1} &= tU_{m-1,n}^- U_{m,n}^+ - 2(2s + 2n + 1)U_{m-1,n}U_{m,n}, \\ 4U_{m-1,n+1}^- U_{m,n-1}^+ &= tU_{m-1,n}^- U_{m,n}^+ - 4sU_{m-1,n}U_{m,n}, \\ 4U_{m-1,n}^- U_{m+1,n-1}^+ &= tU_{m,n-1}^+ U_{m,n}^- + 4sU_{m,n-1}U_{m,n}, \\ 4U_{m-1,n-1}U_{m+1,n} &= tU_{m,n-1}^+ U_{m,n}^- + 2(2s - 2m - 1)U_{m,n-1}U_{m,n}, \\ 4U_{m-1,n-1}U_{m,n+1}^- &= tU_{m,n}^- U_{m-1,n} - 2(2s - 2m - 1)U_{m,n}U_{m-1,n}^-, \\ 4U_{m,n-1}U_{m-1,n+1}^- &= tU_{m,n}^- U_{m-1,n} - 4(s - m + n)U_{m,n}U_{m-1,n}^-, \\ 4U_{m+1,n-1}U_{m-1,n}^- &= tU_{m,n}U_{m,n-1}^- + 4(s - m + n - 1)U_{m,n}^- U_{m,n-1}, \\ 4U_{m+1,n}U_{m-1,n-1}^- &= tU_{m,n}U_{m,n-1}^- + 2(2s + 2n - 1)U_{m,n}^- U_{m,n-1}, \end{aligned}$$

with

$$(3.16) \quad U_{-1,-1} = U_{-1,0} = U_{0,-1} = U_{0,0} = 1,$$

where we denote  $U_{m,n}^\pm = U_{m,n}(t, s \pm 1)$ . Then,

$$(3.17) \quad y = -\frac{U_{m,n-1}(t, s)U_{m-1,n}(t, s)}{U_{m-1,n}(t, s-1)U_{m,n-1}(t, s+1)},$$

gives the rational solutions of  $P_V$  (1.1) with parameters

$$(3.18) \quad \kappa_\infty = s, \quad \kappa_0 = s - m + n, \quad \theta = m + n - 1.$$

#### §4. Proof of Theorem 1.1

In this section, we give the proof of Theorem 1.1.

DEFINITION 2. Let  $p_k^{(r)} = p_k^{(r)}(x)$  and  $q_k^{(r)} = q_k^{(r)}(x)$  be polynomials defined by

$$(4.1) \quad \sum_{k=0}^{\infty} p_k^{(r)} \eta^k = (1 - \eta)^{-r} \exp\left(-\frac{x\eta}{1 - \eta}\right), \quad p_k^{(r)} = 0 \text{ for } k < 0,$$

$$(4.2) \quad q_k^{(r)}(x) = p_k^{(r)}(-x),$$

respectively. For  $m, n \in \mathbb{Z}_{\geq 0}$ , we define a family of polynomials  $R_{m,n}^{(r)} = R_{m,n}^{(r)}(x)$  by

$$(4.3) \quad R_{m,n}^{(r)}(x) = \begin{vmatrix} q_1^{(r)} & q_0^{(r)} & \cdots & q_{-m+2}^{(r)} & q_{-m+1}^{(r)} & \cdots & q_{-m-n+3}^{(r)} & q_{-m-n+2}^{(r)} \\ q_3^{(r)} & q_2^{(r)} & \cdots & q_{-m+4}^{(r)} & q_{-m+3}^{(r)} & \cdots & q_{-m-n+5}^{(r)} & q_{-m-n+4}^{(r)} \\ \vdots & \vdots & \ddots & \vdots & \vdots & \ddots & \vdots & \vdots \\ q_{2m-1}^{(r)} & q_{2m-2}^{(r)} & \cdots & q_m^{(r)} & q_{m-1}^{(r)} & \cdots & q_{-n+1}^{(r)} & q_{-n}^{(r)} \\ p_{n-m}^{(r)} & p_{n-m+1}^{(r)} & \cdots & p_{n-1}^{(r)} & p_n^{(r)} & \cdots & p_{2n-2}^{(r)} & p_{2n-1}^{(r)} \\ \vdots & \vdots & \ddots & \vdots & \vdots & \ddots & \vdots & \vdots \\ p_{-n-m+4}^{(r)} & p_{-n-m+5}^{(r)} & \cdots & p_{-n+3}^{(r)} & p_{-n+4}^{(r)} & \cdots & p_2^{(r)} & p_3^{(r)} \\ p_{-n-m+2}^{(r)} & p_{-n-m+3}^{(r)} & \cdots & p_{-n+1}^{(r)} & p_{-n+2}^{(r)} & \cdots & p_0^{(r)} & p_1^{(r)} \end{vmatrix}.$$

For  $m, n \in \mathbb{Z}_{< 0}$ , we define  $R_{m,n}^{(r)}$  through

$$(4.4) \quad R_{m,n}^{(r)} = (-1)^{m(m+1)/2} R_{-m-1,n}^{(r)}, \quad R_{m,n}^{(r)} = (-1)^{n(n+1)/2} R_{m,-n-1}^{(r)}.$$

*Remark.* The polynomials  $p_k$  and  $q_k$  ( $k \geq 0$ ) are essentially the Laguerre polynomials, namely,  $p_k^{(r)}(x) = L_k^{(r-1)}(x)$ . Moreover,  $R_{m,n}^{(r)}$  is related to  $S_{m,n}$  in Theorem 1.1 as

$$(4.5) \quad R_{m,n}^{(r)}(x) = S_{m,n}(t, s), \quad x = \frac{t}{2}, \quad r = 2s - m + n.$$

PROPOSITION 4.1. For  $m, n \in \mathbb{Z}$ ,  $R_{m,n}^{(r)}$  satisfy the following bilinear equations.

$$\begin{aligned}
& -(2n+1)R_{m,n+1}^{(r+1)}R_{m-1,n-1}^{(r)} \\
& \quad = xR_{m-1,n}^{(r-1)}R_{m,n}^{(r+2)} - (r+m+n+1)R_{m-1,n}^{(r+1)}R_{m,n}^{(r)}, \\
& -(2n+1)R_{m-1,n+1}^{(r)}R_{m,n-1}^{(r+1)} \\
& \quad = xR_{m-1,n}^{(r-1)}R_{m,n}^{(r+2)} - (r+m-n)R_{m-1,n}^{(r+1)}R_{m,n}^{(r)}, \\
& (2m+1)R_{m-1,n}^{(r-1)}R_{m+1,n-1}^{(r)} \\
& \quad = xR_{m,n-1}^{(r+1)}R_{m,n}^{(r-2)} + (r+m-n)R_{m,n-1}^{(r-1)}R_{m,n}^{(r)}, \\
& (2m+1)R_{m-1,n-1}^{(r)}R_{m+1,n}^{(r-1)} \\
& \quad = xR_{m,n-1}^{(r+1)}R_{m,n}^{(r-2)} + (r-m-n-1)R_{m,n-1}^{(r-1)}R_{m,n}^{(r)}, \\
& -(2n+1)R_{m-1,n-1}^{(r)}R_{m,n+1}^{(r-1)} \\
& \quad = xR_{m,n}^{(r-2)}R_{m-1,n}^{(r+1)} - (r-m-n-1)R_{m,n}^{(r)}R_{m-1,n}^{(r-1)}, \\
& -(2n+1)R_{m,n-1}^{(r-1)}R_{m-1,n+1}^{(r)} \\
& \quad = xR_{m,n}^{(r-2)}R_{m-1,n}^{(r+1)} - (r-m+n)R_{m,n}^{(r)}R_{m-1,n}^{(r-1)}, \\
& (2m+1)R_{m+1,n-1}^{(r-2)}R_{m-1,n}^{(r-1)} \\
& \quad = xR_{m,n}^{(r)}R_{m,n-1}^{(r-3)} + (r-m+n-2)R_{m,n}^{(r-2)}R_{m,n-1}^{(r-1)}, \\
& (2m+1)R_{m+1,n}^{(r-1)}R_{m-1,n-1}^{(r-2)} \\
& \quad = xR_{m,n}^{(r)}R_{m,n-1}^{(r-3)} + (r+m+n-1)R_{m,n}^{(r-2)}R_{m,n-1}^{(r-1)}.
\end{aligned} \tag{4.6}$$

Comparing (4.6) with (3.15), we obtain an explicit formula for  $U_{m,n} = U_{m,n}(t, s)$  in Proposition 3.3.

PROPOSITION 4.2. We have

$$(4.7) \quad U_{m,n}(t, s) = c_m d_n S_{m,n}(t, s), \quad m, n \in \mathbb{Z},$$

where  $c_m$  and  $d_n$  are constants determined by

$$\begin{aligned}
(4.8) \quad c_{m+1}c_{m-1} &= \left(m + \frac{1}{2}\right)c_m^2, & c_{-1} = c_0 = 1, \\
d_{n+1}d_{n-1} &= -\left(n + \frac{1}{2}\right)d_n^2, & d_{-1} = d_0 = 1.
\end{aligned}$$

*Proof.* Putting

$$(4.9) \quad U_{m,n}(t, s) = c_m d_n R_{m,n}^{(r)}(x),$$

with

$$(4.10) \quad x = \frac{t}{2}, \quad r = 2s - m + n,$$

we find that the bilinear relations (4.6) become (3.15). Taking (4.5) into account, we obtain Proposition 4.2.  $\square$

Applying  $s_1$  to the solutions (1.8) with (1.9), we get the solutions (1.8) with (1.10). Then, the first half of Theorem 1.1 is a direct consequence of Propositions 3.3 and 4.2. It is easy to find that the latter half of Theorem 1.1 is obtained by applying  $\pi s_1$  to the solutions (1.8) with (1.9). Therefore, now the proof of Theorem 1.1 is reduced to that of Proposition 4.1.

It is possible to reduce the number of bilinear equations to be proved in (4.6) by the following symmetry of  $R_{m,n}^{(r)}(x)$ .

LEMMA 4.3. *We have the relations for  $m, n \in \mathbb{Z}_{\geq 0}$*

$$(4.11) \quad R_{n,m}^{(r)}(-x) = R_{m,n}^{(r)}(x),$$

$$(4.12) \quad R_{n,m}^{(-r)}(x) = (-1)^{m(m+1)/2+n(n+1)/2} R_{m,n}^{(r)}(x).$$

*Proof.* The first relation (4.11) is easily obtained from (4.3). To verify the second relation (4.12), we introduce polynomials  $\bar{q}_k^{(r)} = \bar{q}_k^{(r)}(x)$  by

$$(4.13) \quad \sum_{k=0}^{\infty} \bar{q}_k^{(r)} \eta^k = (1 + \eta)^r \exp\left(\frac{x\eta}{1 + \eta}\right), \quad \bar{q}_k^{(r)} = 0 \text{ for } k < 0.$$

Comparing the generating function for  $q_k$  with that for  $\bar{q}_k$ , we see that each  $\bar{q}_k^{(r)}(x)$  is a linear combination of  $q_j^{(r)}(x)$ ,  $j = k, k - 2, k - 4, \dots$ . Therefore



we can express  $R_{m,n}^{(r)}$  for  $m, n \in \mathbb{Z}_{\geq 0}$  in terms of  $p_k$  and  $\bar{q}_k$  as

$$(4.14) \quad R_{m,n}^{(r)}(x) = \begin{vmatrix} \bar{q}_1^{(r)} & \bar{q}_0^{(r)} & \cdots & \bar{q}_{-m+2}^{(r)} & \bar{q}_{-m+1}^{(r)} & \cdots & \bar{q}_{-m-n+3}^{(r)} & \bar{q}_{-m-n+2}^{(r)} \\ \bar{q}_3^{(r)} & \bar{q}_2^{(r)} & \cdots & \bar{q}_{-m+4}^{(r)} & \bar{q}_{-m+3}^{(r)} & \cdots & \bar{q}_{-m-n+5}^{(r)} & \bar{q}_{-m-n+4}^{(r)} \\ \vdots & \vdots & \ddots & \vdots & \vdots & \ddots & \vdots & \vdots \\ \bar{q}_{2m-1}^{(r)} & \bar{q}_{2m-2}^{(r)} & \cdots & \bar{q}_m^{(r)} & \bar{q}_{m-1}^{(r)} & \cdots & \bar{q}_{m-n+1}^{(r)} & \bar{q}_{m-n}^{(r)} \\ p_{n-m}^{(r)} & p_{n-m+1}^{(r)} & \cdots & p_{n-1}^{(r)} & p_n^{(r)} & \cdots & p_{2n-2}^{(r)} & p_{2n-1}^{(r)} \\ \vdots & \vdots & \ddots & \vdots & \vdots & \ddots & \vdots & \vdots \\ p_{-n-m+4}^{(r)} & p_{-n-m+5}^{(r)} & \cdots & p_{-n+3}^{(r)} & p_{-n+4}^{(r)} & \cdots & p_2^{(r)} & p_3^{(r)} \\ p_{-n-m+2}^{(r)} & p_{-n-m+3}^{(r)} & \cdots & p_{-n+1}^{(r)} & p_{-n+2}^{(r)} & \cdots & p_0^{(r)} & p_1^{(r)} \end{vmatrix}.$$

Noticing that  $\bar{q}_k$  and  $p_k$  are related as

$$(4.15) \quad \bar{q}_k^{(r)}(x) = (-1)^k p_k^{(-r)}(x),$$

we obtain the relation (4.12).  $\square$

From the symmetries of  $R_{m,n}^{(r)}(x)$  described by (4.4) and Lemma 4.3, it is sufficient to prove the first two equations in (4.6) for  $m, n \in \mathbb{Z}_{\geq 0}$ , which are equivalent to

$$(4.16) \quad R_{m-1,n+1}^{(r)} R_{m,n-1}^{(r+1)} - R_{m,n+1}^{(r+1)} R_{m-1,n-1}^{(r)} - R_{m-1,n}^{(r+1)} R_{m,n}^{(r)} = 0,$$

$$(4.17) \quad \begin{aligned} & -(2n+1) R_{m-1,n+1}^{(r)} R_{m,n-1}^{(r+1)} \\ & = x R_{m-1,n}^{(r-1)} R_{m,n}^{(r+2)} - (r+m-n) R_{m-1,n}^{(r+1)} R_{m,n}^{(r)}. \end{aligned}$$

In the following, we show that these bilinear equations are reduced to Jacobi's identity of determinants. Let  $D$  be an  $(m+n+1) \times (m+n+1)$  determinant and  $D \begin{bmatrix} i_1 & i_2 & \cdots & i_k \\ j_1 & j_2 & \cdots & j_k \end{bmatrix}$  the minor which are obtained by deleting the rows with indices  $i_1, \dots, i_k$  and the columns with indices  $j_1, \dots, j_k$ . Then we have Jacobi's identity

$$(4.18) \quad \begin{aligned} & D \cdot D \begin{bmatrix} m & m+1 \\ 1 & m+n+1 \end{bmatrix} \\ & = D \begin{bmatrix} m \\ 1 \end{bmatrix} D \begin{bmatrix} m+1 \\ m+n+1 \end{bmatrix} - D \begin{bmatrix} m+1 \\ 1 \end{bmatrix} D \begin{bmatrix} m \\ m+n+1 \end{bmatrix}. \end{aligned}$$

We first choose proper determinants as  $D$  ( $D$  itself should be expressed in terms of  $R_{m,n}^{(r)}$ ). Secondly, we construct such formulas that express the minor determinants by  $R_{m,n}^{(r)}$ . Then, Jacobi's identity yields bilinear equations for  $R_{m,n}^{(r)}$  which are nothing but (4.16) and (4.17).

We have the following lemmas.

LEMMA 4.4. *We set*

$$(4.19) \quad D \equiv \begin{vmatrix} -q_1^{(r+1)} & q_1^{(r)} & \cdots & q_{-m-n+3}^{(r)} & q_{-m-n+2}^{(r)} \\ -q_3^{(r+1)} & q_3^{(r)} & \cdots & q_{-m-n+5}^{(r)} & q_{-m-n+4}^{(r)} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ -q_{2m-1}^{(r+1)} & q_{2m-1}^{(r)} & \cdots & q_{m-n+1}^{(r)} & q_{m-n}^{(r)} \\ p_{n-m+1}^{(r+1)} & p_{n-m+2}^{(r)} & \cdots & p_{2n}^{(r)} & p_{2n+1}^{(r)} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ p_{-n-m+3}^{(r+1)} & p_{-n-m+4}^{(r)} & \cdots & p_2^{(r)} & p_3^{(r)} \\ p_{-n-m+1}^{(r+1)} & p_{-n-m+2}^{(r)} & \cdots & p_0^{(r)} & p_1^{(r)} \end{vmatrix}.$$

Then, we have

$$(4.20) \quad \begin{aligned} D &= (-1)^m R_{m,n+1}^{(r+1)}, & D \begin{bmatrix} m \\ 1 \end{bmatrix} &= R_{m-1,n+1}^{(r)}, \\ D \begin{bmatrix} m+1 \\ 1 \end{bmatrix} &= R_{m,n}^{(r)}, & D \begin{bmatrix} m \\ m+n+1 \end{bmatrix} &= (-1)^{m-1} R_{m-1,n}^{(r+1)}, \\ D \begin{bmatrix} m+1 \\ m+n+1 \end{bmatrix} &= (-1)^m R_{m,n-1}^{(r+1)}, \\ D \begin{bmatrix} m & m+1 \\ 1 & m+n+1 \end{bmatrix} &= R_{m-1,n-1}^{(r)}. \end{aligned}$$

LEMMA 4.5. *We set*

$$(4.21) \quad D \equiv \begin{vmatrix} \tilde{q}_1^{(r-m-n+2)} & q_1^{(r-m-n+1)} & q_0^{(r-m-n+2)} & \cdots & q_{-m-n+2}^{(r)} \\ \tilde{q}_3^{(r-m-n+2)} & q_3^{(r-m-n+1)} & q_2^{(r-m-n+2)} & \cdots & q_{-m-n+4}^{(r)} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \tilde{q}_{2m-1}^{(r-m-n+2)} & q_{2m-1}^{(r-m-n+1)} & q_{2m-2}^{(r-m-n+2)} & \cdots & q_{m-n}^{(r)} \\ (-1)^{m+n} \tilde{p}_{2n}^{(r-m-n+2)} & (-1)^{m+n} p_{2n+1}^{(r-m-n+1)} & (-1)^{m+n-1} p_{2n+1}^{(r-m-n+2)} & \cdots & (-1)^1 p_{2n+1}^{(r)} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ (-1)^{m+n} \tilde{p}_2^{(r-m-n+2)} & (-1)^{m+n} p_3^{(r-m-n+1)} & (-1)^{m+n-1} p_3^{(r-m-n+2)} & \cdots & (-1)^1 p_3^{(r)} \\ (-1)^{m+n} \tilde{p}_0^{(r-m-n+2)} & (-1)^{m+n} p_1^{(r-m-n+1)} & (-1)^{m+n-1} p_1^{(r-m-n+2)} & \cdots & (-1)^1 p_1^{(r)} \end{vmatrix},$$

where  $\hat{p}_{2k}^{(r)}$  and  $\tilde{q}_{2k-1}^{(r)}$  are defined by

$$(4.22) \quad \hat{p}_{2k}^{(r)} = \frac{p_{2k}^{(r)}}{2k+1}, \quad \tilde{q}_{2k-1}^{(r)} = \frac{q_{2k-1}^{(r)}}{r+2k-2}.$$

Then, we have

$$(4.23) \quad \begin{aligned} D &= \frac{x^{m+n}}{\prod_{j=0}^n (2j+1) \prod_{k=1}^m (r-m-n+2k)} R_{m,n}^{(r+2)}, \\ D \begin{bmatrix} m \\ 1 \end{bmatrix} &= (-1)^{n+1} R_{m-1,n+1}^{(r)}, \quad D \begin{bmatrix} m+1 \\ 1 \end{bmatrix} = (-1)^n R_{m,n}^{(r)}, \\ D \begin{bmatrix} m \\ m+n+1 \end{bmatrix} &= \frac{(-1)^{n+1} x^{m+n-1}}{\prod_{j=0}^n (2j+1) \prod_{k=1}^{m-1} (r-m-n+2k)} R_{m-1,n}^{(r+1)}, \\ D \begin{bmatrix} m+1 \\ m+n+1 \end{bmatrix} &= \frac{(-1)^n x^{m+n-1}}{\prod_{j=0}^{n-1} (2j+1) \prod_{k=1}^m (r-m-n+2k)} R_{m,n-1}^{(r+1)}, \\ D \begin{bmatrix} m & m+1 \\ 1 & m+n+1 \end{bmatrix} &= R_{m-1,n}^{(r-1)}. \end{aligned}$$

It is easy to see that the bilinear relations (4.16) and (4.17) follow immediately from Jacobi's identity (4.18) by using Lemmas 4.4 and 4.5, respectively. We give the proof of Lemmas 4.4 and 4.5 in Appendix B. This completes the proof of Proposition 4.1 and thus our main result Theorem 1.1.

## §5. Discussion

As we mentioned in Section 2.3, the  $\tau$ -cocycles  $\phi_{k,l,m,n}$  are polynomials in  $\alpha_i$  and  $f_i$  and admit a determinant expression in terms of a generalized Jacobi-Trudi formula [21]. When specialized to the seed solution (3.1), we obtain a determinant formula for  $\phi_{0,0,m,n} = \phi_{m,n}$ , which are given as follows.

PROPOSITION 5.1. *Let  $g_k^{(l)}$  ( $k, l \in \mathbb{Z}$ ) be functions defined by*

$$(5.1) \quad \begin{aligned} g_{2k}^{(2l)} &= \frac{(-1)^k}{\xi_k} L_k^{(2s-1)}(t/2), & g_{2k+1}^{(2l)} &= \frac{\sqrt{t}}{2} \frac{(-1)^k}{\xi_{k+1}^\dagger} L_k^{(2s^\dagger)}(t/2), \\ g_{2k}^{(2l+1)} &= \frac{(-1)^k}{\xi_k^\dagger} L_k^{(2s^\dagger-1)}(t/2), & g_{2k+1}^{(2l+1)} &= \frac{\sqrt{t}}{2} \frac{(-1)^k}{\xi_{k+1}} L_k^{(2s)}(t/2), \end{aligned}$$

with

$$(5.2) \quad \xi_k = \xi_k(s) = \prod_{j=1}^k \left( s + \frac{j-1}{2} \right), \quad \xi_k^\dagger = \xi_k(s^\dagger),$$

where  $L_k^{(r)}(x)$  are the Laguerre polynomials and  $s^\dagger = 1/2 - s$ . Then,  $\phi_{m,n}$  under the specialization (3.1) are given by

$$(5.3) \quad \phi_{m,n} = N_{m,n} \det \left( g_{\lambda_j - j + i}^{(m+n+1-i)} \right)_{i,j=1}^{m+n},$$

where the partition  $\lambda$  and the normalization factor  $N_{m,n}$  are given by

$$(5.4) \quad \lambda = \begin{cases} (3m - n - 1, 3m - n - 4, \dots, 2n + 5, 2n + 2, 2n, 2n, \dots, 4, 4, 2, 2), & (m > n), \\ (3n - m, 3n - m - 3, \dots, 2m + 3, 2m, 2m, \dots, 4, 4, 2, 2), & (m \leq n), \end{cases}$$

and

$$(5.5) \quad N_{m,n} = \begin{cases} (-1)^{n(n+1)/2} c_m d_n \prod_{k=1}^n \hat{\zeta}_k \prod_{k=1}^m \zeta_k^\dagger \prod_{k=1}^{m-n-1} \hat{\zeta}_k^\dagger, & (m > n), \\ (-1)^{n(n+1)/2} c_m d_n \prod_{k=1}^n \hat{\zeta}_k \prod_{k=1}^m \zeta_k^\dagger \prod_{k=1}^{n-m} \zeta_k, & (m \leq n), \end{cases}$$

with

$$(5.6) \quad \zeta_k = \prod_{j=1}^k (s + j - 1), \quad \hat{\zeta}_k = \prod_{j=1}^k \left( s + \frac{2j-1}{2} \right),$$

respectively.

This gives a different expression for the rational solutions discussed in this paper. Studying the relationship between this formula and our result might be an interesting problem.

**Acknowledgements.** The authors would like to thank Prof. S. Okada for leading their attention to the universal characters. They also thank Prof. M. Noumi, Prof. Y. Yamada and Prof. H. Watanabe for useful suggestions and discussions.

## Appendix A. Determinant formula for the Umemura polynomials

In [9], Noumi and Yamada gave a determinant formula of Jacobi-Trudi type for the Umemura polynomials in terms of 2-reduced Schur functions. In this appendix, we give a brief review on this determinant formula, and show that it is recovered as a special case of our formula.

We normalize the polynomials  $U_{m,n}$  in Section 3 as

$$(A.1) \quad U_{m,n} = 2^{m(m+1)/2+n(n+1)/2} T_{m,n}.$$

Then we find that the functions  $T_n = T_{0,n}(t, s)$  are monic polynomials generated by the Toda equation

$$(A.2) \quad T_{n+1}T_{n-1} = t \left[ \left( \frac{d^2 T_n}{dt^2} \right) T_n - \left( \frac{dT_n}{dt} \right)^2 \right] + \frac{dT_n}{dt} T_n + \left( \frac{t}{8} - \frac{2s+1+3n}{4} \right) T_n^2,$$

with initial conditions  $T_{-1} = T_0 = 1$ . The polynomials  $T_n$  are called Umemura polynomials. It is easy to see that we have

$$(A.3) \quad T_{-1,n}(t, s) = T_{0,n}(t, s + 1/2).$$

Introducing  $\widehat{T}_n(t, s)$  by  $\widehat{T}_n(t, s) = T_{-n}(t, s)$ , we have the following proposition.

**PROPOSITION A.1.** *Let  $\widehat{T}_n = \widehat{T}_n(t, s)$  be a sequence of polynomials in  $t$  and  $s$  defined through the Toda equation*

$$(A.4) \quad \widehat{T}_{n+1}\widehat{T}_{n-1} = t \left[ \left( \frac{d^2 \widehat{T}_n}{dt^2} \right) \widehat{T}_n - \left( \frac{d\widehat{T}_n}{dt} \right)^2 \right] + \frac{d\widehat{T}_n}{dt} \widehat{T}_n + \left[ \frac{t}{8} - \frac{1}{2} \left( s + \frac{1}{2} \right) + \frac{3}{4} n \right] \widehat{T}_n^2,$$

with initial conditions  $\widehat{T}_0 = \widehat{T}_1 = 1$ . Then, the rational function

$$(A.5) \quad y = - \frac{\widehat{T}_{n+1}(t, s) \widehat{T}_n(t, s + 1/2)}{\widehat{T}_{n+1}(t, s + 1) \widehat{T}_n(t, s - 1/2)},$$

solves  $P_V$  with the parameters

$$(A.6) \quad \kappa_\infty = s, \quad \kappa_0 = s - n, \quad \theta = -n - 1.$$

The explicit formula for  $\widehat{T}_n$  was given by Noumi and Yamada, which is expressed in terms of the 2-reduced Schur functions.

PROPOSITION A.2. *Let  $S_n = S_n(t_1, t_2, \dots)$  for  $n \geq 0$  be the Schur function associated with a partition  $\lambda = (n, n-1, \dots, 2, 1)$ . Then, we have*

$$(A.7) \quad \widehat{T}_{n+1}(t, s) = N_n S_n$$

with

$$(A.8) \quad \begin{aligned} N_n &= 2^{-n(n+1)} (2n-1)!! (2n-3)!! \cdots 3!! 1!! \\ t_j &= \frac{t}{2} + \frac{-2s + n + 1}{j}. \end{aligned}$$

It is easy to verify that Proposition A.2 is recovered by putting  $m = 0$  for the solutions (1.8) with (1.9) in Theorem 1.1.

## Appendix B. Proof of Lemmas 4.4 and 4.5

We first note that the following contiguity relations hold by definition,

$$(B.1) \quad p_k^{(r)} - p_{k-1}^{(r)} = p_k^{(r-1)}, \quad q_k^{(r)} - q_{k-1}^{(r)} = q_k^{(r-1)},$$

and

$$(B.2) \quad (k+1)p_{k+1}^{(r)} = r p_k^{(r+1)} - x p_k^{(r+2)}, \quad (k+1)q_{k+1}^{(r)} = r q_k^{(r+1)} + x q_k^{(r+2)}.$$

Let us prove Lemma 4.4. Noticing that  $p_1^{(r)} = 1$  and  $p_k^{(r)} = 0$  for  $k < 0$ , we see that  $R_{m,n}^{(r)}$  can be rewritten as

$$(B.3) \quad R_{m,n}^{(r)} = \begin{vmatrix} q_1^{(r)} & q_0^{(r)} & \cdots & q_{-m-n+3}^{(r)} & q_{-m-n+2}^{(r)} & q_{-m-n+1}^{(r)} \\ q_3^{(r)} & q_2^{(r)} & \cdots & q_{-m-n+5}^{(r)} & q_{-m-n+4}^{(r)} & q_{-m-n+3}^{(r)} \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ q_{2m-1}^{(r)} & q_{2m-2}^{(r)} & \cdots & q_{m-n+1}^{(r)} & q_{m-n}^{(r)} & q_{m-n-1}^{(r)} \\ p_{n-m}^{(r)} & p_{n-m+1}^{(r)} & \cdots & p_{2n-2}^{(r)} & p_{2n-1}^{(r)} & p_{2n}^{(r)} \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ p_{-n-m+4}^{(r)} & p_{-n-m+5}^{(r)} & \cdots & p_2^{(r)} & p_3^{(r)} & p_4^{(r)} \\ p_{-n-m+2}^{(r)} & p_{-n-m+3}^{(r)} & \cdots & p_0^{(r)} & p_1^{(r)} & p_2^{(r)} \\ p_{-n-m}^{(r)} & p_{-n-m+1}^{(r)} & \cdots & p_{-2}^{(r)} & p_{-1}^{(r)} & p_0^{(r)} \end{vmatrix}.$$

Subtracting the  $(j-1)$ -th column from the  $j$ -th column of  $R_{m,n}^{(r+1)}$  for  $(j = m+n, m+n-1, \dots, 2)$  and using (B.1), we get

$$(B.4) \quad R_{m,n}^{(r+1)} = (-1)^m \begin{vmatrix} -q_1^{(r+1)} & q_1^{(r)} & \cdots & q_{-m-n+4}^{(r)} & q_{-m-n+3}^{(r)} \\ -q_3^{(r+1)} & q_3^{(r)} & \cdots & q_{-m-n+6}^{(r)} & q_{-m-n+5}^{(r)} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ -q_{2m-1}^{(r+1)} & q_{2m-1}^{(r)} & \cdots & q_{m-n+2}^{(r)} & q_{m-n+1}^{(r)} \\ p_{n-m}^{(r+1)} & p_{n-m+1}^{(r)} & \cdots & p_{2n-2}^{(r)} & p_{2n-1}^{(r)} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ p_{-n-m+4}^{(r+1)} & p_{-n-m+5}^{(r)} & \cdots & p_2^{(r)} & p_3^{(r)} \\ p_{-n-m+2}^{(r+1)} & p_{-n-m+3}^{(r)} & \cdots & p_0^{(r)} & p_1^{(r)} \end{vmatrix}.$$

From (B.3) and (B.4), we obtain Lemma 4.4.

We next prove Lemma 4.5. Subtracting the  $(i+1)$ -th column from the  $i$ -th column of  $R_{m,n}^{(r)}$  for  $(i = 1, 2, \dots, j, j = m+n-1, m+n-2, \dots, 1)$  and using (B.1), we get

$$(B.5) \quad R_{m,n}^{(r)} = \begin{vmatrix} q_1^{(r-m-n+1)} & q_0^{(r-m-n+2)} & \cdots & q_{-m-n+3}^{(r-1)} & q_{-m-n+2}^{(r)} \\ q_3^{(r-m-n+1)} & q_2^{(r-m-n+2)} & \cdots & q_{-m-n+5}^{(r-1)} & q_{-m-n+4}^{(r)} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ q_{2m-1}^{(r-m-n+1)} & q_{2m-2}^{(r-m-n+2)} & \cdots & q_{m-n+1}^{(r-1)} & q_{m-n}^{(r)} \\ (-1)^{m+n-1} p_{2n-1}^{(r-m-n+1)} & (-1)^{m+n-2} p_{2n-1}^{(r-m-n+2)} & \cdots & (-1)^1 p_{2n-1}^{(r-1)} & (-1)^0 p_{2n-1}^{(r)} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ (-1)^{m+n-1} p_3^{(r-m-n+1)} & (-1)^{m+n-2} p_3^{(r-m-n+2)} & \cdots & (-1)^1 p_3^{(r-1)} & (-1)^0 p_3^{(r)} \\ (-1)^{m+n-1} p_1^{(r-m-n+1)} & (-1)^{m+n-2} p_1^{(r-m-n+2)} & \cdots & (-1)^1 p_1^{(r-1)} & (-1)^0 p_1^{(r)} \end{vmatrix}.$$

Noticing that  $p_0^{(r)} = 1$  and  $p_k^{(r)} = 0$  for  $k < 0$ , we see that  $R_{m,n}^{(r)}$  can be rewritten as

$$(B.6) \quad R_{m,n}^{(r)} = \begin{vmatrix} q_1^{(r-m-n)} & q_0^{(r-m-n+1)} & \cdots & q_{-m-n+2}^{(r-1)} & q_{-m-n+1}^{(r)} \\ q_3^{(r-m-n)} & q_2^{(r-m-n+1)} & \cdots & q_{-m-n+4}^{(r-1)} & q_{-m-n+3}^{(r)} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ q_{2m-1}^{(r-m-n)} & q_{2m-2}^{(r-m-n+1)} & \cdots & q_{m-n}^{(r-1)} & q_{m-n-1}^{(r)} \\ (-1)^{m+n} p_{2n}^{(r-m-n)} & (-1)^{m+n-1} p_{2n}^{(r-m-n+1)} & \cdots & (-1)^1 p_{2n}^{(r-1)} & (-1)^0 p_{2n}^{(r)} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ (-1)^{m+n} p_2^{(r-m-n)} & (-1)^{m+n-1} p_2^{(r-m-n+1)} & \cdots & (-1)^1 p_2^{(r-1)} & (-1)^0 p_2^{(r)} \\ (-1)^{m+n} p_0^{(r-m-n)} & (-1)^{m+n-1} p_0^{(r-m-n+1)} & \cdots & (-1)^1 p_0^{(r-1)} & (-1)^0 p_0^{(r)} \end{vmatrix}.$$

We add the  $j$ -th column multiplied by  $(r-2+m+n-j)/x$  to the  $(j+1)$ -th column of (B.6) for  $(j = m+n, m+n-1, \dots, 1)$ . Then using (B.1) and

(B.2), we obtain

$$(B.7) \quad R_{m,n}^{(r)} = \prod_{j=0}^n (2j+1) \prod_{k=1}^m (r-m-n+2k-2)x^{-(m+n)} \\ \times \begin{vmatrix} q_1^{(r-m-n)} & q_1^{(r-m-n-1)} & \cdots & q_{-m-n+3}^{(r-3)} & q_{-m-n+2}^{(r-2)} \\ q_3^{(r-m-n)} & q_3^{(r-m-n-1)} & \cdots & q_{-m-n+5}^{(r-3)} & q_{-m-n+4}^{(r-2)} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ q_{2m-1}^{(r-m-n)} & q_{2m-1}^{(r-m-n-1)} & \cdots & q_{m-n+1}^{(r-3)} & q_{m-n}^{(r-2)} \\ (-1)^{m+n} p_{2n}^{(r-m-n)} & (-1)^{m+n} p_{2n+1}^{(r-m-n-1)} & \cdots & (-1)^2 p_{2n+1}^{(r-3)} & (-1)^1 p_{2n+1}^{(r-2)} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ (-1)^{m+n} p_2^{(r-m-n)} & (-1)^{m+n} p_3^{(r-m-n-1)} & \cdots & (-1)^2 p_3^{(r-3)} & (-1)^1 p_3^{(r-2)} \\ (-1)^{m+n} p_0^{(r-m-n)} & (-1)^{m+n} p_1^{(r-m-n-1)} & \cdots & (-1)^2 p_1^{(r-3)} & (-1)^1 p_1^{(r-2)} \end{vmatrix}.$$

Lemma 4.5 follows from (B.5), (B.6) and (B.7).

#### REFERENCES

- [1] M. Jimbo and T. Miwa, *Solitons and infinite dimensional Lie algebras*, Publ. RIMS. Kyoto Univ., **19** (1983), 943–1001.
- [2] K. Kajiwara and Y. Ohta, *Determinant structure of the rational solutions for the Painlevé II equation*, J. Math. Phys., **37** (1996), 4693–4704.
- [3] K. Kajiwara and T. Masuda, *On the Umemura polynomials for the Painlevé III equation*, Phys. Lett., **A 260** (1999), 462–467.
- [4] K. Kajiwara and Y. Ohta, *Determinant structure of the rational solutions for the Painlevé IV equation*, J. Phys. A: Math. Gen., **31** (1998), 2431–2446.
- [5] A. V. Kitaev, C. K. Law and J. B. McLeod, *Rational solutions of the fifth Painlevé equation*, Differential and Integral Equations, **7** (1994), 967–1000.
- [6] K. Koike, *On the decomposition of tensor products of the representations of the classical groups: by means of the universal characters*, Adv. Math., **74** (1989), 57–86.
- [7] M. Noumi, S. Okada, K. Okamoto and H. Umemura, *Special polynomials associated with the Painlevé equations II*, Proceedings of the Taniguchi Symposium 1997, Integrable Systems and Algebraic Geometry (M. H. Saito, Y. Shimizu and K. Ueno, eds.), World Scientific, Singapore (1998), pp. 349–372.
- [8] M. Noumi and Y. Yamada, *Symmetries in the fourth Painlevé equation and Okamoto polynomials*, Nagoya Math. J., **153** (1999), 53–86.
- [9] ———, *Umemura polynomials for the Painlevé V equation*, Phys. Lett., **A247** (1998), 65–69.
- [10] ———, *Higher order Painlevé equations of type  $A_l^{(1)}$* , Funkcial. Ekvac., **41** (1998), 483–503.
- [11] ———, *Affine Weyl groups, discrete dynamical systems and Painlevé equations*, Commun. Math. Phys., **199** (1998), 281–295.
- [12] K. Okamoto, *Studies on the Painlevé equations I, sixth Painlevé equation  $P_{VI}$* , Annali di Matematica pura ed applicata, **CXLVI** (1987), 337–381.



- [13] ———, *Studies on the Painlevé equations II, fifth Painlevé equation  $P_V$* , Japan J. Math., **13** (1987), 47–76.
- [14] ———, *Studies on the Painlevé equations III, second and fourth Painlevé equations,  $P_{II}$  and  $P_{IV}$* , Math. Ann., **275** (1986), 222–254.
- [15] ———, *Studies on the Painlevé equations IV, third Painlevé equation  $P_{III}$* , Funkcial. Ekvac., **30** (1987), 305–332.
- [16] M. Taneda, *Polynomials associated with an algebraic solution of the sixth Painlevé equation*, to appear in Jap. J. Math., **27** (2002).
- [17] H. Umemura, *Special polynomials associated with the Painlevé equations I*, to appear in the Proceedings of the Workshop on “Painlevé Transcendents” (CRM, Montreal, Canada, 1996).
- [18] ———, *Irreducibility of the Painlevé equations—Evolution in the past 100 years*, to appear in the Proceedings of the Workshop on “Painlevé Transcendents” (CRM, Montreal, Canada, 1996).
- [19] A. P. Vorob’ev, *On rational solutions of the second Painlevé equation*, Diff. Uravn., **1** (1965), 58–59.
- [20] H. Watanabe, *Solutions of the fifth Painlevé equation I*, Hokkaido Math. J., **24** (1995), 231–267.
- [21] Y. Yamada, *Determinant formulas for the generalized Painlevé equations of type A*, Nagoya Math. J., **156** (1999), 123–134.

Tetsu Masuda  
*Department of Mathematics*  
*Kobe University*  
*Rokko, Kobe, 657-8501*  
*Japan*  
masuda@math.kobe-u.ac.jp

Yasuhiro Ohta  
*Department of Applied Mathematics, Faculty of Engineering*  
*Hiroshima University*  
*1-4-1 Kagamiyama*  
*Higashi-Hiroshima, 739-8527*  
*Japan*  
ohta@kurims.kyoto-u.ac.jp

Kenji Kajiwara  
*Graduate School of Mathematics*  
*Kyushu University*  
*6-10-1 Hakozaki*  
*Higashi-ku, Fukuoka, 812-8512*  
*Japan*  
kaji@math.kyushu-u.ac.jp