

RIEMANNIAN FOLIATIONS WITH PARALLEL CURVATURE

ROBERT A. BLUMENTHAL

§ 1. Introduction

Let M be a smooth compact manifold and let \mathcal{F} be a smooth codimension q Riemannian foliation of M . Let $T(M)$ be the tangent bundle of M and let $E \subset T(M)$ be the subbundle tangent to \mathcal{F} . We may regard the normal bundle $Q = T(M)/E$ of \mathcal{F} as a subbundle of $T(M)$ satisfying $T(M) = E \oplus Q$. Let g be a smooth Riemannian metric on Q invariant under the natural parallelism along the leaves of \mathcal{F} . This is equivalent to the existence of a bundle-like metric [16] and to the existence of a transverse $O(q)$ -structure [5]. Recall that a connection ∇ on Q is basic if the induced parallel translation along a path lying in a leaf of \mathcal{F} agrees with the natural parallelism along the leaves and that such a connection is characterized by the condition that $\nabla_x Y = [X, Y]_q$ for all vector fields X tangent to E and Y tangent to Q where $[X, Y]_q$ denotes the Q -component of the Lie bracket of X and Y [3]. The torsion of ∇ is the tensor field of type $(1, 2)$ on M defined by $T(X, Y) = \nabla_x Y_q - \nabla_y X_q - [X, Y]_q$ where X and Y are vector fields on M . There is a unique torsion-free metric-preserving basic connection ∇ on Q [9], [11] defined as follows. Let $x \in M$. Let $f: U \rightarrow V$ be a submersion whose level sets are the leaves of $\mathcal{F}|U$ where U is a neighborhood of x in M and V is an open set in R^q . There is a unique Riemannian metric \bar{g} on V such that $f^*(\bar{g}) = g|U$. Let $\bar{\nabla}$ be the Riemannian connection on V . Then $\nabla|U = f^{-1}(\bar{\nabla})$. It is natural to study the relationship between the curvature of ∇ and the structure of the foliated manifold (M, \mathcal{F}) .

In the present work we study the case of parallel curvature, that is $\nabla R = 0$ where $R(X, Y)Z$ denotes the curvature tensor of ∇ .

Let \mathcal{F} be a Riemannian foliation with parallel curvature of a compact manifold M .

THEOREM 1. *Let \tilde{M} be the universal cover of M and let $\tilde{\mathcal{F}}$ be the lift*

Received March 15, 1982.

of \mathcal{F} to \tilde{M} . Then \tilde{M} fibers over a simply connected Riemannian symmetric space with the leaves of $\tilde{\mathcal{F}}$ as fibers.

Let $p \in M$. Let π_p be a two-dimensional subspace of Q_p and let $\{X, Y\}$ be an orthonormal basis of π_p . The (transverse) sectional curvature of π_p is defined by $K(\pi_p) = -g_p(R(X, Y)X, Y)$ and depends only on π_p . If $K(\pi_p) > 0$ (respectively, $\leq 0, \geq 0$) for all two-dimensional subspaces $\pi_p \subset Q_p$ and all $p \in M$, we say that (M, \mathcal{F}) has positive (respectively, non-positive, non-negative) sectional curvature.

COROLLARY 1. *If (M, \mathcal{F}) has non-positive sectional curvature, then \tilde{M} is diffeomorphic to a product $\tilde{L} \times R^q$ where \tilde{L} is the (common) universal cover of the leaves of \mathcal{F} and the leaves of $\tilde{\mathcal{F}}$ are identified with the sets $\tilde{L} \times \{x\}$, $x \in R^q$.*

COROLLARY 2. *If $\pi_1(M)$ is finite, then (M, \mathcal{F}) has non-negative sectional curvature and all the leaves of \mathcal{F} are compact with finite holonomy.*

Remark. If \mathcal{F} is a flat Riemannian foliation of the compact manifold M , then (M, \mathcal{F}) has zero curvature and zero sectional curvature and so, by Corollary 1, $\tilde{M} \cong \tilde{L} \times R^q$ and $\tilde{\mathcal{F}}$ is the product foliation. A theorem of G. Reeb [14] states that if \mathcal{F} is a codimension one foliation of a compact manifold M defined by a closed nonsingular one-form, then $\tilde{M} \cong \tilde{L} \times R$ and $\tilde{\mathcal{F}}$ is the product foliation. It is easy to see that such a codimension one foliation admits a flat Riemannian structure and so we obtain Reeb's theorem from Corollary 1.

A differential r -form ω on M is called base-like if on each coordinate neighborhood U with coordinates $(x^1, \dots, x^k, y^1, \dots, y^q)$ respecting the foliation \mathcal{F} , the local expression of ω is of the form

$$\sum_{1 \leq i_1 < \dots < i_r \leq q} a_{i_1, \dots, i_r}(y^1, \dots, y^q) dy^{i_1} \wedge \dots \wedge dy^{i_r}$$

[16], [17]. Since the exterior derivative of a base-like form is again base-like, one can construct the base-like cohomology algebra $H^*(M, \mathcal{F}) = \sum_{r \geq 0} H^r(M, \mathcal{F})$ of the foliated manifold (M, \mathcal{F}) . For each $r \geq 0$, let $\beta_r(M, \mathcal{F})$ be the dimension of $H^r(M, \mathcal{F})$.

THEOREM 2. *If (M, \mathcal{F}) has positive sectional curvature, then $\beta_1(M, \mathcal{F}) = 0$.*

Recall from [13] the definition of the growth of a leaf L of \mathcal{F} . Pick a Riemannian metric on M (bundle-like or not) and restrict to obtain a

Riemannian metric on L . Let $p \in L$ and define the growth function of L at p by $g_p(r) = \text{vol}(B_p(r))$ where $B_p(r)$ denotes the open ball in L of radius r centered at p . The growth type of L is then defined to be the growth type of the function $g_p: R^+ \rightarrow R^+$ and is independent of the choice of metric on M and of $p \in L$ [13].

THEOREM 3. *The growth type of each leaf of \mathcal{F} is dominated by the growth type of $\pi_1(M)$.*

§ 2. Proofs

Let $\pi: F(Q) \rightarrow M$ be the frame bundle of Q , let ω be the connection form on $F(Q)$ associated to ∇ , and let $H \subset T(F(Q))$ be the corresponding horizontal distribution. Let $\{(U_\alpha, f_\alpha, g_{\alpha\beta})\}_{\alpha, \beta \in A}$ be an R^q -cocycle defining \mathcal{F} . Let $F(R^q)$ be the frame bundle of R^q . Then $\{(\pi^{-1}(U_\alpha), f_{\alpha*}, g_{\alpha\beta*})\}_{\alpha, \beta \in A}$ is an $F(R^q)$ -cocycle on $F(Q)$ and defines a codimension $q(q+1)$ foliation \mathcal{F}' of $F(Q)$. Let $E' \subset T(F(Q))$ be the integrable distribution whose integral manifolds are the leaves of \mathcal{F}' . Since ∇ is basic we have $E' \subset H$ [12]. Let θ be the R^q -valued one-form on $F(Q)$ defined by $\theta_u(Y) = u^{-1}(\pi_{*u}(Y)_Q)$ for $u \in F(Q)$, $Y \in T_u(F(Q))$ where $u: R^q \rightarrow Q_{\pi(u)}$ denotes the vector space isomorphism which sends the standard basis of R^q to the frame u of $Q_{\pi(u)}$. The torsion form of ∇ is the R^q -valued two-form Θ on $F(Q)$ defined by $\Theta_u(X, Y) = (d\theta)_u(X_H, Y_H)$ for $u \in F(Q)$ and $X, Y \in T_u(F(Q))$. Since ∇ is the Riemannian basic connection, we have $\Theta = 0$.

For each $h \in R^q$ let $B(h)$ be the unique horizontal vector field on $F(Q)$ satisfying $\pi_{*u}(B(h)_u) = u(h)$ for all $u \in F(Q)$. Let $E^i = B(e_i)$ for $i = 1, \dots, q$ where $\{e_1, \dots, e_q\}$ is the standard basis of R^q and let $Q' \subset T(F(Q))$ be the q -plane bundle spanned by E^1, \dots, E^q . Then $H = E' \oplus Q'$ and so $T(F(Q)) = E' \oplus Q' \oplus V$ where V is the bundle of vertical vectors. Hence we may regard $Q' \oplus V$ as the normal bundle of \mathcal{F}' . Let E_h^k be the $q \times q$ matrix with a 1 in the h^{th} column and k^{th} row and 0 elsewhere and let $\sigma(E_h^k)$ be the corresponding fundamental vector field on $F(Q)$. Then $\{E^i, \sigma(E_h^k): i, h, k = 1, \dots, q\}$ is a trivialization of the normal bundle of \mathcal{F}' . Recall that a vector field Y on $F(Q)$ which is normal to \mathcal{F}' is said to be parallel along the leaves to \mathcal{F}' if $(f_{\alpha*})_*(Y|\pi^{-1}(U_\alpha))$ is a well-defined vector field on $f_{\alpha*}(\pi^{-1}(U_\alpha)) \subset F(R^q)$ for each $\alpha \in A$. This is equivalent to Y being invariant under the natural parallelism along the leaves and is characterized by the condition that $[X, Y]$ is tangent to \mathcal{F}' whenever X is a vector field tangent to \mathcal{F}' [5]. Since the fundamental vector fields

on $F(R^q)$ are preserved by the maps $g_{\alpha\beta}$, it follows that the vector fields $\sigma(E_h^k)$ are parallel along the leaves to \mathcal{F}' . Since ∇ is the Riemannian basic connection, it is transversely projectable and hence $[X, E^i]$ is tangent to \mathcal{F}' for $i = 1, \dots, q$ whenever X is a vector field tangent to \mathcal{F}' [10], and so E^1, \dots, E^q are parallel along the leaves to \mathcal{F}' . Hence $\{E^i, \sigma(E_h^k): i, h, k = 1, \dots, q\}$ is an e -structure for \mathcal{F}' [5].

Let $u \in F(Q)$ and let $X \in T_u(F(Q))$. Then there is a unique expression $X = X_{E'} + X_{Q'} + X_V$. Thus $\theta_u(X) = \theta_u(X_{Q'}) = \theta_u(B(h)_u) = h$ for some $h \in R^q$ and so $X_{Q'} = B(\theta_u(X))_u$. Also $\omega_u(X) = \omega_u(X_V) = \omega_u(\sigma(A)_u) = A$ for some $A \in \mathfrak{gl}(q, R)$ and so $X_V = \sigma(\omega_u(X))_u$. Letting $X = [E^i, E^j]_u$, we obtain

$$[E^i, E^j]_{Q'_u} = B(\theta_u([E^i, E^j]_u))_u \quad \text{and} \quad [E^i, E^j]_{V_u} = \sigma(\omega_u([E^i, E^j]_u))_u.$$

Since

$$\begin{aligned} -\theta_u([E^i, E^j]_u) &= E_u^i \theta(E^j) - E_u^j \theta(E^i) - \theta_u([E^i, E^j]_u) \\ &= (d\theta)_u(E_u^i, E_u^j) = \Theta_u(E_u^i, E_u^j) = 0, \end{aligned}$$

we have $[E^i, E^j]_{Q'} = 0$. Let Ω be the curvature form of ∇ and write $\Omega = \sum_{h,k=1}^q \Omega_k^h E_h^k$ where the Ω_k^h are two-forms on $F(Q)$. Since

$$\begin{aligned} -\omega_u([E^i, E^j]_u) &= E_u^i \omega(E^j) - E_u^j \omega(E^i) - \omega_u([E^i, E^j]_u) \\ &= (d\omega)_u(E_u^i, E_u^j) = \Omega_u(E_u^i, E_u^j), \end{aligned}$$

we have that

$$[E^i, E^j]_V = - \sum_{h,k=1}^q \Omega_k^h(E^i, E^j) \sigma(E_h^k).$$

Let $u_0 \in F(Q)$ and let $P(u_0) = \{u \in F(Q) : u \text{ can be joined to } u_0 \text{ by a horizontal curve}\}$, the holonomy bundle through u_0 . Let $u \in P(u_0)$ and let $p = \pi(u) \in M$. Let $c : (-\varepsilon, \varepsilon) \rightarrow M$ be a smooth curve with $c(0) = p$, and let $c^* : (-\varepsilon, \varepsilon) \rightarrow F(Q)$ be the unique horizontal lift of c satisfying $c^*(0) = u$. Fix $1 \leq i, j, l \leq q$ and let $X(t) = c^*(t)_i$, $Y(t) = c^*(t)_j$, and $Z(t) = c^*(t)_l$. Then X, Y and Z are normal vector fields along c which are parallel along c and hence, since R is parallel, $\nabla_{\dot{c}(t)} R(X, Y)Z = 0$. Defining $f : (-\varepsilon, \varepsilon) \rightarrow R^q$ by $f(t) = c^*(t)^{-1}(R(X, Y)Z)_{c^*(t)}$, we have that $\nabla_{\dot{c}(t)} R(X, Y)Z = u(f'(t))$ and so $f'(0) = 0$. But

$$(R(X, Y)Z)_{c^*(t)} = c^*(t)(\Omega_{c^*(t)}(E_{c^*(t)}^i, E_{c^*(t)}^j)c^*(t)^{-1}(c^*(t)_l))$$

and hence $f(t) = \Omega_{c^*(t)}(E_{c^*(t)}^i, E_{c^*(t)}^j) \cdot e_l = l^{\text{th}}$ column of $\Omega_{c^*(t)}(E_{c^*(t)}^i, E_{c^*(t)}^j)$. Thus $\dot{c}^*(0)(l^{\text{th}}$ column of $\Omega(E^i, E^j) = f'(0) = 0$ and hence

$$W(l^{\text{th}} \text{ column of } \Omega(E^i, E^j)) = 0$$

for all $W \in H_u$. Thus $\Omega_k^h(E^i, E^j)$ is constant on $P(u_0)$ for each $1 \leq h, k \leq q$.

Let $\Phi(u_0)$ be the holonomy group with reference point u_0 ; that is, $\Phi(u_0) = \{A \in \text{gl}(q, R) : u_0 \text{ and } u_0A \text{ can be joined by a horizontal curve}\}$. Then $P(u_0)$ is a reduced bundle with structure group $\Phi(u_0)$ such that the natural parallelism along the leaves of \mathcal{F} carries elements of $P(u_0)$ to elements of $P(u_0)$ and so $P(u_0)$ is a transverse $\Phi(u_0)$ -structure for \mathcal{F} [5] and ω is reducible to a basic connection in $P(u_0)$. Let $V' \subset T(P(u_0))$ be the subbundle consisting of vectors tangent to the fibers of $P(u_0)$. Then $T(P(u_0)) = E' \oplus Q' \oplus V'$ and \mathcal{F}' is a foliation of $P(u_0)$ whose tangent bundle is E' . Let A_1, \dots, A_r be a basis of the Lie algebra of $\Phi(u_0)$. Then $\{E^1, \dots, E^q, \sigma(A_1), \dots, \sigma(A_r)\}$ is an e -structure for \mathcal{F}' on $P(u_0)$. On $P(u_0)$ we have

$$\begin{aligned} [E^i, E^j]_{Q'} &= 0 \\ [E^i, E^j]_{V'} &= \sum_{k=1}^r f_{ij}^k \sigma(A_k) \\ [\sigma(A_i), \sigma(A_j)] &= \sum_{k=1}^r c_{ij}^k \sigma(A_k) \\ [\sigma(A_i), E^j] &= B(A_i \cdot e_j) = \sum_{k=1}^q b_{ij}^k E^k \end{aligned}$$

where f_{ij}^k, c_{ij}^k , and b_{ij}^k are constants.

Let G be the unique simply connected Lie group with Lie algebra g spanned by elements $Z_1, \dots, Z_q, B_1, \dots, B_r$ satisfying

$$\begin{aligned} [Z_i, Z_j] &= \sum_{k=1}^r f_{ij}^k B_k \\ [B_i, B_j] &= \sum_{k=1}^r c_{ij}^k B_k \\ [B_i, Z_j] &= \sum_{k=1}^q b_{ij}^k Z_k. \end{aligned}$$

Let h be the subalgebra of g spanned by B_1, \dots, B_r and let m be the subspace of g spanned by Z_1, \dots, Z_q . Then $g = h \oplus m$, $[h, h] \subset h$, $[h, m] \subset m$, and $[m, m] \subset h$. Let $X \in g$ and write X uniquely as $X = Y + Z$ where $Y \in h$, $Z \in m$. Let $\tau(X) = Y - Z$. Then τ is an automorphism of g and τ^2 is the identity. Since G is simply connected there is an automorphism $F : G \rightarrow G$ such that $F_* = \tau$. Let H be the identity component of the subgroup of G fixed by F . Then H is a closed Lie subgroup of G and the triple (G, H, F) is a symmetric space.

Let $u \in P(u_0)$. Since $E^1, \dots, E^q, \sigma(A_1), \dots, \sigma(A_r)$ are parallel along the leaves to \mathcal{F}' , there is a neighborhood W of u in $P(u_0)$ and a smooth submersion $\bar{f}: W \rightarrow G$ such that

$$\begin{aligned} \text{kernel}(\bar{f}_{*y}) &= E'_y \\ \bar{f}_{*y}(E_y^i) &= Z_{i\bar{f}(y)}, \quad i = 1, \dots, q \\ \bar{f}_{*y}(\sigma(A_j)_y) &= B_{j\bar{f}(y)}, \quad j = 1, \dots, r \end{aligned}$$

for all $y \in W$. Let $U = \pi(W)$, a neighborhood of $\pi(u)$ in M . Then \bar{f} induces a smooth submersion $f: U \rightarrow G/H$ such that $\text{kernel}(f_{*p}) = E_p$ for all $p \in U$ and the diagram

$$\begin{array}{ccc} W & \xrightarrow{\bar{f}} & G \\ \pi \downarrow & & \downarrow \\ U & \xrightarrow{f} & G/H \end{array}$$

commutes. Let $\bar{\theta}$ be the canonical left-invariant \mathfrak{g} -valued one-form on G and let $\bar{\theta}_h$ be the \mathfrak{h} -component of $\bar{\theta}$. Then $\bar{\theta}_h$ defines a G -invariant connection in the principal H -bundle $G \rightarrow G/H$ which induces the canonical linear connection on the symmetric space G/H [8], and $\bar{f}^*\bar{\theta}_h = \omega$. Thus if $F(G/H)$ is the frame bundle of G/H and $\bar{\omega}$ is the connection form on $F(G/H)$ corresponding to the canonical linear connection on G/H , we have that $(f_*)^*\bar{\omega} = \omega$ on $F(Q)|_U$. Thus we can find a G/H -cocycle $\{(U_\alpha, f_\alpha, g_{\alpha\beta})\}_{\alpha, \beta \in A}$ defining \mathcal{F} such that $(f_{\alpha_*})^*\bar{\omega} = \omega$ on $F(Q)|_{U_\alpha}$. If $U_\alpha \cap U_\beta \neq \emptyset$ then, since $(f_{\alpha_*})^*\bar{\omega} = \omega = (f_{\beta_*})^*\bar{\omega}$ on $F(Q)|_{U_\alpha \cap U_\beta}$, it follows that $(g_{\alpha\beta_*})^*\bar{\omega} = \bar{\omega}$ on $F(G/H)|_{f_\beta(U_\alpha \cap U_\beta)}$. Without loss of generality we may assume that $U_\alpha \cap U_\beta$ is connected whenever it is non-empty. Hence, since $\bar{\omega}$ is a complete analytic linear connection on the simply connected analytic manifold G/H , $g_{\alpha\beta}$ extends to an affine isomorphism of G/H [8].

Hence \mathcal{F} is transversely homogeneous. The foliated manifold (\tilde{M}, \mathcal{F}) admits a complete bundle-like metric, and so we have that \mathcal{F} is regular [1]. Hence the space of leaves \tilde{M}/\mathcal{F} is a complete, Riemannian, Hausdorff manifold and the natural projection $f: \tilde{M} \rightarrow \tilde{M}/\mathcal{F}$ is a fibration [16]. Let N denote the Riemannian manifold \tilde{M}/\mathcal{F} . Since the metric on N is induced by the bundle-like metric on \tilde{M} , it follows that the curvature tensor field of N is parallel. Thus N is a complete, simply connected, Riemannian locally symmetric space and hence is Riemannian symmetric [8] and so Theorem 1 is proved.

If (M, \mathcal{F}) has non-positive sectional curvature, then N has non-positive sectional curvature. Since N is complete and simply connected we have that N is diffeomorphic to R^q [8] and hence the fibration $f: \tilde{M} \rightarrow N$ is a product, proving Corollary 1. See [2] for similar results in the flat non-Riemannian case.

If $\pi_1(M)$ is finite, then \tilde{M} is compact and so the leaves of $\tilde{\mathcal{F}}$ are compact. Thus all the leaves of \mathcal{F} are compact. Hence \mathcal{F} is a closed metric foliation and so the holonomy group of each leaf is finite [15]. Since N is compact it has non-negative sectional curvature [18] and so (M, \mathcal{F}) has non-negative sectional curvature, proving Corollary 2.

We now prove Theorem 2. Each covering transformation σ of \tilde{M} induces an isometry $\Psi(\sigma)$ of N . We thus obtain a homomorphism $\Psi: \pi_1(M) \rightarrow I(N)$ where $I(N)$ denotes the isometry group of N such that $\Psi(\sigma) \circ f = f \circ \sigma$ for all $\sigma \in \pi_1(M)$. Let Σ be the image of Ψ in $I(N)$ and let K be the closure of Σ . For each $r \geq 0$ let $A_K^r(N)$ be the space of K -invariant r -forms on N and let $A^r(M, \mathcal{F})$ be the space of base-like r -forms on M . Let $\eta \in A_K^r(N)$. Then $f^*\eta$ is an r -form on \tilde{M} which is base-like with respect to $\tilde{\mathcal{F}}$. Since η is Σ -invariant it follows that $f^*\eta$ is $\pi_1(M)$ -invariant and hence there exists a unique $\omega \in A^r(M, \mathcal{F})$ such that $f^*\eta = p^*\omega$ where $p: \tilde{M} \rightarrow M$ is the covering projection. Conversely, suppose $\omega \in A^r(M, \mathcal{F})$. Then $p^*\omega$ is base-like and hence there exists a unique r -form η on N such that $p^*\omega = f^*\eta$. Since $p^*\omega$ is $\pi_1(M)$ -invariant it follows that η is Σ -invariant and hence K -invariant. Thus $\eta \in A_K^r(N)$. We have constructed an isomorphism of cochain complexes

$$\begin{array}{ccccc} A^0(M, \mathcal{F}) & \xrightarrow{d} & A^1(M, \mathcal{F}) & \xrightarrow{d} & \dots \\ \updownarrow & & \updownarrow & & \\ A_K^0(N) & \xrightarrow{d} & A_K^1(N) & \xrightarrow{d} & \dots \end{array}$$

and hence we obtain an isomorphism in cohomology $H^*(M, \mathcal{F}) \rightarrow H_K^*(N)$. Since (M, \mathcal{F}) has positive sectional curvature it follows that N has positive sectional curvature. Thus N is compact [18] and so K is compact. Hence the inclusion of the algebra of K -invariant forms on N into the algebra of differential forms on N induces an injection $H_K^*(N) \rightarrow H^*(N)$ [4], [6]. Thus $H^*(M, \mathcal{F})$ is isomorphic to a subalgebra of $H^*(N)$. Since N is simply connected we have that $H^1(N) = 0$ and hence $H^1(M, \mathcal{F}) = 0$.

Let $\Sigma \backslash N$ be the orbit space of Σ . To prove Theorem 3 we construct a map $h: M/\mathcal{F} \rightarrow \Sigma \backslash N$ and apply a result in [1]. Let L be a leaf of \mathcal{F} .

Choose a leaf \tilde{L} of \mathcal{F} such that $p(\tilde{L}) = L$. Let $x = f(\tilde{L}) \in N$. Then the orbit of x under Σ depends only on L and we denote it by $h(L)$. The growth type of a leaf L of \mathcal{F} is dominated by the growth type of $h(L)$ [1] which in turn is dominated by the growth type of Σ . Hence, since Σ is a homomorphic image of $\pi_1(M)$, we have that the growth type of L is dominated by the growth type of $\pi_1(M)$.

§ 3. Examples

EXAMPLE 1. Let G be a compact connected Lie group of dimension q and let \mathfrak{g} be the Lie algebra of G . Let M be a compact manifold and suppose ω is a smooth \mathfrak{g} -valued one-form of rank q on M satisfying $d\omega + \frac{1}{2}[\omega, \omega] = 0$. Then ω defines a smooth codimension q foliation \mathcal{F} on M which is a Lie foliation modeled on G [7]. Let \langle, \rangle be a bi-invariant Riemannian metric on G . Then \langle, \rangle induces on (M, \mathcal{F}) a Riemannian structure with parallel curvature and non-negative sectional curvature.

EXAMPLE 2. Let H be a Lie subgroup of the n -dimensional torus T^n . Then the foliation \mathcal{F} of T^n by the cosets of H admits a Riemannian structure with vanishing curvature and $\beta_1(T^n, \mathcal{F}) \neq 0$.

EXAMPLE 3. Let M be the unit tangent bundle of the two-holed torus T_2 . Let \mathcal{F} be the foliation of M by the circle fibers. Then \mathcal{F} admits a Riemannian structure with parallel curvature and negative sectional curvature and $\beta_1(M, \mathcal{F}) \neq 0$.

EXAMPLE 4. Let $\Phi : \pi_1(T_2) \rightarrow SO(3)$ be a homomorphism whose image is dense in $SO(3)$. This defines a left action of $\pi_1(T_2)$ on S^2 . Let H be the universal cover of T_2 . Then H is a principal $\pi_1(T_2)$ -bundle over T_2 . Let $M = H \times_{\pi_1(T_2)} S^2$ be the associated bundle with fiber S^2 . The foliation of $H \times S^2$ whose leaves are the sets $H \times \{x\}$, $x \in S^2$ passes to a foliation \mathcal{F} of M all of whose leaves are dense. Since $\pi_1(T_2)$ acts on S^2 by isometries, (M, \mathcal{F}) admits a Riemannian structure with parallel curvature and positive sectional curvature. Also $\beta_1(M, \mathcal{F}) = 0$.

EXAMPLE 5. Define a left action of the integers Z on S^2 by letting the generator act as

$$\begin{pmatrix} \cos 2\pi\alpha & \sin 2\pi\alpha & 0 \\ -\sin 2\pi\alpha & \cos 2\pi\alpha & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

where $0 < \alpha < 1$ is irrational. Let $M = R \times_z S^2$. Then M is an S^2 -bundle over S^1 with a codimension 2 foliation \mathcal{F} . The foliated manifold (M, \mathcal{F}) admits a Riemannian structure with parallel curvature and positive sectional curvature and $\beta_1(M, \mathcal{F}) = 0$. There are exactly two compact leaves. If L is a non-compact leaf then \bar{L} is diffeomorphic to T^2 . The foliation of \bar{L} by the leaves of \mathcal{F} is an irrational slope foliation and hence is a flat Riemannian foliation and $\beta_1(\bar{L}, \mathcal{F}) \neq 0$.

REFERENCES

- [1] R. A. Blumenthal, Transversely homogeneous foliations, *Ann. Inst. Fourier*, **29** (1979), 143–158.
- [2] —, Foliated manifolds with flat basic connection, *J. Differential Geom.*, **16** (1981), 401–406.
- [3] R. Bott, Lectures on characteristic classes and foliations (notes by L. Conlon), *Lecture Notes in Math.*, no. 279, Springer-Verlag, New York, 1972, 1–80.
- [4] C. Chevalley and S. Eilenberg, Cohomology theory of Lie groups and Lie algebras, *Trans. Amer. Math. Soc.*, **63** (1948), 85–124.
- [5] L. Conlon, Transversally parallelizable foliations of codimension two, *Trans. Amer. Math. Soc.*, **194** (1974), 79–102.
- [6] W. Greub, S. Halperin and R. Vanstone, *Connections, curvature, and cohomology*, vol. II, Academic Press, New York, 1973.
- [7] R. Hermann, On the differential geometry of foliations, *Ann. of Math.*, **72** (1960), 445–457.
- [8] S. Kobayashi and K. Nomizu, *Foundations of differential geometry*, vol. I, II. Interscience Tracts in Pure and Appl. Math., 15, Interscience, New York, 1963.
- [9] C. Lazarov and J. Pasternack, Secondary characteristic classes for Riemannian foliations, *J. Differential Geom.*, **11** (1976), 365–385.
- [10] F. Molino, Connexions et G-structures sur les variétés feuilletées, *Bull. Sci. Math.*, **92** (1968), 59–63.
- [11] —, Etude des feuilletages transversalement complets et applications, *Ann. Sci. École Norm. Sup.*, **10** (1977), 289–307.
- [12] —, Feuilletages et classes caractéristiques, *Symposia Mathematica*, **X** (1972), 199–209.
- [13] J. F. Plante, Foliations with measure preserving holonomy, *Ann. of Math.*, **102** (1975), 327–361.
- [14] G. Reeb, Sur certaines propriétés topologiques des variétés feuilletées, *Actualités Sci. Indust.*, no. 1183, Hermann, Paris, 1952.
- [15] B. Reinhart, Closed metric foliations, *Michigan Math. J.*, **8** (1961), 7–9.
- [16] —, Foliated manifolds with bundle-like metrics, *Ann. of Math.*, **69** (1959), 119–132.
- [17] —, Harmonic integrals on foliated manifolds, *Amer. J. Math.*, **81** (1959), 529–536.
- [18] J. Wolf, *Spaces of constant curvature*, McGraw-Hill, New York, 1967.

Department of Mathematics
Saint Louis University
St. Louis, Missouri 63103

