

**CORRECTIONS TO 'CHARACTER THEORY OF FINITE
GROUPS WITH TRIVIAL INTERSECTION
SUBSETS (Vol 27, 515—524)**

JOHN H. WALTER

1. The conditions (TI 1) and (TI 2) are stated for $H = \mathfrak{N}_G(\mathbf{D})$ and henceforth in the paper H is understood to be $\mathfrak{N}_G(\mathbf{D})$ when \mathbf{D} is taken to be a T.I. subset of G . Also in the definition of T.I. subset the condition is that $\mathbf{D} \cap \mathbf{D}^g \neq \phi$ where ϕ is the empty set.

2. Just before formula (11), the symbol should read

$$\{\varepsilon(\tau_i)\xi_i \mid \varepsilon(\tau_i)\xi_i^{\tau_i} = \varepsilon\}.$$

3. In the statement of Proposition 8, the penultimate sentence should read: '*If \mathbf{D} contains a section $\mathfrak{S}_H(P)$ of a p -element P belonging to a defect group V of \mathfrak{B}^G , then $\mathfrak{S}_G(\mathfrak{C}_G, (\mathbf{D}^G, \mathfrak{B}^G))$ contains all characters of zero height in \mathfrak{B}^G .*' It is required to know that $\varepsilon(R) \neq 0$ in the proof for a character ε of \mathfrak{B}^G for an appropriate p -regular element R in order to have $\varepsilon(PR) \neq 0$ where $PR \in \mathbf{D}$. The assumption of zero height is need to justify this step.

The author is grateful to Professor J.G. Thompson and Warren Wong for their comments. There is a substantial overlap with the article and Wong's 'Exceptional Character Theory and the Theory of Blocks', *Math. Zeit.*, 91, 363-379 (1966).