

GAUSSIAN MEASURE ON A BANACH SPACE AND ABSTRACT WINER MEASURE

HIROSHI SATO

In this paper, we shall show that any Gaussian measure on a separable or reflexive Banach space is an abstract Wiener measure in the sense of L. Gross [1] and, for the proof of that, establish the Radon extensibility of a Gaussian measure on such a Banach space. In addition, we shall give some remarks on the support of an abstract Wiener measure.

An abstract Wiener measure is a σ -extension in a Banach space X of the canonical Gaussian cylinder measure $\mu_{\mathfrak{X}}$ of a real separable Hilbert space \mathfrak{X} which is contained in X densely. The idea of the abstract Wiener measure coincides with that of *the White Noise* (T. Hida [13]) and plays an important role not only in the theory of probability but in the theory of functional analysis (T. Hida [13], Y. Umemura [12], I.E. Segal [4, 5], L. Gross [3] and Yu. L. Daletskii [16]).

We shall show first that any Gaussian measure on a separable or reflexive Banach space can be extended to a Radon measure on the strong topological σ -algebra (Theorem 1). With the same idea of the proof of Theorem 1, we can prove that this result is true for any probability measure on a Banach space, the finite dimensional distribution of which is Radon.

Utilizing the above result, we shall restrict the support of a Gaussian measure to a separable subspace which is explicitly constructed. Furthermore, constructing a suitable Hilbert subspace of the support, we shall show that any Gaussian measure on such a Banach space is an abstract Wiener measure (Theorem 2). L. Gross [1] showed that there exists an abstract Wiener measure on any separable Banach space. Our result shows that any *given* Gaussian measure on a separable or reflexive Banach space is an abstract Wiener measure. This means that the study of a Gaussian measure

on such a Banach space can be reduced to that of an abstract Wiener measure on a separable Banach space, and clears a new way for the investigation of a Gaussian measure on a Banach space, and makes the study of an abstract Wiener measure more meaningful.

As a corollary of Theorem 2, we shall show that the canonical Gaussian cylinder measure of a *nonseparable* Hilbert space can not be extended to a σ -additive measure in any Banach space.

Before stating the remaining results in this paper, we establish terminology and notation.

Let X be a real Banach space, X^* be its topological dual space and $\xi(x)$, ($\xi \in X^*$, $x \in X$), be the natural linear form. A cylinder set in X is a set of the form

$$C = \{x \in X: (\xi_1(x), \dots, \xi_n(x)) \in D\}$$

where $\xi_1, \xi_2, \dots, \xi_n$ are in X^* and D is a Borel set in the n -dimensional Euclidean space R_n . \mathfrak{A}_X is the family of all cylinder sets in X and $\overline{\mathfrak{A}}_X$ is the minimal σ -algebra including \mathfrak{A}_X . τ_X is the weak topological σ -algebra in X and $\hat{\tau}_X$ is the strong topological σ -algebra in it. Evidently we have

$$\mathfrak{A}_X \subset \overline{\mathfrak{A}}_X \subset \tau_X \subset \hat{\tau}_X$$

and if X is separable, then $\overline{\mathfrak{A}}_X = \hat{\tau}_X$ (E. Mourier [8]).

Let \mathfrak{X} be a real Hilbert space. *The canonical Gaussian cylinder measure* $\mu_{\mathfrak{X}}$ of \mathfrak{X} is a finitely additive nonnegative set function on $(\mathfrak{X}, \mathfrak{A}_{\mathfrak{X}})$ such that

$$\mu_{\mathfrak{X}}[x \in \mathfrak{X}: \xi(x) \leq \alpha] = \frac{1}{\sqrt{2\pi} |\xi|} \int_{-\infty}^{\alpha} \exp\left[-\frac{u^2}{2|\xi|^2}\right] du, \quad (1.1)$$

for any $\xi \in \mathfrak{X}^*$ and real number α , where $|\xi|$ is the norm in \mathfrak{X}^* . It is well-known that $\mu_{\mathfrak{X}}$ does not have σ -additive extension to $(\mathfrak{X}, \overline{\mathfrak{A}}_{\mathfrak{X}})$, (see Corollary of Lemma 6).

Let $\|x\|$ be a continuous norm on \mathfrak{X} , and X be the Banach space obtained by the completion of \mathfrak{X} in the norm $\|x\|$. Since we may consider X^* as a subspace of \mathfrak{X}^* through the natural imbedding, $\mu_{\mathfrak{X}}$ induces a Gaussian cylinder measure μ on (X, \mathfrak{A}_X) as follows. If $\xi_1, \xi_2, \dots, \xi_n$ are in X^* and D is a Borel set in R_n , define

$$\begin{aligned} & \mu[x \in X; (\xi_1(x), \dots, \xi_n(x)) \in D] \\ &= \mu_{\mathfrak{X}}[x \in \mathfrak{X}; (\xi_1(x), \dots, \xi_n(x)) \in D]. \end{aligned} \quad (1.2)$$

μ is well-defined. Furthermore, if μ has a σ -additive extension on $(X, \overline{\mathfrak{A}}_X)$, then we call it *the σ -extension of $\mu_{\mathfrak{X}}$ on the Banach space X* and the norm $\|x\|$ *admissible* on \mathfrak{X} . If a norm on \mathfrak{X} is induced by an inner product, namely, a continuous symmetric bilinear form on \mathfrak{X} , then we call it *Hilbertian*. A measurable norm is defined by L. Gross [1, 2] as follows. A norm $\|x\|_1$ on \mathfrak{X} is a *measurable norm* if for every positive real number ε there exists a finite dimensional projection P_0 of \mathfrak{X} such that for every finite dimensional projection P orthogonal to P_0 we have

$$\mu_{\mathfrak{X}}[x \in \mathfrak{X}: \|Px\|_1 > \varepsilon] < \varepsilon.$$

L. Gross [1] showed that the measurable norm is admissible.

In the last section, we shall give some remarks on the admissible norm. We shall give a necessary and sufficient condition for a Hilbertian norm to be admissible (Theorem 3) and show that there exists a measurable norm such that there is no Hilbertian admissible norm stronger than it (Example 2). This means that as a support of an abstract Wiener measure we can choose a Banach subspace which includes no Hilbert subspace of full measure. We shall also show that there exists an admissible norm which is not a measurable norm. This means that for a norm to be an admissible norm it is not necessary to be a measurable norm.

2. Gaussian measure and Radon measure.

Let X be a Banach space with norm $\|x\|$ and X^* be the topological dual for X with norm $\|\xi\|$. A probability measure μ on $(X, \overline{\mathfrak{A}}_X)$ is *Gaussian* if for every $\xi \in X^*$, $\xi(x)$ is a Gaussian random variable with mean zero on the probability space $(X, \overline{\mathfrak{A}}_X, \mu)$. In other words, for every $\xi \in X^*$ and real number α ,

$$\mu[x \in X: \xi(x) \leq \alpha] = \frac{1}{\sqrt{2\pi v(\xi)}} \int_{-\infty}^{\alpha} \exp\left[-\frac{u^2}{2v(\xi)}\right] du, \tag{2. 1}$$

where $v(\xi)$ is the variance of $\xi(x)$.

Theorem 1. Every Gaussian measure μ on a separable or reflexive Banach space $(X, \overline{\mathfrak{A}}_X)$ can be extended to a Radon measure on (X, τ_X) .

Proof. If X is separable, $\overline{\mathfrak{A}}_X = \tau_X$ and the proof is trivial. Let X be a reflexive Banach space and let X^{**} be the topological dual space of X^* . Let $\overline{\mathfrak{A}}^*$ be the minimal σ -algebra of subsets of X^{**} with respect to which

all the functions $\xi(x)$, $\xi \in X^*$, are measurable, where $\xi(x)$ ($\xi \in X^*$, $x \in X^{**}$) denotes the continuous linear form and τ^* is the topological σ -algebra with respect to X^* -topology in X^{**} (W. Dunford and J.T. Schwartz [15], p. 419). Define a measure μ^* on $(X^{**}, \overline{\mathfrak{A}}^*)$ as follows:

$$\begin{aligned} \mu^*[x \in X^{**}: (\xi_1(x), \dots, \xi_n(x)) \in D] \\ = \mu[x \in X: (\xi_1(x), \dots, \xi_n(x)) \in D]. \end{aligned} \quad (2.2)$$

where $\xi_1, \xi_2, \dots, \xi_n$ are in X^* and D is a Borel set in R_n . The measure μ^* is well defined and is Gaussian. Since all the open sets in $\overline{\mathfrak{A}}^*$ form an open basis which determines X^* -topology and since X^{**} is the topological dual for the Banach space X^* , μ^* can be extended to a Radon measure $\tilde{\mu}^*$ on (X^{**}, τ^*) uniquely (Yu. V. Prohorov [10], Theorem 1, Lemma 3 and Example 1). Since X is reflexive, we have $X = X^{**}$ and $\tau^* = \tau_X$. Therefore $\tilde{\mu}^*$ is a Gaussian Radon measure on (X, τ_X) . Since X is a Banach space, the weak Radon measure $\tilde{\mu}^*$ can be extended to a strong Radon measure $\tilde{\mu}$ on (X, τ_X) and, it is easy to see from (2.2), that $\tilde{\mu}$ is an extension of μ . Thus we have proved the theorem.

Remark. Without any change in the proof, we can prove Theorem 1 not only for a Gaussian measure but for any probability measure on a Banach space, the finite dimensional distribution of which is Radon.

We can therefore consider a Gaussian measure on a Banach space X as a Radon measure on (X, τ_X) .

3. Gaussian measure and abstract Wiener measure.

Let μ be a Gaussian measure on a separable or reflexive Banach space X . We use the same notations used in Section 2. Choose the maximal subset $\{\xi_\alpha; \alpha \in A\}$ of X^* such that

$$\begin{aligned} \xi_\alpha \in X^* \text{ and } \|\xi_\alpha\| = 1, \quad \alpha \in A \\ \int_X \xi_\alpha(x) \xi_\beta(x) d\mu(x) = 0 \quad \text{if } \alpha \neq \beta, \alpha, \beta \in A. \end{aligned} \quad (3.1)$$

LEMMA 1. *Let $A_0 = \{\alpha \in A; v(\xi_\alpha) \neq 0\}$, then A_0 is an at most countable subset of A .*

Proof. Let $\{\alpha_n\}_{n=1,2,\dots}$, be an arbitrary countable subset of \mathcal{A} . Since it holds that

$$\sup_n |\xi_{\alpha_n}(x)| \leq \sup_{\substack{\|\xi\|=1 \\ \xi \in X^*}} |\xi(x)| = \|x\| < +\infty, \text{ for every } x \in X, \quad (3.2)$$

we can choose a positive number M such that

$$\mu[x \in X: \sup_n |\xi_{\alpha_n}(x)| \leq M] > \frac{1}{2}. \quad (3.3)$$

On the other hand, we have

$$\begin{aligned} & \mu[x \in X; \sup_n |\xi_{\alpha_n}(x)| \leq M] \\ &= \lim_{N \rightarrow +\infty} \mu[x \in X: \sup_{1 \leq n \leq N} |\xi_{\alpha_n}(x)| \leq M] \\ &= \lim_{N \rightarrow +\infty} \mu[\cap_{1 \leq n \leq N} \{x \in X: |\xi_{\alpha_n}(x)| \leq M\}]. \end{aligned}$$

Since the collection $\{\xi_{\alpha_n}(x)\}$ is Gaussian, from (3.1), $\xi_{\alpha_n}(x)$ and $\xi_{\alpha_m}(x)$ are mutually independent if $n \neq m$. Therefore,

$$\begin{aligned} & \mu[x \in X: \sup_n |\xi_{\alpha_n}(x)| \leq M] \\ &= \lim_{N \rightarrow +\infty} \prod_{1 \leq n \leq N} \mu[x \in X: |\xi_{\alpha_n}(x)| \leq M] \\ &= \lim_{N \rightarrow +\infty} \prod_{1 \leq n \leq N} \frac{1}{\sqrt{2\pi v(\xi_{\alpha_n})}} \int_{-M}^M \exp\left[-\frac{u^2}{2v(\xi_{\alpha_n})}\right] du \\ &= \lim_{N \rightarrow +\infty} \prod_{1 \leq n \leq N} \frac{1}{\sqrt{2\pi}} \int_{-\frac{M}{\sqrt{v(\xi_{\alpha_n})}}}^{\frac{M}{\sqrt{v(\xi_{\alpha_n})}}} \exp\left[-\frac{u^2}{2}\right] du. \end{aligned}$$

Together with (3.3), we have

$$\lim_{N \rightarrow +\infty} v(\xi_{\alpha_n}) = 0. \quad (3.4)$$

Since the choice of the countable subset $\{\alpha_n\}$ is arbitrary, the set

$$\mathcal{A}_N = \left\{ \alpha \in \mathcal{A}; v(\xi_{\alpha_n}) \geq \frac{1}{N} \right\}$$

must be a finite subset of Λ for every positive integer N . Otherwise we have a contradiction to (3.4). Therefore,

$$A_0 = \bigcup_{N=1}^{+\infty} A_N$$

must be a countable subset of Λ .

LEMMA 2. Define X_α , $\alpha \in \Lambda$, by

$$X_\alpha = \{x \in X; \xi_\alpha(x) = 0\}, \quad \alpha \in \Lambda,$$

and set $\tilde{X} = \bigcap_{\alpha \in \Lambda - A_0} X_\alpha$. Then we have

$$\mu[\tilde{X}] = 1. \quad (3.5)$$

Proof. Let Γ be the family of all finite subsets of $\Lambda - A_0$ and define $X_J = \bigcap_{\alpha \in J} X_\alpha$; $J \in \Gamma$. Obviously X_J is a strongly closed linear subspace of X and the family $\{X_J; J \in \Gamma\}$ is directed. Since $v(\xi_\alpha) = 0$, $\xi_\alpha(x)$ is a Dirac measure for every $\alpha \in \Lambda - A_0$, we have

$$\mu[X_J] = 1 \quad \text{for every } J \in \Gamma.$$

Therefore,

$$\begin{aligned} \mu[\tilde{X}] &= \mu\left[\bigcap_{J \in \Gamma} X_J\right] \\ &= \inf_{J \in \Gamma} \mu[X_J] = 1, \end{aligned}$$

(L. Schwartz [11]). Thus we have proved the lemma.

This lemma means that the measure μ is concentrated in some closed linear subspace \tilde{X} . \tilde{X} is also a Banach space with the norm $\|x\|$. Let \mathfrak{C} be the closed linear manifold spanned by $\{\xi_\alpha; \alpha \in \Lambda - A_0\}$. Then the topological dual \tilde{X}^* for \tilde{X} is isomorphic to X^*/\mathfrak{C} .

It is easy to see that in \tilde{X}^*

$$v(\xi) = 0 \quad \text{implies} \quad \xi = 0. \quad (3.6)$$

Let $\|\xi\|$ be the norm in \tilde{X}^* again.

Hereafter, we restrict the measure μ to \tilde{X} . For every $\xi, \eta \in X^*$ define

$$(\xi, \eta) = \int_{\tilde{X}} \xi(x)\eta(x) d\mu(x), \quad (3.7)$$

$$|\xi| = \sqrt{(\xi, \xi)} = \sqrt{v(\xi)} \quad (3.8)$$

Then, according to (3. 6),

$$|\xi| = 0 \text{ if and only if } \|\xi\| = 0, \quad (3. 9)$$

in \tilde{X}^* . Therefore the bilinear form (ξ, η) is an inner product and $|\xi|$ is a norm on \tilde{X}^* . Next we shall show that the norm $|\xi|$ is continuous.

LEMMA 3. *There exists a positive constant C such that*

$$|\xi| \leq C \|\xi\| \quad \text{for every } \xi \in \tilde{X}^*. \quad (3. 10)$$

Proof. It is sufficient to show

$$C = \sup_{\substack{\|\xi\|=1 \\ \xi \in \tilde{X}^*}} |\xi| < +\infty.$$

Suppose not, then there exists a sequence $\{\xi_n\}$ in \tilde{X}^* such that

$$\begin{aligned} \|\xi_n\| &= 1, \quad n = 1, 2, 3, \dots \\ \lim_{n \rightarrow +\infty} |\xi_n| &= +\infty. \end{aligned}$$

By choosing a sufficiently large number M , we have

$$\mu[x \in \tilde{X}: \sup_n |\xi_n(x)| \leq M] > \frac{1}{2}, \quad (3. 11)$$

(see the proof of Lemma 1). On the other hand,

$$\begin{aligned} & \mu[x \in \tilde{X}: \sup_n |\xi_n(x)| \leq M] \\ &= \lim_{n \rightarrow +\infty} \mu[x \in \tilde{X}: \sup_{1 \leq \nu \leq n} |\xi_\nu(x)| \leq M] \\ &\leq \lim_{n \rightarrow +\infty} \mu[x \in \tilde{X}: |\xi_n(x)| \leq M] \\ &= \lim_{n \rightarrow +\infty} \frac{1}{\sqrt{2\pi} |\xi_n|} \int_{-M}^M \exp\left[-\frac{u^2}{2|\xi_n|^2}\right] du \\ &= \lim_{n \rightarrow +\infty} \frac{1}{\sqrt{2\pi}} \int_{-\frac{M}{|\xi_n|}}^{\frac{M}{|\xi_n|}} \exp\left[-\frac{u^2}{2}\right] du = 0. \end{aligned}$$

This contradicts (3. 11) and concludes the proof.

Let \mathfrak{X}^* be the Hilbert space obtained by the completion of \tilde{X}^* with respect to the inner product (ξ, η) , and let \mathfrak{X} be its topological dual space. By the definition (3. 8) of the norm $|\xi|$, the relation (1. 2) is valid for μ and the canonical Gaussian cylinder measure $\mu_{\mathfrak{X}}$ of the Hilbert space \mathfrak{X} .

This means that μ is a σ -extension of $\mu_{\mathfrak{X}}$ in \tilde{X} . On the other hand, it is easy to see that the system $\{\xi_\alpha / \|\xi_\alpha\| : \alpha \in A_0\}$ is a C.O.N.S. (complete orthonormal system) in \mathfrak{X}^* . Since A_0 is at most countable, \mathfrak{X} is a separable Hilbert space.

LEMMA 4. \mathfrak{X} is a subspace of \tilde{X} .

Proof. The measure μ extends to a Gaussian measure μ^* on \tilde{X}^{**} by (2. 2), where \tilde{X}^{**} is the topological dual for \tilde{X} . Then \tilde{X} is a measurable subset of \tilde{X}^{**} and $\mu^*(\tilde{X}^{**}) = \mu^*(\tilde{X}) = 1$ is true (see the proof of Theorem 1). Since \tilde{X}^* is included in \mathfrak{X}^* its dual \mathfrak{X} is included in $(\tilde{X}^*)^* = \tilde{X}^{**}$. The relation (1. 2) is also valid for μ^* and $\mu_{\mathfrak{X}}$. Therefore, by identifying \mathfrak{X}^* and \mathfrak{X} , for every $x_0 \in \mathfrak{X} (= \mathfrak{X}^*)$

$$\mu^*[\tilde{X} + x_0] = \mu^*[\tilde{X}] = 1, \quad (3. 12)$$

due to the fact that μ^* is quasi-invariant. (Y. Umemura [12]). On the other hand, if \mathfrak{X} is not a subspace of \tilde{X} , namely, if there exists x_0 in \mathfrak{X} which is not in \tilde{X} , then we have

$$[\tilde{X} + x_0] \cap \tilde{X} = \phi. \quad (3. 13)$$

For, if there exists y in $[\tilde{X} + x_0] \cap \tilde{X}$, then there exists y' in \tilde{X} such that $y = y' + x_0$. Since \tilde{X} is a linear space, $x_0 = y - y'$ is in \tilde{X} . This is a contradiction to the assumption on x_0 and (3. 13) is true. Thus we have

$$1 = \mu^*[\tilde{X}^{**}] \geq \mu^*[[\tilde{X} + x_0] \cup \tilde{X}] = 2.$$

This contradicts (3. 12), which proves the lemma.

LEMMA 5. \mathfrak{X} is dense in \tilde{X} .

Proof. Let $\bar{\mathfrak{X}}$ be the closure of \mathfrak{X} in \tilde{X} . If there exists x_0 in $\tilde{X} - \bar{\mathfrak{X}}$, then, by the Hahn-Banach theorem, there exists $\xi \neq 0$ in \tilde{X}^* such that $\xi(x) = 0$ on \mathfrak{X} . On the other hand, let $\|x\|_0$ be the norm on \mathfrak{X} . Then we have

$$\|\xi\| = \sup_{\substack{\|x\|_0=1 \\ x \in \mathfrak{X}}} |\xi(x)| = 0.$$

According to (3. 9), this means $\xi = 0$ in \tilde{X}^* and contradicts the choice of ξ . Therefore $\tilde{X} = \bar{\mathfrak{X}}$, that is, \mathfrak{X} is dense in \tilde{X} .

COROLLARY. \tilde{X} is separable.

Proof. The space \mathfrak{X} is a separable Hilbert space and, by Lemma 5, is dense in \tilde{X} . Furthermore, the norm $|x|_0$ on \mathfrak{X} is stronger than that on \tilde{X} . Therefore \tilde{X} is separable.

Summing up these results, we can derive the following theorem.

THEOREM 2. (A). *Let μ be a Gaussian measure on a separable or reflexive Banach space. Then there exists a separable closed linear subspace \tilde{X} such that $\mu[\tilde{X}] = 1$ and (3. 6) is valid in \tilde{X}^* .*

(B). *Let μ be a Gaussian measure on a separable Banach space \tilde{X} , and assume that (3. 6) is valid in \tilde{X}^* . Then there exists a dense Hilbert subspace \mathfrak{X} of \tilde{X} such that μ is an abstract Wiener measure, that is, μ is a σ -extension in \tilde{X} of the canonical Gaussian cylinder measure $\mu_{\mathfrak{X}}$ of \mathfrak{X} . The norm $\|x\|$ is admissible on \mathfrak{X} .*

COROLLARY. *There is no admissible norm on a nonseparable Hilbert space \mathfrak{X} .*

Proof. Suppose that a norm $\|x\|$ on \mathfrak{X} is admissible, X be the completion of \mathfrak{X} in the norm $\|x\|$, and let μ be the σ -extension in X of the canonical Gaussian cylinder measure $\mu_{\mathfrak{X}}$ of \mathfrak{X} . Since \mathfrak{X} is dense in X and $\|x\| = 0$ implies $x = 0$ in \mathfrak{X} , we can show that X^* is a dense subspace of \mathfrak{X}^* and (3. 6) is valid in X^* in the manner similar to that used in the proof of Lemma 5. Therefore, we can choose a C.O.N.S. $\{\xi_\alpha^0: \alpha \in A\}$ of \mathfrak{X}^* from X^* . A is an uncountable set since \mathfrak{X}^* is nonseparable. Let $\xi_\alpha = \xi_\alpha^0 / \|\xi_\alpha^0\|$; $\alpha \in A$. Then (3. 1) is valid for $\{\xi_\alpha: \alpha \in A\}$. On the other hand, considering (3. 6), $v(\xi_\alpha) = \frac{1}{\|\xi_\alpha^0\|} \neq 0$ for every $\alpha \in A$. This contradicts Lemma 1.

4. Admissible norm.

Let \mathfrak{X} be a separable Hilbert space with norm $|x|$ and inner product (x, y) . We study the condition under which a Hilbertian norm on \mathfrak{X} is admissible.

LEMMA 6^(*). *Let H be a separable Hilbert space and let μ be a Gaussian cylinder measure on (H, \mathfrak{A}_H) , that is, for every $\xi \in H^*$, $\xi(x)$ is a Gaussian random variable on (H, \mathfrak{A}_H, μ) with mean $m(\xi)$ and variance $v(\xi)$. (In this lemma, we do not assume zero mean.)*

^(*) This lemma was suggested by Prof. K. Ito.

Then μ has a σ -additive extension to $(H, \overline{\mathfrak{A}}_H)$ if and only if the characteristic functional of μ is of the form

$$\int_H e^{i\xi(x)} d\mu(x) = \exp \left[i \langle \xi, m \rangle - \frac{1}{2} \|S\xi\|^2 \right], \quad \xi \in H^*, \quad (4.1)$$

where m is an element of H , S is a nonnegative self-adjoint Hilbert-Schmidt operator and $\|\xi\|$ is the norm on H^* .

Proof. The sufficiency is derived from V.V. Sazonav [6].

We have only to prove the necessity. Assume that there exists a σ -additive extension to $(H, \overline{\mathfrak{A}}_H)$ and denote it by μ again. Identify H^* and H and let $\langle \cdot, \cdot \rangle$ be its inner product and $\|\cdot\|$ be its norm. Then $\langle \xi, x \rangle$; $\xi \in H^*(=H)$, $x \in H$ denotes the natural linear form.

Let $\{\xi_n\}$ be a sequence in H convergent to zero. Then $\langle \xi_n, x \rangle$ converges to zero for all x in H . Since $\{\langle \xi_n, x \rangle\}$ is a Gaussian random sequence on $(H, \overline{\mathfrak{A}}_H, \mu)$,

$$m(\xi_n) = \int_H \langle \xi_n, x \rangle d\mu(x) \quad (4.2)$$

converges to zero (§33, Lemma 1 of K. Ito [14]). Therefore $m(\xi)$ is a continuous linear functional on H^* and there exists $m \in H$ such that

$$m(\xi) = \langle \xi, m \rangle \quad \text{for any } \xi \in H. \quad (4.3)$$

Next, let $\{\varphi_j\}$ be a C.O.N.S. in H , and, for m and for every ξ, x in H , set

$$\begin{aligned} m_j &= \langle \varphi_j, m \rangle, \\ x_j &= x_j(x) = \langle \varphi_j, x \rangle, \quad j = 1, 2, 3, \dots, \\ \xi_j &= \xi_j(\xi) = \langle \varphi_j, \xi \rangle. \end{aligned} \quad (4.4)$$

Then obviously

$$\mu[x \in H: \sum_{j=1}^{+\infty} x_j(x)^2 < +\infty] = \mu[H] = 1. \quad (4.5)$$

On the other hand, let

$$\begin{aligned} \xi^N &= \sum_{j=1}^N \xi_j \varphi_j, \quad N = 1, 2, 3, \dots \\ v_{ij} &= \int_H (x_j(x) - m_j)(x_i(x) - m_i) d\mu(x), \\ & \quad i, j = 1, 2, 3, \dots \end{aligned} \quad (4.6)$$

Then

$$\begin{aligned} & \int_H \exp [i \langle \xi^N, x \rangle] d\mu(x) \\ &= \int_H \exp \left[i \sum_{j=1}^N \xi_j x_j(x) \right] d\mu(x) \\ &= \exp \left[i \sum_{j=1}^N m_j \xi_j - \frac{1}{2} \sum_{k,j=1}^N v_{kj} \xi_k \xi_j \right]. \end{aligned} \tag{4.7}$$

Averaging both sides of (4.7) with respect to the measure

$$(2\pi)^{-\frac{N}{2}} \exp \left[-\frac{1}{2} \sum_{j=1}^N \xi_j^2 \right] d\xi_1 d\xi_2 \cdots d\xi_N,$$

we have

$$\int_H \exp \left[-\frac{1}{2} \sum_{j=1}^N x_j(x)^2 \right] d\mu(x) \leq \frac{1}{\sqrt{1 + \sum_{j=1}^N v_{jj}}}. \tag{4.8}$$

If $\sum_{j=1}^{+\infty} v_{jj}$ is divergent, then from (4.8) we have

$$\int_H \exp \left[-\frac{1}{2} \sum_{j=1}^{+\infty} x_j(x)^2 \right] d\mu(x) = 0,$$

and

$$\exp \left[-\frac{1}{2} \sum_{j=1}^{+\infty} x_j(x)^2 \right] = 0, \quad \text{a.e.}$$

Therefore

$$\mu \left[\sum_{j=1}^{+\infty} x_j(x)^2 = +\infty \right] = 1.$$

This contradicts (4.5) and we have,

$$\sum_{j=1}^{+\infty} v_{jj} < +\infty. \tag{4.9}$$

Define a linear operator V on H by

$$\langle V\varphi_i, \varphi_j \rangle = v_{ij}, \quad i, j = 1, 2, 3, \dots \tag{4.10}$$

Then V is a nonnegative self-adjoint operator on H and further, it is nuclear, since

$$\sum_{j=1}^{+\infty} \langle V\varphi_j, \varphi_j \rangle = \sum_{j=1}^{+\infty} v_{jj} < +\infty.$$

Let S be \sqrt{V} . Then it is easy to see that S is the required Hilbert-Schmidt operator. Thus we have proved the lemma.

COROLLARY 1. *The canonical Gaussian cylinder measure $\mu_{\mathfrak{X}}$ on a Hilbert space \mathfrak{X} does not have a σ -additive extension to $(\mathfrak{X}, \overline{\mathfrak{X}})$.*

Proof. The characteristic functional of $\mu_{\mathfrak{X}}$ is

$$\begin{aligned} \int_{\mathfrak{X}} \exp[i\xi(x)] d\mu_{\mathfrak{X}}(x) &= \exp\left[-\frac{1}{2} |\xi|^2\right] \\ &= \exp\left[-\frac{1}{2} |I\xi|^2\right], \end{aligned} \quad (4.11)$$

where $|\xi|$ is the norm on \mathfrak{X}^* and I is the identity. But I is not of Hilbert-Schmidt type. Therefore, by Lemma 6, $\mu_{\mathfrak{X}}$ does not have a σ -additive extension to $(\mathfrak{X}, \overline{\mathfrak{X}})$.

COROLLARY 2. *In Lemma 6, if μ has a σ -additive extension to $(H, \overline{\mathfrak{X}}_H)$ and mean zero, then for every $\xi, \eta \in H^*(=H)$*

$$\int_H \xi(x)\eta(x) d\mu(x) = \langle S\xi, S\eta \rangle, \quad (4.12)$$

where S is the Hilbert-Schmidt operator determined by (4.1).

Utilizing Lemma 6, we have the following theorem.

THEOREM 3. *A Hilbertian norm $\|x\|$ on a separable Hilbert space \mathfrak{X} is admissible if and only if there exists a one to one Hilbert-Schmidt operator S_0 such that*

$$\|x\| = |S_0x|, \quad x \in \mathfrak{X}, \quad (4.13)$$

where $|x|$ is the initial norm on \mathfrak{X} .

Proof. The sufficiency is well-known (for example, see Y. Umemura [12]).

We prove the necessity. Let $\|x\|$ be a Hilbertian admissible norm induced by an inner product $\langle x, y \rangle$ on \mathfrak{X} and let H be the completion of \mathfrak{X} in the norm $\|x\|$. Then H is also a Hilbert space with the inner product $\langle x, y \rangle$. Let μ be the σ -extension in H of the canonical Gaussian cylinder measure $\mu_{\mathfrak{X}}$ of \mathfrak{X} . Then μ is a Gaussian measure on the Hilbert space H . Therefore, by Lemma 6, there exists a nonnegative Hilbert-Schmidt opera-

tor S on H^* determined by (4.1). Since we are assuming mean zero, (4.12) is also valid (Corollary 2 of Lemma 6).

Identifying \mathfrak{X} and \mathfrak{X}^* , and remembering H^* is a subspace of $\mathfrak{X}^*(=\mathfrak{X})$, we have

$$\begin{aligned} \|S\xi\|^2 &= \int_H \xi(x)^2 d\mu(x) \\ &= \int_{\mathfrak{X}} (\xi, x)^2 d\mu_{\mathfrak{X}}(x) = |\xi|^2, \end{aligned} \tag{4.14}$$

for every ξ in H^* where $\|\xi\|$ is the norm on H^* . Consequently,

$$\|S\xi\| = |\xi|, \quad \text{for every } \xi \in H^*. \tag{4.15}$$

Since $\|x\| = 0$ implies $x = 0$ in \mathfrak{X} and so $|\xi| = 0$ implies $\xi = 0$ in H^* . Therefore, by (4.15), $S\xi = 0$ implies $\xi = 0$ in H^* and S is a one to one operator.

Let $\{\lambda_j\}$ and $\{\varphi_j\}$ be eigenvalues and eigenvectors of S , respectively. Then $\lambda_j > 0$, $j = 1, 2, \dots$, and $\sum_{j=1}^{+\infty} \lambda_j^2 < +\infty$ because S is a one to one Hilbert-Schmidt operator.

Further, since μ is the σ -extension of the canonical Gaussian cylinder measure $\mu_{\mathfrak{X}}$, we have

$$\begin{aligned} (\varphi_i, \varphi_j) &= \int_{\mathfrak{X}} \varphi_j(x) \varphi_i(x) d\mu_{\mathfrak{X}}(x) \\ &= \int_H \langle \varphi_i, x \rangle \langle \varphi_j, x \rangle d\mu(x) \\ &= \langle S\varphi_i, S\varphi_j \rangle = \lambda_i \lambda_j \delta_{ij}, \\ &\quad i, j = 1, 2, 3, \dots \end{aligned}$$

Let $\phi_j = \lambda_j^{-1} \varphi_j$, $j = 1, 2, 3, \dots$. Then $\{\phi_j\}$ is a C.O.N.S. in \mathfrak{X} and

$$\begin{aligned} \sum_{j=1}^{+\infty} |S\phi_j|^2 &= \sum_{j=1}^{+\infty} \|S^2\phi_j\|^2 = \sum_{j=1}^{+\infty} \|\lambda_j \varphi_j\|^2 \\ &= \sum_{j=1}^{+\infty} \lambda_j^2 < +\infty. \end{aligned}$$

Therefore S can be extended to a Hilbert-Schmidt operator on $\mathfrak{X}^*(=\mathfrak{X})$ and we denote it by S again. Let S_0 be the dual operator of S in \mathfrak{X} . Then S_0 is the required operator. In fact, since SH^* is dense in H^* and H^* is dense in $\mathfrak{X}^*(=\mathfrak{X})$, for every x in $\mathfrak{X}(\subset H)$,

$$\begin{aligned}
\|x\| &= \sup_{\substack{\|\xi\|=1 \\ \xi \in H^*}} |\xi(x)| = \sup_{\substack{\|S\xi\|=1 \\ \xi \in H^*}} |(S\xi)(x)| \\
&= \sup_{\substack{\|\xi\|=1 \\ \xi \in H^*}} |(S\xi, x)| = \sup_{\substack{\|\xi\|=1 \\ \xi \in H^*}} |(\xi, S^*x)| \\
&= \sup_{\substack{\|\xi\|=1 \\ \xi \in \mathfrak{X}^*}} |(\xi, S_0x)| = |S_0x|.
\end{aligned}$$

The proof is now complete.

COROLLARY. *Let $\|x\|$ be an admissible norm on \mathfrak{X} . If there exists a Hilbertian admissible norm stronger than $\|x\|$ then for any C.O.N.S. $\{\varphi_j\}$ in \mathfrak{X} we have*

$$\sum_{j=1}^{+\infty} \|\varphi_j\|^2 < +\infty. \quad (4.16)$$

Proof. Suppose that $\|x\|'$ is a Hilbertian admissible norm stronger than $\|x\|$, say, $\|x\| \leq \|x\|'$. By Theorem 3, there exists a Hilbert-Schmidt operator S such that $\|x\|' = |Sx|$, $x \in \mathfrak{X}$. Then for any C.O.N.S. $\{\varphi_j\}$ in \mathfrak{X} ,

$$\begin{aligned}
\sum_{j=1}^{+\infty} \|\varphi_j\|^2 &\leq \sum_j \|\varphi_j\|'^2 \\
&= \sum_j |S\varphi_j|^2 < +\infty.
\end{aligned}$$

This was to be proved.

Next we give some examples of admissible norms on a separable Hilbert space \mathfrak{X} .

EXAMPLE 1. Define

$$\|x\|_1 = |Sx|, \quad x \in \mathfrak{X},$$

where S is a one to one Hilbert-Schmidt operator on \mathfrak{X} . Then $\|x\|_1$ is a measurable norm (Section 1). Therefore, by Theorem 3, every Hilbertian admissible norm is a measurable norm.

EXAMPLE 2. Define

$$\|x\|_2 = \sup_n \frac{1}{\sqrt{n}} |(\varphi_n, x)|, \quad x \in \mathfrak{X}$$

where $\{\varphi_n\}$ is a C.O.N.S. in \mathfrak{X} . Then $\|x\|_2$ is a measurable norm but there is no Hilbertian admissible norm stronger than $\|x\|_2$.

In fact it is evident that $\|x\|_2$ is a norm on \mathfrak{X} . To prove that $\|x\|_2$ is a measurable norm, we imbed \mathfrak{X} in a measurable space $(\Omega, \overline{\mathfrak{A}})$ in which \mathfrak{X} is an $\overline{\mathfrak{A}}$ -measurable subspace and all functions (φ_n, x) , $n = 1, 2, 3, \dots$ are extended to $\overline{\mathfrak{A}}$ -measurable functions on Ω , further, there exists a σ -additive extension μ of $\mu_{\mathfrak{X}}$. As an example of such a space, we can choose the space of all sequences.

Then since μ is a σ -extension of $\mu_{\mathfrak{X}}$, we have

$$\begin{aligned}
 & \mu[x \in \Omega: \|x\|_2 < +\infty] \\
 &= \mu\left[x \in \Omega: \sup_n \frac{1}{\sqrt{n}} |(\varphi_n, x)| < +\infty\right] \\
 &= \lim_{N \rightarrow +\infty} \lim_{M \rightarrow +\infty} \mu\left[x \in \Omega: \sup_{1 \leq n \leq N} \frac{1}{\sqrt{n}} |(\varphi_n, x)| \leq M\right] \\
 &= \lim_{N \rightarrow +\infty} \lim_{M \rightarrow +\infty} \mu_{\mathfrak{X}}\left[x \in \mathfrak{X}: \sup_{1 \leq n \leq N} \frac{1}{\sqrt{n}} |(\varphi_n, x)| \leq M\right] \\
 &= \lim_N \lim_M \prod_{n=1}^N \left\{ \frac{1}{\sqrt{2\pi}} \int_{-M\sqrt{n}}^{M\sqrt{n}} \exp\left[-\frac{u^2}{2}\right] du \right\} \\
 &= \lim_N \lim_M \prod_{n=1}^N \left\{ 1 - \sqrt{\frac{2}{\pi}} \int_{M\sqrt{n}}^{+\infty} \exp\left[-\frac{u^2}{2}\right] du \right\} \\
 &\geq \lim_N \lim_M \prod_{n=1}^N \left\{ 1 - \sqrt{\frac{2}{\pi}} \frac{1}{M\sqrt{n}} \exp\left[-\frac{M^2}{2} n\right] \right\} \\
 &\geq \lim_N \lim_M \prod_{n=1}^N \left\{ 1 - \exp\left[-\frac{M^2}{2} n\right] \right\} \\
 &\geq \lim_N \lim_M \left\{ 1 - \sum_{n=1}^N \exp\left[-\frac{M^2}{2} n\right] \right\} \\
 &= \lim_N \lim_M \left\{ 1 - \frac{1 - \exp\left[-\frac{M^2}{2} N\right]}{1 - \exp\left[-\frac{M^2}{2}\right]} \exp\left[-\frac{M^2}{2}\right] \right\} \\
 &= 1,
 \end{aligned}$$

and for any positive number ε

$$\begin{aligned}
 & \mu[x \in \Omega: \|x\|_2 < \varepsilon] \\
 &= \lim_{N \rightarrow +\infty} \mu_{\mathfrak{X}}\left[x \in \mathfrak{X}: \sup_{1 \leq n \leq N} \frac{1}{\sqrt{n}} |(\varphi_n, x)| < \varepsilon\right] \\
 &\geq \prod_{n=1}^{+\infty} \left\{ 1 - \frac{1}{\varepsilon\sqrt{n}} \exp\left[-\frac{\varepsilon^2}{2} n\right] \right\} > 0,
 \end{aligned}$$

because

$$\sum_{n=1}^{+\infty} \frac{1}{\varepsilon \sqrt{n}} \exp\left[-\frac{\varepsilon^2}{2} n\right] \leq \frac{1}{\varepsilon} \frac{\exp\left[-\frac{\varepsilon^2}{2}\right]}{1 - \exp\left[-\frac{\varepsilon^2}{2}\right]} < +\infty.$$

Therefore, by Corollary 4.5 of L. Gross [2], $\|x\|_2$ is a measurable norm.

While for the C.O.N.S. $\{\varphi_n\}$ in \mathfrak{X}

$$\begin{aligned} \sum_{n=1}^{+\infty} \|\varphi_n\|_2^2 &= \sum_{n=1}^{+\infty} \left\{ \sup_{\nu} \frac{1}{\sqrt{\nu}} |(\varphi_{\nu}, \varphi_n)| \right\}^2 \\ &= \sum_{n=1}^{+\infty} \frac{1}{n} = +\infty. \end{aligned}$$

By Corollary of Theorem 3, there is no Hilbertian admissible norm stronger than $\|x\|_2$. This means that there is no Hilbert space of full measure which is included in the Banach space obtained by the completion of \mathfrak{X} in the norm $\|x\|_2$.

EXAMPLE 3. Define

$$\|x\|_3 = \left[\sup_n \frac{1}{n} \sum_{\nu=1}^n |(\varphi_{\nu}, x)|^2 \right]^{\frac{1}{2}}, \quad x \in \mathfrak{X}$$

where $\{\varphi_n\}$ is a C.O.N.S. in \mathfrak{X} . Then $\|x\|_3$ is an admissible norm on \mathfrak{X} but not a measurable norm.

Proof. Imbed \mathfrak{X} in the measurable space $(\Omega, \overline{\mathfrak{A}}, \mu)$ as in Example 2. Then by the law of large number, we have

$$\begin{aligned} &\mu[x \in \Omega: \|x\|_3 < +\infty] \\ &\geq \mu\left[x \in \Omega: \limsup_n \frac{1}{n} \sum_{\nu=1}^n |(\varphi_{\nu}, x)|^2 = 1\right] = 1. \end{aligned}$$

Therefore $\|x\|_3$ is an admissible norm; but, according to Corollary 4.5 of L. Gross [2], it is not a measurable norm. This means that for a norm on a separable Hilbert space to be admissible, it is not necessary to be a measurable norm in the sense of L. Gross [1].

BIBLIOGRAPHY

- [1] Gross, L., Abstract Wiener spaces. Proceedings of the Fifth Berkley Symposium, Vol. 2, Part 1, pp. 31–42, (1965).
- [2] Gross, L., Measurable functions on Hilbert space. Trans. A.M.S., Vol. 105, pp. 372–390, (1962).
- [3] Gross, L., Potential theory on Hilbert space. J. of Functional Analysis, Vol. 1, pp. 123–181, (1967).
- [4] Segal, I.E., Tensor algebras over Hilbert spaces, I. Trans. A.M.S., Vol. 81, pp. 106–134, (1956).
- [5] Segal, I.E., Distributions in Hilbert space and canonical systems of operators. Trans. A.M.S., Vol. 88, pp. 12–41, (1958).
- [6] Sazonov, V.V., A remark on characteristic functionals. Theory of Prob. and its Appl., Vol. 3, pp. 188–192, (1958). (English translation).
- [7] Getoor, P.K., On characteristic functions of Banach space-valued random variables. Pacific J. Math., Vol. 7, pp. 885–897, (1957).
- [8] Mourier, E., Éléments aleatoires dans un espaces de Banach. Ann. d'Inst. H. Poincare, Vol. 13, pp. 161–244, (1953).
- [9] Mourier, E., Random elements in linear spaces. Proceedings of the Fifth Berkley Symposium, Vol. 2, Part 1, pp. 43–53, (1965).
- [10] Prohorov, Yu.V. The method of characteristic functionals. Proceedings of the Fourth Berkley Symposium, Vol. 2, pp. 403–419, (1960).
- [11] Schawrtz, L., Mesures de Radon sur des espaces topologiques arbitraires. 3rd cycle 1964–1965, Inst. H. Poincare, Paris.
- [12] Umemura, Y., Measures on infinite dimensional vector spaces. Publ. Research Inst. for Math. Sci. Kyoto Univ., Vol. 1, pp. 1–47, (1965).
- [13] Hida, T., Stationary stochastic processes. White noise. Lecture note in Princeton University, (1968). (to appear).
- [14] Ito, K., Probability theory. Iwanami, Tokyo, (1953). (in Japanese).
- [15] Dunford, N. and Schwartz, J.T., Linear operators, Part 1. Interscience, N.Y., (1958).
- [16] Далецукый, Ю.Л., Бесконечномерные эллиптические операторы и связанные с ними параболические уравнения. Успехи Матем. Наук, Том. 22, стр. 3–54, (1967).

Tokyo Metropolitan University

