

GROUP CHARACTERS AND NORMAL HALL SUBGROUPS

P. X. GALLAGHER

TO RICHARD BRAUER ON HIS 60TH BIRTHDAY

1. Introduction

Let G be a finite group and let ψ be an (ordinary) irreducible character of a normal subgroup N . If ψ extends to a character of G then ψ is invariant under G , but the converse is false. In section 3 it is shown that if ψ extends coherently to the intermediate groups H for which H/N is elementary, then ψ extends to G . If N is a Hall subgroup, then in order for ψ to extend to G it is sufficient that ψ be invariant under G . This leads to a construction of the characters of G from the characters of N and the characters of the subgroups of G/N in this case.

Let $1(H)_G$ denote the character of G induced by the 1-character of the π -Hall subgroup H . If H is normal in G then the degree of each irreducible component of $1(H)_G$ divides the index of H . In section 4, the converse is proved in the case in which G is π -solvable and in the case in which G has a nilpotent π' -Hall subgroup.

2. Notation and Preliminary Results

In what follows, group means finite group. For a character ψ of a subgroup H of a group G , ψ_G denotes the character of G induced by ψ (for a definition, see [1]). Induction has the following properties:

- (i) If $H \subset K \subset G$, then $(\psi_K)_G = \psi_G$.
- (ii) If χ is a character of G , then $(\psi_G, \chi)_G = (\psi, \chi)_H$.

If χ_1 and χ_2 are characters of G , then $\chi_2 \in \chi_1$ means that there is a character χ_3 such that $\chi_1 = \chi_2 + \chi_3$. For example, $\psi \in \psi_G|H$ for each character ψ of the subgroup H of G .

The 1-character of G is denoted by $1(G)$.

Received November 18, 1961.

The degree of the representation with character χ is denoted by f_χ . A linear character is a character of degree 1.

If ϕ is a matrix representation, then $\det \phi$ is a linear character. Equivalent representations have the same determinant, so we may put $\det \phi = N(\chi)$, where χ is the character of ϕ . For each linear character ω , we have $N(\omega\chi) = \omega^{f_\chi} N(\chi)$.

If π is a set of prime numbers and f is a rational integer, then f divides π means that each prime factor of f is in π . A π -group is a group whose order divides π . A group is π -solvable if each of its composition factors is either a π -group or a π' -group, where π' is the complimentary set of prime numbers. A π -Hall subgroup of a group G is a π -subgroup whose index in G divides π' .

In what follows, character means irreducible character.

We will need the following results due to Clifford [3]:

THEOREM 1. *Let ψ be a character of the normal subgroup N of G and let T be the subgroup fixing ψ . Then for each character $\chi \in \psi_G$ there is a unique character ζ of T such that $\zeta \in \psi_T$ and $\chi \in \zeta_G$. For this ζ , $\zeta|N = a\psi$ for some positive integer a , and $\zeta_G = \chi$.*

Proof. Given $\chi \in \psi_G$, choose a character ζ of T such that $\zeta \in \psi_T$ and $\chi \in \zeta_G$. Then $\zeta|N \in \psi_T|N = (T:N)\psi$, so $\zeta|N = a\psi$ for some positive integer a .

For $\rho \in T$, $(\zeta^\rho, \zeta)_N = a_2(\psi^\rho, \psi)_N = 0$. Therefore $(\zeta_G, \zeta_G)_G = (\zeta_G, \zeta)_T = \sum_{\rho \in T} (\zeta^\rho, \zeta)_{T \cap T^\rho} (T:T \cap T^\rho)^{-1} = 1$. It follows that $\zeta_G = \chi$.

Since $\chi|T = \zeta + \zeta_1$, where $\psi \notin \zeta_1|N$, the only character of T with the required properties is ζ .

COROLLARY. *With the same hypotheses, $(G:T)f_\psi$ divides f_χ .*

THEOREM 2. *With the same hypotheses, if ψ extends to a character ψ_1 of T , then as ω ranges over the characters of T/N , $(\omega\psi_1)_G$ ranges over distinct characters of G , and*

$$(1) \quad \psi_G = \sum_{\omega} f_\omega \cdot (\omega\psi_1)_G$$

Proof. Both ψ_T and $\sum_{\omega} f_\omega \cdot \omega\psi_1$ are 0 off N and restrict on N to $(T:N)\psi$. Therefore they are equal. Inducing to G , we have (1). From (1) it follows that

$$(T:N) = (\psi_G, \psi)_N = (\psi_G, \psi_G)_G = \sum_{\omega, \omega'} f_\omega f_{\omega'} ((\omega\psi_1)_G, (\omega'\psi_1)_G)_G.$$

The diagonal terms alone contribute at least $\sum f_w^2 = (T:N)$, so the $(\omega\psi_1)_G$ are irreducible and distinct.

COROLLARY. *If N is normal in G and ψ_1 is a fixed extension of the character ψ of N to G , then each extension of ψ to G is uniquely of the form $\omega\psi_1$ where ω is a linear character of G/N .*

THEOREM 3. *Let N be a normal subgroup of prime index p in G . Then each invariant character of N extends to a character of G .*

Proof. Let ψ be an invariant character of N . Let χ be a character in ψ_G . By Theorem 1, $\chi|N = a\psi$. For each character ω of G/N , $\omega\chi$ is a character of G and $\omega\chi|N = a\psi$. The characters $\omega\chi$ are distinct. In fact, if $\omega\chi = \omega'\chi$ for $\omega \neq \omega'$, then $\chi = 0$ off N , so $a^2 = (\chi, \chi)_N = p(\chi, \chi)_G = p$, a contradiction. Therefore

$$\sum_{\omega} a \cdot \omega\chi \in \psi_G.$$

Since $(\psi_G, \psi_G)_G = (\psi_G, \psi)_N = p$, this implies that $a = 1$. Thus χ is an extension of ψ to G .

3. Extension of Characters from Normal Subgroups

An elementary group is a direct product of a cyclic group with a group of prime-power order. According to a theorem of Brauer [1], each character of a group G may be written in the form $\sum a_{\sigma} \xi_{\sigma}$ where ξ ranges over all linear characters of all elementary subgroups H of G and the coefficients a_{σ} are rational integers. Using Brauer's theorem we will find conditions under which an invariant character of a normal subgroup can be extended to a character of the whole group.

THEOREM 4. *Let N be a normal subgroup of G and let ψ be a character of N . Suppose that for each intermediate group H for which H/N is elementary ψ may be extended to a character $\psi(H)$ of H in such a way that*

- (i) $\psi(H)^{\rho} = \psi(H^{\rho}) \quad (\rho \in G)$,
- (ii) $\psi(H') = \psi(H'')|_{H'} \quad (H' \subset H'')$.

Then ψ may be extended to a character of G .

Proof. By Brauer's theorem we may write

$$(2) \quad 1(G/N) = \sum a_{\sigma} \xi_{\sigma}$$

where ξ ranges over all linear characters of H/N , for all intermediate groups H for which H/N is elementary, and the coefficients a_{ξ} are rational integers. Put

$$(3) \quad \chi = \sum a_{\xi} (\xi \cdot \psi(H_{\xi}))_{\sigma},$$

where H_{ξ} is the group of which ξ is a character. Then χ is a generalized character of G for which $\chi|N = \psi$.

From (2) and (3) we have

$$1 = (1(G/N), 1(G/N))_{\sigma} = \sum a_{\xi} a_{\xi'} (\xi_{\sigma}, \xi'_{\sigma})_{\sigma},$$

and

$$(\chi, \chi)_{\sigma} = \sum a_{\xi} a_{\xi'} ((\xi\psi(H_{\xi}))_{\sigma}, (\xi'\psi(H_{\xi'}))_{\sigma})_{\sigma}$$

We shall show that corresponding inner products in these two sums are equal, proving that $(\chi, \chi)_{\sigma} = 1$ and hence that χ or $-\chi$ is a character, from which since $\chi|N = \psi$, it will follow that χ is a character.

For any two characters ζ and ζ' of subgroups H and H' of G ,

$$(\zeta_{\sigma}, \zeta'_{\sigma})_{\sigma} = (\zeta_{\sigma}, \zeta')_{H'} = \sum_{\rho \bmod H} (\zeta^{\rho}, \zeta')_{H' \cap H^{\rho}} (H' : H' \cap H^{\rho})^{-1}.$$

Thus it is enough, in view of (i) and (ii), to show that for each intermediate subgroup H for which H/N is elementary, and each pair of linear characters ω and ω' of H/N ,

$$(\omega\psi(H), \omega'\psi(H))_H = (\omega, \omega')_H.$$

This is true since by the corollary to Theorem 2, $\omega\psi(H)$ and $\omega'\psi(H)$ are equal only for $\omega = \omega'$.

As an application of Theorem 4, we will prove the following special result which may also be proved using Schur's lemma and factor sets.

THEOREM 5. *Let N be a normal subgroup of G and let ψ be a character of N such that*

- (a) ψ is invariant under G ,
- (b) $N(\psi)$ extends to a character ξ of G ,
- (c) f_{ψ} is prime to $(G:N)$.

Then there is a unique character χ of G such that $\chi|N = \psi$ and $N(\chi) = \xi$.

Proof. If ψ extends to a character of G , there is a unique extension χ such that $N(\chi) = \xi$. In fact, all extensions are given by $\omega\chi_1$ where χ_1 is any

one extension and ω ranges over the linear characters of G/N . Since $\xi N(\chi_1)^{-1}$ is a linear character of G/N and f_ψ is prime to $(G:N)$, there is a unique linear character ω of G/N for which $\omega^{f_\psi} = \xi N(\chi_1)^{-1}$, i.e., for which $N(\omega\chi_1) = \omega^{f_\psi} N(\chi_1) = \xi$.

To prove the existence of an extension, suppose first that G/N is supersolvable. If $G = N$, put $\chi = \psi$. If $G \neq N$, let K/N be a normal subgroup of prime order of G/N . Since ψ is invariant under K , Theorem 3 shows that ψ has an extension to a character of K . By the last paragraph, ψ has a unique extension ψ_1 for which $N(\psi_1) = \xi|K$. For each $\rho \in G$, ψ_1^ρ is an extension of ψ to K for which $N(\psi_1^\rho) = \xi|K$. Thus ψ_1 is invariant under G . By induction, ψ_1 extends to a character of G .

To complete the proof in the general case, we note that elementary subgroups are supersolvable. Hence there are unique extensions $\psi(H)$ of ψ to the intermediate groups H for which H/N is elementary such that $N(\psi(H)) = \xi|H$. Properties (i) and (ii) of these extensions follow easily from this uniqueness and the invariance of ξ under conjugation.

LEMMA 1. *Let $G = HN$ where H is a subgroup of G and N is a normal subgroup of G and $H \cap N = 1$. Then each linear character of N which is invariant under G extends to a character of G .*

Proof. Let δ be an invariant linear character of N . Put $\xi(\sigma\tau) = \delta(\tau)$ for $\sigma \in H$, $\tau \in N$. Then $\xi|N = \delta$ and ξ is a linear character of G , since for $\sigma_1, \sigma_2 \in H$ and $\tau_1, \tau_2 \in N$,

$$\begin{aligned} \xi(\sigma_1\tau_1\sigma_2\tau_2) &= \xi(\sigma_1\sigma_2\tau_1^{\sigma_2}\tau_2) = \delta(\tau_1^{\sigma_2}\tau_2) \\ &= \delta(\tau_1^{\sigma_2})\delta(\tau_2) = \delta(\tau_1)\delta(\tau_2) = \xi(\sigma_1\tau_1)\xi(\sigma_2\tau_2). \end{aligned}$$

THEOREM 6. *If N is a normal Hall subgroup of G , then each invariant character of N extends to a character of G .*

Proof. Let ψ be an invariant character of N . Since f_ψ divides $(N:1)$, f_ψ is prime to $(G:N)$. By a theorem of Schur [5, p. 162], there is a subgroup H such that $G = HN$ and $H \cap N = 1$. Therefore $N(\psi)$, which is also invariant, extends to a linear character of G . Consequently ψ extends to a character of G , by Theorem 5.

THEOREM 7. *Let N be a normal Hall subgroup of G . Pick a character ψ*

of N , let ψ_1 be an extension of ψ to the group T fixing ψ , and let ω be a character of T/N . Then $(\omega\psi_1)_G$ is a character of G , and each character of G is obtained in this way.

Proof. By Theorem 6, ψ_1 exists. By Theorem 2, each $(\omega\psi_1)_G$ is a character of G and each character $\chi \in \psi_G$ is obtained in this way. For each character χ of G there is a ψ such that $\chi \in \psi_G$ so χ is of the form $(\omega\psi_1)_G$ for some ψ and some ω .

4. Conditions Implying a Hall Subgroup is Normal

It was proved by N. Ito [4] that if the degree of each character of a solvable group is prime to p , then the group has an abelian normal p -Sylow subgroup. In this section we prove an analogous result in which the conclusion is “ G has a normal π -Hall subgroup” and the hypothesis “ $p \nmid f_\chi$, for all characters χ ” is replaced by

α : G has a π -Hall subgroup H and the degree of each character in $1(H)_G$ divides π' .

LEMMA 2. *If G satisfies α and $N \triangleleft G$, then N and G/N also satisfy α .*

Proof. Let G satisfy α relative to the π -Hall subgroup H and let $N \triangleleft G$. Then $H \cap N$ is a π -Hall subgroup of N and HN/N is a π -Hall subgroup of G/N .

Since $\psi \in \psi_G \mid N$ for each character ψ of N , $\psi \in 1(H \cap N)_N$ implies $1(H \cap N) \in \psi \mid H \cap N \in \psi_G \mid H \cap N$ and hence

$$(\psi, 1(H)_G)_N = (\psi_G, 1(H)_G)_G = (\psi_G, 1(H))_H = (\psi_G, 1(H \cap N))_{H \cap N} (H : H \cap N)^{-1} \neq 0.$$

Therefore for each character $\psi \in 1(H \cap N)_N$, there is a character $\chi \in 1(H)_G$ for which $\psi \in \chi \mid N$ and hence for which f_ψ divides f_χ . By hypothesis, f_χ divides π' , so f_ψ also divides π' . Thus N satisfies α relative to $H \cap N$.

Since $1(HN) \in 1(H)_{HN}$, we have

$$1(HN)_G \in (1(H)_{HN})_G = 1(H)_G.$$

If χ is a character of G/N in $1(HN)_G$, then χ is a character of G in $1(H)_G$. By hypothesis f_χ divides π' . Thus G/N satisfies α relative to HN/N .

THEOREM 8. *If G is a π -solvable group which satisfies α , then the π -Hall subgroup H is normal in G .*

Proof. Let M be a minimal normal subgroup of G . Then G/M is also a π -solvable group which satisfies α , so by induction G/M has a normal π -Hall subgroup K/M .

If M is a π -group, then K is a normal π -Hall subgroup of G .

If $K \neq G$, then K is a π -solvable group which satisfies α , so by induction K has a normal π -Hall subgroup, which is then a normal π -Hall subgroup of G .

Finally, suppose M is a π' -group and $K = G$. Then $G = HM$ and $H \cap M = 1$. Consequently, $1(H)_G | M = \sum f_\psi \psi$, where ψ ranges over the characters of M . Let ψ be a character of M . Then there is a character $\chi \in 1(H)_G$ for which $\psi \in \chi | M$. Let T be the subgroup fixing ψ . Then $(G:T)$ divides f_χ . By the hypothesis, f_χ divides π' , so $(G:T)$ divides π' .

Since $(G:T)$ divides $(H:1)$, which divides π , $(G:T) = 1$. Thus each character ψ of M is invariant under G . It follows that H , in its action on M by conjugation, preserves the M -classes of M . By a theorem of Burnside [2, p. 89], H centralizes M . Thus $G = H \times M$, so $H \triangleleft G$.

LEMMA 3. *Let A be an abelian group, B a subgroup of A , and χ a possibly reducible character of A such that $\chi = 0$ off B . Then $(A:B)$ divides f_χ .*

Proof. Let $\chi = \sum a(\xi)\xi$ be the decomposition of χ into linear characters. For each character ω of A/B , we have $\omega\chi = \chi$, so $a(\omega\xi) = a(\xi)$. Also, $\omega\xi = \omega'\xi$ only for $\omega = \omega'$. Hence

$$\chi = \sum_{\xi}' a(\xi) \sum_{\omega} \omega\xi = \sum_{\omega} \omega \sum_{\xi}' a(\xi)\xi,$$

where the prime indicates that the summation is only over some ξ . It follows that $f_\chi = (A:B) \sum' a(\xi)$.

For a character χ of a group G , denote by $Z(\chi)$ the normal subgroup on which $|\chi| = f_\chi$. Denote by $Z(K)$ the center of a group K .

LEMMA 4. *Suppose that the group G has a Hall subgroup K and let χ be a character of G such that f_χ divides $(K:1)$. Then $(Z(K):Z(K) \cap Z(\chi))$ divides f_χ .*

Proof. For $\sigma \in Z(K)$, the centralizer of σ contains K , so the number of elements in the class of σ is prime to $(K:1)$ and hence to f_χ . By a theorem of Burnside [2, p. 322], either $\sigma \in Z(\chi)$ or $\chi(\sigma) = 0$. Thus $\chi | Z(K)$ is 0 off $Z(K) \cap Z(\chi)$, and the result now follows by Lemma 3.

THEOREM 9. *If G satisfies α relative to a π -Hall subgroup H and G has a nilpotent π' -Hall subgroup K , then H is normal in G .*

Proof. It is enough by Theorem 8 to prove that G is π -solvable. Since the hypotheses of Theorem 9 carry over to normal subgroups and factor groups, it is enough to show that if G has no proper normal subgroups, then $K = 1$.

If G has no proper normal subgroups, then $Z(\chi) = 1$ for all characters χ of G except $1(G)$. Lemma 4 then shows that $(Z(K):1)$ divides f_χ for all $\chi \in 1(H)_G$ except $1(G)$. Putting

$$1(H)_G = 1(G) + \sum a_\chi \chi,$$

we then have

$$(K:1) = (G:H) = 1 + \sum a_\chi f_\chi \equiv 1 \pmod{(Z(K):1)}.$$

This implies that $Z(K) = 1$. Since K is nilpotent, $K = 1$.

REFERENCES

- [1] Brauer, R. and Tate, J.: On the characters of finite groups. *Ann. of Math.* **62**, 1-7 (1955).
- [2] Burnside, W.: *Theory of Groups of Finite Order*. Second Edition, Cambridge Univ. Press. 1911.
- [3] Clifford, A. H.: Representations induced in an invariant subgroup. *Ann. of Math.* **38**, 533-550 (1937).
- [4] Ito, N.: Some studies on group characters. *Nagoya Math. Journal* **2**, 17-28 (1951).
- [5] Zassenhaus, H.: *The Theory of Groups*. Second Edition, Chelsea Publishing Co. 1958.

*Massachusetts Institute of Technology
and Paris, France*