

$$\alpha_{11}a'_1 + (\mu_1\alpha_{11} - \alpha_{12})b_1 + \alpha_{12}a'_2 + \mu_2\alpha_{12}b_2 \text{ on } T_2.$$

Suppose that  $k_1 \sim \varepsilon k_2$  in  $V_2$  for  $\varepsilon = 1$  or  $-1$ . Since  $k_1 \sim 0$  on  $T_2$ , one of the following systems of equations holds.

$$\begin{cases} \mu_1\alpha_{11} = 0, \\ \alpha_{11} - \mu_2\alpha_{12} = 0. \end{cases} \quad \begin{cases} \mu_1\alpha_{11} - \alpha_{12} = 0, \\ \mu_2\alpha_{12} = 0. \end{cases}$$

Using  $\mu_1\mu_2 \neq 0$ , we can show that it is impossible. This completes the proof.

Now we return to the proof for Case 2. Without loss of generality, we may assume that  $\varphi_1 F_3 \cap F_3$  contains  $c^*$ . Suppose that there exists a curve of type IV on  $\varphi_1 F_3$  in  $\varphi_1 F_3 \cap F_3$ . Let  $c$  be a simple closed curve of type IV on  $\varphi_1 F_3$  which bounds a Möbius band  $B$  such that  $B \cap F_3 = c \cup c^*$ . Since  $B \cap V_2$  is an annulus, it follows from Assertion B that  $c$  is of type IV on  $F_3$ . Hence  $c$  bounds a Möbius band  $B'$  on  $F_3$ . Let  $F'_3$  denote the surface obtained by deforming  $F_3 - B' \cup B$  slightly so that it is disjoint from  $B$ . Then, as is similar to Case 1 [Fig. 4.1],  $\varphi_1 F'_3 \cap F'_3$  contains fewer curves of type IV on  $\varphi_1 F'_3$  than  $\varphi_1 F_3 \cap F_3$ . Repeating these procedures, we can show that  $\varphi_1$  is equivalent to  $\varphi_2$  such that  $\varphi_2 F_3 \cap F_3$  does not contain a curve of type IV on  $\varphi_2 F_3$  and  $F_3$ .

Suppose that  $\varphi_2 F_3 \cap F_3$  contains at least two curves of type III on  $\varphi_2 F_3$ . Then there exists an annulus  $A$  on  $\varphi_2 F_3$  such that  $A \cap F_3 = \partial A$ . Let  $A'$  be an annulus on  $F_3$  which bounds  $\partial A$ . Deforming  $F_3 - A' \cup A$  slightly until it is disjoint from  $A$ , we obtain  $A'$  such that  $\varphi_2 F'_3 \cap F'_3$  has fewer components than  $\varphi_2 F_3 \cap F_3$ . Hence we can find an involution  $\varphi$  which is equivalent to  $\varphi_2$  such that  $\varphi F_3 \cap F_3$  consists of  $c^*$  and at most one curve of type III on  $\varphi F_3$ . If  $\varphi F_3 \cap F_3$  contains a curve  $c$  of type III, then  $c$  is  $\varphi$ -invariant. Since any two-sided curve in  $\varphi F_3 \cap F_3$  is not  $\varphi$ -invariant [12], the proof is completed.

*Case 3.* We will show that this case can not occur except for  $\mu_1\mu_2 = -2$ .

**Assertion C.** Suppose that  $\mu_1\mu_2 \neq -2$ . Let  $l_1$  and  $l_2$  be disjoint simple closed curves on  $T_2$  such that  $\pi l_1$  is of type II or V, and  $\pi l_2$  is of type III on  $F_3$ . Then  $l_1$  is not homologous to  $\varepsilon l_2$ , for  $\varepsilon = 1$  and  $-1$ , in  $V_2$ .

*Proof.* Let  $\rho$  be an autohomeomorphism of  $F_3$  such that  $\rho\pi l_1$  coincides with  $\partial N(c_1)$  or  $c_1$  and  $\rho\pi l_2 = b^*$ . By  $\begin{pmatrix} \alpha_{11} & \alpha_{12} \\ \alpha_{21} & \alpha_{22} \end{pmatrix}$  we denote a matrix corresponding to  $\rho^{-1}$ . Then, by Lemma 3.2,  $l_1$  is homologous to  $\alpha_{22}b_1 - \alpha_{21}b_2$  and  $l_2$  is homologous to either

$$\varepsilon(-\alpha_{21}a'_1 - \mu_1\alpha_{21}b_1 - \alpha_{22}a'_2 + (\alpha_{21} - \mu_2\alpha_{22})b_2)$$

or

$$\varepsilon(\alpha_{21}a'_1 + (\mu_1\alpha_{21} - \alpha_{22})b_1 + \alpha_{22}a'_2 + \mu_2\alpha_{22}b_2), \text{ for } \varepsilon = 1 \text{ or } -1, \text{ on } T_2.$$

and  $\rho c = \partial N(c_\mu)$ ,  $\mu = 1, 2$  or  $3$ . An annulus  $B \cap V_2$  has the boundaries  $k_1$  and  $k_2$  such that  $\pi k_1 = c'$  and  $\pi k_2 = c$ . If we suppose that  $B' \supset c'$ , then  $\rho c = \partial N(c_2)$  or  $\partial N(c_3)$ . Hence, in order to show that  $B' \supset c'$ , it suffices to prove that  $k_1 \sim \bar{\rho}^{-1}c_2$ ,  $-\bar{\rho}^{-1}c_3$ ,  $\bar{\rho}^{-1}c_3$  and  $-\bar{\rho}^{-1}c_3$  in  $V_2$ , where  $\bar{\rho}$  denotes the lifting of  $\rho$ . Let  $\begin{pmatrix} \alpha_{11} & \alpha_{12} \\ \alpha_{21} & \alpha_{22} \end{pmatrix}$  be a matrix in  $GL(2, Z)$  corresponding to the isotopy class of  $\rho^{-1}$ . Then it follows from Lemma 3.2 that

$$\bar{\rho}^{-1}\tilde{c}_1 \sim \alpha_{22}b_1 - \alpha_{21}b_2, \quad \bar{\rho}^{-1}\tilde{c}_2 \sim -(\alpha_{22} + \alpha_{21})b_1 + (\alpha_{21} + \alpha_{11})b_2$$

and

$$\bar{\rho}^{-1}\tilde{c}_3 \sim -\alpha_{12}b_1 + \alpha_{11}b_2 \text{ in } T_2.$$

Since  $\alpha_{11}\alpha_{22} - \alpha_{12}\alpha_{21} = \pm 1$ , it can be easily shown that  $k_1 = \bar{\rho}^{-1}\tilde{c}_1 \sim \bar{\rho}^{-1}\tilde{c}_2$ ,  $-\bar{\rho}^{-1}\tilde{c}_2$ ,  $\bar{\rho}^{-1}\tilde{c}_3$  and  $-\bar{\rho}^{-1}\tilde{c}_3$  in  $V_2$ . Let  $F'_3$  be the surface obtained by deforming  $F_3 - B' \cup B$  slightly keeping the exterior of  $N(B)$  fixed until it intersects  $B$  in  $c'$  [Fig. 4.1].




Fig. 4.1

Then  $\varphi_1 F'_3 \cap F'_3$  has fewer components than  $\varphi_1 F_3 \cap F_3$ . Since we can deform  $F'_3$  onto  $F_3$  by an ambient isotopy, we obtain an involution  $\varphi_2$  such that  $\varphi_2 F_3 \cup F_3$  is isotopic to  $\varphi_1 F'_3 \cup F'_3$  in  $L(2\alpha, \beta)$ . Repeating these procedures, we can show that  $\varphi_1$  is equivalent to  $\varphi$  such that  $\varphi F_3 \cap F_3$  consists of three curves of type II on  $\varphi F_3$  and  $F_3$ .

**Case 2.** In this case each curve of  $\varphi_1 F_3 \cap F_3$  is of either type I, type III or type IV.

**Assertion B.** Let  $k_1$  and  $k_2$  be simple closed curves on  $T_2$  such that  $\pi k_1$  is of type I or IV, and  $\pi k_2$  is of type III. Then  $k_1$  is not homologous to  $\varepsilon k_2$ , for  $\varepsilon = 1$  and  $-1$ , in  $V_2$ .

**Proof.** Let  $\rho$  be an autohomeomorphism of  $F_3$  such that  $\rho \pi k_1$  coincides with  $c^*$  or  $\partial N(c^*)$ , and  $\rho \pi k_2 = a^*$ . By  $\begin{pmatrix} \alpha_{11} & \alpha_{12} \\ \alpha_{21} & \alpha_{22} \end{pmatrix}$  we denote a matrix in  $GL(2, Z)$  corresponding to the isotopy class of  $\rho^{-1}$ . Then, by using Lemma 3.4,  $k_2$  or  $-k_2$  is homologous to either

$$-\alpha_{11}a'_1 - \mu_1\alpha_{11}b_1 - \alpha_{12}a'_2 + (\alpha_{11} - \mu_2\alpha_{12})b_2$$

or

$$Id \times i: M \times S(E \oplus F) \rightarrow M \times V_m^\Lambda(E \oplus F \oplus \Lambda^{m-1})$$

induces a bundle embedding  $\tilde{j}: B' \rightarrow B$ . There is a one-to-one correspondence between  $G$ -maps from  $M$  to  $V_m^\Lambda(E \oplus F \oplus \Lambda^{m-1})$  and cross sections of  $B$ , and there is also a one-to-one correspondence between their homotopies. This shows that the following two lemmas are equivalent:

**Lemma 3.** *Let*

$$f: M \rightarrow V_m^\Lambda(E \oplus F \oplus \Lambda^{m-1})$$

*be a  $G$ -map, and let*

$$P: \partial M \times [0, 1] \rightarrow V_m^\Lambda(E \oplus F \oplus \Lambda^{m-1})$$

*be a  $G$ -homotopy with  $P_0 = f|_{\partial M}$  and  $P_1(\partial M) \subset i(S(E \oplus F))$ . Then  $P$  extends to a  $G$ -homotopy*

$$Q: M \times [0, 1] \rightarrow V_m^\Lambda(E \oplus F \oplus \Lambda^{m-1})$$

*with  $Q_0 = f$  and  $Q_1(M) \subset j(S(E \oplus F))$ .*

**Lemma 4.** *Let  $N = M/G$ . Let  $s: N \rightarrow B$  be a cross section of  $B$ , and let  $P: \partial N \times [0, 1] \rightarrow B|_{\partial N}$  be a homotopy of cross section of  $B|_{\partial N}$  with  $P_0 = s|_{\partial N}$  and  $P_1(\partial N) \subset \tilde{j}(B')$ . Then  $P$  extends to a homotopy of cross section of  $B$ ,  $Q: N \times [0, 1] \rightarrow B$ , with  $Q_0 = s$  and  $Q_1(N) \subset \tilde{j}(B')$ .*

We give a proof of Lemma 4 making use of the obstruction theory. Refer to [4; Part III] for the obstruction theory.

**Proof of Lemma 4.** Since  $N$  is a smooth manifold, we obtain a triangulation of  $N$ . Let  $n = \dim S(E \oplus F)$ . Then  $\dim N \leq n$ , and  $S(E \oplus F)$ , which is the fibre of  $B'$ , is  $(n-1)$ -connected. So the cross section  $\tilde{j}^{-1}P_1$  of  $B'|_{\partial N}$  extends to a cross section  $s_1: N \rightarrow B'$  of  $B'$ . We see from Proposition 1 that  $V_m^\Lambda(E \oplus F \oplus \Lambda^{m-1})$  is also  $(n-1)$ -connected. Let  $N^{n-1}$  denote the  $(n-1)$ -skeleton of  $N$ , which contains  $\partial N$ . Then  $P$  extends to a homotopy of cross section,

$$R: N^{n-1} \times [0, 1] \rightarrow B|_{N^{n-1}},$$

with  $R_0 = s|_{N^{n-1}}$  and  $R_1 = \tilde{j}s_1|_{N^{n-1}}$ . So, if  $\dim N < n$ , the lemma is proved.

Now let  $\dim N = n$ . Let  $B(\pi_n)$  and  $B'(\pi_n)$  be the bundles of coefficients associated with the bundles  $B$  and  $B'$  by the  $n$ -th homotopy group, respectively. Also let  $C^n(N; B(\pi_n))$  and  $C^n(N; B'(\pi_n))$  be the groups of  $n$ -cochains of  $N$  with coefficients in  $B(\pi_n)$  and  $B'(\pi_n)$ , respectively. The bundle embedding  $\tilde{j}: B' \rightarrow B$  induces a group homomorphism

$$\tilde{j}_*: C^n(N; B'(\pi_n)) \rightarrow C^n(N; B(\pi_n)).$$

We see from Proposition 1 that  $\tilde{j}_*$  is an epimorphism. Let

homotopy set.

Our result is

**Theorem 2.** *Let  $E, F$  be representations of a compact Lie group  $G$  over  $\Lambda$ . Let*

$$j_*: [S(E), S(E \oplus F)]_G \rightarrow [S(E), V_m^\Lambda(E \oplus F \oplus \Lambda^{m-1})]_G$$

*be the transformation induced from the  $G$ -map*

$$j: S(E \oplus F) \rightarrow V_m^\Lambda(E \oplus F \oplus \Lambda^{m-1}).$$

*Then*

- (a)  $j_*$  is surjective,
- (b)  $j_*$  is bijective in particular in each case of the followings (i), (ii):
- (i)  $\Lambda = \mathbf{R}$ ,  $\dim_{\mathbf{R}} E^H$  is odd for any  $H \in \mathfrak{N}(E, F) = \{H \in \mathfrak{M}_r(E) \mid \dim_{\mathbf{R}} F^H = 0\}$ ,

*and*

$$r = \prod_{H \in \mathfrak{N}(E, F)} r_H: [S(E), S(E \oplus F)]_G \rightarrow \prod_{H \in \mathfrak{N}(E, F)} [S(E^H), S(E^H \oplus F^H)]$$

*is injective,*

- (ii)  $\Lambda = \mathbf{C}$  or  $\mathbf{Q}$ , and  $r$  is injective,
- (c) if  $\dim_{\mathbf{R}} E^G \geq 2$  then  $[S(E), V_m^\Lambda(E \oplus F \oplus \Lambda^{m-1})]_G$  has a group structure and  $j_*$  is a group homomorphism.

NOTE. The injectivity of  $r$  is studied by several authors, e.g., Hauschild [1; Satz 4.5].

In the subsequent sections we prove Theorem 2. Section 2 is devoted to preliminary lemmas. Section 3 is devoted to proving the surjectivity of  $j_*$ , and section 4 is devoted to proving the injectivity of  $j_*$ . In section 5 we give a group structure to  $[S(E), V_m^\Lambda(E \oplus F \oplus \Lambda^{m-1})]_G$  so that  $j_*$  is a group homomorphism.

## 2. Preliminary lemmas

Let  $E, F$  be representations of a compact Lie group  $G$  over  $\Lambda$ , and let  $M$  be a compact, smooth, free  $G$ -manifold with  $\dim M \leq \dim S(E \oplus F)$ . Consider the fibre bundles

$$B = M \times_G V_m^\Lambda(E \oplus F \oplus \Lambda^{m-1}) \rightarrow M/G$$

with fibre  $V_m^\Lambda(E \oplus F \oplus \Lambda^{m-1})$ , and

$$B' = M \times_G S(E \oplus F) \rightarrow M/G$$

with fibre  $S(E \oplus F)$ . The  $G$ -map

$$d = d(s, R, \tilde{j}s_1) \in C^n(N; B(\pi_n))$$

be the deformation  $n$ -cochain. (See [4; p 172].) There is  $d' \in C^n(N; B'(\pi_n))$  with  $\tilde{j}_*(d') = d$ . By [4; 33.9] there is a cross section  $s_2$  of  $B'$  such that  $s_2$  agrees with  $s_1$  on  $N^{n-1}$  and  $d(s_1, s_2) = -d'$ , where  $d(s_1, s_2)$  is the difference  $n$ -cochain. (Also see [4; p 172].) We see

$$d(\tilde{j}s_1, \tilde{j}s_2) = \tilde{j}_*(d(s_1, s_2)) = -d.$$

We define a homotopy of cross section of  $B|N^{n-1}$ ,

$$S: N^{n-1} \times [0, 1] \rightarrow B|N^{n-1},$$

by

$$S(x, t) = \begin{cases} R(x, 2t) & \text{if } 0 \leq t \leq 1/2 \\ \tilde{j}s_2(x) & \text{if } 1/2 \leq t \leq 1. \end{cases}$$

By [4; 33.7],

$$\begin{aligned} d(s, S, \tilde{j}s_2) &= d(s, R, \tilde{j}s_1) + d(\tilde{j}s_1, \tilde{j}s_2) \\ &= d - d \\ &= 0. \end{aligned}$$

By this and [4; 33.8],  $S$  extends to a homotopy of cross section of  $B, T: N \times [0, 1] \rightarrow B$ , with  $T_0 = s$  and  $T_1 = \tilde{j}s_2$ . Let  $\partial N \times [0, 1] \subset N$  be a collar of  $\partial N$  in  $N$ . Then we define a homotopy  $Q: N \times [0, 1] \rightarrow B$  as, for  $x \in N$  and  $t \in [0, 1]$ ,

$$\begin{aligned} Q(x, t) &= T(x, t) \quad \text{if } x \in N - \partial N \times [0, 1], \\ Q(x, t) &= T(x, t/(2-r)) \quad \text{if } x = (y, r) \in \partial N \times [0, 1] \quad \text{and } 2t+r \leq 2, \\ Q(x, t) &= T(x, (t+r-1)/r) \quad \text{if } x = (y, r) \in \partial N \times [0, 1], \quad 2t+r \geq 2 \\ &\text{and } r \neq 0. \end{aligned}$$

$Q$  is well-defined, and is an extension of  $P$  with the desired property.

Q.E.D.

### 3. The surjectivity of $j_*$

Let  $E, F$  be representations of a compact Lie group  $G$  over  $\Lambda$ , and let

$$j_*: [S(E), S(E \oplus F)]_G \rightarrow [S(E), V_m^\Lambda(E \oplus F \oplus \Lambda^{m-1})]_G$$

be the transformation induced from the  $G$ -map

$$j: S(E \oplus F) \rightarrow V_m^\Lambda(E \oplus F \oplus \Lambda^{m-1}).$$

The purpose of this section is to prove the surjectivity of  $j_*$ . Since  $j$  is an embedding, it suffices to prove the following fact:

**Lemma 5.** *Let*

$$f: S(E) \rightarrow V_m^\Delta(E \oplus F \oplus \Lambda^{m-1})$$

*be a  $G$ -map, and let  $N$  be a compact smooth  $G$ -submanifold of  $S(E)$  with  $\dim N = \dim S(E)$ . Let*

$$T: N \times [0, 1] \rightarrow V_m^\Delta(E \oplus F \oplus \Lambda^{m-1})$$

*be a  $G$ -homotopy with  $T_0 = f|N$  and  $T_1(N) \subset j(S(E \oplus F))$ . Then  $T$  extends to a  $G$ -homotopy*

$$R: S(E) \times [0, 1] \rightarrow V_m^\Delta(E \oplus F \oplus \Lambda^{m-1})$$

*with  $R_0 = f$  and  $R_1(S(E)) \subset i(S(E \oplus F))$ .*

*Proof.*  $\mathfrak{M}(E)$  is a finite set. Let us number its elements

$$\mathfrak{M}(E) = \{(H_1), (H_2), \dots, (H_a)\}$$

in such a way that if  $i < k$  then  $(H_i) \subset (H_k)$ . Consider the following Assertion:

**ASSERTION.** *There are compact smooth  $G$ -submanifolds  $M_0, M_1, \dots, M_a$  of  $S(E)$  such that*

$$\begin{aligned} \dim M_i &= \dim S(E) \quad \text{for } i = 0, 1, \dots, a, \quad M_0 \supset N, \quad \text{and} \\ \text{Int } M_i &\supset M_{i-1} \cup S(E)_{(H_i)} \quad \text{for } i = 1, 2, \dots, a. \end{aligned}$$

*Furthermore there are  $G$ -homotopies  $R^{(0)}, R^{(1)}, \dots, R^{(a)}$  such that*

$$\begin{aligned} R^{(i)}: M_i \times [0, 1] &\rightarrow V_m^\Delta(E \oplus F \oplus \Lambda^{m-1}) \quad \text{for } i = 0, 1, \dots, a, \\ R_0^{(i)} &= f|_{M_i} \quad \text{for } i = 0, 1, \dots, a, \\ R_1^{(i)}(M_i) &\subset j(S(E \oplus F)) \quad \text{for } i = 0, 1, \dots, a, \\ R^{(0)}|_{N \times [0, 1]} &= T, \quad \text{and} \\ R^{(i)}|_{M_{i-1} \times [0, 1]} &= R^{(i-1)} \quad \text{for } i = 1, 2, \dots, a. \end{aligned}$$

Lemma 5 follows from the Assertion since  $M_a = S(E)$ . In the following we prove the Assertion.

$N$  and  $T$  satisfy the conditions for  $M_0$  and  $R^{(0)}$ , respectively. Suppose that  $M_0, \dots, M_{i-1}$ , and  $R^{(0)}, \dots, R^{(i-1)}$  are constructed. Put

$$M = (S(E) - \text{Int } M_{i-1})^{H_i} = S(E^{H_i}) - \text{Int } M_{i-1}^{H_i}.$$

Then  $M$  is a compact smooth manifold with boundary  $\partial M = M \cap \partial M_{i-1}$ . Moreover  $M$  is  $N(H_i)$ -invariant, and all isotropy subgroups on  $M$  are  $H_i$ . So  $M$  becomes a free  $N(H_i)/H_i$ -manifold. Regard  $E^{H_i}$  and  $F^{H_i}$  as representations of  $N(H_i)/H_i$ . By Lemma 3 there is an  $N(H_i)/H_i$ -homotopy

$$Q: M \times [0, 1] \rightarrow V_m^\Delta(E^{H_i} \oplus F^{H_i} \oplus \Lambda^{m-1})$$

such that

$$\begin{aligned} Q_0 &= f|_M, \\ Q_1(M) &\subset i(S(E^{H_i} \oplus F^{H_i})), \quad \text{and} \\ Q|_{\partial M \times [0, 1]} &= R^{(i-1)}|_{\partial M \times [0, 1]}. \end{aligned}$$

Since  $G(M) = G \times_{N(H_i)} M$ , we may extend  $Q$  to a  $G$ -homotopy

$$Q': G(M) \times [0, 1] \rightarrow G(V_m^\Delta(E^{H_i} \oplus F^{H_i} \oplus \Lambda^{m-1})) \subset V_m^\Delta(E \oplus F \oplus \Lambda^{m-1})$$


such that

$$\begin{aligned} Q'_0 &= f|_{G(M)}, \\ Q'_1(G(M)) &\subset i(S(E \oplus F)), \quad \text{and} \\ Q'|_{\partial G(M) \times [0, 1]} &= R^{(i-1)}|_{\partial G(M) \times [0, 1]}. \end{aligned}$$

Applying [3; Lemma 1.1] to the  $G$ -manifold  $A = S(E) - \text{Int } M_{i-1}$  and the submanifold  $G(M)$  of  $A$ , we obtain compact  $G$ -submanifolds  $K, L$  of  $A$  such that

- (i)  $K \cup L = A$ ,
- (ii)  $\partial L = L \cap K$ ,  
 $\partial K = \partial L \cup \partial A = \partial L \cup \partial M_{i-1}$ ,  
 $\partial M_{i-1} \cap \partial L = \emptyset$ ,
- (iii)  $\partial M_{i-1} \cup G(M) \subset K$ , and
- (iv)  $K$  is a mapping cylinder of some  $G$ -map

$$\psi: \partial L \rightarrow \partial M_{i-1} \cup G(M).$$



Put  $M_i = M_{i-1} \cup K$  in  $S(E)$ . Then  $M_i$  is a compact smooth  $G$ -submanifold of  $S(E)$  with  $\dim M_i = \dim S(E)$ , and with  $\text{Int } M_i \supset M_{i-1} \cup S(E)_{(H_i)}$ . According to (iv), let us denote a point of  $K$  by the form  $[y, s]$ , where  $y \in \partial L$  and  $s \in [0, 1]$ . Under this form  $[y, 1] = y$  and  $[y, 0] = \psi(y)$ . We define a  $G$ -homotopy

$$R^{(i)}: M_i \times [0, 1] \rightarrow V_m^\Delta(E \oplus F \oplus \Lambda^{m-1})$$

as the following: For  $(x, t) \in M_i \times [0, 1]$ ,

$$\begin{aligned} R^{(i)}(x, t) &= R^{(i-1)}(x, t) \quad \text{if } x \in M_{i-1}, \\ R^{(i)}(x, t) &= f([y, s-2t]) \quad \text{if } x = [y, s] \in K \quad \text{and } 2t \leq s, \\ R^{(i)}(x, t) &= R^{(i-1)}(\psi(y), (2t-s)/(2-s)) \quad \text{if } x = [y, s] \in K, \\ &\quad \psi(y) \in \partial M_{i-1} \quad \text{and } s \leq 2t, \\ R^{(i)}(x, t) &= Q'(\psi(y), (2t-s)/(2-s)) \quad \text{if } x = [y, s] \in K, \\ &\quad \psi(y) \in G(M) \quad \text{and } s \leq 2t. \end{aligned}$$

$M_i$  and  $R^{(i)}$  constructed above satisfy the conditions in the Assertion. Thus this completes the proof. Q.E.D.

Now let  $X, Y$  be  $G$ -spaces, and let  $x_0 \in X^G, y_0 \in Y^G$ . Denote by

$$[(X, x_0), (Y, y_0)]_G$$

the set of  $G$ -homotopy classes rel.  $x_0$  of  $G$ -maps  $f: X \rightarrow Y$  with  $f(x_0) = y_0$ .

The following Proposition is required in section 5.

**Proposition 6.** *Let  $E, F$  be representations of  $G$ , and let  $x_0 \in S(E^G), y_0 \in S(E^G \oplus F^G)$ . Then*

$$j_*: [(S(E), x_0), (S(E \oplus F), y_0)]_G \rightarrow [(S(E), x_0), (V_m^\Lambda(E \oplus F \oplus \Lambda^{m-1}), j(y_0))]_G$$

*is surjective.*

Proof. Let

$$f: S(E) \rightarrow V_m^\Lambda(E \oplus F \oplus \Lambda^{m-1})$$

be a  $G$ -map with  $f(x_0) = j(y_0)$ . Let  $D$  be a  $G$ -invariant, top-dimensional, small disc in  $S(E)$  with  $x_0$  as its center. We may deform  $f$  to a  $G$ -map  $f'$  such that  $f'(D) = j(y_0)$  and  $f' \simeq f$  rel.  $x_0$ . By Lemma 5 there is a  $G$ -homotopy

$$R: S(E) \times [0, 1] \rightarrow V_m^\Lambda(E \oplus F \oplus \Lambda^{m-1})$$

such that

$$\begin{aligned} R_0 &= f', \\ R_1(S(E)) &\subset j(S(E \oplus F)), \quad \text{and} \\ R(D \times [0, 1]) &= j(y_0). \end{aligned}$$

Then  $f'$  is  $G$ -homotopic to  $R_1$  rel.  $x_0$ . This proves the Proposition. Q.E.D.

#### 4. The injectivity of $j_*$

Let  $E, F$  be representations of a compact Lie group  $G$  over  $\Lambda$ , and let



$$j_*: [S(E), S(E \oplus F)]_G \rightarrow [S(E), V_m^\Lambda(E \oplus F \oplus \Lambda^{m-1})]_G$$

be the transformation induced from the  $G$ -map

$$j: S(E \oplus F) \rightarrow V_m^\Lambda(E \oplus F \oplus \Lambda^{m-1}).$$

The purpose of this section is to prove the injectivity of  $j_*$  under the assumption (i) or (ii) in Theorem 2.

For any closed subgroup  $H$  of  $G$ , let

$$j^H = j|_{S(E^H \oplus F^H)}: S(E^H \oplus F^H) \rightarrow V_m^\Lambda(E^H \oplus F^H \oplus \Lambda^{m-1}).$$

The following diagram is commutative:

$$\begin{array}{ccc} [S(E), S(E \oplus F)]_G & \xrightarrow{j_*} & [S(E), V_m^\Lambda(E \oplus F \oplus \Lambda^{m-1})]_G \\ r_H \downarrow & & \downarrow r'_H \\ [S(E^H), S(E^H \oplus F^H)] & \xrightarrow{j_*^H} & [S(E^H), V_m^\Lambda(E^H \oplus F^H \oplus \Lambda^{m-1})] \end{array}$$

where  $r_H$  and  $r'_H$  are the transformations restricting to the fixed point set by  $H$ .

Now suppose

$$j_*(\alpha) = j_*(\beta)$$

for  $\alpha, \beta \in [S(E), S(E \oplus F)]_G$ . Then, by the commutativity of the above diagram,

$$j_*^H r_H(\alpha) = j_*^H r_H(\beta)$$

for any closed subgroup  $H$  of  $G$ . Proposition 1 implies that  $j_*^H$  is an isomorphism under the assumption (i) or (ii) in Theorem 2. Thus

$$r_H(\alpha) = r_H(\beta)$$

for any  $H$ . Hence  $r(\alpha) = r(\beta)$ . By the assumption  $r$  is injective, hence  $\alpha = \beta$ . Thus  $j_*$  is injective.

## 5. The group structure

Let  $E, F$  be representations of a compact Lie group  $G$  over  $\Lambda$ . Suppose  $\dim_{\mathbb{R}} E^G \geq 2$ . Then, according to [3; Section 6],  $[S(E), S(E \oplus F)]_G$  has a group structure. In the similar way we may give a group structure to

$$[S(E), V_m^\Lambda(E \oplus F \oplus \Lambda^{m-1})]_G$$

so that

$$j_*: [S(E), S(E \oplus F)]_G \rightarrow [S(E), V_m^\Lambda(E \oplus F \oplus \Lambda^{m-1})]_G$$

is a group homomorphism. To show this is the purpose of this section.

**Lemma 7.** Suppose  $\dim_R E^G = 2$  and  $x_0 \in S(E^G)$ . Let

$$\omega: [0, 1] \rightarrow S(E^G) \subset S(E)$$

be a path with  $\omega(0) = \omega(1) = x_0$ . Then there is a  $G$ -homotopy

$$H: S(E) \times [0, 1] \rightarrow S(E)$$

such that

$$\begin{aligned} H_0 = H_1 = Id, \quad \text{and} \\ H(x_0, t) = \omega(t) \quad \text{for any } t \in [0, 1]. \end{aligned}$$

*Proof.* Choose a homotopy

$$J: S(E^G) \times [0, 1] \rightarrow S(E^G) \subset S(E)$$

such that

$$\begin{aligned} J_0 = J_1 = \text{the inclusion}, \quad \text{and} \\ J(x_0, t) = \omega(t) \quad \text{for any } t \in [0, 1]. \end{aligned}$$

Denote by  $(E^G)^\perp$  the orthogonal complement of  $E^G$  in  $E$ , and denote a point of  $E$  by the form  $x + y$  where  $x \in E^G$  and  $y \in (E^G)^\perp$ . Define

$$H: S(E) \times [0, 1] \rightarrow S(E)$$

as

$$\begin{aligned} H(x + y, t) &= \|x\| J(x/\|x\|, t) + y \quad \text{if } x \neq 0, \quad \text{and} \\ H(x + y, t) &= y \quad \text{if } x = 0. \end{aligned}$$

Then  $H$  is a  $G$ -homotopy with the desired property. Q.E.D.

**Lemma 8.** Suppose  $\dim_R E^G \geq 2$ ,  $x_0 \in S(E^G)$  and  $y_0 \in S(E^G \oplus F^G)$ . Then the natural transformations

$$\psi_1: [(S(E), x_0), (S(E \oplus F), y_0)]_G \rightarrow [S(E), S(E \oplus F)]_G$$

and

$$\psi_2: [(S(E), x_0), (V_m^\Delta(E \oplus F \oplus \Lambda^{m-1}), j(y_0))]_G \rightarrow [S(E), V_m^\Delta(E \oplus F \oplus \Lambda^{m-1})]_G$$

are bijective.

*Proof.* Consider the commutative diagram:

$$\begin{array}{ccc} [(S(E), x_0), (S(E \oplus F), y_0)]_G & \xrightarrow{\psi_1} & [S(E), S(E \oplus F)]_G \\ j_* \downarrow & & \downarrow j_* \\ [(S(E), x_0), (V_m^\Delta(E \oplus F \oplus \Lambda^{m-1}), j(y_0))]_G & \xrightarrow{\psi_2} & [S(E), V_m^\Delta(E \oplus F \oplus \Lambda^{m-1})]_G \end{array}$$

In [3; Section 6],  $\psi_1$  is already seen to be bijective. The two  $j_*$  are surjective by the arguments in section 3. So it follows that  $\psi_2$  is surjective.

It only remains to show that  $\psi_2$  is injective. Suppose

$$\psi_2(\alpha) = \psi_2(\beta)$$

for  $\alpha, \beta \in [(S(E), x_0), (V_m^\Delta(E \oplus F \oplus \Lambda^{m-1}), j(y_0))]_G$ . Since  $j_*$  is surjective, there are  $G$ -maps

$$f, g: S(E) \rightarrow S(E \oplus F)$$

such that  $f(x_0) = y_0$ ,  $g(x_0) = y_0$ , and  $jf, jg$  are representatives of  $\alpha, \beta$ , respectively. There also is a  $G$ -homotopy

$$K: S(E) \times [0, 1] \rightarrow V_m^\Delta(E \oplus F \oplus \Lambda^{m-1})$$

with  $K_0 = jf$  and  $K_1 = jg$ . Define a path

$$\omega: [0, 1] \rightarrow V_m^\Delta(E^G \oplus F^G \oplus \Lambda^{m-1})$$

by  $\omega(t) = K(x_0, t)$  for  $t \in [0, 1]$ . Then

$$\omega(0) = \omega(1) = j(y_0).$$

By Proposition 1 there is a path

$$\omega': [0, 1] \rightarrow S(E^G \oplus F^G)$$

such that

$$\begin{aligned} \omega'(0) &= \omega'(1) = y_0, & \text{and} \\ \omega &\simeq j\omega' \text{ rel. } \{0, 1\}. \end{aligned}$$

Let  $D$  be a  $G$ -invariant, top-dimensional, small disc in  $S(E)$  with  $x_0$  as its center, and let  $D' = \frac{1}{2}D$ . By radius contraction we may deform  $K$  to a  $G$ -homotopy

$$K': S(E) \times [0, 1] \rightarrow V_m^\Delta(E \oplus F \oplus \Lambda^{m-1})$$

such that  $K'(x, t) = j\omega'(t)$  for  $x \in D'$  and  $t \in [0, 1]$ . Moreover, if we put  $f' = K'_0$  and  $g' = K'_1$ , then

$$\begin{aligned} f'(S(E)) &\subset i(S(E \oplus F)), \\ g'(S(E)) &\subset i(S(E \oplus F)), \end{aligned}$$

and  $f', g'$  are  $G$ -homotopic to  $jf, jg$  rel.  $x_0$ , respectively.

So, to show  $\alpha = \beta$  we must show that  $f'$  is  $G$ -homotopic to  $g'$  rel.  $x_0$ .

(i) Suppose  $\dim_{\mathbb{R}} E^G \oplus F^G > 2$ . By Proposition 1,  $j\omega'$  is homotopic to the constant path at  $j(y_0)$  rel.  $\{0, 1\}$ . From this we may deform  $K'$  to a  $G$ -homotopy

$$K'': S(E) \times [0, 1] \rightarrow V_m^\Delta(E \oplus F \oplus \Lambda^{m-1})$$

such that

$$\begin{aligned} K'_0 &= f', \quad K'_1 = g', \quad \text{and} \\ K''(x_0, t) &= j(y_0) \quad \text{for any } t \in [0, 1]. \end{aligned}$$

Therefore  $f'$  is  $G$ -homotopic to  $g'$  rel.  $x_0$ .

(ii) Suppose  $\dim_R E^G \oplus F^G = 2$ . Define a path

$$\omega'': [0, 1] \rightarrow S(E^G \oplus F^G)$$

by  $\omega'' = (\omega')^{-1}$ , i.e.,  $\omega''(t) = \omega'(1-t)$ . Applying Lemma 7 to the path  $\omega''$ , there is a  $G$ -homotopy

$$H: S(E \oplus F) \times [0, 1] \rightarrow S(E \oplus F)$$

such that

$$\begin{aligned} H_0 &= H_1 = Id, \quad \text{and} \\ H(y_0, t) &= \omega''(t) \quad \text{for any } t \in [0, 1]. \end{aligned}$$

Define a  $G$ -homotopy

$$L: S(E) \times [0, 1] \rightarrow V_m^\Lambda(E \oplus F \oplus \Lambda^{m-1})$$

as, for  $x \in S(E)$  and  $t \in [0, 1]$ ,

$$\begin{aligned} L(x, t) &= K'(x, 2t) \quad \text{if } 0 \leq t \leq 1/2, \quad \text{and} \\ L(x, t) &= jH(j^{-1}g'(x), 2t-1) \quad \text{if } 1/2 \leq t \leq 1. \end{aligned}$$

Then

$$\begin{aligned} L_0 &= f', \quad L_1 = g', \quad \text{and} \\ L(x, t) &= j\omega' * j\omega''(t) \quad \text{for } x \in D' \quad \text{and } t \in [0, 1]. \end{aligned}$$

$j\omega' * j\omega''$  is homotopic to the constant path at  $j(y_0)$  rel.  $\{0, 1\}$ . So we may deform  $L$  to a  $G$ -homotopy

$$L': S(E) \times [0, 1] \rightarrow V_m^\Lambda(E \oplus F \oplus \Lambda^{m-1})$$

such that

$$\begin{aligned} L'_0 &= f', \quad L'_1 = g', \quad \text{and} \\ L'(x_0, t) &= j(y_0) \quad \text{for any } t \in [0, 1]. \end{aligned}$$

Therefore  $f'$  is  $G$ -homotopic to  $g'$  rel.  $x_0$ .

Q.E.D.

Now suppose  $\dim_R E^G \geq 2$ ,  $x_0 \in S(E^G)$  and  $y_0 \in S(E^G \oplus F^G)$ . Let  $\lambda$  be the real one-dimensional subspace of  $E$  spanned by  $x_0$ , and let  $\lambda^\perp$  be the orthogonal complement of  $\lambda$  in  $E$ . We may identify  $S(E)$  with a nonreduced suspension

$$\Sigma S(\lambda^\perp) = [0, 1] \times S(\lambda^\perp) / \sim.$$



Under this identification  $x_0=[0, x]$  and  $-x_0=[1, x]$  for  $x \in S(\lambda^\perp)$ . Let  $Y$  be one of  $S(E \oplus F)$  and  $V_m^\Delta(E \oplus F \oplus \Lambda^{m-1})$ . Put  $z_0=y_0$  if  $Y$  is the former, and  $z_0=j(y_0)$  if  $Y$  is the latter. Then we may give a group structure to  $[S(E), Y]_G$  as follows. Let  $[f], [g] \in [S(E), Y]_G$ . By Lemma 8 we may choose  $f$  and  $g$  in such a way that  $f(-x_0)=z_0$  and  $g(x_0)=z_0$ . Define  $h: S(E) \rightarrow Y$  as, for  $[t, x] \in \Sigma S(\lambda^\perp) = S(E)$ ,

$$\begin{aligned} h([t, x]) &= f([2t, x]) \quad \text{if } 0 \leq t \leq 1/2, \quad \text{and} \\ h([t, x]) &= g([2t-1, x]) \quad \text{if } 1/2 \leq t \leq 1. \end{aligned}$$

Define  $[f] + [g] = [h]$ . This gives a group structure to  $[S(E), Y]_G$ , and the transformation

$$j_*: [S(E), S(E \oplus F)]_G \rightarrow [S(E), V_m^\Delta(E \oplus F \oplus \Lambda^{m-1})]_G$$

becomes a group homomorphism. We note that this group structure does not depend on the choice of  $x_0 \in S(E^G)$  and  $y_0 \in S(E^G \oplus F^G)$ .

#### References

- [1] H. Hauschild: *Zerspaltung äquivarianter Homotopiemengen*, Math. Ann. **230** (1977), 279–292.
- [2] D. Husemoller: *Fibre bundles* (second edition), Springer-Verlag, New York-Heidelberg-Berlin, 1975.
- [3] R.L. Rubinsztein: *On the equivariant homotopy of spheres*, Dissertationes Math. (Rozprawy Mat.), **134** (1976).
- [4] N. Steenrod: *The topology of fibre bundles*, Princeton University Press, Princeton, 1951.

Department of Mathematics  
Yamaguchi University  
Yamaguchi 753, Japan