

Theorem 14 is false as stated. We will show elsewhere that a positive parabolic function v on $(-1, 1) \times (0, 1)$ has the integral representation given, for a finite measure α , if and only if $\lim_{t \rightarrow 1^-} v(0, t) < \infty$. Here we take $p_0 = (0, 1)$.

We also note the following errors. Theorem 5 should read " $B|_{\Gamma} = C(\Gamma)$ " instead of " $B|_{\Gamma}$ dense in $C(\Gamma)$ ". On p. 347 the definition B_n should read: "all C^2 functions which satisfy $u_{xx}(x, y) - u_{yy}(x, y) = 0$ and $u_y(x, 0) = 0$ ".

Corrections to

SUBSEQUENCES AND REARRANGEMENTS OF SEQUENCES IN FK SPACES

ROBERT DEVOS

Volume 64 (1976), 129-135

In Lemma 1 and all subsequent results, whenever we take a sequence in $E \setminus l^p$ we need take it in $E \setminus (l^p \oplus \{e\})$. This error was pointed out by R. A. Shoop.

Corrections to

EXACT FUNCTORS AND MEASURABLE CARDINALS

ANDREAS BLASS

Volume 63 (1977), 335-346

Professor V. Trnková and J. Reiterman have informed me that the main results in [1] are contained in or easily deducible from [3] and that the example constructed in the last paragraph of [1] was also obtained in [2].

1. A. Blass, *Exact functors and measurable cardinals*, Pacific J. Math., **63** (1976), 335-346.
2. J. Reiterman, *An example concerning set-functors*, Comm. Math. Univ. Carolinae, **12** (1971), 227-233.
3. V. Trnková, *On descriptive classification of setfunctors*, Comm. Math. Univ. Carolinae, **12** (1971), 143-174 and 345-357.