

NILPOTENCY AND SPECTRAL OPERATORS

JAMES E. SIMPSON

1. **Introduction.** The present paper is concerned with conditions under which the quasi-nilpotent part of a spectral operator is actually nilpotent of some order k . As might be expected, the case of a spectral operator on a Hilbert space has been settled longest. (See [4].) The case of a Banach space has been treated quite thoroughly by C. A. McCarthy [7] who showed that with a certain rate of growth condition on Q , the nilpotent part of the spectral operator $T = S + Q$, satisfies $Q^{m+2} = 0$, where the m is a positive integer involved in the rate of growth condition. He also discusses more special cases in which $Q^{m+1} = 0$ and provides examples to show that these exponents are the lowest possible in each case. The question of extending these results to general locally convex spaces could not even be formulated until a theory of spectral operators in these spaces had been devised. The work of C. Ionescu Tulcea [5] having laid the foundations in this area, we may now attempt to solve the problem of generalizing McCarthy's results. It is shown below that his theorems, and indeed some part of the proofs, may be carried over to the locally convex case, with a suitable reformulation of some of the conditions and reworking of some of the supporting theory.

The basic assumptions are as follows. E denotes a locally convex linear topological space over the field, C , of complex numbers. Moreover, E is assumed to be separated, barrelled and quasi-complete. The strong dual of E is denoted by E' . The space of continuous linear mappings of E into itself is $\mathcal{L}(E, E)$, which we shall always assume to be given the topology \mathcal{T}_b of uniform convergence on the bounded subsets of E . We denote the adjoint of T by $'T$, for each $T \in \mathcal{L}(E, E)$. The resolvent set of T , $\text{res } T$, is a certain subset of \hat{C} , the one-point compactification of C . Specifically, $\lambda \in \text{res } T$ provided there is a neighborhood V_λ of λ in \hat{C} and a function R_r with domain $V_\lambda \cap C$ and range in $\mathcal{L}(E, E)$ such that

(a) the set $\{R_r(z)x: z \in V_\lambda \cap C\}$ is a bounded subset of E for each $x \in E$, and

(b) $R_r(z)(zI - T) = (zI - T)R_r(z) = I$ for all $z \in V_\lambda \cap C$. The complement of $\text{res } T$, in \hat{C} , is the spectrum of T , denoted $\text{sp } T$. If $\infty \notin \text{sp } T$, then $\text{sp } T$ is compact in C and we say T is regular. We

Received June 6, 1963. Most of the work appearing in this paper was done while the author was a National Science Foundation Graduate Fellow and appears in somewhat modified form in his dissertation, submitted to Yale University while he was a candidate for the doctoral degree.

denote by $B^\infty(C)$ the algebra of bounded complex-valued Baire functions on C , with norm $\|f\|_\infty = \sup_{z \in C} |f(z)|$, and by $S(C)$ the Baire sets, i.e., those subsets, A , of C whose characteristic function, φ_A , is in $B^\infty(C)$. We denote by M^1 the set of bounded Radon measures on C , with norm $\|\mu\| = \sup \left\{ \left| \int_C f d\mu \right| : f \in B^\infty(C), \|f\|_\infty = 1 \right\}$. By $\text{supp } \mu$ we mean the support of the measure μ . Further information about locally convex spaces and Radon measures may be found in the well-known Bourbaki books [1] and [2]. Concerning the resolvent and spectrum of T , see L. Waelbroeck [10] and [11].

Turning now to spectral operators, we review some of the definitions and theorems to be found in the above mentioned work of Ionescu Tulcea [5]. See also F. Maeda [6] and H. Schaeffer [8]. The latter paper contains a monumental quantity of information about spectral measures and extensions and proofs of many of the observations listed in this paragraph. With E as above, let $\mathcal{F} = \{\mu_{x,x'} : x \in E, x' \in E'\}$ be a set of bounded Radon measures on C , indexed as indicated by $E \times E'$. We say that \mathcal{F} is a spectral family of measures if there is a continuous algebraic representation of $B^\infty(C)$ in $\mathcal{L}(E, E)$, denoted by $f \rightarrow U_{\mathcal{F},f}$ (or U_f , if no confusion will ensue) such that $U_I = I$, and $\langle U_f x, x' \rangle = \int_C f d\mu_{x,x'}$ for all $x \in E, x' \in E'$, and $f \in B^\infty(C)$. The function l is defined by $l(z) = 1$ for all $z \in C$. By $\text{sup } \mathcal{F}$ we mean $\overline{\bigcup_{\substack{x \in E \\ x' \in E'}} \text{supp } \mu_{x,x'}}$. We say that $T \in \mathcal{L}(E, E)$ commutes with a spectral family \mathcal{F} , provided $TU_f = U_f T$ for all $f \in B^\infty(C)$. Denote by $P_{\mathcal{F}}$ the $\mathcal{L}(E, E)$ -valued function defined on $S(C)$ by $P_{\mathcal{F}}(\sigma) = U_{\varphi_\sigma}$. Then $P_{\mathcal{F}}$ has the following properties:

- (i) $P_{\mathcal{F}}(\phi) = 0$.
- (ii) $P_{\mathcal{F}}(\sigma \cap \delta) = P_{\mathcal{F}}(\sigma)P_{\mathcal{F}}(\delta)$ for all $\sigma, \delta \in S(C)$.
- (iii) The set function m_x , defined on $S(C)$ with values in E by $m_x(\sigma) = P_{\mathcal{F}}(\sigma)x$, is countably additive for each $x \in E$.
- (iv) $P_{\mathcal{F}}(C) = I$.

We shall call $P_{\mathcal{F}}$ the spectral measure associated with \mathcal{F} . It is quite common to write $\int_C f dP_{\mathcal{F}}$ for U_f . For each $\sigma \in S(C)$, let $E_\sigma = P_{\mathcal{F}}(\sigma)E$. If $T \in \mathcal{L}(E, E)$ commutes with \mathcal{F} , then $T_\sigma : E_\sigma \rightarrow E_\sigma$ may be defined by $T_\sigma x = Tx$ for all $x \in E_\sigma$. An element T of $\mathcal{L}(E, E)$ is said to be a spectral operator if there is a spectral family \mathcal{F} ($= \mathcal{F}_T$, if necessary) on C such that T commutes with \mathcal{F} and $\text{sp } T_\sigma \subset \sigma$ for every compact subset σ of C . An element Q of $\mathcal{L}(E, E)$ is said to be quasi-nilpotent if $\lim_n |\langle Q^n x, x' \rangle|^{1/n} = 0$ for all $x \in E, x' \in E'$. An element S of $\mathcal{L}(E, E)$ is said to be scalar relative to a spectral family \mathcal{F} if the function $f: f(z) = z$ is $\mu_{x,x'}$ -measurable for all $\mu_{x,x'} \in \mathcal{F}$ and $\langle Sx, x' \rangle = \int_C f d\mu_{x,x'}$ for all $x \in E, x' \in E'$. Finally we mention con-

dition $P\mathcal{E}$), which is described in [5] and [6]. The central decomposition theorem of spectral theory then is that if T is a spectral operator whose associated spectral family \mathcal{F}_T satisfies condition $P\mathcal{E}$), there is a unique decomposition $T = S + Q$ where S is scalar relative to \mathcal{F}_T and Q is quasi-nilpotent.

One last tool will be needed below, namely an “operational calculus” for regular operators. Suppose T is regular. Let $\mathcal{G}(T)$ be the class of all complex-valued functions f , analytic on an open set $D(f)$ which contains $\text{sp } T$. Let D be any Cauchy domain satisfying $\text{sp } T \subset D \subset \bar{D} \subset D(f)$. Then for each $f \in \mathcal{G}(T)$, define $f(T) = 1/2\pi i \int_{\Gamma} f(z)R_T(z)dz$, where Γ is the boundary of D . We then have the following theorem.

THEOREM 1. *For any $f \in \mathcal{G}(T)$, $f(T)$ is a well-defined element of $\mathcal{L}(E, E)$ independent of the choice of D (provided it satisfies the above conditions). Moreover, if f and g are both in $\mathcal{G}(T)$, then $(f + g)(T) = f(T) + g(T)$ and $(fg)(T) = f(T)g(T)$. If Γ is a circle of sufficiently large radius to contain $\text{sp } T$ in its interior, then $T = 1/2\pi i \int_{\Gamma} zR_T(z)dz$ and $I = 1/2\pi i \int_{\Gamma} R_T(z)dz$. Finally,*

$$\left\langle \int_{\Gamma} f(z)R_T(z)dz x, x' \right\rangle = \int_{\Gamma} \langle f(z)R_T(z)x, x' \rangle dz$$

for every $x \in E$, $x' \in E'$, and $f \in \mathcal{G}(T)$, $\Gamma \subset D(f)$.

The definition of Cauchy domain is to be found in Taylor’s paper [9], which also contains a proof of a theorem very similar to the above which may easily be adjusted to fit the present situation. The theorem might also be considered a special case of some of the work of H. Cartan [3]. With this background we are prepared to discuss conditions under which $Q^k = 0$ for some positive integer k .

2. The general case. Let E be a separated, locally convex space which is barrelled and quasi-complete. Let T be a spectral operator on E whose spectral family \mathcal{F} satisfies condition $P\mathcal{E}$) so that we may write $T = S + Q$. We now state our rate of growth condition:

DEFINITION. With E and T as described, we say that R_T satisfies an m th-order rate of growth condition (m being a positive integer) if the set $\{d(z, \sigma)^m R_{T_\sigma}(z)P_{\mathcal{F}}(\sigma) : z \notin \sigma, \sigma \text{ compact}\}$ is bounded in $(\mathcal{L}(E, E), \mathcal{F}_b)$. Here $d(z, \sigma)$ is the distance from z to σ .

For the rest of this section we assume that R_T satisfies an m th-order rate of growth of condition.

LEMMA 1. *Let n be a fixed positive integer, σ a compact subset of C , V a neighborhood of zero in $\mathcal{L}(E, E)$. Then there is a finite partition of σ by Borel sets $\{\sigma_j; j = 1, 2, \dots, k\}$ with the property that if $\{\lambda_j; j = 1, 2, \dots, k\}$ is any choice of k complex numbers with each $\lambda_j \in \sigma_j$, then*

$$(1) \quad Q^n P_{\mathcal{F}}(\sigma) - \sum_j (T - \lambda_j)^n P_{\mathcal{F}}(\sigma_j) \in V.$$

Proof. Denoting by f_i the function $f_i(z) = z^i \varphi_{\sigma}(z)$, for $i = 1, 2, \dots, n$, we observe that for every $\varepsilon > 0$ there is a partition $\{\sigma_j; j = 1, 2, \dots, k\}$ of σ such that $\|\sum_j \lambda_j^i \varphi_{\sigma_j} - f_i\|_u < \varepsilon$ for all $i = 1, 2, \dots, n$, where the λ_j are arbitrary in σ_j . Next, since the mapping $f \rightarrow U_f$ is continuous, it follows that $S^i P_{\mathcal{F}}(\sigma) = U_{f_i}$ may be approximated in the topology of $\mathcal{L}(E, E)$ by operators of the form $\sum_j \lambda_j^i P_{\mathcal{F}}(\sigma_j)$, uniformly for $i = 1, 2, \dots, n$, and for $\lambda_j \in \sigma_j$. The theorem may now be proved by considering

$$Q^n P_{\mathcal{F}}(\sigma) = (T - S)^n P_{\mathcal{F}}(\sigma) = \sum_{i=0}^n (-1)^i \binom{n}{i} T^{n-i} S^i P_{\mathcal{F}}(\sigma).$$

Given V , choose U , an equilibrated neighborhood of zero in $\mathcal{L}(E, E)$, such that $\sum_{i=0}^n \binom{n}{i} U \subset V$. For each $i = 0, 1, \dots, n$, choose a neighborhood W_i of zero in $\mathcal{L}(E, E)$ such that $A \in W_i$ implies $T^{n-i} A \in U$. Finally, choose $\{\sigma_j\}$ so that $\sum_j \lambda_j^i P_{\mathcal{F}}(\sigma_j) - S^i P_{\mathcal{F}}(\sigma) \in \cap_i W_i$. Then,

$$\sum_i (-1)^i \binom{n}{i} \sum_j T^{n-i} \lambda_j^i P_{\mathcal{F}}(\sigma_j) - \sum_i (-1)^i \binom{n}{i} T^{n-i} S^i P_{\mathcal{F}}(\sigma) \in V.$$

But the first term in this last expression is just $\sum_j (T - \lambda_j)^n P_{\mathcal{F}}(\sigma_j)$ so that (1) is proved.

LEMMA 2. *For every bounded subset B of E , every equicontinuous subset B' of E' , and every positive integer n , there is a positive real number $M = M(B, B', n)$ such that*

$$(2) \quad |\langle Q^n P_{\mathcal{F}}(\sigma)x, x' \rangle| \leq M\varepsilon^{n-m+1}$$

for all $x \in B$, $x' \in B'$, provided $0 < \varepsilon \leq 1$ and σ is a Borel set of diameter $\leq \varepsilon$.

Proof. If σ is empty there is no problem. Next consider the case where σ is a nonvoid compact subset of C , and fix $\eta > 0$. By Lemma 1 there is a partition $\{\sigma_j\}$ of σ such that

$$(3) \quad |\langle Q^n P_{\mathcal{F}}(\sigma)x, x' \rangle - \langle \sum_j (T - \lambda_j)^n P_{\mathcal{F}}(\sigma_j)x, x' \rangle| < \eta$$

for all $x \in B$, $x' \in B'$. Using any point in σ as center, construct a

circle, Γ , of radius 2ε . Then $|z - \lambda| \geq \varepsilon$ for all $z \in \Gamma$, $\lambda \in \sigma$. Then, for all $x \in B$, $x' \in B'$,

$$\begin{aligned} & |\langle \sum_j (T - \lambda_j)^n P_{\mathcal{F}}(\sigma_j)x, x' \rangle| \\ &= \left| \left\langle \sum_j \frac{1}{2\pi i} \int_{\Gamma} (z - \lambda_j)^n R_{T_\sigma}(z) dz P_{\mathcal{F}}(\sigma_j)x, x' \right\rangle \right| \\ &= \frac{1}{2\pi} \left| \int_{\Gamma} \langle \sum_j (z - \lambda_j)^n R_{T_\sigma}(z) P_{\mathcal{F}}(\sigma_j)x, x' \rangle dz \right| \\ &\leq \frac{1}{2\pi} L(\Gamma) \sup_{z \in \Gamma} |\langle \sum_j (z - \lambda_j)^n P_{\mathcal{F}}(\sigma_j) R_{T_\sigma}(z) P_{\mathcal{F}}(\sigma)x, x' \rangle| \\ &= 2\varepsilon \sup_{z \in \Gamma} \left| \left\langle \int_{\sigma} \sum_j (z - \lambda_j)^n \varphi_{\sigma_j}(\lambda) dP_{\mathcal{F}}(\lambda) R_{T_\sigma}(z) P_{\mathcal{F}}(\sigma)x, x' \right\rangle \right|. \end{aligned}$$

Let $g_z(\lambda) = \sum_j (z - \lambda_j)^n \varphi_{\sigma_j}(\lambda)$ be the integrand function in the spectral integral in this last expression. Then the above computation implies

$$(4) \quad |\langle \sum_j (T - \lambda_j)^n P_{\mathcal{F}}(\sigma_j)x, x' \rangle| \leq 2\varepsilon \sup_{z \in \Gamma} |\langle U_{g_z} R_{T_\sigma}(z) P_{\mathcal{F}}(\sigma)x, x' \rangle|.$$

Since $\sup_{z \in \Gamma} \|g_z\|_u \leq (3\varepsilon)^n$ and the set $\{U_f: \|f\|_u \leq 1\}$ is equicontinuous, corresponding to B' we may find a neighborhood W of zero in E such that

$$(5) \quad |\langle U_{g_z} y, x' \rangle| \leq (3\varepsilon)^n$$

for all $y \in W$, and $x' \in B'$. It is apparent that W is independent of ε , of σ , of the choice of $\{\sigma_j\}$, and of $z \in \Gamma$. But B and W determine a neighborhood of zero in $\mathcal{L}(E, E)$. Consequently, from the rate of growth condition, there is $\alpha > 0$ such that

$$(6) \quad d(z, \sigma)^m \alpha^{-1} R_{T_\sigma}(z) P_{\mathcal{F}}(\sigma)x \in W$$

for all $x \in B$, α being dependent only on B and B' (by way of W).

Substituting (6) in (5) gives, for all $z \in \Gamma$,

$$(7) \quad |\langle U_{g_z} R_{T_\sigma}(z) P_{\mathcal{F}}(\sigma)x, x' \rangle| \leq \alpha (3\varepsilon)^n d(z, \sigma)^{-m} \leq \alpha 3^m \varepsilon^{n-m}.$$

Letting $M = 2 \cdot 3^n \alpha$ and substituting (7) in (4) and (3) we have

$$\begin{aligned} |\langle Q^n P_{\mathcal{F}}(\sigma)x, x' \rangle| &\leq |\langle \sum_j (T - \lambda_j)^n P_{\mathcal{F}}(\sigma_j)x, x' \rangle| + \eta \\ &\leq M\varepsilon^{n-m+1} + \eta \end{aligned}$$

for all $x \in B$, $x' \in B'$, with M dependent only on B , B' , and n . Since η is arbitrary, the theorem is proved under the additional assumption that σ be compact. However, the general case may be readily deduced from this one.

THEOREM 2. *Let σ be a Borel set in C whose Hausdorff p -mea-*

sure is zero for some p . Then, for all $k \geq p + m - 1$, $Q^k P_{\mathcal{F}}(\sigma) = 0$.

Proof. The hypothesis asserts that for every $\varepsilon > 0$ there is a partition $\{\sigma_j\}$ of σ by finitely or countably many Borel sets of diameters ε_j respectively such that $\varepsilon_j \leq 1$ for all j and $\sum_j \varepsilon_j^p < \varepsilon$. Let $k = p + m - 1$. Then

$$|\langle Q^k P_{\mathcal{F}}(\sigma)x, x' \rangle| \leq \sum_j |\langle Q^k P_{\mathcal{F}}(\sigma_j)x, x' \rangle| \leq \sum_j M \varepsilon_j^{k+1-m} \leq M \varepsilon,$$

where M depends on x, x' , and k . Since ε, x , and x' are arbitrary, we are done.

THEOREM 3. $Q^{m+2} = 0$.

Proof. Let $p = 3$, $\sigma = \sup \mathcal{F}$, and $k = 3 + m - 1 = m + 2$ in the previous theorem.

3. Variations on the theme. In the case where E is a Banach space, McCarthy has pointed out a number of variations on Theorem 3. The simplest of these, equally valid in our locally convex setting, flow directly from Theorem 2 when the p -measure of $\text{sp } T$ or of $\sup \mathcal{F}$ is 0 for $p = 1$ or 2. An entirely different type of variation (also considered in [7]) may be discovered by observing that certain well-known Banach spaces may be embedded in E or E' . As in part 2 we assume that $T = S + Q$ is a spectral operator whose resolvent satisfies an m th-order rate of growth condition.

THEOREM 4. For every $x \in E$, and $x' \in E'$, the measure $\mu_{Q^{m+1}x, x'}$ has base Lebesgue planar measure, λ . In fact, for every bounded set B in E , and equicontinuous set B' in E' , there is $N = N(B, B')$ such that for all Borel sets σ , all $x \in B$, and all $x' \in B'$, we have

$$\mu_{Q^{m+1}x, x'}(\sigma) \leq N \lambda(\sigma).$$

Proof. Actually we prove this for μ_2 , Hausdorff 2-measure in the plane, but this is equivalent to proving it for λ . Let σ be a Borel set in the plane of finite μ_2 measure, and $\varepsilon > 0$. Partition σ by Borel sets $\{\sigma_i\}$ such that the diameter of σ_i is less than ε_i , with $0 < \varepsilon_i \leq 1$, and $\sum_i \varepsilon_i^2 < \mu_2(\sigma) + \varepsilon$. Letting $n = m + 1$ in Lemma 2, we find that for each B, B' as described in the Theorem, there is $M > 0$ such that $|\langle Q^{m+1} P_{\mathcal{F}}(\sigma_i)x, x' \rangle| \leq M \varepsilon_i^2$ for all $x \in B, x' \in B'$, and i . Consequently, $|\langle Q^{m+1} P_{\mathcal{F}}(\sigma)x, x' \rangle| \leq \sum_i M \varepsilon_i^2 \leq M(\mu_2(\sigma) + \varepsilon)$ for all $x \in B$, and $x' \in B'$. Since ε is arbitrary, we are done.

We now denote by $g_{x, x'}$ the Radon-Nikodym derivative of $\mu_{Q^{m+1}x, x'}$ with respect to λ . If f and g are two elements of $B^\infty(C)$ which agree

λ -almost everywhere, then for all $x \in E, x' \in E'$, we have

$$|\langle U_{f-g}Q^{m+1}x, x' \rangle| = \left| \int_{\sigma} (f - g)g_{x,x'}d\lambda \right| = 0.$$

Hence $U_fQ^{m+1} = U_gQ^{m+1}$. Moreover, if $\{f_n\}$ is a sequence of simple functions which converges to f in $L_1(C)$, then $U_{f_n}Q^{m+1}$ is a sequence which converges in $\mathcal{L}(E, E)$ to an operator which we denote by U_fQ^{m+1} . In this case $\langle U_fQ^{m+1}x, x' \rangle = \int_{\sigma} fd\mu_{Q^{m+1}x,x'}$.

We have already seen that $Q^{m+2} = 0$. If we now assume $Q^{m+1} \neq 0$, by letting $\sigma = \text{sup } \mathcal{F}$, we see that for some $x, x', \mu_{Q^{m+1}x,x'}(\sigma) \neq 0$. Consequently there is a set τ , compact if necessary, with nonvoid interior and a number $\alpha > 0$, such that $\lambda(\tau) \neq 0$ and $|g_{x,x'}(z)| > \alpha$ for all $z \in \tau$.

These two constructions are the basis for the embedding procedures mentioned above. The basic idea is to assume E has some property which is inherited by all of its closed linear subspaces. Then, if $Q^{m+1} \neq 0$, we may embed a suitable space (perhaps an L_p -space) in E or E' which does not have the property, thus obtaining a contradiction. In a written communication, McCarthy has suggested that $C(\tau)$ would be better than the $L_{\infty}(\tau)$ used in his paper for the case where E is assumed weakly complete. As an indication of some of the details involved in such a construction, we prove here the following:

THEOREM 5. *If E is semi-reflexive, then $Q^{m+1} = 0$.*

Proof. As indicated, we assume $Q^{m+1} \neq 0$ and construct τ and α corresponding to some x_0, x'_0 . Define $\Phi: L_1(\tau) \rightarrow E$ by the formula

$$\Phi(f) = U_{f\varphi_{\tau}}Q^{m+1}x_0.$$

Fix $\varepsilon > 0$ and let B' be an arbitrary equicontinuous subset of E' . Then for any $x' \in B'$,

$$\begin{aligned} |\langle \Phi(f), x' \rangle| &= |\langle U_{f\varphi_{\tau}}Q^{m+1}x_0, x' \rangle| \\ &= \left| \int_{\tau} fd\mu_{Q^{m+1}x_0,x'} \right| \leq N \|f\|_{L_1(\tau)}, \end{aligned}$$

where N is chosen according to Theorem 4 with $B = \{x_0\}$. Thus $|\langle \Phi(f), x' \rangle| < \varepsilon$ for all $x' \in B'$ whenever $\|f\|_{L_1} < \varepsilon/N$. Since B' is arbitrary, Φ is continuous. To see that Φ is one-to-one, first define f^* for each f in $L_1^0(\tau)$ by

$$f^*(z) = \begin{cases} \overline{f(z)}/|f(z)| & \text{when } f(z) \neq 0, \\ 0 & \text{when } f(z) = 0. \end{cases}$$

Then f^* is in $B^\infty(C)$ so that we may define $y' = {}^t\left(\int_\tau f^*/g_{x_0, x'_0} dP_{\mathcal{F}}\right)x'_0$. Then $\langle \Phi(f), y' \rangle = \int_\tau f f^* d\lambda = \int_\tau |f| d\lambda = \mathbf{||}f\mathbf{||}_{L_1(\tau)}$. Thus the kernel of Φ is zero. To show that Φ^{-1} is continuous on the range of Φ it is sufficient to find, for each $\varepsilon > 0$, an equicontinuous set B' in E' such that $\mathbf{||}f\mathbf{||}_{L_1} < \varepsilon$ whenever $|\langle \Phi(f), x' \rangle| \leq \varepsilon$ for all $x' \in B'$. The set

$$B = \left\{ \int_\tau f^*/g_{x_0, x'_0} dP_{\mathcal{F}} : f \in L_1(\tau) \right\}$$

is an equicontinuous subset of $\mathcal{L}(E, E)$. Consequently the set $B' = \{{}^tAx'_0 : A \in B\}$ is equicontinuous in E' . Finally,

$$\begin{aligned} \mathbf{||}f\mathbf{||}_{L_1(\tau)} &= \left| \int_\tau |f|/g_{x_0, x'_0} d\mu_{Q^{m+1}x_0, x'_0} \right| \\ &= \left| \left\langle U_{f, \varphi_\tau} Q^{m+1}x_0, {}^t\left(\int_\tau f^*/g_{x_0, x'_0} dP_{\mathcal{F}}\right)x'_0 \right\rangle \right| \\ &\leq \sup_{A \in B} |\langle \Phi(f), {}^tAx'_0 \rangle| = \sup_{x' \in B'} |\langle \Phi(f), x' \rangle|. \end{aligned}$$

Consequently, $\Phi(L_1(\tau))$ is a closed linear subspace of E which is isomorphic in both algebraic and topological sense to $L_1(\tau)$. This contradicts the assumption that E is semi-reflexive, hence the Theorem is proved.

BIBLIOGRAPHY

1. N. Bourbaki, *Espaces vectoriels topologiques, Elements de Mathematique*, Paris, (1953-55).
2. ———, *Intégration. Elements de Mathematique*, Paris (1952-60).
3. H. Cartan, *Theorie spectrale des C-algèbres commutatives*, Seminaire Bourbaki, Exposé **125** (1956).
4. N. Dunford, *Spectral operators*, Pacific J. Math., **4** (1954), 321-354.
5. C. Ionescu Tulcea, *Spectral operators on locally convex spaces*, Bull. Amer. Math. Soc., **67** (1961), 125-128.
6. F. Maeda, *A characterization of spectral operators on locally convex spaces*, Math. Annalen, **143** (1961), 59-74.
7. C. A. McCarthy, *The nilpotent part of a spectral operator*, Pacific J. Math., **9** (1959), 1223-1231.
8. H. Schaeffer, *Spectral measures in locally convex algebras*, Acta Math., **107** (1962), 125-173.
9. A. E. Taylor, *Spectral theory of closed distributive operators*, Acta Math., **84** (1950), 189-224.
10. L. Waelbroeck, *Le calcul symbolique dans les algèbres commutatives*, J. Math. Pures Appl., 9e Série, **33** (1954), 147-186.
11. ———, *Locally convex algebras: Spectral Theory*, Seminar on complex analysis, Institute for Advanced Study, 1958, unpub.