

Erratum: A Parabolic Quasilinear Problem for Linear Growth Functionals

(REV. MAT. IBEROAMERICANA **18** (2002), no. 1, 135–185)

Fuensanta Andreu, Vicent Caselles and José Manuel Mazón

Abstract

We give the correct proof of Lemma 3.6 of the paper *A Parabolic Quasilinear Problem for Linear Growth Functionals* (Rev. Mat. Iberoamericana **18** (2002), no. 1, 135-185).

Following [2], let

$$(1.1) \quad X(\Omega) = \{z \in L^\infty(\Omega, \mathbb{R}^N) : \operatorname{div}(z) \in L^1(\Omega)\}.$$

In [2], the weak trace on $\partial\Omega$ of the normal component of $z \in X(\Omega)$ is defined. More precisely, it is proved that there exists a linear operator $\gamma : X(\Omega) \rightarrow L^\infty(\partial\Omega)$ such that

$$\|\gamma(z)\|_\infty \leq \|z\|_\infty$$

$$\gamma(z)(x) = z(x) \cdot \nu(x) \quad \text{for all } x \in \partial\Omega \text{ if } z \in C^1(\bar{\Omega}, \mathbb{R}^N).$$

We shall denote $\gamma(z)(x)$ by $[z, \nu](x)$.

Assuming that $\partial\Omega$ is of class C^1 , it is proved in [3] that if $x_0 \in \partial\Omega$ is a Lebesgue point of the function $[z, \nu]$, then

$$(1.2) \quad [z, \nu](x_0) = \lim_{\rho \rightarrow 0} \lim_{r \rightarrow 0} \frac{1}{2rw_{N-1}\rho^{N-1}} \int_{C_{r,\rho}(x_0, \nu(x_0))} z(y) \cdot \nu(x_0) dy,$$

$C_{r,\rho}(x, \alpha)$ being the cylinder defined by

$$C_{r,\rho}(x, \alpha) = \{y \in \mathbb{R}^N : |(y-x) \cdot \alpha| < r, |(y-x) - [(y-x) \cdot \alpha] \cdot \alpha| < \rho\}.$$

2000 Mathematics Subject Classification: 35K65, 35K55, 47H06, 47H20.

Keywords: Linear growth functionals, nonlinear parabolic equations, accretive operators, nonlinear semigroups.

Our purpose is to give the correct proof of the following Lemma.

Lemma 1.1 ([1, Lemma 3.6])

i) Let $u_n \in BV(\Omega) \cap L^2(\Omega)$ and $z \in X(\Omega)$. Suppose that

$$(1.3) \quad \mathbf{a}(x, \nabla u_n) \rightharpoonup z \quad \text{weakly}^* \text{ in } L^\infty(\Omega, \mathbb{R}^N)$$

and

$$(1.4) \quad \operatorname{div}(\mathbf{a}(x, \nabla u_n)) \rightharpoonup \operatorname{div} z \quad \text{weakly in } L^2(\Omega).$$

Then

$$(1.5) \quad [\mathbf{a}(x, \nabla u_n), \nu(x)] \rightarrow [z, \nu(x)] \quad \text{weakly in } L^2(\partial\Omega) \text{ and}$$

$$(1.6) \quad |z(x) \cdot \nu(x)| \leq f^0(x, \nu(x)) \quad \text{a.e. in } \partial\Omega.$$

ii) Let $u_n \in W^{1,2}(\Omega)$. Let $\mathbf{a}_n(x, \xi) = \mathbf{a}(x, \xi) + \frac{1}{n}\xi$. Suppose that

$$(1.7) \quad \|u_n\|_2 \quad \text{is bounded in } L^2(\Omega),$$

$$(1.8) \quad \frac{1}{n}|\nabla u_n| \rightarrow 0 \quad \text{in } L^2(\Omega),$$

$$(1.9) \quad \mathbf{a}_n(x, \nabla u_n) \rightharpoonup z \quad \text{weakly in } L^2(\Omega, \mathbb{R}^N)$$

and

$$(1.10) \quad \operatorname{div}(\mathbf{a}_n(x, \nabla u_n)) \rightharpoonup \operatorname{div} z \quad \text{weakly in } L^2(\Omega).$$

Then

$$(1.11) \quad [\mathbf{a}_n(x, \nabla u_n), \nu(x)] \rightharpoonup [z, \nu(x)] \quad \text{weakly in } W^{1/2,2}(\partial\Omega)^* \text{ and}$$

$$(1.12) \quad |[z(x), \nu(x)]| \leq f^0(x, \nu(x)) \quad \text{a.e. in } \partial\Omega.$$

To prove it, let us recall the following result, which corresponds to Lemma 3.7 in [1]. We notice that its proof is independent of Lemma 3.6.

Lemma 1.2 ([1, Lemma 3.7]) *Suppose that any of the assumptions of Lemma 3.6 hold. Moreover we assume that*

$$(1.13) \quad u_n \rightarrow u \text{ in } L^2(\Omega) \text{ and } \|u_n\|_{BV} \text{ is bounded,}$$

Then

$$(1.14) \quad z(x) = \mathbf{a}(x, \nabla u(x)) \quad \text{a.e. } x \in \Omega.$$

Proof of Lemma 1.1. Since both proofs are based on similar arguments, we shall only prove *ii*). For the proof of (1.11) we refer to [1]. Let us prove (1.12). By (1.2), we have

$$(1.15) \quad [z, \nu](x) = \lim_{\rho \rightarrow 0} \lim_{r \rightarrow 0} \frac{1}{2rw_{N-1}\rho^{N-1}} \int_{C_{r,\rho}(x,\nu(x))} z(y) \cdot \nu(x) \, dy \quad \mathcal{H}^{N-1}\text{-a.e. on } \partial\Omega.$$

On the other hand, by assumption (H_5) ,

$$\mathbf{a}(x, \xi) \cdot \eta \leq f^0(x, \eta)$$

for all $\xi, \eta \in \mathbb{R}^N$, and all $x \in \overline{\Omega}$. Then, by 1.2, we have

$$(1.16) \quad z(y) \cdot \nu(x) = \mathbf{a}(y, \nabla u(y)) \cdot \nu(x) \leq f^0(y, \nu(x)).$$

Finally, since $f^0(\cdot, \xi)$ is continuous in $\overline{\Omega}$ for all $\xi \in \mathbb{R}^N$, using (1.15), (1.16), we get

$$\begin{aligned} |[z, \nu](x)| &\leq \lim_{\rho \rightarrow 0} \lim_{r \rightarrow 0} \frac{1}{2rw_{N-1}\rho^{N-1}} \int_{C_{r,\rho}(x,\nu(x))} |z(y) \cdot \nu(x)| \, dy \\ &\leq \lim_{\rho \rightarrow 0} \lim_{r \rightarrow 0} \frac{1}{2rw_{N-1}\rho^{N-1}} \int_{C_{r,\rho}(x,\nu(x))} f^0(y, \nu(x)) \, dy = f^0(x, \nu(x)), \end{aligned}$$

\mathcal{H}^{N-1} – a.e. on $\partial\Omega$.

■

References

- [1] ANDREU, F., CASELLES, V. AND MAZÓN, J. M.: A Parabolic Quasilinear Problem for Linear Growth Functionals. *Rev. Mat. Iberoamericana* **18** (2002), no. 1, 135–185.
- [2] ANZELLOTTI, G.: Pairings between measures and bounded functions and compensated compactness. *Ann. Mat. Pura Appl. (4)* **135** (1983), 293–318.

- [3] ANZELLOTTI, G.: Traces of bounded vector fields and the Divergence Theorem. Unpublished paper.

Recibido: 22 de enero de 2008

Fuensanta Andreu
Dept. de Matemática Aplicada
Universitat de Valencia
Dr. Moliner 50, 46100 Burjassot, Spain
Fuensanta.Andreu@uv.es

Vicent Caselles
Universitat Pompeu-Fabra
Dept. de Tecnologia
La Rambla 30-32, 08002 Barcelona, Spain
Vicent.Caselles@tecn.upf.es

José M. Mazón
Dept. de Análisis Matemático
Universitat de Valencia
Dr. Moliner 50, 46100 Burjassot, Spain
Mazon@uv.es