

ANALYSIS & PDE

Volume 5

No. 5

2012

HAMID HEZARI AND CHRISTOPHER D. SOGGE

A NATURAL LOWER BOUND FOR THE SIZE OF NODAL SETS

A NATURAL LOWER BOUND FOR THE SIZE OF NODAL SETS

HAMID HEZARI AND CHRISTOPHER D. SOGGE

We prove that, for an n -dimensional compact Riemannian manifold (M, g) , the $(n - 1)$ -dimensional Hausdorff measure $|Z_\lambda|$ of the zero-set Z_λ of an eigenfunction e_λ of the Laplacian having eigenvalue $-\lambda$, where $\lambda \geq 1$, and normalized by $\int_M |e_\lambda|^2 dV_g = 1$ satisfies

$$C|Z_\lambda| \geq \lambda^{\frac{1}{2}} \left(\int_M |e_\lambda| dV_g \right)^2$$

for some uniform constant C . As a consequence, we recover the lower bound $|Z_\lambda| \gtrsim \lambda^{(3-n)/4}$.

The purpose of this brief note is to prove a natural lower bound for the $(n - 1)$ -dimensional Hausdorff measure of nodal sets of eigenfunctions. To wit:

Theorem 1. *Let (M, g) be a compact manifold of dimension n and e_λ an eigenfunction satisfying*

$$-\Delta_g e_\lambda = \lambda e_\lambda, \text{ and } \int_M |e_\lambda|^2 dV_g = 1.$$

Then if $Z_\lambda = \{x \in M : e_\lambda(x) = 0\}$ is the nodal set and $|Z_\lambda|$ its $(n - 1)$ -dimensional Hausdorff measure, we have

$$\lambda^{\frac{1}{2}} \left(\int_M |e_\lambda| dV_g \right)^2 \leq C|Z_\lambda|, \quad \lambda \geq 1, \tag{1}$$

for some uniform constant C . Consequently,

$$\lambda^{\frac{3-n}{4}} \lesssim |Z_\lambda|, \quad \lambda \geq 1. \tag{2}$$

Inequality (2) follows from (1) and the lower bounds in [Sogge and Zelditch 2011a]

$$\lambda^{\frac{1-n}{8}} \lesssim \int_M |e_\lambda| dV_g. \tag{3}$$

The lower bound (2) is due to Colding and Minicozzi [2011]. Yau [1982] conjectured that $\lambda^{\frac{1}{2}} \approx |Z_\lambda|$. This lower bound $\lambda^{\frac{1}{2}} \lesssim |Z_\lambda|$ was verified in the two-dimensional case by Brüning [1978] and independently by Yau (unpublished). The bounds in (2) seem to be the best known ones for higher dimensions, although Donnelly and Fefferman [1988; 1990] showed that, as conjectured, $|Z_\lambda| \approx \lambda^{\frac{1}{2}}$, if (M, g) is assumed to be real analytic.

The authors were supported in part by NSF grants DMS-0969745 and DMS-1069175.
 MSC2010: 35P15.

Keywords: eigenfunctions, nodal lines.

The first “polynomial type” lower bounds appear to be those given in [Colding and Minicozzi 2011] and [Sogge and Zelditch 2011a] (see also [Mangoubi 2011]). As we shall point out, inequality (1) cannot be improved and it to some extent unifies the approaches in [Colding and Minicozzi 2011] and [Sogge and Zelditch 2011a]. As was shown in the latter paper, the L^1 -lower bounds in (3) follow from Hölder’s inequality and the L^p eigenfunction estimates of [Sogge 1988] for the range where $2 < p \leq 2(n + 1)/(n - 1)$. These too cannot be improved, but it is thought better L^p -bounds hold for a typical eigenfunction or if one makes geometric assumptions such as negative curvature (cf. [Sogge and Zelditch 2010; 2011b]). Thus, it is natural to expect to be able to improve (3) and hence the lower bounds (2) for all eigenfunctions on manifolds with negative curvature, or for “typical” eigenfunctions on any manifold. Of course, Yau’s conjecture that $|Z_\lambda| \approx \lambda^{\frac{1}{2}}$ would be the ultimate goal, but understanding when (3) can be improved is a related problem of independent interest.

Let us now turn to the proof of Theorem 1. We shall use an identity from [Sogge and Zelditch 2011a]:

$$\int_M |e_\lambda| (\Delta_g + \lambda) f \, dV_g = 2 \int_{Z_\lambda} |\nabla_g e_\lambda| f \, dS_g, \tag{4}$$

Here dS_g is the Riemannian surface measure on Z_λ , and ∇_g is the gradient coming from the metric and $|\nabla_g u|$ is the norm coming from the metric, meaning that in local coordinates

$$|\nabla_g u|_g^2 = \sum_{jk=1}^n g_{jk}(x) \partial_j u \partial_k u. \tag{5}$$

Identity (4) follows from the Gauss–Green formula and a related earlier identity was proved by Dong [1992].

As in [Hezari and Wang 2011], if we take $f \equiv 1$ and apply Schwarz’s inequality we get

$$\lambda \int_M |e_\lambda| \, dV_g \leq 2|Z_\lambda|^{1/2} \left(\int_{Z_\lambda} |\nabla_g e_\lambda|^2 \, dS_g \right)^{1/2}. \tag{6}$$

Thus we would have (1) if we could prove that the energy of e_λ on its nodal set satisfies the natural bounds

$$\int_{Z_\lambda} |\nabla_g e_\lambda|^2 \, dS_g \lesssim \lambda^{\frac{3}{2}}. \tag{7}$$

We shall do this by choosing a different auxiliary function f . This time we want to use

$$f = (1 + \lambda e_\lambda^2 + |\nabla_g e_\lambda|_g^2)^{\frac{1}{2}}. \tag{8}$$

If we plug this into (4) we get that

$$2 \int_{Z_\lambda} |\nabla_g e_\lambda|_g^2 \, dS_g \leq \int_M |e_\lambda| (\Delta_g + \lambda) (1 + \lambda e_\lambda^2 + |\nabla_g e_\lambda|_g^2)^{\frac{1}{2}} \, dV_g.$$

Since we have the L^2 -Sobolev bounds

$$\|e_\lambda\|_{H^s(M)} = O(\lambda^{\frac{s}{2}}), \tag{9}$$

it is clear that

$$\lambda \int_M |e_\lambda| (1 + \lambda e_\lambda^2 + |\nabla_g e_\lambda|_g^2)^{\frac{1}{2}} dV_g = O(\lambda^{\frac{3}{2}}),$$

and thus to prove (7), it suffices to show that

$$\int_M |e_\lambda| \Delta_g (1 + \lambda e_\lambda^2 + |\nabla_g e_\lambda|_g^2)^{\frac{1}{2}} dV_g = O(\lambda^{\frac{3}{2}}). \tag{10}$$

To prove this we first note that

$$\partial_k (1 + \lambda e_\lambda^2 + |\nabla_g e_\lambda|_g^2)^{\frac{1}{2}} = \frac{\lambda e_\lambda \partial_k e_\lambda + \frac{1}{2} \partial_k |\nabla_g e_\lambda|_g^2}{(1 + \lambda e_\lambda^2 + |\nabla_g e_\lambda|_g^2)^{\frac{1}{2}}},$$

from this and (9) we deduce that

$$\int_M |e_\lambda| |\nabla_g (1 + \lambda e_\lambda^2 + |\nabla_g e_\lambda|_g^2)^{\frac{1}{2}}| dV_g = O(\lambda).$$

This means that the contribution of the first order terms of the Laplace–Beltrami operator (written in local coordinates) to (10) are better than required, and so it suffices to show that in a compact subset K of a local coordinate patch we have

$$\int_K |e_\lambda| \left| \partial_j \partial_k (1 + \lambda e_\lambda^2 + |\nabla_g e_\lambda|_g^2)^{\frac{1}{2}} \right| dV_g = O(\lambda^{\frac{3}{2}}). \tag{11}$$

A calculation shows that $\partial_j \partial_k (\lambda e_\lambda^2 + |\nabla_g e_\lambda|_g^2)^{\frac{1}{2}}$ equals

$$-\frac{(\lambda e_\lambda \partial_j e_\lambda + \frac{1}{2} \partial_j |\nabla_g e_\lambda|_g^2)(\lambda e_\lambda \partial_k e_\lambda + \frac{1}{2} \partial_k |\nabla_g e_\lambda|_g^2)}{(1 + \lambda e_\lambda^2 + |\nabla_g e_\lambda|_g^2)^{\frac{3}{2}}} + \frac{\lambda \partial_j e_\lambda \partial_k e_\lambda + \lambda e_\lambda \partial_j \partial_k e_\lambda + \frac{1}{2} \partial_j \partial_k |\nabla_g e_\lambda|_g^2}{(1 + \lambda e_\lambda^2 + |\nabla_g e_\lambda|_g^2)^{\frac{1}{2}}}.$$

If $|D^m f| = \sum_{|\alpha|=m} |\partial^\alpha f|$, then by (5)

$$\partial_k |\nabla_g e_\lambda|_g^2 = O(|D^2 e_\lambda| |De_\lambda| + |De_\lambda|^2),$$

and

$$\partial_j \partial_k |\nabla_g e_\lambda|_g^2 = O(|D^3 e_\lambda| |De_\lambda| + |D^2 e_\lambda|^2 + |D^2 e_\lambda| |De_\lambda| + |De_\lambda|^2).$$

Therefore,

$$\begin{aligned} & \partial_j \partial_k (\lambda e_\lambda^2 + |\nabla_g e_\lambda|_g^2)^{\frac{1}{2}} \\ &= O\left(\frac{\lambda^2 |e_\lambda|^2 |De_\lambda|^2 + |D^2 e_\lambda|^2 |De_\lambda|^2 + |De_\lambda|^4}{(1 + \lambda e_\lambda^2 + |\nabla_g e_\lambda|_g^2)^{\frac{3}{2}}} \right) \\ & \quad + O\left(\frac{\lambda |De_\lambda|^2 + \lambda |e_\lambda| |D^2 e_\lambda| + |D^3 e_\lambda| |De_\lambda| + |D^2 e_\lambda|^2 + |D^2 e_\lambda| |De_\lambda| + |De_\lambda|^2}{(1 + \lambda e_\lambda^2 + |\nabla_g e_\lambda|_g^2)^{\frac{1}{2}}} \right). \end{aligned}$$

This implies that the integrand in the left side of (11) is dominated by

$$\begin{aligned} & (\lambda^{\frac{1}{2}} |De_\lambda|^2 + \lambda^{-\frac{1}{2}} |D^2 e_\lambda|^2 + |De_\lambda|^2) \\ & + (\lambda^{\frac{1}{2}} |De_\lambda|^2 + \lambda^{\frac{1}{2}} |e_\lambda| |D^2 e_\lambda| + |e_\lambda| |D^3 e_\lambda| + \lambda^{-\frac{1}{2}} |D^2 e_\lambda|^2 + |D^2 e_\lambda| |e_\lambda| + |De_\lambda| |e_\lambda|), \end{aligned}$$

leading to (11) after applying (9). \square

Remarks.

- We could also have taken f to be $(\lambda + \lambda e_\lambda^2 + |\nabla_g e_\lambda|^2)^{\frac{1}{2}}$ and obtained the same upper bounds, but there does not seem to be any advantage to doing this.
- Inequality (1) cannot be improved. There are many cases when the L^1 and L^2 -norms of eigenfunctions are comparable. For instance, for the sphere the zonal functions have this property and it is easy to check that their nodal sets satisfy $|Z_\lambda| \approx \lambda^{\frac{1}{2}}$, which means that for zonal functions (1) cannot be improved.
- There are many cases where inequality (1) can be improved. For instance, the L^2 -normalized highest weight spherical harmonics Q_k have eigenvalues $\lambda = \lambda_k \approx k^2$, and L^1 -norms $\approx k^{-\frac{n-1}{4}}$ (see e.g., [Sogge 1986]). This means that for the highest weight spherical harmonics the left side is proportional to $\lambda^{\frac{3-n}{4}}$ even though here too $|Z_\lambda| \approx \lambda^{\frac{1}{2}}$. Similarly, the highest weight spherical harmonics saturate (7). It is because of functions like the highest weight spherical harmonics that the current techniques only seem to yield (2). Note that inequality (2) gives the correct lower bound in the trivial case where the dimension n is one. As the dimension increases, the bound gets worse and worse due to the fact that (3) is saturated by functions like the highest weight spherical harmonics (“Gaussian beams”) whose mass is supported on a $\lambda^{-\frac{1}{4}}$ neighborhood of a geodesic and the volume of such a tube decreases geometrically as n increases. (See [Bourgain 2009; Sogge 2011] for related work on this phenomena.)
- W. Minicozzi pointed out to us that (7) also follows from the identity

$$2 \int_{Z_\lambda} |\nabla_g e_\lambda|^2 dS_g = - \int_M \operatorname{sgn}(e_\lambda) \operatorname{div}_g (|\nabla_g e_\lambda| \nabla_g e_\lambda) dV_g. \quad (12)$$

and (9). Like the proof of (4) in [Sogge and Zelditch 2011a], the identity (12) follows from an application of the divergence theorem applied to each of the nodal domains of e_λ .

Acknowledgments

The authors wish to thank W. Minicozzi and S. Zelditch for several helpful and interesting discussions.

References

- [Bourgain 2009] J. Bourgain, “Geodesic restrictions and L^p -estimates for eigenfunctions of Riemannian surfaces”, pp. 27–35 in *Linear and complex analysis*, edited by A. Alexandrov et al., Amer. Math. Soc. Transl. Ser. 2 **226**, American Mathematical Society, Providence, RI, 2009. [MR 2011b:58066](#) [Zbl 1189.58015](#)

- [Brüning 1978] J. Brüning, “Über Knoten von Eigenfunktionen des Laplace–Beltrami-Operators”, *Math. Z.* **158**:1 (1978), 15–21. [MR 57 #17732](#) [Zbl 0349.58012](#)
- [Colding and Minicozzi 2011] T. H. Colding and W. P. Minicozzi, II, “Lower bounds for nodal sets of eigenfunctions”, *Comm. Math. Phys.* **306**:3 (2011), 777–784. [MR 2825508](#) [Zbl 1238.58020](#)
- [Dong 1992] R.-T. Dong, “Nodal sets of eigenfunctions on Riemann surfaces”, *J. Differential Geom.* **36**:2 (1992), 493–506. [MR 93h:58159](#) [Zbl 0776.53024](#)
- [Donnelly and Fefferman 1988] H. Donnelly and C. Fefferman, “Nodal sets of eigenfunctions on Riemannian manifolds”, *Invent. Math.* **93**:1 (1988), 161–183. [MR 89m:58207](#) [Zbl 0659.58047](#)
- [Donnelly and Fefferman 1990] H. Donnelly and C. Fefferman, “Nodal sets for eigenfunctions of the Laplacian on surfaces”, *J. Amer. Math. Soc.* **3**:2 (1990), 333–353. [MR 92d:58209](#) [Zbl 0702.58077](#)
- [Hezari and Wang 2011] H. Hezari and Z. Wang, “Lower bounds for volumes of nodal sets: an improvement of a result of Sogge–Zelditch”, preprint, 2011. [arXiv 1107.0092](#)
- [Mangoubi 2011] D. Mangoubi, “A remark on recent lower bounds for nodal sets”, *Comm. Partial Differential Equations* **36**:12 (2011), 2208–2212. [MR 2852075](#) [Zbl 1232.58025](#) [arXiv 1010.4579](#)
- [Sogge 1986] C. D. Sogge, “Oscillatory integrals and spherical harmonics”, *Duke Math. J.* **53**:1 (1986), 43–65. [MR 87g:42026](#) [Zbl 0636.42018](#)
- [Sogge 1988] C. D. Sogge, “Concerning the L^p norm of spectral clusters for second-order elliptic operators on compact manifolds”, *J. Funct. Anal.* **77**:1 (1988), 123–138. [MR 89d:35131](#) [Zbl 0641.46011](#)
- [Sogge 2011] C. D. Sogge, “Kakeya–Nikodym averages and L^p -norms of eigenfunctions”, *Tohoku Math. J. (2)* **63**:4 (2011), 519–538. [MR 2872954](#) [Zbl 1234.35156](#) [arXiv 0907.4827](#)
- [Sogge and Zelditch 2010] C. D. Sogge and S. Zelditch, “Concerning the L^4 norms of typical eigenfunctions on compact surfaces”, preprint, 2010. [arXiv 1011.0215](#)
- [Sogge and Zelditch 2011a] C. D. Sogge and S. Zelditch, “Lower bounds on the Hausdorff measure of nodal sets”, *Math. Res. Lett.* **18**:1 (2011), 25–37. [MR 2012c:58055](#) [Zbl 06026600](#)
- [Sogge and Zelditch 2011b] C. D. Sogge and S. Zelditch, “On eigenfunction restriction estimates and L^4 -bounds for compact surfaces with nonpositive curvature”, preprint, 2011. [arXiv 1108.2726](#)
- [Yau 1982] S. T. Yau, “Survey on partial differential equations in differential geometry”, pp. 3–71 in *Seminar on Differential Geometry*, edited by S. T. Yau, Ann. of Math. Stud. **102**, Princeton University Press, Princeton, NJ, 1982. [MR 83i:53003](#) [Zbl 0478.53001](#)

Received 12 Aug 2011. Accepted 24 Oct 2011.

HAMID HEZARI: hezari@math.uci.edu

Department of Mathematics, University of California, Irvine, CA 92697, United States

CHRISTOPHER D. SOGGE: sogge@jhu.edu

Department of Mathematics, Johns Hopkins University, Baltimore, MD 21093, United States

Analysis & PDE

msp.berkeley.edu/apde

EDITORS

EDITOR-IN-CHIEF

Maciej Zworski
University of California
Berkeley, USA

BOARD OF EDITORS

Michael Aizenman	Princeton University, USA aizenman@math.princeton.edu	Nicolas Burq	Université Paris-Sud 11, France nicolas.burq@math.u-psud.fr
Luis A. Caffarelli	University of Texas, USA caffarel@math.utexas.edu	Sun-Yung Alice Chang	Princeton University, USA chang@math.princeton.edu
Michael Christ	University of California, Berkeley, USA mchrist@math.berkeley.edu	Charles Fefferman	Princeton University, USA cf@math.princeton.edu
Ursula Hamenstaedt	Universität Bonn, Germany ursula@math.uni-bonn.de	Nigel Higson	Pennsylvania State University, USA higson@math.psu.edu
Vaughan Jones	University of California, Berkeley, USA vfr@math.berkeley.edu	Herbert Koch	Universität Bonn, Germany koch@math.uni-bonn.de
Izabella Laba	University of British Columbia, Canada ilaba@math.ubc.ca	Gilles Lebeau	Université de Nice Sophia Antipolis, France lebeau@unice.fr
László Lempert	Purdue University, USA lempert@math.purdue.edu	Richard B. Melrose	Massachusetts Institute of Technology, USA rbm@math.mit.edu
Frank Merle	Université de Cergy-Pontoise, France Frank.Merle@u-cergy.fr	William Minicozzi II	Johns Hopkins University, USA minicozz@math.jhu.edu
Werner Müller	Universität Bonn, Germany mueller@math.uni-bonn.de	Yuval Peres	University of California, Berkeley, USA peres@stat.berkeley.edu
Gilles Pisier	Texas A&M University, and Paris 6 pisier@math.tamu.edu	Tristan Rivière	ETH, Switzerland riviere@math.ethz.ch
Igor Rodnianski	Princeton University, USA irod@math.princeton.edu	Wilhelm Schlag	University of Chicago, USA schlag@math.uchicago.edu
Sylvia Serfaty	New York University, USA serfaty@cims.nyu.edu	Yum-Tong Siu	Harvard University, USA siu@math.harvard.edu
Terence Tao	University of California, Los Angeles, USA tao@math.ucla.edu	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu
Gunther Uhlmann	University of Washington, USA gunther@math.washington.edu	András Vasy	Stanford University, USA andras@math.stanford.edu
Dan Virgil Voiculescu	University of California, Berkeley, USA dvv@math.berkeley.edu	Steven Zelditch	Northwestern University, USA zelditch@math.northwestern.edu

PRODUCTION

production@msp.org

Silvio Levy, Scientific Editor

Sheila Newbery, Senior Production Editor


See inside back cover or msp.berkeley.edu/apde for submission instructions.

The subscription price for 2012 is US \$140/year for the electronic version, and \$240/year for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to Mathematical Sciences Publishers, Department of Mathematics, University of California, Berkeley, CA 94720-3840, USA.

Analysis & PDE, at Mathematical Sciences Publishers, Department of Mathematics, University of California, Berkeley, CA 94720-3840 is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

APDE peer review and production are managed by EditFLOW™ from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
<http://msp.org/>

A NON-PROFIT CORPORATION

Typeset in L^AT_EX

Copyright ©2012 by Mathematical Sciences Publishers

ANALYSIS & PDE

Volume 5 No. 5 2012

An inverse problem for the wave equation with one measurement and the pseudorandom source	887
TAPIO HELIN, MATTI LASSAS and LAURI OKSANEN	
Two-dimensional nonlinear Schrödinger equation with random radial data	913
YU DENG	
Schrödinger operators and the distribution of resonances in sectors	961
TANYA J. CHRISTIANSEN	
Weighted maximal regularity estimates and solvability of nonsmooth elliptic systems, II	983
PASCAL AUSCHER and ANDREAS ROSÉN	
The two-phase Stefan problem: regularization near Lipschitz initial data by phase dynamics	1063
SUNHI CHOI and INWON KIM	
C^∞ spectral rigidity of the ellipse	1105
HAMID HEZARI and STEVE ZELDITCH	
A natural lower bound for the size of nodal sets	1133
HAMID HEZARI and CHRISTOPHER D. SOGGE	
Effective integrable dynamics for a certain nonlinear wave equation	1139
PATRICK GÉRARD and SANDRINE GRELLIER	
Nonlinear Schrödinger equation and frequency saturation	1157
RÉMI CARLES	