

ANALYSIS & PDE

Volume 7

No. 5

2014

GUSTAV HOLZEGEL AND JACQUES SMULEVICI

**QUASIMODES AND A LOWER BOUND ON THE UNIFORM
ENERGY DECAY RATE FOR KERR-ADS SPACETIMES**

QUASIMODES AND A LOWER BOUND ON THE UNIFORM ENERGY DECAY RATE FOR KERR–ADS SPACETIMES

GUSTAV HOLZEGEL AND JACQUES SMULEVICI

We construct quasimodes for the Klein–Gordon equation on the black hole exterior of Kerr–AdS (anti-de Sitter) spacetimes. Such quasimodes are associated with time-periodic approximate solutions of the Klein–Gordon equation and provide natural candidates to probe the decay of solutions on these backgrounds. They are constructed as the solutions of a semiclassical nonlinear eigenvalue problem arising after separation of variables, with the (inverse of the) angular momentum playing the role of the semiclassical parameter. Our construction results in *exponentially small* errors in the semiclassical parameter. This implies that general solutions to the Klein Gordon equation on Kerr–AdS cannot decay faster than logarithmically. The latter result completes previous work by the authors, where a logarithmic decay rate was established as an upper bound.

1. Introduction	1057
2. Preliminaries	1063
3. Separation of variables and reduced equations	1066
4. Bound states	1069
5. Agmon estimates	1078
6. The construction of quasimodes	1083
7. Proof of Corollary 1.6	1084
Appendix: The improved boundedness statement	1086
Acknowledgments	1088
References	1088

1. Introduction

There is currently a lot of mathematical activity concerning the analysis of waves on the exterior of black hole backgrounds.¹ The main motivation is the black hole stability problem, i.e., the conjectured nonlinear asymptotic stability of the two-parameter family of asymptotically flat Kerr spacetimes $(\mathcal{M}, g_{M,a})$, the latter being stationary solutions of the vacuum Einstein equations $\text{Ric}[g] = 0$. From the point of view of nonlinear partial differential equations, the analysis of linear scalar waves on black holes is a prerequisite to successfully understanding the nonlinear hyperbolic Einstein equations in a neighborhood of the Kerr family.

MSC2010: primary 58J50; secondary 83C57.

Keywords: wave equation, black holes, decay estimates, Kerr – anti-de Sitter.

¹See [Dafermos and Rodnianski 2013] for an introduction and a review of recent results.

A major insight that crystallized in the last decade [Dafermos and Rodnianski 2005; 2009; 2010a; 2010b; Marzuola et al. 2010; Tataru and Tohaneanu 2011; Andersson and Blue 2009; Dyatlov 2011; 2012] is that the fundamental geometric obstacles to the decay of waves, namely superradiance and trapped null-geodesics, can be overcome by exploiting the normal hyperbolicity of the trapping, the redshift effect near the event horizon and the natural dispersion of waves in asymptotically flat (and asymptotically de Sitter) spacetimes. In particular, polynomial decay rates have been established for solutions to the scalar wave equation on the exterior of any member of the subextremal Kerr family of spacetimes [Dafermos and Rodnianski 2010b].

Changing the black hole geometry can have dramatic effects on the behavior of linear waves through the subtle interplay of the redshift, the superradiance and the trapping. Aretakis [2012; 2013] showed that for *extremal* black holes (whose vanishing surface gravity leads to a degeneration of the redshift effect) the transversal derivatives of general solutions to the wave equation will grow along the event horizon. In [Shlapentokh-Rothman 2014], it is proven that for the *massive* wave equation on a subextremal ($|a| < M$) Kerr spacetime, exponentially growing solutions can be constructed on the exterior, exploiting an amplification of the superradiance caused by the confining properties of the mass term.

In this paper, we shall be interested in a black hole geometry for which a strong trapping phenomenon leads to a very slow (only logarithmic) decay of waves. More precisely, we will study the behavior of solutions to the massive wave equation

$$\left(\square_g + \frac{\alpha}{l^2}\right)\psi = 0 \tag{1}$$

in the exterior of asymptotically anti-de Sitter (AdS) black holes with spacetime metric g .

Due to their AdS asymptotics, these spacetimes are not globally hyperbolic. Nonetheless, Equation (1) is well-posed in suitably weighted Sobolev spaces, denoted here by H_{AdS}^k , provided α satisfies the Breitenlohner–Friedmann bound $\alpha < \frac{9}{4}$. See [Holzegel 2012] as well as [Vasy 2012; Bachelot 2008; 2011; Ishibashi and Wald 2004; Warnick 2013] for a complete treatment of general boundary conditions.

The *global* properties of solutions to (1) on the exterior of nonsuperradiant² Kerr–AdS black holes were studied in [Holzegel 2010; Holzegel and Smulevici 2013a; Holzegel and Warnick 2014]. In particular, boundedness was obtained in [Holzegel 2010; Holzegel and Warnick 2014] and logarithmic decay in time for general H_{AdS}^2 solutions in [Holzegel and Smulevici 2013a]. We summarize these results in the following theorem. We refer to Section 2A1 for the precise definitions of the Kerr–AdS spacetimes, the area-radius r_+ of the event horizon and the Σ_{t^*} -foliation and to Section 2B for the definitions of the norms and energies used in the statement below. At this point we only remark that $e_1[\psi]$ is an energy density involving all first derivatives of ψ while $e_2[\psi]$ and $\tilde{e}_2[\psi]$ involve all second derivatives (with appropriate weights):

Theorem 1.1. *Let (g, \mathcal{R}) denote the black hole exterior of a Kerr–AdS spacetime with mass $M > 0$, angular momentum per unit mass a and cosmological constant $\Lambda = -3/l^2$. Assume that the parameters satisfy $\alpha < \frac{9}{4}$, $|a| < l$. Fix a spacelike slice $\Sigma_{t_0}^*$ intersecting \mathcal{H}^+ . Then:*

²This means that the parameters of the black hole satisfy $r_+^2 > |a|l$. See Remark 1.3 and Section 2.

1. *Equation (1) is well-posed in CH^k_{AdS} on (g, \mathbb{R}) for any $k \geq 2$ for initial data prescribed on $\Sigma_{t_0^*}$. (See [Holzegel 2012].)*
2. *The solutions of (1) arising from data prescribed on $\Sigma_{t_0^*}$ remain uniformly bounded on the black hole exterior provided $r_+^2 > |a|l$ holds. In particular,*

$$\int_{\Sigma_{t^*}} e_1[\psi](t^*) r^2 \sin \theta \, dr \, d\theta \, d\phi \lesssim \int_{\Sigma_{t_0^*}} e_1[\psi](t_0^*) r^2 \sin \theta \, dr \, d\theta \, d\phi. \tag{2}$$

Analogous statements hold for all higher H^k_{AdS} -norms. In particular,

$$\int_{\Sigma_{t^*}} e_2[\psi](t^*) r^2 \sin \theta \, dr \, d\theta \, d\phi \lesssim \int_{\Sigma_{t_0^*}} e_2[\psi](t_0^*) r^2 \sin \theta \, dr \, d\theta \, d\phi, \tag{3}$$

and the same statement for $\tilde{e}_2[\psi](t^*)$. (See [Holzegel 2010; Holzegel and Warnick 2014].³)

3. *If the parameters satisfy $r_+^2 > |a|l$, the solutions of (1) satisfy the global decay estimate*

$$\int_{\Sigma_{t^*}} e_1[\psi](t^*) r^2 \sin \theta \, dr \, d\theta \, d\phi \lesssim \frac{1}{[\log(2 + t^*)]^2} \int_{\Sigma_{t_0^*}} e_2[\psi](t_0^*) r^2 \sin \theta \, dr \, d\theta \, d\phi \tag{4}$$

for all $t^* \geq t_0^* > 0$. (See [Holzegel and Smulevici 2013a].)

Remark 1.2. The constant implicit in the symbol “ \lesssim ” appearing in items 2 and 3 depends only on the fixed parameters M, ℓ, a and α .

Remark 1.3. The condition $r_+^2 > |a|l$ on the parameters in items 2 and 3 guarantees the existence of a globally causal Killing vector field on the black hole exterior, the Hawking–Reall vector field [Hawking and Reall 2000], which explains why such black holes are sometimes referred to as “nonsuperradiant”. If one restricts to *axisymmetric* solutions of (1), this condition can be dropped for statements 2 and 3 in Theorem 1.1.

Remark 1.4. The boundedness statement (2) does not lose derivatives while the decay statement (4) does. This is the familiar loss of derivatives caused by the existence of trapped null-geodesics [Ralston 1969; Sbierski 2013].

This logarithmic decay rate (4) was conjectured to be sharp in [Holzegel and Smulevici 2013a] in view of the discovery of a new stable trapping phenomenon, itself a consequence of the coupling between the lack of dispersion at the asymptotic end and the usual (unstable) trapping on black hole exteriors [ibid.].

1A. The main results. In this paper, we shall prove that the logarithmic decay estimate of Theorem 1.1 is indeed sharp. Recall that for the obstacle problem, it is classical [Ralston 1969] that lower bounds on the rate of energy decay can be obtained from the construction of approximate eigenfunctions, also called *quasimodes*, of the associated elliptic operator, obtained by formally taking the Fourier transform in time

³These papers and [Holzegel 2012] are only concerned with the $\tilde{e}_2[\psi]$ energy. It is remarked that by commutation with angular momentum operators one can prove boundedness for the $e_2[\psi]$ energy (which differs from the $\tilde{e}_2[\psi]$ energy through the weights of the angular derivatives). For completeness, we provide in an Appendix an explicit proof of this statement.

of the wave operator. Our main theorem establishes the existence of such quasimodes for Kerr–AdS with exponentially small errors.⁴

The statement of the following theorem will involve the quantity

$$r_{\max} \in \left(r_+, \frac{3M}{1 - a^2/l^2} \right],$$

which is determined in Lemma 3.1 as the location of the unique maximum of a simple radial function. For Schwarzschild–AdS, $r_{\max} = 3M$ will be the location of the well-known photon sphere.

Theorem 1.5 (quasimodes for Kerr–AdS). *Let (g, \mathcal{R}) denote the black hole exterior of a Kerr–AdS spacetime, with mass $M > 0$, angular momentum per unit mass a and cosmological constant $\Lambda = -3/l^2$. Assume that the parameters satisfy $\alpha < \frac{9}{4}$, $|a| < l$. Let (t, r, θ, φ) denote standard Boyer–Lindquist coordinates on \mathcal{R} . Then, for $\delta > 0$ sufficiently small (depending only on the parameters l, M, a, α), there exists a family of nonzero functions $\psi_\ell \in H^k_{\text{AdS}}$ for any $k \geq 0$ satisfying the following conditions:*

- (1) $\psi_\ell(t, r, \theta, \varphi) = e^{i\omega_\ell t} \varphi_\ell(r, \theta)$ (axisymmetric and time-periodic).
- (2) $0 < c < \frac{\omega_\ell^2}{\ell(\ell+1)} < C$, for constants c and C independent of ℓ (uniform bounds on the frequencies).
- (3) For all $t^* \geq t_0^*$ and all $k \geq 0$, we have $\|(\square_g + \alpha/l^2)\psi_\ell\|_{H^k_{\text{AdS}}(\Sigma_{t^*})} \leq C_k e^{-C_k \ell} \|\psi_\ell\|_{H^0_{\text{AdS}}(\Sigma_{t_0^*})}$, for some $C_k > 0$ independent of ℓ (approximate solutions to the wave equation).
- (4) The support of $F_\ell := (\square_g + \alpha/l^2)\psi_\ell$ is contained in $\{r_{\max} \leq r \leq r_{\max} + \delta\}$ (spatial localization of the error).
- (5) The support of $\varphi_\ell(r, \theta)$ is contained in $\{r \geq r_{\max}\}$ (spatial localization of the solution).

Note that the ψ_ℓ have constant H^k_{AdS} -norms and hence exhibit no decay. On the other hand, a standard application of Duhamel’s formula shows that the ψ_ℓ are good approximations to the solution of $(\square_g + \alpha/l^2)\psi = 0$ arising from the data induced by ψ_ℓ , at least up to a time $t \sim e^{C_k \ell}$.

Corollary 1.6. *Let (\mathcal{R}, g) denote the black hole exterior of a Kerr–AdS spacetime as in Theorem 1.5. Denote by SCH^2_{AdS} the set of CH^2_{AdS} solutions to (1) with $\alpha < \frac{9}{4}$. Let $t_0^* \geq 0$ be fixed and define for any nonzero ψ and $t^* \geq 0$*

$$Q[\psi](t^*) := \log(2 + t^*) \left(\frac{\int_{\Sigma_{t^*} \cap \{r \geq r_{\max}\}} e_1[\psi](t^*) r^2 \sin \theta \, dr \, d\theta \, d\varphi}{\int_{\Sigma_{t_0^*}} e_2[\psi](t_0^*) r^2 \sin \theta \, dr \, d\theta \, d\varphi} \right)^{\frac{1}{2}}.$$

Then there exists a universal (depending only on $M, \alpha, |a|$ and l) constant $C > 0$ such that

$$\limsup_{t^* \rightarrow +\infty} \sup_{\substack{\psi \in SCH^2_{\text{AdS}} \\ \psi \neq 0}} Q[\psi](t^*) > C > 0.$$

Remark 1.7. Corollary 1.6 implies that the semilocal energy in $\{r \geq r_{\max}\}$ cannot decay universally faster than $(\log t^*)^{-2}$, unless one loses more derivatives.

⁴Note that in order to deduce the sharpness of the logarithmic decay rate from the quasimodes, polynomial errors would a priori not be sufficient.

Remark 1.8. We emphasize that no smallness assumption on the angular momentum a is needed, apart from the condition $|a| < l$ which ensures that the metric is a regular black hole metric.

Remark 1.9. The ψ_ℓ constructed in [Theorem 1.5](#) are axisymmetric, while the decay estimate of [Theorem 1.1](#) holds for general solutions (provided $r_+^2 > |a|l$ holds in the nonaxisymmetric case). Since we are concerned here with a *lower* bound on the uniform decay rate, an analysis within axisymmetry is sufficient. This allows us to drop the nonsuperradiant condition $r_+^2 > |a|l$ in the analysis. On the other hand, for (sufficiently) superradiant black holes $r_+^2 < |a|l$ one can adapt the proof of [\[Shlapentokh-Rothman 2014\]](#) to construct exponentially growing solutions. Hence in this case, the quasimodes we construct are not the “worst” solutions on these backgrounds.

Remark 1.10. Stable trapping occurs only in the region $r \geq r_{\max}$ and is associated to certain frequencies. As a consequence, stronger local energy decay in $r \leq r_{\max}$ or for some frequency projections of solutions is a priori compatible with the results of this paper.

1B. *Related works and discussion.*

1B1. *Nonlinear analysis on asymptotically AdS spacetimes.* In [\[Holzegel and Smulevici 2012a; 2013b\]](#), the nonlinear spherically symmetric Einstein–Klein–Gordon system for asymptotically AdS initial data was studied, and, in particular, asymptotic stability of Schwarzschild–AdS was proven within this model.

For a discussion connecting the logarithmic decay to the nonlinear stability or instability of asymptotically AdS black holes, we refer to Section 1.4 of [\[Holzegel and Smulevici 2013a\]](#). We also mention the recent heuristic analysis of [\[Dias et al. 2012\]](#) drawing attention to a potential stability mechanism caused by the lack of exact nonlinear resonances in this setting. For AdS itself, instability was conjectured in [\[Dafermos and Holzegel 2006; Anderson 2006; Dafermos 2006\]](#). More recently, both numerical and additional heuristic evidence has been presented [\[Bizon and Rostworowski 2011\]](#). Finally, let us note that asymptotically AdS solutions to the Einstein equations have been constructed in [\[Friedrich 1995\]](#).


1B2. *Quasinormal modes of the asymptotically AdS black holes.* Quasinormal modes, also called resonances, are complex frequencies generalizing the well-known normal modes to systems which dissipate energy. There is a strong connection between quasimodes and resonances [\[Tang and Zworski 1998\]](#). One way to mathematically define them is as poles of the meromorphic continuation of a truncated resolvent. In the case of asymptotically de Sitter black holes, this theory has been very successfully developed; see [\[Bony and Häfner 2008; Dyatlov 2011; 2012\]](#). In a recent paper, Gannot [\[2014\]](#) has established, in the case of Schwarzschild–AdS, the existence of a sequence of quasinormal modes (based on an independent construction of quasimodes), indexed by angular momentum ℓ , and with imaginary parts of size $\mathcal{O}(\exp(-\ell/C))$. In particular, his construction confirms the numerical results of [\[Festuccia and Liu 2009\]](#) and provides an independent proof of [Theorem 1.5](#) and [Corollary 1.6](#), albeit restricted to the Schwarzschild–AdS case. While we restrict ourselves here to the construction of quasimodes, we strongly believe that our results can be used as a basis for the construction of resonances in the Kerr–AdS case for the whole range of parameters satisfying $|a| < l$ and $r_+^2 > |a|l$.

1B3. Universal minimal decay rates of waves outside stationary black holes. In the context of the obstacle problem in Minkowski space, a celebrated result of Burq [1998] establishes a logarithmic decay rate for the local energy of waves, independently of the geometry of the obstacle causing the trapping. For waves outside black holes, in view of the results of [Holzegel and Smulevici 2013a], a natural conjecture is: Given any black hole exterior (\mathcal{R}, g) of a stationary spacetime, a logarithmic decay of energy similar to that of [ibid.] will hold, provided a uniform boundedness statement is true for solutions to the wave equation $\square_g \psi = 0$ on \mathcal{R} .

1C. Outline and overview of the proof. Section 2 introduces the family of Kerr–AdS spacetimes as well as the norms required to state our estimates. In Section 3, we exploit the classical fact that the wave equation separates on Kerr–AdS. An ingredient which considerably simplifies our analysis here is the important observation that we can restrict ourselves to *axisymmetric* solutions. With axisymmetry, the separation of variables leads to relatively simple, one-dimensional, second order ordinary differential equations for the radial functions. In the case of Schwarzschild, they are roughly of the form of the semiclassical problem

$$-u''_\ell \frac{1}{\ell(\ell+1)} + V_\sigma u_\ell = \frac{\omega^2}{\ell(\ell+1)} u_\ell \tag{5}$$

for a potential $V_\sigma(r)$, whose general form is depicted below.⁵ In Section 3C, we shall describe in detail the analytic properties of the potentials appearing in these equations.



To construct the quasimodes, we first construct eigenfunctions for the problem (5) with Dirichlet conditions $u = 0$ imposed on u at $r = r_{\max}$ and $r \rightarrow \infty$ (Section 4). In particular, we prove a version of Weyl’s law, ensuring that for any energy between $1/l^2 < E < V_{\max} = V_\sigma(r_{\max})$ we can find (lots of) eigenvalues

⁵For the purpose of this exposition, we neglect terms of lower order in $\frac{1}{\ell(\ell+1)}$ in the potential, as well as the mass term. The latter is actually unbounded and needs to be absorbed with a Hardy inequality. We suppress such technical difficulties in the present discussion.

$\omega_\ell^2/(\ell(\ell + 1))$ of (5) in a strip $[E - \delta, E + \delta]$. In the Kerr case, the eigenvalue problem (5) turns into a problem of the form⁶

$$-u_\ell'' \frac{1}{\mu_\ell(a^2\omega^2)} + V_\sigma u_\ell = \frac{\omega^2}{\mu_\ell(a^2\omega^2)} u_\ell, \tag{6}$$

which is *nonlinear* in ω^2 . An application of the implicit function theorem together with global estimates on the behavior of the eigenvalues still allow us to conclude the existence of eigenfunctions u_ℓ of (6) with corresponding eigenvalues in the range $(1/l^2, E + \delta]$. *These estimates, together with the analysis of the potential in the Kerr–AdS case, constitute the core of our paper.*

In Section 5, we recall the so-called Agmon estimates, their proof being included to make the paper self-contained. These estimates quantify that the solutions u_ℓ constructed from the above eigenvalue problem decay exponentially in ℓ in a region $[r_{\max}, r_{\max} + \delta']$.

In Section 6, the quasimodes are constructed by cutting off the solution u_ℓ of the eigenvalue problem in $[r_{\max}, r_{\max} + \delta']$ so that it vanishes with all derivatives at r_{\max} and then continuing it to be identically zero in $[r_+, r_{\max}]$. The function ϕ_ℓ thus constructed will be defined on $[r_+, \infty)$ and the corresponding wave function $\psi_\ell = e^{i\omega_\ell t} \phi_\ell$ will satisfy the wave equation everywhere except in the small strip $[r_{\max}, r_{\max} + \delta']$, where the error is exponentially small by the Agmon estimate.

In the last section, we prove Corollary 1.6 using the Duhamel formula. Finally, the Appendix contains a proof of boundedness for the second energy used in this paper. This boundedness statement differs from that obtained in [Holzegel 2010; Holzegel and Warnick 2014] in that it allows for stronger radial weights near infinity for the angular derivatives.

2. Preliminaries

2A. The Kerr–AdS family of spacetimes. We recall here some basic facts about the family of Kerr–AdS spacetimes required in the paper. We refer the reader to the detailed discussion in our work [Holzegel and Smulevici 2013a].

2A1. The fixed manifold with boundary \mathcal{R} . Let \mathcal{R} denote the manifold with boundary

$$\mathcal{R} = [0, \infty) \times \mathbb{R} \times \mathbb{S}^2.$$

We define standard coordinates y^* for $[0, \infty)$, t^* for \mathbb{R} and (θ, ϕ) for \mathbb{S}^2 . This defines a coordinate system on \mathcal{R} which is global up to the well-known degeneration of the spherical coordinates. We define the event horizon \mathcal{H}^+ to be the boundary of \mathcal{R} :

$$\mathcal{H}^+ = \partial\mathcal{R} = \{y^* = 0\}.$$

The manifold \mathcal{R} will coincide with the domain of outer communication of the black hole spacetimes including the future event horizon \mathcal{H}^+ .

⁶The $\mu_\ell(a^2\omega^2)$ generalize the familiar spherical eigenvalues $\ell(\ell + 1)$ of the Schwarzschild case to Kerr. See Section 3A.

2A2. *The parameter space and the radial function.* A Kerr–AdS spacetime is characterized by its mass $M > 0$, its angular momentum per unit mass a and the value of the cosmological constant $\Lambda < 0$. For convenience, we shall use mostly $l = \sqrt{3/|\Lambda|}$, instead of Λ . For the spacetime to be regular, we will require $|a| < l$. Let us thus fix $M, l > 0$ and $|a| < l$. We then define $r_+(M, l, a) > 0$ as the unique real solution of $\Delta_-(x) = 0$, where

$$\Delta_-(x) = \left(1 + \frac{x^2}{l^2}\right)(x^2 + a^2) - 2Mx. \tag{7}$$

We now define a function r on \mathcal{R} as follows. As a function of $(t^*, y^*, \varphi, \theta)$, r only depends on y^* and is a diffeomorphism from $[0, \infty)$ to $[r_+, \infty)$. The collection $(t^*, r(y^*), \varphi, \theta)$ then forms a coordinate system on \mathcal{R} , global up to the degeneration of the spherical coordinates. Moreover, the horizon \mathcal{H}^+ coincides with $\{r = r_+\}$.

2A3. *More coordinates: $r^*, t, \tilde{\phi}$.* Let us define r^* by

$$\frac{dr^*}{dr}(r) = \frac{r^2 + a^2}{\Delta_-(r)}, \quad r^*(r = +\infty) = \frac{\pi}{2},$$

where $\Delta_-(r)$ is given by (7). Note that $r^*(r_+) = -\infty$.

By a small abuse of notation, we shall often write for functions f and g , $f(r^*) = g(r)$, instead of $f(r^*) = g(r(r^*))$ or $f(r^*(r)) = g(r)$.

Finally, let $r_{\text{cut}} = r_+ + \frac{1}{2}(r_{\text{max}} - r_+)$, with $r_{\text{max}} > r_+$ defined in Lemma 3.1 depending only on the parameters M, a and l , and let $\chi(r)$ be a smooth cut-off function with the property

$$\chi(r) = \begin{cases} 1 & \text{if } r \in [r_+, \frac{1}{2}(r_{\text{cut}} - r_+)], \\ 0 & \text{if } r \geq r_{\text{cut}}. \end{cases} \tag{8}$$

We introduce the time coordinate t and another angular coordinate $\tilde{\phi}$ as

$$t = t^* - A(r)\chi(r) \quad \text{and} \quad \tilde{\phi} = \phi - B(r)\chi(r), \tag{9}$$

where

$$\frac{dA}{dr} = \frac{2Mr}{\Delta_-(1 + r^2/l^2)}, \quad \frac{dB}{dr} = \frac{a(1 - a^2/l^2)}{\Delta_-},$$

and A and B vanish at infinity.

Note that $t, \tilde{\phi}$ and r^* are not well-behaved functions at the horizon $r = r_+$. As a consequence, the coordinate systems $(t, r, \theta, \tilde{\phi})$ and (t, r^*, θ, ϕ) only cover $\text{int}(\mathcal{R})$. Observe also that the two coordinate systems $(t, r, \theta, \tilde{\phi})$ and (t^*, r, θ, ϕ) are identical for $r \geq r_{\text{cut}}$.

2A4. *The Kerr–AdS metric for fixed (a, M, l) .* We may now introduce the Kerr–AdS metric as the unique smooth extension to \mathcal{R} of the tensor given in the Boyer–Lindquist chart by

$$g_{\text{KAdS}} = \frac{\Sigma}{\Delta_-} dr^2 + \frac{\Sigma}{\Delta_\theta} d\theta^2 + \frac{\Delta_\theta(r^2 + a^2)^2 - \Delta_- a^2 \sin^2 \theta}{\Xi^2 \Sigma} \sin^2 \theta d\tilde{\phi}^2 - 2 \frac{\Delta_\theta(r^2 + a^2) - \Delta_-}{\Xi \Sigma} a \sin^2 \theta d\tilde{\phi} dt - \frac{\Delta_- - \Delta_\theta a^2 \sin^2 \theta}{\Sigma} dt^2, \tag{10}$$

where Δ_- is defined by (7) and

$$\Sigma = r^2 + a^2 \cos^2 \theta, \quad \Delta_\theta = 1 - \frac{a^2}{l^2} \cos^2 \theta, \quad \Xi = 1 - \frac{a^2}{l^2}. \tag{11}$$

See [Holzegel and Smulevici 2013a] for explicit expressions for the inverse of (10) and its determinant in Boyer–Lindquist coordinates. That the tensor (10) indeed extends to a smooth metric on \mathcal{R} is clear from expressing the metric in (t^*, r, θ, ϕ) coordinates, which is carried out explicitly in [ibid.]. Note that for $a = 0$ the metric (10) reduces to the well-known Schwarzschild–AdS spacetime

$$g = -\left(1 - \frac{2M}{r} + \frac{r^2}{l^2}\right) dt^2 + \left(1 - \frac{2M}{r} + \frac{r^2}{l^2}\right)^{-1} dr^2 + r^2 d\sigma_{S^2},$$

where $d\sigma_{S^2}$ is the standard metric on the unit sphere.

The Boyer–Lindquist coordinates degenerate at the horizon $r = r_+$, but the importance of these coordinates is that it is in these coordinates that the Klein–Gordon operator separates (see Section 3).

2A5. The Klein–Gordon operator in Boyer–Lindquist coordinates. For any Lorentzian metric g and any scalar function ψ , we define as usual

$$\square_g \psi = \frac{1}{\sqrt{|g|}} \partial_\alpha (\sqrt{|g|} g^{\alpha\beta} \partial_\beta \psi);$$

the Klein–Gordon operator acting on scalar functions in Boyer–Lindquist coordinates is given by

$$\begin{aligned} \left(\square_g + \frac{\alpha}{l^2}\right) \psi &= \left(-\frac{(r^2 + a^2)^2}{\Sigma \Delta_-} + \frac{a^2 \sin^2 \theta}{\Sigma \Delta_\theta}\right) \partial_t^2 \psi + \frac{1}{\Sigma} \partial_r (\Delta_- \partial_r \psi) + 2\left(\frac{\Xi(r^2 + a^2)a}{\Delta_- \Sigma} - \frac{\Xi a}{\Delta_\theta \Sigma}\right) \partial_t \partial_\phi \psi \\ &+ \left(\frac{\Xi^2}{\Delta_\theta \Sigma} \frac{1}{\sin^2 \theta} - \frac{\Xi^2 a^2}{\Sigma \Delta_-}\right) \partial_\phi \partial_\phi \psi + \frac{1}{\Sigma \sin \theta} \partial_\theta (\Delta_\theta \sin \theta \partial_\theta \psi) + \frac{\alpha}{l^2} \psi. \end{aligned} \tag{12}$$

2B. The norms. Let g and ∇ denote the induced metric and the covariant derivative on the spheres $S_{t^*, r}^2$ of constant t^* and r in \mathcal{R} .

We write $|\nabla \cdots \nabla \psi|^2 = g^{AA'} \cdots g^{BB'} \nabla_A \cdots \nabla_B \psi \nabla_{A'} \cdots \nabla_{B'} \psi$ to denote the induced norms on these spheres. We denote by Ω_i with $i = 1, 2, 3$ the standard basis of angular momentum operators on the unit sphere in θ, ϕ coordinates.

With these conventions, we define the energy densities

$$\begin{aligned} e_1[\psi] &= \frac{1}{r^2} (\partial_{t^*} \psi)^2 + r^2 (\partial_r \psi)^2 + |\nabla \psi|^2 + \psi^2, \\ \tilde{e}_2[\psi] &= e_1[\psi] + e_1[\partial_{t^*} \psi] + r^4 (\partial_r \partial_r \psi)^2 + r^2 |\partial_r \nabla \psi|^2 + |\nabla \nabla \psi|^2, \\ e_2[\psi] &= \tilde{e}_2[\psi] + \sum_i e_1[\Omega_i \psi]. \end{aligned}$$

Similarly, we define the following energy norms for the scalar field ψ (see [Holzegel 2012]):

$$\begin{aligned} \|\psi\|_{H_{\text{AdS}}^{0,s}(\Sigma_{t^*})}^2 &= \int_{\Sigma_{t^*}} r^s \psi^2 r^2 dr \sin \theta d\theta d\phi \\ \|\psi\|_{H_{\text{AdS}}^{1,s}(\Sigma_{t^*})}^2 &= \int_{\Sigma_{t^*}} r^s (r^2 (\partial_r \psi)^2 + |\nabla \psi|^2 + \psi^2) r^2 dr \sin \theta d\theta d\phi \\ \|\psi\|_{H_{\text{AdS}}^{2,s}(\Sigma_{t^*})}^2 &= \|\psi\|_{H_{\text{AdS}}^{1,s}(\Sigma_{t^*})}^2 + \int_{\Sigma_{t^*}} r^s (r^4 (\partial_r \partial_r \psi)^2 + r^2 |\nabla \partial_r \psi|^2 + |\nabla \nabla \psi|^2) r^2 dr \sin \theta d\theta d\phi. \end{aligned}$$

Note in particular the relations

$$\int_{\Sigma_{t^*}} e_1[\psi] r^2 dr \sin \theta d\theta d\phi = \|\psi\|_{H_{\text{AdS}}^{1,0}(\Sigma_{t^*})}^2 + \|\partial_{t^*} \psi\|_{H_{\text{AdS}}^{0,-2}(\Sigma_{t^*})}^2, \tag{13}$$

$$\int_{\Sigma_{t^*}} e_2[\psi] r^2 dr \sin \theta d\theta d\phi = \|\Omega_i \psi\|_{H_{\text{AdS}}^{1,0}(\Sigma_{t^*})}^2 + \|\partial_{t^*} \psi\|_{H_{\text{AdS}}^{1,0}(\Sigma_{t^*})}^2 + \|\psi\|_{H_{\text{AdS}}^{2,0}(\Sigma_{t^*})}^2 + \|\partial_{t^*} \partial_{t^*} \psi\|_{H_{\text{AdS}}^{0,-2}(\Sigma_{t^*})}^2. \tag{14}$$

Higher-order norms may be defined similarly. We denote by $H_{\text{AdS}}^{k,s}(\Sigma_{t^*})$ the space of functions ψ such that $\nabla^i \psi \in L_{loc}^2(\Sigma_{t^*})$ for $i = 0, \dots, k$ and $\|\psi\|_{H_{\text{AdS}}^{k,s}(\Sigma_{t^*})} < \infty$. We denote by $CH_{\text{AdS}}^{k,s}$ the set of functions ψ defined on \mathcal{R} such that

$$\psi \in \bigcap_{q=0,\dots,k} C^q(\mathbb{R}_{t^*}; H_{\text{AdS}}^{k-q,s_q}(\Sigma_{t^*})), \quad \text{where } s_k = -2, s_{k-1} = 0 \text{ and } s_j = s \text{ for } j = 0, \dots, k-2.$$

When $s = 0$, we will feel free to drop the s in the notation, i.e., $H_{\text{AdS}}^{k,0} := H_{\text{AdS}}^k$ and $CH_{\text{AdS}}^k := CH_{\text{AdS}}^{k,0}$.

2C. A final remark. In [Holzegel and Smulevici 2013a], the coordinates t and $\tilde{\phi}$ in (9) are defined with the $\chi(r)$ of (8) being globally equal to 1. Here, for convenience in the subsequent analysis (which happens mostly away from the horizon, in $r \geq r_{\text{max}}$), we have altered these coordinates away from the horizon to agree with the Boyer–Lindquist coordinates. Note that these two coordinate systems are equivalent in the sense that the statement $\|\psi\|_{H_{\text{AdS}}^{k,s}(\Sigma_{t^*})}^2$ decays logarithmically in t^* is independent of whether the coordinates (and Σ_{t^*} -slices) of our earlier paper or the cut-off coordinates (9) are used.

3. Separation of variables and reduced equations

3A. The (modified) oblate spheroidal harmonics. For each $\xi \in \mathbb{R}$, define the unbounded $L^2(\sin \theta d\theta d\tilde{\phi})$ -self-adjoint operator $P_{\mathbb{S}^2}(\xi)$ with domain the space of $H^2(\mathbb{S}^2)$ -complex valued functions (see Section 7 of [Dafermos and Rodnianski 2010a] for a more detailed discussion), as

$$-P_{\mathbb{S}^2}(\xi) f = \frac{1}{\sin \theta} \partial_\theta (\Delta_\theta \sin \theta \partial_\theta f) + \frac{\Xi^2}{\Delta_\theta} \frac{1}{\sin^2 \theta} \partial_\phi^2 f + \Xi \frac{\xi^2}{\Delta_\theta} \cos^2 \theta f - 2i\xi \frac{\Xi}{\Delta_\theta} \frac{a^2}{l^2} \cos^2 \theta \partial_{\tilde{\phi}} f. \tag{15}$$

We also define the operator P_α , which is equal to

$$P_{\mathbb{S}^2,\alpha}(\xi) = \begin{cases} P(\xi) + \frac{\alpha}{l^2} a^2 \sin^2 \theta & \text{if } \alpha > 0, \\ P(\xi) + \frac{|\alpha|}{l^2} a^2 \cos^2 \theta & \text{if } \alpha \leq 0. \end{cases} \tag{16}$$

The motivation for introducing these operators is that, formally replacing ξ by $ai \partial_t \psi$, they appear naturally in the Klein–Gordon operator (12) when trying to separate variables; see the next section.

When $l \rightarrow \infty$ the operator $P_{\mathbb{S}^2}(\xi)$ reduces to an oblate spheroidal operator on \mathbb{S}^2 as considered for instance in [ibid.]. If also $\xi = a = 0$ we recover the Laplacian on the round sphere. Both operators $P_{\mathbb{S}^2}(\xi)$ and $P_{\mathbb{S}^2, \alpha}(\xi)$ have discrete spectra. We will use the following notation:

$$P_{\mathbb{S}^2, \alpha}(\xi) \text{ has eigenvalues } \lambda_{m\ell}(\xi) \text{ with eigenfunctions } S_{m\ell}(\xi, \cos \theta)e^{im\tilde{\phi}}.$$

Later we will restrict attention to axisymmetric solutions and hence to the eigenvalues $\lambda_{0\ell}$ of the operators

$$-P_{\theta, \alpha}(\xi^2) := -P_{\mathbb{S}^2, \alpha}(\xi)|_{m=0}. \tag{17}$$

In fact, it will be convenient (in view of their manifest positivity) to work with the eigenvalues of the shifted operators, which are, for $\alpha \leq 0$,

$$-P_{\theta, \alpha}(\xi^2) - \xi^2 = \frac{1}{\sin \theta} \partial_\theta (\Delta_\theta \sin \theta \partial_\theta (\cdot)) - \frac{|\alpha|}{l^2} a^2 \cos^2 \theta - \frac{\sin^2 \theta}{\Delta_\theta} \xi^2 \tag{18}$$

(\cos^2 is replaced by \sin^2 in the second term if $\alpha > 0$). In view of these considerations, we shall denote the eigenvalues of the operator $P_{\theta, \alpha}(\xi^2) + \xi^2$ by $\mu_\ell(\xi^2)$. By min-max and comparison with the spherical Laplacian [Holzegel and Smulevici 2013a, Lemma 5.1] in the axisymmetric case, we have

$$\mu_\ell(\xi^2) \geq \mu_\ell(0) \geq \Xi \ell(\ell + 1) > c_{a,l} \ell(\ell + 1). \tag{19}$$

3B. The separation of variables. We now present the reduced equations obtained after separation of variables. For this purpose, we consider the Klein–Gordon operator in Boyer–Lindquist coordinates (12).

For the construction of quasimodes, it would be sufficient to start directly from the reduced equations. However, we will instead derive them from the Klein–Gordon equation (1) to show this relation. Thus, in this section ψ will denote any regular solution to (1). Let us introduce the time-Fourier transform⁷

$$\psi(t, r, \theta, \tilde{\phi}) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{-i\omega t} \hat{\psi}(\omega, r, \theta, \tilde{\phi}) d\omega. \tag{20}$$

We decompose the $\hat{\psi}$ of (20) as

$$\hat{\psi}(\omega, r, \theta, \tilde{\phi}) = \sum_{m\ell} (\hat{\psi})_{m\ell}^{(a\omega)}(r) S_{m\ell}(a\omega, \cos \theta) e^{im\tilde{\phi}}, \tag{21}$$

where $S_{m\ell}$ are the modified spheroidal harmonics introduced above and

$$(\hat{\psi})_{m\ell}^{(a\omega)}(r) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} dt \int_{\mathbb{S}^2(t,r)} d\tilde{\sigma} e^{i\omega t} S_{m\ell}(a\omega, \cos \theta) e^{-im\tilde{\phi}} \psi(t, r, \theta, \tilde{\phi}),$$

with $d\tilde{\sigma} = \sin \theta d\theta d\tilde{\phi}$.

After the renormalization

$$u_{m\ell}^{(a\omega)}(r) = (r^2 + a^2)^{1/2} (\hat{\psi})_{m\ell}^{(a\omega)}(r), \tag{22}$$

⁷Note that in general ψ is not an L^2 -function in time and therefore, $\hat{\psi}$ is defined only as a tempered distribution. Since here we are merely trying to justify the origin of the reduced equations, it will be sufficient to understand all computations formally.

we finally obtain from (1) and (12) the equation

$$(u_{m\ell}^{(a\omega)}(r))'' + (\omega^2 - V_{m\ell}^{(a\omega)}(r))u_{m\ell}^{(a\omega)} = 0, \tag{23}$$

where ' denotes differentiation with respect to r^* and where the potential $V_{m\ell}^{(a\omega)}(r)$ is defined as

$$V_{m\ell}^{(a\omega)}(r) = V_{+,m\ell}^{(a\omega)}(r) + V_{0,m\ell}^{(a\omega)}(r) + V_\alpha(r), \tag{24}$$

where

$$V_{+,m\ell}^{(a\omega)}(r) = -\Delta_-^2 \frac{3r^2}{(r^2+a^2)^4} + \Delta_- \frac{5r^4/l^2 + 3r^2(1+a^2/l^2) - 4Mr + a^2}{(r^2+a^2)^3} = (r^2+a^2)^{-1/2}(\sqrt{r^2+a^2})'',$$

$$V_{0,m\ell}^{(a\omega)}(r) = \frac{\Delta_-(\lambda_{m\ell} + \omega^2 a^2) - \Xi^2 a^2 m^2 - 2m\omega a \Xi(\Delta_- - (r^2+a^2))}{(r^2+a^2)^2}, \tag{25}$$

$$V_\alpha(r) = -\frac{\alpha}{l^2} \frac{\Delta_-}{(r^2+a^2)^2} (r^2 + \Theta(\alpha)a^2), \tag{26}$$

with $\Theta(\alpha) = 1$ if $\alpha > 0$ and $\Theta(\alpha) = 0$ if $\alpha \leq 0$ (recall that the $\lambda_{m\ell}$ also depend on α through (16)). Note that V_+ grows like $2r^2/l^4$ near infinity, while the V_0 part remains bounded.

3C. The axisymmetric reduced equations. We now look at the axisymmetric case; that is, we consider the above equations under the assumption that ψ is independent of the azimuthal variable $\tilde{\phi}$. The reduced equations are then obtained by setting $m = 0$ in the decomposed equations. Hence, we will consider the following set of equations:

$$(u_{0\ell}^{(a\omega)}(r))'' + (\omega^2 - V_{0\ell}^{(a\omega)}(r))u_{0\ell}^{(a\omega)}(r) = 0, \tag{27}$$

where the potential $V_{0\ell}^{(a\omega)}(r)$ is defined as

$$V_{0\ell}^{(a\omega)}(r) = V_{\text{junk}}(r) + V_{\text{mass}}(r) + V_\sigma(r) \cdot \mu_\ell(\omega^2 a^2), \tag{28}$$

where

$$V_{\text{junk}}(r) = -\Delta_-^2 \frac{3r^2}{(r^2+a^2)^4} + \Delta_- \frac{5r^4/l^2 + 3r^2(1+a^2/l^2) - 4Mr + a^2}{(r^2+a^2)^3} - \frac{2}{l^2} \frac{\Delta_- r^2}{(r^2+a^2)^2} - \frac{\alpha}{l^2} \frac{\Delta_-}{(r^2+a^2)^2} \Theta(\alpha)a^2,$$

$$V_{\text{mass}}(r) = \frac{2-\alpha}{l^2} \frac{\Delta_- r^2}{(r^2+a^2)^2}, \quad V_\sigma(r) = \frac{\Delta_-}{(r^2+a^2)^2}. \tag{29}$$

Here we rearranged the terms in the different potentials so that $V_{\text{mass}} = 0$ corresponds to the conformal case $\alpha = 2$. In particular, we have

$$|V_{\text{junk}}(r)| \leq C_{M,l,a} \frac{\Delta_-}{(r^2+a^2)^2} \lesssim V_\sigma;$$

hence V_{junk} is uniformly bounded.

Lemma 3.1 (properties of V_σ). *For all $|a| < l$, the potential V_σ enjoys the following properties:*

- $V_\sigma(r^*) \rightarrow 1/l^2$ as $r^* \rightarrow \pi/2$.

- V_σ has a unique local and global maximum V_{\max} at r_{\max}^* in $[r_+^*, \pi/2)$. Also, V_σ is monotonically decreasing in $[r_{\max}^*, \pi/2)$.
- $V_{\max} = V_\sigma(r_{\max}^*) \geq \frac{1}{l^2} + \frac{3M/\Xi + \Xi a^2}{((3M/\Xi)^2 + a^2)^2}$.

Remark 3.2. In particular, for any $0 < a_0 < l$ and all $|a| \leq a_0$, the size of the interval $[1/l^2, V_\sigma(r_{\max}^*)]$ is bounded uniformly (in a) from below by a strictly positive uniform constant.

Proof. The first claim can be trivially checked. For the second and third claims, let us write

$$V_\sigma(r) = \frac{\Delta_-}{(r^2 + a^2)^2} = \frac{(r^2 + a^2)(r^2/l^2 + a^2/l^2 + \Xi) - 2Mr}{(r^2 + a^2)^2} = \frac{1}{l^2} + \frac{\Xi(r^2 + a^2) - 2Mr}{(r^2 + a^2)^2}. \tag{30}$$

The r -derivative of V_σ is

$$\begin{aligned} \partial_r V_\sigma(r) &= \frac{(r^2 + a^2)^2(2\Xi r - 2M) - 4r(r^2 + a^2)(\Xi(r^2 + a^2) - 2Mr)}{(r^2 + a^2)^4} \\ &= \frac{(r^2 + a^2)(2\Xi r - 2M) - 4r(\Xi(r^2 + a^2) - 2Mr)}{(r^2 + a^2)^3} = \frac{-2\Xi r^3 + 6Mr^2 - 2\Xi a^2 r - 2Ma^2}{(r^2 + a^2)^3}. \end{aligned} \tag{31}$$

If $a = 0$, then one has as usual that the only zero in $[r_+, \infty)$ is at $r = 3M$. Assume $a \neq 0$. Observe that the derivative is positive near $-\infty$, negative at $r = 0$, positive at $r = r_+$ and negative near ∞ . This tells us that there are three real roots for $\partial_r V_\sigma$. The one of interest to us is the one corresponding to the (unique) maximum of V_σ in the interval $[r_+, \infty)$. Define $r_{\text{guess}} = 3M/\Xi$. We claim $r_+ < r_{\text{guess}} < \infty$. This follows from $r_+ < 2M$, which is in turn a consequence of the fact that $\Delta_- = r(r - 2M) + r^4/l^2 + a^2(1 + r^2/l^2)$ is positive for $r \geq 2M$. Finally, at $r_{\text{guess}} = 3M/\Xi$ we have from (30)

$$V_\sigma(r_{\text{guess}}) = \frac{1}{l^2} + \frac{3M/\Xi + \Xi a^2}{(r_{\text{guess}}^2 + a^2)^2} > \frac{1}{l^2}. \quad \square$$

4. Bound states

As proven in [Holzegel and Smulevici 2013a], there exist no periodic solutions of the massive wave equation on Kerr–AdS. In this section, we will introduce an additional boundary, located at $r = r_{\max}$ (the location of the top of the potential V_σ , as defined in Lemma 3.1), enabling us to construct periodic solutions whose associated energies lie below the top of V_σ .

In order to avoid confusion between the mode number ℓ and the real number l determining the cosmological constant, we introduce the semiclassical parameter $h > 0$ by defining

$$h^{-2}(\ell, \omega) := \lambda_{0\ell}(a^2\omega^2) + a^2\omega^2 = \mu_\ell(a^2\omega^2), \tag{32}$$

as well as the shorthand

$$h_0^{-2} = h^{-2}(\ell, 0) = \mu_\ell(0). \tag{33}$$

In the rest of this paper, the notation u_ℓ or u_h will be used when we want to make explicit that a solution u depends on ℓ or h .

Having in mind a semiclassical-type analysis with semiclassical parameter h , we then rewrite (27) as the *nonlinear eigenvalue problem*

$$h^2\omega^2 u = P(h)u := -h^2u'' + V_\sigma u + h^2(V_{\text{junk}} + V_{\text{mass}})u, \tag{34}$$

with boundary conditions $u(r_{\text{max}}^*) = 0$ and $\int_{r_{\text{max}}^*}^{\pi/2} (|u'|^2 + r^2u^2) dr^* < \infty$.

Remark 4.1. We will refer to these boundary conditions as Dirichlet conditions. They imply that $u(\pi/2) = 0$. See Remark 4.2 below.

Unless $a = 0$, (34) is a nonlinear eigenvalue problem. Indeed, a solution to the eigenvalue problem

$$\kappa u = P(h)u \tag{35}$$

with Dirichlet boundary conditions is a solution of (34) if and only if $\kappa = h^2\omega^2$. If $a = 0$, (34) reduces to the linear problem (35), since the $P(h)$ operator becomes independent of ω , and, therefore, given any solution to (35), one simply obtains a solution to (34) by defining $\omega^2 = h^{-2}\kappa$.

What we would like to prove is that, given fixed parameters M, a, l and α , we can find, for any sufficiently small h (or, equivalently, any sufficiently large ℓ), an ω_ℓ^2 such that (34) is solved for some u_ℓ , and that we can control the size of this ω_ℓ^2 (see Proposition 4.8 below).

In order to understand (34), it will be useful to first look at the following *linear eigenvalue problem*:

$$h_0^2\omega^2 u = P_{\text{base}}(h_0)u := -h_0^2u'' + V_\sigma u + h_0^2(V_{\text{junk}} + V_{\text{mass}})u, \tag{36}$$

with boundary conditions $u(r_{\text{max}}^*) = 0$ and $\int_{r_{\text{max}}^*}^{\pi/2} [|u'|^2 + r^2u^2] dr^* < \infty$.

As explained above, (36) can be seen as a linear eigenvalue problem because h_0 , and therefore $P_{\text{base}}(h_0)$, depends only on ℓ (but not on ω).

Remark 4.2. By Proposition 4 of [Holzegel and Warnick 2014], (36) is a well-posed eigenvalue problem. This is a nontrivial statement because the potential V_{mass} is unbounded on the domain $(r_{\text{max}}^*, \pi/2i)$ unless $\alpha = 2$. The condition $\int_{r_{\text{max}}^*}^{\pi/2} [|u'|^2 + r^2u^2] dr^* < \infty$ implies that $\psi = \sqrt{r^2 + a^2} u \cdot S_{0\ell}(\theta) \in H_{\text{ADS}}^1$ (in particular, $r^{1/2-\epsilon}u(\pi/2) = 0$ for any $\epsilon > 0$) and ensures the existence of a positive discrete spectrum with eigenfunctions in the energy space.⁸

To prove that eigenvalues $h_0^2\omega^2$ for (36) exist in a suitable range, we will perform a semiclassical-type analysis for a semiclassical operator whose principal part should be $-h_0^2u'' + V_\sigma u$. Since $h_0^2V_{\text{junk}}$ is controlled by $h_0^2V_\sigma$, this term will be of lower order and hence negligible. On the other hand, unless one considers the conformal case $\alpha = 2$, the a priori lower-order (in powers of h_0) potential term $h_0^2V_{\text{mass}}$ is unbounded near $r^* = \pi/2$, so that some care (a Hardy inequality) is required.

Observe finally that if we set $a = 0$ in $V_\sigma, V_{\text{junk}}$ and V_{mass} in (36), then

$$P_{\text{base}}^{a=0}(h_0)u = \kappa \cdot u \tag{37}$$

⁸Using the twisted derivatives of [Holzegel and Warnick 2014], i.e., writing $u'' + V_{\text{mass}}u = r^n(r^{-2n}(r^n u)')' + V_{\text{twist}} \cdot u$ with $n = \frac{1}{2}(1 - \sqrt{9 - 4\alpha})$, so V_{twist} is uniformly bounded, one could generalize our construction to other boundary conditions.

would be precisely the eigenvalue problem one needs to study for Schwarzschild–AdS. In any case, in the next section we will establish the existence of eigenvalues $\kappa = h_0^2 \omega^2$ of the more general eigenvalue problem (36) as the latter is easier to connect to the full problem (34).

4A. Weyl’s law for the linear eigenvalue problem (36). The aim of this section is to prove a Weyl’s-law-type result for the linear problem (36). This is a classical problem which we will approach using a slight modification of the usual Dirichlet–Neumann bracketing argument (see for instance [Reed and Simon 1978], Section XIII.15, for an introduction to this method).

For the purpose of this section, it will be convenient to introduce the following notation. For all $c < d$, we define $P_{DD}^{\text{base}}(c, d)$ as the eigenvalue problem

$$P_{\text{base}}(h_0)u = \kappa u,$$

with Dirichlet boundary conditions $u(c) = u(d) = 0$, plus the condition $\int_c^d dr^*(|u'|^2 + r^2 \cdot u^2) < \infty$ if $d = \pi/2$ (see Remark 4.2); here $P_{\text{base}}(h_0)$ is the operator defined by (36). Similarly, we will write $P_{NN}^{\text{base}}(c, d)$ for the Neumann problem (which will never be considered with $d = \pi/2$). Finally, we write $P_{ND}^{\text{base}}(c, d)$ for the Neumann boundary at c and Dirichlet at d . In the latter case we again impose $\int_c^d dr^*(|u'|^2 + r^2 \cdot u^2) < \infty$ if $d = \pi/2$ to ensure that all eigenfunctions live in the energy space; see Remark 4.2. Note that $P_{DD}^{\text{base}}(r_{\text{max}}^*, \pi/2)$ is precisely the linear eigenvalue problem (36).

Proposition 4.3. *Let $\alpha < \frac{9}{4}$, $M > 0$ and $|a| < l$ be fixed, and $E \in (1/l^2, V_{\text{max}})$ be given. Then, for any $\delta > 0$ such that $[E - \delta, E + \delta] \subset (1/l^2, V_{\text{max}})$, there exists an $H_0 > 0$, such that, for any $0 < h_0 \leq H_0$, there exists a smooth solution u_{h_0} of the eigenvalue problem $P_{DD}^{\text{base}}(r_{\text{max}}^*, \pi/2)$, with corresponding eigenvalue κ lying in $[E - \delta, E + \delta]$. In particular, there exists a sequence $((h_0)_n, u_{(h_0)_n})$ such that the associated eigenvalues $\kappa((h_0)_n) \rightarrow E$ as $(h_0)_n \rightarrow 0$.*

In the rest of this section, (a, M, l, α) are fixed parameters satisfying the assumptions of the proposition.

We shall in fact prove in this section a stronger result than Proposition 4.3, namely a version of Weyl’s law adapted to our problem. This is the statement of Lemma 4.5, from which Proposition 4.3 immediately follows. The proof of Lemma 4.5 in turn requires the following auxiliary lemma, which ensures nonexistence of eigenvalues below a certain threshold.

Lemma 4.4. *Let $E > 0$ be given. Then there exists an $H_0 > 0$ so that for all $0 < h_0 \leq H_0$, there exists a $r_K^*(E, h_0, \alpha)$ such that the problems $P_{DD}^{\text{base}}(r_K^*, \pi/2)$ and $P_{ND}^{\text{base}}(r_K^*, \pi/2)$ have no solutions with $\kappa \leq E$. Moreover,*

$$r_{\text{max}} < r_K \leq \frac{C}{h_0} \cdot E, \tag{38}$$

where C depends only on M, a, l and α .

Proof. Assume there was a solution u of $P_{DD}^{\text{base}}(r_K^*, \pi/2)$ or $P_{ND}^{\text{base}}(r_K^*, \pi/2)$ with $\kappa \leq E$. Then we would have

$$\int_{r_K^*}^{\pi/2} dr^* [h_0^2 |u'|^2 + (V_\sigma + h_0^2 V_{\text{junk}} + h_0^2 V_{\text{mass}} - E) |u|^2] \leq 0 \tag{39}$$

for this u . On the other hand, since u solves $P_{DD}^{\text{base}}(r_K^*, \pi/2)$ or $P_{ND}^{\text{base}}(r_K^*, \pi/2)$, we have $r^{1/2}u(r) = o(1)$,

so that the Hardy inequality

$$\int_{r_K^*}^{\pi/2} dr^* \frac{\Delta_-}{r^2 + a^2} |u|^2 \leq 4l^2 \int_{r_K^*}^{\pi/2} dr^* |u'|^2 \tag{40}$$

proven in [Holzegel and Smulevici 2013a] holds for u . This implies that

$$\int_{r_K^*}^{\pi/2} dr^* \left[\left(h_0^2 \frac{1}{4l^2} \frac{\Delta_-}{r^2 + a^2} + V_\sigma + h_0^2 V_{\text{junk}} + h_0^2 V_{\text{mass}} - E \right) |u|^2 \right] \leq 0.$$

The dominant term in the integrand near infinity is $h_0^2(\frac{9}{4} - \alpha)r^2/l^2$, which is positive, while all other terms remain bounded. Hence by choosing r_K^* sufficiently large ($r_K \geq C/h_0 \cdot E$ for some constant C) we obtain a contradiction as the parenthetical in the integrand eventually becomes positive. \square

Consider now the eigenvalue problem $P_{DD}^{\text{base}}(r_{\text{max}}^*, \pi/2)$ and fix an energy level $\mathcal{E} \in (1/l^2, V_{\text{max}})$. Lemma 4.4 produces an $r_K^*(\mathcal{E}, h_0, \alpha)$, to which we associate the phase space volume

$$\begin{aligned} \mathcal{P}_{\mathcal{E}, h_0, \alpha} &= \text{Vol}\{(\xi, r^*) \in \mathbb{R} \times [r_{\text{max}}^*, r_K^*] \mid \xi^2 + V_\sigma + h_0^2 V_{\text{mass}} + h_0^2 V_{\text{junk}} \leq \mathcal{E}\} \\ &= 2 \int_{r_{\text{max}}^*}^{r_K^*} dr^* \sqrt{\mathcal{E} - V(r^*)} \cdot \chi_{\{V(r^*) \leq \mathcal{E}\}}. \end{aligned} \tag{41}$$

Note that for fixed \mathcal{E} this expression converges uniformly in h_0 as $h_0 \rightarrow 0$. This is already immediate for $\alpha \leq 2$: $V(r^*)$ is then bounded below and hence the integrand itself is obviously uniformly bounded in h_0 . For $2 < \alpha < \frac{9}{4}$, the integral (41) also converges uniformly in h_0 since

$$\int_{r_{\text{max}}^*}^{r_K^*} dr^* h_0 r \leq C \int_{r_{\text{max}}^*}^{r_K^*} dr h_0 \frac{1}{r} \leq Ch_0 \log\left(\frac{C}{h_0} E\right)$$

goes to zero as $h_0 \rightarrow 0$. Here we have used the estimate (38) on r_K .

Finally, to state and prove Weyl’s law, we also introduce an expression for the phase space volume between two energy levels, say $[E - \delta, E + \delta] \subset (1/l^2, V_{\text{max}})$:

$$\mathcal{Q}_{E, \alpha} = \lim_{h_0 \rightarrow 0} \mathcal{P}_{E+\delta, h_0, \alpha} - \lim_{h_0 \rightarrow 0} \mathcal{P}_{E-\delta, h_0, \alpha} = \text{Vol}\{(\xi, r^*) \mid E - \delta \leq \xi^2 + V_\sigma \leq E + \delta\}.$$

By an elementary computation, we have a lower bound $\mathcal{Q}_{E, \alpha} \geq C_{E, M, l, \alpha} \delta$ for a constant independent of h_0 .

Lemma 4.5. *Consider the eigenvalue problem $P_{DD}^{\text{base}}(r_{\text{max}}^*, \pi/2)$. Fix an energy level $V_{\text{max}} > E > 1/l^2$ and prescribe a small $\delta > 0$. Then the number of eigenvalues of $P_{DD}^{\text{base}}(r_{\text{max}}^*, \pi/2)$ lying in the interval $[E - \delta, E + \delta] \subset (1/l^2, V_{\text{max}})$, denoted by $N[E - \delta, E + \delta]$, satisfies Weyl’s law*

$$N[E - \delta, E + \delta] \underset{h_0 \rightarrow 0}{\sim} \frac{1}{2\pi h_0} \mathcal{Q}_{E, \alpha}. \tag{42}$$

Proof. Choose $r_K^*(E + \delta, h_0)$ such that by Lemma 4.4 there are no eigenvalues below $E + \delta$ of $P_{DD}^{\text{base}}(r_K^*, \pi/2)$ and $P_{ND}^{\text{base}}(r_K^*, \pi/2)$. We equipartition the domain $[r_{\text{max}}^*, r_K^*(E, h_0)]$ into k intervals

of length $\beta = (\pi/2 - r_K^*)/k$. We then consider two comparison problems:

- The Dirichlet problem $P_{DD}^{\text{base}}(r_K^*, \pi/2)$ in conjunction with k Dirichlet problems P_D^{i+} ($i = 1, \dots, k$) arising as follows: They are the problems $P_{DD}^{\text{base}}(r_{\text{max}}^*, r_{\text{max}}^* + \beta), \dots, P_{DD}^{\text{base}}(r_K^* - \beta, r_K^*)$ but with the potential replaced by a constant, which equals the *maximum* of the potential on the interval.
- The mixed problem $P_{ND}^{\text{base}}(r_K^*, \pi/2)$ in conjunction with k Neumann problems P_N^{i-} ($i = 1, \dots, k$) arising as follows: They are the problems $P_{NN}^{\text{base}}(r_{\text{max}}^*, r_{\text{max}}^* + \beta), \dots, P_{NN}^{\text{base}}(r_K^* - \beta, r_K^*)$ but with the potential replaced by a constant, which equals the *minimum* of the potential on the interval.

We can estimate the number of eigenvalues of $P_{DD}^{\text{base}}(r_{\text{max}}^*, \pi/2)$ below a threshold \mathcal{E} by

$$\begin{aligned} \sum_{i=1}^k N_{\leq \mathcal{E}}(P_D^{i+}) + N_{\leq \mathcal{E}}(P_{DD}^{\text{base}}(r_K^*, \pi/2)) &\leq N_{\leq \mathcal{E}}(P_{DD}^{\text{base}}(r_{\text{max}}^*, \pi/2)) \\ &\leq \sum_{i=1}^k N_{\leq \mathcal{E}}(P_N^{i-}) + N_{\leq \mathcal{E}}(P_{ND}^{\text{base}}(r_K^*, \pi/2)). \end{aligned}$$

By our choice of r_K^* , we have $N_{\leq \mathcal{E}}(P_{DD}^{\text{base}}(r_K^*, \pi/2)) = N_{\leq \mathcal{E}}(P_{ND}^{\text{base}}(r_K^*, \pi/2)) = 0$ for $\mathcal{E} = E + \delta$. On the other hand, for each P_D^{i+} and each P_N^{i-} , the number of eigenvalues can be estimated directly (as each problem can be solved explicitly). We have

$$\sum_{i=1}^k N_{\leq \mathcal{E}}(P_D^{i+}) = \sum_{i=1}^k \left\lfloor \frac{\beta}{2\pi h_0} \max(0, \mathcal{E} - V_+^i) \sqrt{\mathcal{E} - V_+^i} \right\rfloor = \sum_{i=1}^k \frac{\beta}{2\pi h_0} \max(0, \mathcal{E} - V_+^i) \sqrt{\mathcal{E} - V_+^i} + \mathcal{O}(k).$$

The estimate for P_N^{i-} is similar, with the potential replaced by V_-^i and the number of eigenvalues in each cell being

$$\left\lfloor \frac{\beta}{2\pi h_0} \max(0, \mathcal{E} - V_-^i) \sqrt{\mathcal{E} - V_-^i} \right\rfloor + 1.$$

To conclude, let us choose the number of cells k such that $k(h_0)$ tends to ∞ as h_0 goes to 0 and moreover $k(h) = o(1/h_0)$. The sums converge as a Riemann sum and the errors are then of order $o(1/h_0)$. Therefore we get

$$\sum_{i=1}^k N_{\leq \mathcal{E}}(P_D^{i+}) \underset{h_0 \rightarrow 0}{\sim} \frac{1}{2\pi h_0} \int_{r_{\text{max}}^*}^{r_K^*} dr^* \sqrt{\mathcal{E} - V(r^*)} \cdot \chi_{V(r^*) \leq \mathcal{E}}. \tag{43}$$

The statement of the lemma then follows from

$$N[E - \delta, E + \delta] = N_{\leq E + \delta} - N_{\leq E - \delta}$$

using the previous formula with $\mathcal{E} = E \pm \delta$. □

4B. Kerr-AdS. In the last section we showed that for any fixed given parameters $M > 0, |a| < l, \alpha < \frac{9}{4}$, the eigenvalue problem (36), $P_{\text{base}}(h_0)u = \kappa \cdot u$ with Dirichlet conditions, admits (lots of) eigenvalues κ in the range $E - \delta \leq \kappa = h_0^2 \omega^2 < E + \delta$, provided h_0 is chosen sufficiently small (i.e., l large).

As an immediate corollary, we obtain the existence of eigenvalues in the desired range for Schwarzschild–AdS, simply by setting $a = 0$; see (37). For the Kerr–AdS case, we still need to relate the above result to the full problem, which we recall is the nonlinear eigenvalue problem (34) given by

$$P(h)u = \kappa \cdot u, \quad \text{with } \kappa = \omega^2 h^2$$

and boundary conditions $u(r_{\max}^*) = 0$ as well as $\int_{r_{\max}^*}^{\pi/2} dr^* [|u'|^2 + r^2 u^2] < \infty$. To achieve this, consider for fixed $|a| < l$, $M > 0$, $\alpha < \frac{9}{4}$ the two-parameter family of linear eigenvalue problems

$$Q_\ell(b^2, \omega^2)u = \Lambda(b^2, \omega^2)u \tag{44}$$

for the operator

$$Q_\ell(b^2, \omega^2)u := -u'' + (V_\sigma \mu_\ell(b^2 a^2 \omega^2)u + (V_{\text{junk}} + V_{\text{mass}} - \omega^2))u, \tag{45}$$

complemented by the above boundary conditions. Here $b^2 \in [0, 1]$ is a dimensionless parameter and $\omega^2 \in \mathbb{R}^+$. Our goal is to show that for $b = 1$ there exists an ω^2 such that the above problem has a zero eigenvalue and, moreover, to suitably control the size of this ω^2 .

By the results of the previous section, we know that for $b = 0$ there exists, for any sufficiently large ℓ , an $\omega_{0,\ell}^2$ (satisfying $E - \delta \leq \omega_{0,\ell}^2 / \mu_\ell(0) \leq E + \delta$) such that $Q_\ell(0, \omega_{0,\ell}^2)$ admits a zero eigenvalue. Moreover, this eigenvalue is nondegenerate, by standard Sturm–Liouville theory. Listing the eigenvalues of $Q_\ell(0, \omega_{0,\ell}^2)$ in ascending order, let us say that it is the n_ℓ -th eigenvalue, which is zero.

The strategy, now, is the following: We will show by an application of the implicit function theorem that for any $b \in [0, 1]$ we can find an $\omega_{b,\ell}^2$ such that the n_ℓ -th eigenvalue of the operator $Q_\ell(b^2, \omega_{b,\ell}^2)$ is zero. As a second step, we will provide a global estimate on the quotient $\omega_{b,\ell}^2 / \mu_\ell(b^2 a^2 \omega_{b,\ell}^2)$. For this last step, an important monotonicity will be exploited.

Lemma 4.6. *Fix parameters $|a| < l$, $M > 0$ and $\alpha < \frac{9}{4}$. Suppose we are given parameters $b_0 \in [0, 1]$ and $\omega_{b_0,\ell}^2 \in \mathbb{R}^+$ such that the n_ℓ -th eigenvalue of $Q_\ell(b_0^2, \omega_{b_0,\ell}^2)$ is zero. Then, there exists an $\varepsilon > 0$ such that:*

1. *For any $b^2 \in (\max(0, b_0^2 - \varepsilon), b_0^2 + \varepsilon)$ one can find an associated $\omega_{b,\ell}^2 \in \mathbb{R}^+$ such that the n_ℓ -th eigenvalue of $Q_\ell(b^2, \omega_{b,\ell}^2)$ is zero,*
2. *$\omega_{b,\ell}^2$ changes differentiably in $b^2 \in (\max(0, b_0^2 - \varepsilon), b_0^2 + \varepsilon)$, and we have the estimate*

$$0 \leq \frac{d\omega_{b,\ell}^2}{db^2} \leq C\omega_{b,\ell}^2,$$

for some constant $C > 0$ which is independent of b_0, ℓ and ε .

3. *The $\varepsilon > 0$ can be taken to be independent of b_0 (but may depend on ℓ).*
4. *$\omega_{b,\ell}^2$ satisfies the estimate*

$$c^{-1} \leq \frac{\omega_{b,\ell}^2}{\ell(\ell + 1)} \leq c,$$

for some $c > 0$ depending only on the parameters a, l, M, α .

Proof. The n -th eigenvalue of $Q_\ell(b^2, \omega_{b,\ell}^2)$, denoted by $\Lambda_n(b^2, \omega^2)$, moves smoothly in the parameters b^2 and ω^2 , and we have the formula

$$\Lambda_n(b^2, \omega^2) = \int_{r_{\max}^*}^{\pi/2} dr^* \psi_n(b^2, \omega^2) Q_\ell(b^2, \omega^2) \psi_n(b^2, \omega^2) \tag{46}$$

for the eigenvalue, provided we normalize the associated eigenfunctions $\psi_n(b^2, \omega^2)$ by

$$\int_{r_{\max}^*}^{\pi/2} dr^* |\psi_n(b^2, \omega^2)|^2 = 1.$$

By assumption, $\Lambda_n(b_0^2, \omega_{b_0,\ell}^2) = 0$. Note that from the normalized condition on the eigenfunctions

$$\int_{r_{\max}^*}^{\pi/2} dr^* \frac{\partial \psi_n}{\partial \omega^2}(b^2, \omega^2) \psi_n(b^2, \omega^2) = 0,$$

which, combined with the eigenvalue equation $\Lambda_n \psi_n = Q_\ell \psi_n$, leads to

$$\int_{r_{\max}^*}^{\pi/2} dr^* \frac{\partial \psi_n}{\partial \omega^2}(b^2, \omega^2) Q_\ell(b^2, \omega^2) \psi_n(b^2, \omega^2) = 0.$$

Thus, differentiating (46), we get

$$\frac{\partial \Lambda_n}{\partial \omega^2} = \int_{r_{\max}^*}^{\pi/2} dr^* \psi_n(b^2, \omega^2) \frac{\partial Q_\ell}{\partial \omega^2}(b^2, \omega^2) \psi_n(b^2, \omega^2),$$

and a similar formula holds replacing $\frac{\partial}{\partial \omega^2}$ with $\frac{\partial}{\partial b^2}$. Using this, we compute

$$\frac{\partial \Lambda_n}{\partial \omega^2}(b_0^2, \omega_{b_0,\ell}^2) = \int_{r_{\max}^*}^{\pi/2} dr^* \psi_n^2(b_0^2, \omega_{b_0,\ell}^2) \cdot \left(V_\sigma \cdot \frac{\partial \mu_\ell}{\partial \omega^2}(b_0^2, \omega_{b_0,\ell}^2) - 1 \right), \tag{47}$$

$$\frac{\partial \Lambda_n}{\partial b^2}(b_0^2, \omega_{b_0,\ell}^2) = \int_{r_{\max}^*}^{\pi/2} dr^* \psi_n^2(b_0^2, \omega_{b_0,\ell}^2) \cdot \left(V_\sigma \cdot \frac{\partial \mu_\ell}{\partial b^2}(b_0^2, \omega_{b_0,\ell}^2) \right). \tag{48}$$

The angular eigenvalue $\mu_\ell(b^2, \omega^2)$ is itself a smooth function of the two parameters b^2 and ω^2 . We have the formula

$$\mu_\ell(b^2, \omega^2) = \int_0^\pi d\theta \sin \theta \phi_\ell(b^2, \omega^2) [P_{\theta,\alpha}(b^2 a^2 \omega^2) + b^2 a^2 \omega^2] \phi_\ell(b^2, \omega^2) \tag{49}$$

for the eigenvalues, provided we normalize the associated eigenfunctions $\phi_\ell(b^2, \omega^2)$ by

$$\int_0^\pi d\theta \sin \theta |\phi_\ell(b^2, \omega^2)|^2.$$

Recalling from (18) that

$$P_{\theta,\alpha}(b^2 a^2 \omega^2) f + b^2 a^2 \omega^2 f = -\frac{1}{\sin \theta} \partial_\theta (\Delta_\theta \sin \theta \partial_\theta f) + \frac{\sin^2 \theta}{\Delta_\theta} b^2 a^2 \omega^2 f + \frac{|\alpha|}{l^2} a^2 \begin{cases} \cos^2 \theta f \\ \sin^2 \theta f \end{cases}, \tag{50}$$

to be read with the upper (lower) line if $\alpha < 0$ ($\alpha \geq 0$), we obtain

$$\frac{\partial \mu_\ell}{\partial \omega^2}(b_0^2, \omega_{b_0, \ell}^2) = b_0^2 l^2 \cdot T_x \quad \text{and} \quad \frac{\partial \mu_\ell}{\partial b^2}(b_0^2, \omega_{b_0, \ell}^2) = \omega_{b_0, \ell}^2 l^2 \cdot T_x, \tag{51}$$

where

$$T_x = \int_0^\pi d\theta \sin \theta \frac{\sin^2 \theta}{\Delta_\theta} \frac{a^2}{l^2} |\phi_\ell(b_0^2, \omega_{b_0, \ell}^2)|^2 \leq \sup_\theta \left(\frac{\sin^2 \theta}{\Delta_\theta} \frac{a^2}{l^2} \right) \int_0^\pi d\theta \sin \theta |\phi_\ell(b_0^2, \omega_{b_0, \ell}^2)|^2 \leq \frac{a^2}{l^2}, \tag{52}$$

where we have used the estimate $\sup_{\theta \in [0, \pi]} (\sin^2 \theta / \Delta_\theta) \leq 1$, which can be easily checked using (11). Going back to (47) and (48), the implicit function theorem allows us to solve for ω^2 as a (smooth) function of b^2 locally near $(b_0^2, \omega_{b_0, \ell}^2)$, provided that the right-hand side of (47) is nonzero. To achieve this, note that

$$1 - V_\sigma \frac{\partial \mu_\ell}{\partial \omega^2} \geq 1 - b_0^2 a^2 V_\sigma \geq 1 - a^2 V_\sigma,$$

using the estimate obtained for $\frac{\partial \mu_\ell}{\partial \omega^2}$ from (51) and (52). On the other hand, using (30),

$$a^2 V_\sigma = \frac{a^2}{l^2} + \frac{\Xi a^2}{r^2 + a^2} = \frac{a^2}{l^2} + \Xi - \frac{\Xi r^2}{r^2 + a^2} < 1 - c_{M, l, a},$$

where $c_{M, l, a}$ is a constant depending only on the parameters M, l, a . It then follows that the right-hand side of (47) is bounded away from zero with the lower bound being independent of b_0 (and ℓ), i.e.,

$$-\frac{\partial \Lambda_n}{\partial \omega^2}(b_0^2, \omega_{b_0, \ell}^2) \geq c_{M, l, a}. \tag{53}$$

This concludes the proof of the first item of the lemma (with ε a priori depending also on b_0^2).

The implicit function theorem also provides a formula for the derivative of the function $\omega^2(b^2)$ just obtained, namely

$$\frac{d\omega_{b, \ell}^2}{db^2}(b^2) = \frac{\frac{\partial \Lambda_n}{\partial b^2}(b^2, \omega_{b, \ell}^2)}{-\frac{\partial \Lambda_n}{\partial \omega^2}(b^2, \omega_{b, \ell}^2)} \quad \text{for any } b^2 \in (\max(0, b_0^2 - \varepsilon), b_0^2 + \varepsilon). \tag{54}$$

We can now repeat the computations following (47) with b_0^2 replaced by any $b^2 \in (\max(0, b_0^2 - \varepsilon), b_0^2 + \varepsilon)$. Proceeding in this way, we obtain the uniform lower bound on the denominator (53) for any such b^2 . The numerator is easily estimated in view of (51), again replacing b_0^2 by b^2 . Note moreover that this derivative has a positive sign since μ_ℓ is an increasing function of b and (48). We thus obtain an estimate of the form

$$0 \leq \frac{d\omega_{b, \ell}^2}{db^2}(b^2) \leq C_{M, l, a} \omega_{b, \ell}^2 \quad \text{for any } b^2 \in (\max(0, b_0^2 - \varepsilon), b_0^2 + \varepsilon), \tag{55}$$

establishing item 2 of the lemma.

We can now apply part 1 of the lemma, first with $b_0^2 = 0$, then with $b_0^2 + \delta$, etc. Integrating the differential inequality (55) from $b_0^2 = 0$ to any point in the interval thus produced will provide the uniform bound $\omega_{b, \ell}^2 \leq C_{M, l, a} \omega_{0, \ell}^2$. Now, $\Lambda_n(b^2, \omega^2)$ is a smooth function on the compact set $[0, 1] \times [0, \omega_{\max, \ell}^2]$, where $\omega_{\max, \ell}^2$ denotes an upper bound for $\omega_{b^2, \ell}^2$ (independent of b !). Since we also have the uniform bound

(53) we obtain uniform control of the errors arising in the implicit function theorem and can conclude that the ε in the implicit function theorem does not depend on b_0 . This is item 3.

To establish item 4, we observe that the identity $\Lambda_n(\omega^2, b^2) = 0$ implies (using the Hardy inequality (40) as well as $V_\sigma \geq 1/l^2$) that $\omega_{b,\ell}^2 \geq (1/2l^2)\mu_\ell(b^2\omega^2) \geq (1/2l^2)\mu_\ell(0) \geq c_{M,l,a}\ell(\ell+1)$ for any $b^2 \in [0, 1]$. Hence the quantity $\omega_{b,\ell}^2$ will stay strictly away from zero and

$$C_{M,l,a} \geq \frac{\omega_{0,\ell}^2}{\ell(\ell+1)} \geq C_{M,l,a}^{-1} \frac{\omega_{b,\ell}^2}{\ell(\ell+1)} \geq c_{M,l,a}, \tag{56}$$

the first inequality following from the analysis of the linear eigenvalue problem (36) in Section 4A, and the second from the uniform estimate (55) on $d\omega_{b,\ell}^2/db^2$. \square

In view of the fact that ε is uniform in b_0^2 , we can apply the implicit function theorem all the way from $b_0 = 0$ to $b = 1$. Note that ε can depend on ℓ . However, for each fixed ℓ it only takes finitely many applications of the implicit function theorem to reach $b = 1$.

To gain quantitative global control beyond (56) on the behavior of $\omega_{b,\ell}^2(b^2)$, let us look at the quotient

$$E_n(b^2) := \frac{\omega_{b,\ell}^2(b^2)}{\mu_\ell(b^2 a^2 \omega_{b,\ell}^2(b^2))} \quad \text{with } b^2 \in [0, 1], \tag{57}$$

which by construction is the n_ℓ -th eigenvalue of the semiclassical operator

$$\tilde{Q}_b(u) = E_n(b^2)u$$

with

$$\tilde{Q}_b(u) = -u''(\mu_\ell(b^2 a^2 \omega_{b,\ell}^2))^{-1} + (V_\sigma + [\mu_\ell(b^2 a^2 \omega_{b,\ell}^2)]^{-1}(V_{\text{junk}} + V_{\text{mass}}))u.$$

Recall that E is an energy level such that $E \in (1/l^2, V_{\text{max}})$ and that, with the notation just introduced, $E_n(0) \in [E - \delta, E + \delta]$ for some small $\delta > 0$.

Lemma 4.7. *For all $\delta' > 0$, there exists an L such that, for all $\ell > L$, we have*

$$\frac{1}{l^2} \leq E_n(1) \leq E + \delta + \delta'. \tag{58}$$

Proof. We first establish the upper bound. Using the Hardy inequality (40) together with (19) (which implies that $\sigma := 1/\mu_\ell(0) - 1/\mu_\ell(b^2 a^2 \omega_{b,\ell}^2) \geq 0$ for any $1 \geq b > 0$), we can estimate

$$\begin{aligned} \int_{r_{\text{max}}^*}^{\pi/2} dr^* [u(\tilde{Q}_0 - \tilde{Q}_b)u] &\geq \sigma \int_{r_{\text{max}}^*}^{\pi/2} dr^* |u|^2 \left(\frac{1}{4l^2} \frac{\Delta_-}{r^2 + a^2} + V_{\text{junk}} + V_{\text{mass}} \right) \\ &\geq \sigma \int_{r_{\text{max}}^*}^{\pi/2} dr^* \left(\left(\frac{9}{4} - \alpha \right) \frac{1}{l^2} \frac{\Delta_-}{r^2 + a^2} |u|^2 - C_{M,l,a} |u|^2 \right), \end{aligned} \tag{59}$$

since V_{junk} is bounded uniformly. In view of $|\sigma| \leq \frac{C_{M,l,a}}{\ell(\ell+1)}$, we conclude that

$$\tilde{Q}_0 \geq \tilde{Q}_b - \frac{C_{M,l,a}}{\ell(\ell+1)}$$

holds for any $1 \geq b > 0$. Hence, by min-max, we infer in particular that (independently of the parameters a, M, ℓ and α)

$$E_n(1) \leq E_n(0) + \delta' \leq E + \delta + \delta', \tag{60}$$

where δ' can be chosen arbitrarily small by choosing ℓ sufficiently large.

For the lower bound, we will establish that $\int dr^* u(\tilde{Q}_b(u)u) > 1/l^2$ holds for any $u \in H^2_{\text{AdS}}$ and ℓ large. To prove the latter, note first that the Hardy inequality (40) reduces the problem to showing that

$$V_\sigma + h^2 \left(\frac{1}{4} \frac{\Delta_-}{r^2 + a^2} + V_{\text{junk}} + V_{\text{mass}} \right) > \frac{1}{l^2}. \tag{61}$$

The square bracket is manifestly positive in a region $[r^*_L, \pi/2)$ for some large r^*_L close to $\pi/2$ (depending only on the parameters $M, |a| < l$ and $\alpha < \frac{9}{4}$). We fix this r^*_L and, in view of $V_\sigma \geq \frac{1}{l^2}$, have established (61) in $[r^*_L, \pi/2)$. In $[r^*_{\text{max}}, r^*_L]$, we have the global estimate

$$V_\sigma \geq \frac{1}{l^2} + \frac{Mr}{(r^2 + a^2)^2}. \tag{62}$$

The second term on the right will dominate the term $h^2 \cdot (V_{\text{junk}})$ pointwise in $[r^*_{\text{max}}, r^*_L]$, provided h is chosen small depending only on M, l, a, α . □

We summarize our results in the following analogue of Proposition 4.3.

Proposition 4.8. *Let $\alpha < \frac{9}{4}, M > 0, |a| < l$ be fixed and $E \in (1/l^2, V_{\text{max}})$ be given. Then, there exists an $L > 0$ such that for any $\ell > L$ there exist an $\omega_\ell^2 \in \mathbb{R}^+$ and a smooth solution u_ℓ of the axisymmetric reduced equation*

$$-u''_\ell + (V_\sigma \cdot \mu_\ell(a^2\omega_\ell^2) + V_{\text{junk}} + V_{\text{mass}})u_\ell = \omega_\ell^2 \cdot u_\ell \tag{63}$$

satisfying $u_\ell(r^*_{\text{max}}) = 0$ and $\int_{r^*}^{\pi/2} (|u'_\ell|^2 + |u_\ell|^2 r^2) dr^* < \infty$. Moreover, the ω_ℓ^2 satisfy the uniform estimates

$$\frac{1}{l^2} < \frac{\omega_\ell^2}{\mu_\ell(a^2\omega_\ell^2)} \leq E + \frac{V_{\text{max}} - E}{2} \quad \text{and} \quad c_{M,l,a} \leq \frac{\omega_\ell^2}{\ell(\ell + 1)} \leq C_{M,l,a}. \tag{64}$$

5. Agmon estimates

In this section, we recall the Agmon estimates. These are (well-known) exponential decay estimates for eigenfunctions for Schrödinger-type operators, in the so-called *forbidden* regions.

5A. Energy inequalities. The Agmon estimates will rely on the following identity.

Lemma 5.1 (energy identity for conjugated operator). *Let $r^*_1 > r^*_0$. Let $h > 0$ and let W, ϕ be smooth real-valued functions on $[r^*_0, r^*_1]$. For all smooth functions u defined on $[r^*_0, r^*_1]$, we have the identity*

$$\begin{aligned} & \int_{r_0^*}^{r_1^*} \left(\left| \frac{d}{dr^*} \left(e^{\frac{\phi}{h}} u \right) \right|^2 + h^{-2} \left(W - \left(\frac{d\phi}{dr^*} \right)^2 \right) e^{2\frac{\phi}{h}} |u|^2 \right) dr^* \\ &= \int_{r_0^*}^{r_1^*} \left(-\frac{d^2 \bar{u}}{dr^{*2}} + h^{-2} W \bar{u} \right) u e^{2\frac{\phi}{h}} dr^* + \int_{x_0}^{x_1} h^{-1} \frac{d\phi}{dr^*} e^{2\frac{\phi}{h}} 2i \Im \left(\bar{u} \frac{du}{dr^*} \right) dr^* \\ & \quad + \left(e^{2\frac{\phi}{h}} \frac{d\bar{u}}{dr^*} u \right) (r_1^*) - \left(e^{2\frac{\phi}{h}} \frac{d\bar{u}}{dr^*} u \right) (r_0^*). \end{aligned}$$

In particular, if u is real-valued and vanishes at r_0^* and r_1^* , then

$$\int_{r_0^*}^{r_1^*} \left(\left| \frac{d}{dr^*} \left(e^{\frac{\phi}{h}} u \right) \right|^2 + h^{-2} \left(W - \left(\frac{d\phi}{dr^*} \right)^2 \right) e^{2\frac{\phi}{h}} |u|^2 \right) dr^* = \int_{r_0^*}^{r_1^*} \left(-\frac{d^2 u}{dr^{*2}} + h^{-2} W u \right) u e^{2\frac{\phi}{h}} dr^*.$$

The same identity holds if instead of assuming that W is smooth on $[r_0^*, r_1^*]$, we assume only that $W|u|^2 \in L^1(r_0^*, r_1^*)$. By density, we may also replace smoothness of u and ϕ by the conditions that $u \in H_0^1[r_0^*, r_1^*]$ and ϕ is a Lipschitz function.

Proof. This follows easily from the computations

$$\begin{aligned} & \int_{r_0^*}^{r_1^*} \left(-\frac{d^2}{dr^{*2}} + h^{-2} W \right) \overline{\left(e^{\frac{\phi}{h}} u \right)} u e^{\frac{\phi}{h}} dr^* \\ &= \int_{r_0^*}^{r_1^*} \left| \frac{d}{dr^*} e^{\frac{\phi}{h}} u \right|^2 + h^{-2} W e^{2\frac{\phi}{h}} |u|^2 dr^* - \left(\frac{d}{dr^*} \left(e^{\frac{\phi}{h}} u \right) u e^{\frac{\phi}{h}} \right) (r_1^*) + \frac{d}{dr^*} \left(e^{\frac{\phi}{h}} u \right) u e^{\frac{\phi}{h}} (r_0^*) \end{aligned}$$

and

$$\begin{aligned} & \int_{r_0^*}^{r_1^*} \left(-\frac{d^2}{dr^{*2}} \right) \overline{\left(e^{\frac{\phi}{h}} u \right)} u e^{\frac{\phi}{h}} dr^* \\ &= \int_{r_0^*}^{r_1^*} -\frac{d}{dr^*} \left(h^{-1} \frac{d\phi}{dr^*} \bar{u} e^{\frac{\phi}{h}} + \frac{d\bar{u}}{dr^*} e^{\frac{\phi}{h}} \right) u e^{\frac{\phi}{h}} dr^* \\ &= \int_{r_0^*}^{r_1^*} \left[-h^{-1} \frac{d\phi}{dr^*} \frac{d\bar{u}}{dr^*} u e^{2\frac{\phi}{h}} - \frac{d^2 \bar{u}}{dr^{*2}} u e^{2\frac{\phi}{h}} \right] dr^* + \int_{r_0^*}^{r_1^*} \left[h^{-1} \frac{d\phi}{dr^*} \frac{du}{dr^*} \bar{u} e^{2\frac{\phi}{h}} + h^{-2} \left(\frac{d\phi}{dr^*} \right)^2 \bar{u} u e^{2\frac{\phi}{h}} \right] dr^* \\ & \quad - h^{-1} \left(\frac{d\phi}{dr^*} |u|^2 e^{2\frac{\phi}{h}} \right) (r_1^*) + h^{-1} \left(\frac{d\phi}{dr^*} |u|^2 e^{2\frac{\phi}{h}} \right) (r_0^*) \\ &= \int_{r_0^*}^{r_1^*} \left(h^{-2} \left(\frac{d\phi}{dr^*} \right)^2 |u|^2 e^{2\frac{\phi}{h}} - \frac{d^2 \bar{u}}{dr^{*2}} u e^{2\frac{\phi}{h}} \right) dr^* + \int_{r_0^*}^{r_1^*} h^{-1} \frac{d\phi}{dr^*} e^{2\frac{\phi}{h}} 2i \Im(\bar{u} u_{r^*}) dr^* \\ & \quad - h^{-1} \left(\frac{d\phi}{dr^*} |u|^2 e^{2\frac{\phi}{h}} \right) (r_1^*) + h^{-1} \left(\frac{d\phi}{dr^*} |u|^2 e^{2\frac{\phi}{h}} \right) (r_0^*). \quad \square \end{aligned}$$

5B. The Agmon distance. We will rely on the Agmon distance to establish our exponential decay estimates.⁹ Given any energy level $\mathcal{E} > 0$ and a potential $V = V(r^*)$ (which may also depend on a

⁹The Agmon distance is actually typically used to obtain *optimal* exponential decay estimates; see for instance [Fournais and Helffer 2010]. For the main purpose of this paper (the construction of quasimodes), we could have used smooth cut-off constructions to prove slightly weaker exponential decay estimates. However, the Agmon distance (despite leading only to Lipschitz cut-offs) has a nice interpretation, which is why we choose to use it here.

parameter h), we define the Agmon distance d between r_1^* and r_2^* as

$$d = d_{(V-\mathcal{E})_+}(r_1^*, r_2^*) = \left| \int_{r_1^*}^{r_2^*} \chi_{\{V \geq \mathcal{E}\}}(r^*) (V(r^*) - \mathcal{E})^{1/2} dr^* \right|,$$

where $\chi_{\{V \geq \mathcal{E}\}}$ is the characteristic function of the set of r^* satisfying $V(r^*) \geq \mathcal{E}$. In other words, d is the distance associated to the Agmon metric $(V - \mathcal{E})_+ dr^*$, where $f_+ = \max(0, f)$ for any function f .

It is easily checked that d satisfies the triangular inequality and that

$$|\nabla_{r^*} d(r^*, r_2^*)|^2 \leq (V - \mathcal{E})_+(r^*).$$

The distance to a set can also be defined as usual. In particular, we define


$$d_{\mathcal{E}}(r^*) := \inf_{r_0^* \in \{\mathcal{E} \geq V\}} d(r^*, r_0^*),$$

which measures the distance to the classical region. We have again

$$|\nabla_{r^*} d_{\mathcal{E}}(r^*)|^2 \leq (V - \mathcal{E})_+(r^*).$$

For a given small $\epsilon \in (0, 1)$ we define the two complementary r^* -regions

$$\begin{aligned} \Omega_{\epsilon}^+ &= \Omega_{\epsilon}^+(\mathcal{E}) := \{r^* \mid V(r^*) > \mathcal{E} + \epsilon\}, \\ \Omega_{\epsilon}^- &= \Omega_{\epsilon}^-(\mathcal{E}) := \{r^* \mid V(r^*) \leq \mathcal{E} + \epsilon\}. \end{aligned}$$



5C. The main estimate. We would like to apply [Lemma 5.1](#) between r_{\max}^* and $\pi/2$ for u , a solution to the eigenvalue problem [\(34\)](#), and for suitable ϕ .

Lemma 5.2. *Let u be a solution to the eigenvalue problem [\(34\)](#); i.e., $\kappa \cdot u = P(h)u$ for some $\kappa = h^2\omega^2$. Define, for any $\epsilon \in (0, 1)$,*

$$\phi_{\kappa, \epsilon} := (1 - \epsilon)d_{\kappa}. \tag{65}$$

Then, for all ϵ sufficiently small, u satisfies

$$\int_{r_{\max}^*}^{\pi/2} h^2 \left| \frac{d}{dr^*} e^{\phi_{\kappa, \epsilon}/h} u \right|^2 dx + \epsilon^2 \int_{\Omega_{\epsilon}^+} e^{2\phi_{\kappa, \epsilon}/h} |u|^2 dr^* \leq D(\kappa + \epsilon) e^{2a(\epsilon)/h} \|u\|_{L^2(r_{\max}^*, \pi/2)}^2, \tag{66}$$

where $a(\epsilon) = \sup_{\Omega_{\epsilon}^-} d_{\kappa}$ and $D > 0$ is a constant depending only on the parameters a, M, l and α .

Remark 5.3. Note

$$a_\kappa(\epsilon) = \sup_{\Omega_\epsilon^-} d_\kappa \rightarrow 0 \quad \text{as } \epsilon \rightarrow 0,$$

uniformly in h (and κ) for h sufficiently small. In view of the exponential weight in the second term on the left, the estimate (66) quantifies that u is exponentially small in the forbidden region, provided we can show a uniform lower bound for $\phi_{\kappa,\epsilon}$ in a suitable subset of Ω_ϵ^+ . This will be achieved in Lemma 5.4.

Proof. Applying Lemma 5.1 between r_{\max}^* and $\pi/2$, we get

$$\int_{r_{\max}^*}^{\pi/2} h^2 \left| \frac{d}{dr^*} e^{\frac{\phi}{h}} u \right|^2 dr^* + \int_{\Omega_\epsilon^+} \left(V - \kappa - \left| \frac{d\phi}{dr^*} \right|^2 \right) e^{2\frac{\phi}{h}} |u|^2 dr^* = \int_{\Omega_\epsilon^-} \left(\kappa - V + \left| \frac{d\phi}{dr^*} \right|^2 \right) e^{2\frac{\phi}{h}} |u|^2 dr^*. \quad (67)$$

In view of our choice $\phi = \phi_{\kappa,\epsilon}$, we have in Ω_ϵ^+ the estimate

$$V - \kappa - \left| \frac{d\phi_{\kappa,\epsilon}}{dr^*} \right|^2 \geq (1 - (1 - \epsilon)^2)(V - \kappa) \geq \epsilon^2 \quad (68)$$

for ϵ sufficiently small, which we will use to estimate the left-hand side of (67).

For the right-hand side of (67), we note that if $V \geq 0$ (which occurs if $\alpha \leq 2$), then we immediately obtain

$$\int_{\Omega_\epsilon^-} \left(\kappa - V + \left| \frac{d\phi_{\kappa,\epsilon}}{dr^*} \right|^2 \right) e^{2\phi_{\kappa,\epsilon}/h} |u|^2 dr^* \leq (\kappa + \epsilon) e^{2a(\epsilon)/h} \|u\|_{L^2(r_{\max}^*, \pi/2)}^2, \quad (69)$$

so that combining (68) and (69) yields (66).

To obtain (66) also in the case $\alpha > 2$ (for which we have $V(r^*) \rightarrow -\infty$ as $r^* \rightarrow \pi/2$), we need once again to appeal to a Hardy-type inequality to absorb the error by the derivative term on the left-hand side of (67). This we do as follows.

Recall that $V = V_\sigma + h^2(V_{\text{junk}} + V_{\text{mass}})$, and the unbounded term is $h^2 V_{\text{mass}} = h^2 \frac{2-\alpha}{l^2} \frac{\Delta_- r^2}{(r^2 + a^2)^2} < 0$ for $\alpha > 2$. Note that

$$\frac{\Delta_- r^2}{(r^2 + a^2)^2} = \frac{\Delta_-}{(r^2 + a^2)} - \frac{\Delta_- a^2}{(r^2 + a^2)^2}.$$

The second term is bounded (and in fact will contribute with the right sign if $\alpha \geq 2$) so its contribution can be treated as before. Thus, we only need to estimate

$$\int_{\Omega_\epsilon^-} \frac{\Delta_-}{(r^2 + a^2)} e^{2\phi_{\kappa,\epsilon}/h} |u|^2 dr^*.$$

By [Holzegel and Smulevici 2013a, Lemma 7.2] (see (40)) we have, for any function v in $H_0^1(r_{\max}^*, \pi/2)$,

$$\int_{R^*}^{\pi/2} \frac{\Delta_-}{(r^2 + a^2)} |v|^2 dr^* \leq 4l^2 \int_{R^*}^{\pi/2} \left| \frac{dv}{dr^*} \right|^2 dr^* \quad \text{for any } R^* \geq r_{\max}^*. \quad (70)$$

Applying the above Hardy inequality to $v = e^{\phi_{\kappa,\epsilon}/h} u$, we obtain that there exists a uniform constant $C > 0$

such that

$$\int_{r_{\max}^*}^{\pi/2} h^2 \left(\frac{9}{4} - \alpha \right) \left| \frac{d}{dr^*} e^{\phi_{\kappa,\epsilon}/h} u \right|^2 dr^* + \int_{\Omega_{\epsilon}^+} \left(V - \kappa - \left| \frac{d\phi_{\kappa,\epsilon}}{dr^*} \right|^2 \right) e^{2\phi_{\kappa,\epsilon}/h} |u|^2 dr^* \leq C(\kappa + \epsilon) e^{2a(\epsilon)/h} \|u\|_{L^2(r_{\max}^*, \pi/2)}^2;$$

i.e., there exists a constant $D > 0$ (which degenerates as $\alpha \rightarrow \frac{9}{4}$) such that

$$\int_{r_{\max}^*}^{\pi/2} h^2 \left| \frac{d}{dr^*} e^{\phi_{\kappa,\epsilon}/h} u \right|^2 dr^* + \int_{\Omega_{\epsilon}^+} \left(V_{\sigma} - \kappa - \left| \frac{d\phi_{\kappa,\epsilon}}{dr^*} \right|^2 \right) e^{2\phi_{\kappa,\epsilon}/h} |u|^2 dr^* \leq D(\kappa + \epsilon) e^{2a(\epsilon)/h} \|u\|_{L^2(r_{\max}^*, \pi/2)}^2. \tag{71}$$

This estimate, when combined with (68), yields again (66) from (67). □

5D. Application of the main estimate. Before we can exploit (66), we need the following lemma, which quantifies the size of the forbidden region for a given energy level.

Lemma 5.4. *Let $E \in (1/l^2, V_{\max})$ and suppose that $\kappa \in (1/l^2, E + \delta]$ for some $\delta > 0$ such that $E + \delta < V_{\max}$. Then there exists $\delta' > 0$ and $C > 0$, both constants independent of h , such that $V_{\sigma} - \kappa > 2C$, in $[r_{\max}, r_{\max} + \delta']$, for all $\kappa \in [E - \delta, E + \delta]$.*

Proof. This is a simple consequence of the continuity of V_{σ} at r_{\max}^* . □

In view of the full potential being $V = V_{\sigma} + h^2(V_{\text{junk}} + V_{\text{mass}})$ we also obtain:

Corollary 5.5. *For h sufficiently small (depending only on M, l and a) we have $V - \kappa > C$ in $[r_{\max}, r_{\max} + \delta']$ for all $\kappa \in [E - \delta, E + \delta]$, with both δ' and C depending only on M, l and a .*

With $E \in (1/l^2, V_{\max})$ given, we now fix $\delta' > 0$ and $C > 0$ as promised by Lemma 5.4. This implies that $\phi_{\kappa,\epsilon} \geq c_{M,l,a}$ in $[r_{\max}, r_{\max} + \delta']$ uniformly in ϵ (the constant $c_{M,l,a}$ being of size $C \cdot \delta'$). Next we fix $\epsilon > 0$ sufficiently small so that, in particular, $a(\epsilon) \leq c_{M,l,a}/2$. We finally conclude from (66) that there exists a $\tilde{C} > 0$ (independent of h) such that


$$\int_{r_{\max}^*}^{r_{\max}^* + \delta'} |u|^2 dr^* \leq \tilde{C} e^{-\tilde{C}/h} \|u\|_{L^2(r_{\max}^*, \pi/2)}^2. \tag{72}$$

Turning to the derivative term on the left of (71), we also have

$$\int_{r_{\max}^*}^{r_{\max}^* + \delta'} h^2 e^{2\phi_{\kappa,\epsilon}/h} \left(\frac{1}{h^2} \left| \frac{d\phi_{\kappa,\epsilon}}{dr^*} \right|^2 |u|^2 + \frac{2}{h} \frac{d\phi_{\kappa,\epsilon}}{dr^*} u \frac{du}{dr^*} + \left| \frac{du}{dr^*} \right|^2 \right) dr^* \lesssim e^{2a(\epsilon)/h} \|u\|_{L^2(r_{\max}^*, \pi/2)}^2.$$

The $|u|^2$ term in the above integral can be ignored since it has the right sign. The cross-term can be absorbed using (72) and $\frac{1}{2}$ of the derivative term. Therefore,

$$\int_{r_{\max}^*}^{r_{\max}^* + \delta'} \left| \frac{du}{dr^*} \right|^2 dr^* \leq \tilde{C} h^{-2} e^{-\tilde{C}/h} \|u\|_{L^2(r_{\max}^*, \pi/2)}^2.$$



Summarizing these decay estimates, we have proven:

Lemma 5.6. *Let $E \in (1/l^2, V_{\max})$ be fixed and let δ be sufficiently small so that $[E - \delta, E + \delta] \subset (1/l^2, V_{\max})$. Then there exist constants $D, \delta' > 0$, depending only on the parameters M, l, a and α , such that the sequence of eigenfunctions $[u_\ell]_{\ell \geq L}^\infty$ arising from Proposition 4.8 satisfies the estimate*

$$\int_{r_{\max}^*}^{r_{\max}^* + \delta'} \left(\left| \frac{du_\ell}{dr^*} \right|^2 + |u_\ell|^2 \right) dr^* \leq D e^{-D/h} \|u_\ell\|_{L^2(r_{\max}^*, \pi/2)}^2,$$

where $h = (\mu_\ell(a^2\omega_\ell^2))^{-1/2}$ and ω_ℓ are as in Proposition 4.8.

We remark that by reusing once again the equation, we can obtain such exponential decay estimates on all higher-order derivatives, with the constants in the above lemma depending on the order of commutation.

6. The construction of quasimodes

By now we have established the existence (Proposition 4.8) of a sequence of functions $[u_\ell]$ such that for each ℓ the corresponding u_ℓ solves

$$\omega_\ell^2 h^2 u_\ell = P(h) u_\ell,$$

where $h = (\mu_\ell(a^2\omega_\ell^2))^{-1/2} \rightarrow 0$ as $\ell \rightarrow \infty$, and such that these u_ℓ obey the estimate of Lemma 5.6 with some constants $D, \delta' > 0$ independent of h (or equivalently ℓ).

Now let χ be a smooth function such that $\chi = 1$ on $[r_{\max}^* + \delta', \pi/2]$ and $\chi = 0$ on $(-\infty, r_{\max}^*]$. We then define $\psi_\ell(t, r, \theta, \tilde{\phi})$ as

$$\psi_\ell(t, r, \theta, \tilde{\phi}) = e^{i\omega_\ell t} \chi(r^*(r))(r^2 + a^2)^{-1/2} u_\ell(r^*(r)) S_{\ell 0}(\theta). \tag{73}$$

Remark 6.1. As defined above, the ψ_ℓ are complex functions but, of course, we could have worked below with $\Re(\psi_\ell)$ or $\Im(\psi_\ell)$.

We now show that the ψ_ℓ satisfy the Klein–Gordon equation up to an exponentially small error.

Lemma 6.2. *For each ℓ and each $k \geq 0$, $\psi_\ell \in CH_{\text{AdS}}^k$. Moreover, there exists $L > 0$ such that we have the following estimates. For all $k \geq 0$, there exists a $C_k > 0$ such that for all $\ell \geq L$ and all $t \geq t_0$,*

$$\left\| \square_g \psi_\ell + \frac{\alpha}{l^2} \psi_\ell \right\|_{H_{\text{AdS}}^k(\Sigma_t)} \leq C_k e^{-C_k \ell} \|\psi_\ell\|_{H_{\text{AdS}}^0(\Sigma_{t_0})}.$$

Moreover, the error $|\square_g \psi_\ell + (\alpha/l^2)\psi_\ell|$ is supported on $[r_{\max}, r_{\max} + \delta']$, for some $\delta' > 0$ independent of ℓ . Finally, all the H^k_{AdS} -norms of ψ_ℓ and of its time derivatives on each Σ_t are constant in t .

Proof. By standard elliptic estimates, any u_ℓ is smooth on $(r^*_{\max}, \pi/2)$. Thus, as far as the regularity of ψ_ℓ is concerned, it is sufficient to check that ψ_ℓ and its derivative decay sufficiently fast near $r = \infty$, which is easy and therefore omitted.

Moreover, in view of our construction, we have $(\square_g + \alpha/l^2)\psi_\ell = 0$ in $[r_+, r_{\max}] \cup [r(r^*_{\max} + \delta'), \infty]$. Hence, the error is supported in a bounded strip in which we have the following naïve estimate: For all (t, r, θ, ϕ) with $r \in [r_{\max}, r_{\max} + \delta']$,

$$\left| \square_g \psi_\ell + \frac{\alpha}{l^2} \psi_\ell \right| \lesssim (\omega^2 |u_\ell| + |u''_\ell| + |u'_\ell| + h^{-2} |u_\ell|) S_{\ell 0}(\theta),$$

which gives the required estimate for $k = 0$ after integration, using the Agmon estimates of the previous section and the equation satisfied by u_ℓ in order to estimate u''_ℓ . For higher k , it suffices to commute the equation and to use the equation for u_ℓ every time two radial derivatives occur, or the equation for $S_{\ell 0}$ every time angular derivatives occur. □

Note that we finally proved [Theorem 1.5](#). Indeed, the ψ_ℓ are of the form claimed in (1) by construction of (73). The estimate on the ω_ℓ in (2) was obtained as part of [Proposition 4.8](#). The error estimate (3) is the statement of [Lemma 6.2](#), while the localization properties (4) and (5) are obvious from (73) itself.

7. Proof of Corollary 1.6

In this section, we prove [Corollary 1.6](#). Given the quasimodes, the proof is standard, but we include it for the paper to be self-contained.

Let us therefore fix a Kerr–AdS spacetime such that the assumption of [Corollary 1.6](#) is satisfied, and also a Klein–Gordon mass $\alpha < \frac{9}{4}$. For convenience, we set $t^*_0 = 0$. Recall also that $t^* = t$ in $r \geq r_{\max}$.

We shall consider solutions ψ to homogeneous and inhomogeneous Klein–Gordon equations with initial data $\psi|_{\Sigma_t}$ and $\partial_t \psi|_{\Sigma_t}$ given on slices of constant t . We shall avoid completely issues regarding the facts that ∂_t is not always timelike and that the coordinate t breaks down at the horizon by considering only axisymmetric data which is compactly supported away from the horizon.

Thus, given any $t, s \in \mathbb{R}$ and given any smooth, axisymmetric initial data set $w = (\bar{\psi}, \overline{T\psi})$ whose support is bounded away from the horizon and which decays sufficiently fast near infinity, we will denote by $P(t, s)w$ the unique solution at time t of the homogeneous problem

$$\left(\square_g + \frac{\alpha}{l^2} \right) \psi = 0, \quad \psi|_{\Sigma_s} = \bar{\psi}, \quad \left. \frac{\partial \psi}{\partial t} \right|_{\Sigma_s} = \overline{T\psi}.$$

Given a smooth axisymmetric function F defined on \mathcal{R} , compactly supported in r away from the horizon and infinity, we can consider the inhomogeneous problem

$$\left(\square_g + \frac{\alpha}{l^2} \right) \psi = F, \quad \psi|_{\Sigma_0} = \bar{\psi}, \quad \left. \frac{\partial \psi}{\partial t} \right|_{\Sigma_0} = \overline{T\psi}.$$

For regular data as above, this problem is well-posed in CH^2_{AdS} and we shall denote its solution by $\psi_F(t)$, suppressing the dependence on r and the angular variables. If the data is axisymmetric, then ψ_F will be axisymmetric and, writing $v(s) = (0, F(s)(g^{tt})^{-1})$, $\psi_F(t)$ is given by the Duhamel formula

$$\psi_F(t) = P(t, 0)w + \int_0^t P(t, s)v(s) ds.$$

We now consider the family of ψ_ℓ given by [Theorem 1.5](#). For each ℓ , ψ_ℓ provides an initial data set

$$w_\ell = \left(\psi_\ell(t = 0), \frac{\partial \psi_\ell}{\partial t}(t = 0) \right)$$

for (1) on the slice $t = 0$. Moreover, each ψ_ℓ satisfies the inhomogeneous Klein–Gordon equation

$$\left(\square_g + \frac{\alpha}{l^2} \right) \psi_\ell = F_\ell,$$

for some F_ℓ satisfying $\|F_\ell\|_{H^{k,-2}_{\text{AdS}}} \leq C_k e^{-C_k \ell} \|\psi_\ell\|_{H^0_{\text{AdS}}}$.

Note that since, from [Lemma 6.2](#), the error F_ℓ is supported away from $r = +\infty$ independently of ℓ , we can safely ignore all powers of r in the following estimates.

Let $\tilde{\psi}_\ell$ denote the solution of the *homogeneous* problem associated with the *same* initial data w_ℓ , i.e., $\tilde{\psi}_\ell = P(t, 0)w_\ell$. From Duhamel’s formula, we then get

$$\begin{aligned} \|\psi_\ell - \tilde{\psi}_\ell\|_{H^1_{\text{AdS}}(\Sigma_t \cap \{r \geq r_{\max}\})} &\leq t \sup_{s \in [0, t]} \|P(t, s)(0, F_\ell)(s)\|_{H^1_{\text{AdS}}(\Sigma_t \cap \{r \geq r_{\max}\})} \\ &\leq tC \|F_\ell\|_{H^{0,-2}(\Sigma_0)} \\ &\leq tC e^{-C\ell} \|\psi_\ell\|_{H^0_{\text{AdS}}(\Sigma_0 \cap \{r \geq r_{\max}\})} \\ &\leq tC e^{-C\ell} \|\psi_\ell\|_{H^1_{\text{AdS}}(\Sigma_0 \cap \{r \geq r_{\max}\})}, \end{aligned} \tag{74}$$

where we use the boundedness statement of [Theorem 1.1](#) to bound $\|P(t, s)(0, F_\ell)(s)\|_{H^0_{\text{AdS}}(\Sigma_t \cap \{r \geq r_{\max}\})}$ in terms of the data, as well as [Lemma 6.2](#). In particular, since the norms of ψ_ℓ are time-invariant, for any $t \leq e^{C\ell}/2C$, the reverse triangle inequality and (74) yield

$$\begin{aligned} \left(\int_{\Sigma_t \cap \{r \geq r_{\max}\}} e_1[\tilde{\psi}_\ell] r^2 dr \sin \theta d\theta d\phi \right)^{\frac{1}{2}} &\geq \|\tilde{\psi}_\ell\|_{H^1_{\text{AdS}}(\Sigma_t \cap \{r \geq r_{\max}\})} \\ &\geq \frac{1}{2} \|\tilde{\psi}_\ell\|_{H^1_{\text{AdS}}(\Sigma_0 \cap \{r \geq r_{\max}\})} \\ &\geq \frac{c}{2\ell} \left[\|\Omega_i \tilde{\psi}_\ell\|_{H^1_{\text{AdS}}(\Sigma_0 \cap \{r \geq r_{\max}\})} + \|\partial_{t^*} \tilde{\psi}_\ell\|_{H^1_{\text{AdS}}(\Sigma_0 \cap \{r \geq r_{\max}\})} \right] \\ &\geq \frac{c}{2\ell} \left(\int_{\Sigma_0} e_2[\tilde{\psi}_\ell] r^2 dr \sin \theta d\theta d\phi \right)^{\frac{1}{2}}. \end{aligned} \tag{75}$$

Here we have used — in the step from the second to the third line — that the data for $\tilde{\psi}_\ell$ is frequency-localized, which allows us to exchange angular and time derivatives with powers of ℓ using the second item of [Theorem 1.5](#), and radial derivatives by angular and time derivatives using the wave equation the

$\tilde{\psi}_\ell$ satisfy. From the third to the fourth line we exploited the fact that the data is localized in $r \geq r_{\max}$. The constant c depends only on the parameters M, l, a and α .

Finally, setting $t_\ell = e^{C\ell}/2C$, we obtain from (75) a family (t_ℓ, ψ_ℓ) such that, for any sufficiently large ℓ , $Q[\tilde{\psi}_\ell](t_\ell) > C > 0$ holds, which proves the corollary.

Appendix: The improved boundedness statement

The boundedness statement at the H^2 -level proven in [Holzegel 2010; 2012] is the estimate (3) for the $\tilde{E}_2[\psi]$ -based energies (see Section 2B). It is remarked in the second of these references that stronger norms can be shown to be uniformly bounded using commutation by angular momentum operators leading to the statement (3). Since the latter statement has been used in this paper and also in [Holzegel and Smulevici 2013a], we provide here a sketch of the proof of this well-known (but absent from the literature) argument. We define the energies

$$E_1[\psi](t^*) = \int_{\Sigma_{t^*}} e_1[\psi](t^*) r^2 \sin \theta \, dr \, d\theta \, d\phi \tag{76}$$

and, with the obvious replacement, $E_2[\psi](t^*)$ and $\tilde{E}_2[\psi](t^*)$. Recall that uniform boundedness for the $\tilde{E}_2[\psi]$ energy is derived by, in addition to using known techniques near the horizon (cf. the red-shift vector field), commuting the Klein–Gordon equation with ∂_t (which yields (2) with ψ replaced by $\partial_t \psi$) followed by elliptic estimates on spacelike slices, which control the H^2_{AdS} norm.

Let us sketch how to prove boundedness (3) for the $E_2[\psi]$ energy. If we commute the Klein–Gordon equation with angular momentum operators we obtain

$$\square_g(\Omega_i \psi) + \frac{\alpha}{l^2}(\Omega_i \psi) = 2^{(\Omega_i)}\pi^{\mu\nu} \cdot \nabla_\mu \nabla_\nu \psi + (2\nabla^\alpha({}^{(\Omega_i)}\pi_{\alpha\mu}) - \nabla_\mu({}^{(\Omega_i)}\pi_\alpha^\alpha))\nabla^\mu \psi, \tag{77}$$

with ${}^{(\Omega_i)}\pi$ the (nonvanishing in Kerr!) deformation tensor of Ω_i . The right-hand side decays suitably in r but not in t . More precisely, in view of the fact that there is no integrated decay estimate available, we cannot close the basic energy estimate on its own. Let us instead commute with localized angular momentum operators $\tilde{\Omega}_i = \chi(r)\Omega_i$, where $\chi(r)$ is equal to 1 for $r \geq 2R$ and equal to 0 for $r \leq R$. Applying the energy estimate for the vector field ∂_{t^*} , we can derive

$$\begin{aligned} & \|\tilde{\Omega}_i \psi_t\|_{H^{0,-2}(\Sigma_{t^*} \cap \{r \geq 2R\})} + \|\tilde{\Omega}_i \psi\|_{H^1_{\text{AdS}}(\Sigma_{t^*} \cap \{r \geq 2R\})} \\ & \leq C_{M,l,a,\alpha} \left(\|\tilde{\Omega}_i \psi_t\|_{H^{0,-2}(\Sigma_{t_0^*})} + \|\tilde{\Omega}_i \psi\|_{H^1_{\text{AdS}}(\Sigma_{t_0^*})} \right) + (\tau_2 - \tau_1) \left(\epsilon \sup_{\tau \in (\tau_1, \tau_2)} E_2[\psi](\tau) + \epsilon \cdot \tilde{E}_2[\psi](0) \right), \end{aligned} \tag{78}$$

where ϵ can be made small by choosing R large. The last term arises from the spacetime error term which decays strongly in r .

The idea is to combine this with an integrated decay estimate for the $\tilde{\Omega}_i \psi$ which loses linearly in τ . Recall that if $\square \Psi + (\alpha/l^2)\Psi = f$, then we have the identity

$$\nabla_a \Psi \nabla^a \psi - \frac{\alpha}{l^2} \Psi^2 = \nabla^\mu (\Psi \nabla_\mu \Psi) - g^{t^*t^*} \nabla_{t^*} \Psi \nabla_{t^*} \Psi - 2g^{t^*b} \nabla_{t^*} \Psi \nabla_b \Psi - f \Psi, \tag{79}$$

where a, b run over r, θ, ϕ only. When integrating this identity (with Ψ replaced by $\tilde{\Omega}_i \psi$ and f the error arising from the commutation in (77)) with the usual spacetime volume we observe that:

- The left-hand side is nonnegative and controls all spatial derivatives after applying the standard Hardy inequality (40).
- The second and third terms on the right are essentially controlled by the $\tilde{E}_2[\psi]$ energy times the length of the time interval $(\tau_2 - \tau_1)$:

$$\begin{aligned}
 g^{t^*t^*} \nabla_{t^*}(\tilde{\Omega}_i \psi) \nabla_{t^*}(\tilde{\Omega}_i \psi) &\sim \frac{1}{r^2} r^2 |\partial_{t^*} \nabla \psi|_g^2 = |\partial_{t^*} \nabla \psi|_g^2 \leq e_1[\partial_{t^*} \psi], \\
 g^{t^*r} \nabla_{t^*}(\tilde{\Omega}_i \psi) \nabla_r(\tilde{\Omega}_i \psi) &\sim \frac{1}{r^3} r^2 (|\partial_{t^*} \nabla \psi|_g^2 + |\partial_r \nabla \psi|_g^2) \leq e_1[\partial_{t^*} \psi], \\
 g^{t^*\phi} \nabla_{t^*}(\tilde{\Omega}_i \psi) \nabla_\phi(\tilde{\Omega}_i \psi) &\sim \frac{1}{r^2} r^2 \left(\frac{1}{\epsilon} |\partial_{t^*} \nabla \psi|_g^2 + \epsilon |\partial_\phi \nabla \psi|_g^2 \right) \leq C_\epsilon \cdot e_1[\partial_{t^*} \psi] + \epsilon \cdot e_2[\psi],
 \end{aligned} \tag{80}$$

and $g^{t^*\theta} = 0$. (We need to borrow an ϵ of $e_2[\psi]$ because the $t^*\phi$ coordinates are not optimal near infinity.) The cross-term $g^{t^*\phi}$ would have much stronger decay in coordinates adapted to the asymptotically AdS end, which would allow us to estimate all terms by the weaker energy $\tilde{E}_2[\psi]$.

- The first term on the right-hand side is a boundary term, which can be estimated by

$$\left| \int_{\mathcal{D}(\tau_1, \tau_2)} \nabla^\mu(\tilde{\Omega}_i \psi \nabla_\mu(\tilde{\Omega}_i \psi)) \right| \leq \sup_{t^*} \int_{\Sigma_{t^*}} \tilde{e}_2[\psi] r^2 \sin \theta \, dr \, d\theta \, d\phi. \tag{81}$$

- The last term in (79) is controlled as previously by the last line in the energy estimate (78).

It follows that integrating (79) furnishes the estimate

$$\begin{aligned}
 \int_{\mathcal{D}(\tau_1, \tau_2) \cap \{r \geq 2R\}} r^2 \sin \theta \, dt^* \, dr \, d\theta \, d\phi \left(r^2 |\partial_r \Omega_i \psi|^2 + |\nabla \Omega_i \psi|^2 \right) \\
 \leq \max(1, \tau_2 - \tau_1) \left(\epsilon \sup_{\tau \in (\tau_1, \tau_2)} E_2[\psi](\tau) + C \cdot \tilde{E}_2[\psi](0) \right). \tag{82}
 \end{aligned}$$

Now note that

$$\tilde{E}_2[\psi](t^*) + \|\Omega_i \psi_t\|_{H_{\text{AdS}}^{0,-2}(\Sigma_{t^*} \cap \{r \leq 2R\})}^2 + \|\Omega_i \psi\|_{H_{\text{AdS}}^1(\Sigma_{t^*} \cap \{r \leq 2R\})}^2 \leq C_R \cdot \tilde{E}_2[\psi](t^*) \leq C_R \cdot \tilde{E}_2[\psi](0) \tag{83}$$

follows right from the boundedness statement for the $\tilde{E}_2[\psi]$ energy and estimating the weights away from infinity. We can integrate (83) in time and add it to (82) which yields (first without the boxed terms)

$$\begin{aligned}
 \boxed{E_2[\psi](\tau_2)} + \int_{\tau_1}^{\tau_2} E_2[\psi](\tau) \, d\tau \\
 \leq \boxed{C_{M,l,a,\alpha} \cdot E_2[\psi](\tau_1)} + \max(1, \tau_2 - \tau_1) \left(\epsilon \sup_{\tau \in (\tau_1, \tau_2)} E_2[\psi](\tau) + C_\epsilon \cdot \tilde{E}_2[\psi](0) \right). \tag{84}
 \end{aligned}$$

The estimate also holds with the boxed terms included, as follows from adding (78) and (83). We claim that (84) implies $E_2[\psi](t^*) \lesssim E_2[\psi](0)$ provided ϵ is sufficiently small depending only on the parameters (the constant $C_{M,l,a,\alpha}$), and leave the verification to the reader.

Remark. An easier proof is available if one is willing to go to H_{AdS}^3 . The Carter operator

$$Q\psi = \Delta_{\mathbb{S}^2}\psi - \partial_\phi^2\psi + (a^2 \sin^2 \theta)\partial_t^2\psi \quad (85)$$

commutes with the wave operator. Since ∂_ϕ^2 and ∂_t^2 trivially commute, we have

$$\begin{aligned} E_1[\Delta_{\mathbb{S}^2}\psi](t^*) &\lesssim E_1[Q\psi](t^*) + E_1[\partial_\phi^2\psi](t^*) + E_1[\partial_t^2\psi](t^*) \\ &\lesssim E_1[Q\psi](0) + E_1[\partial_\phi^2\psi](0) + E_1[\partial_t^2\psi](0), \end{aligned} \quad (86)$$

and we can control all derivatives on S^2 from controlling the Laplacian via elliptic estimates. This yields the desired gain, albeit at the level of three derivatives. This is analogous to commuting with angular momentum operators twice.

Acknowledgments

Theorem 1.5 and **Corollary 1.6** were announced for the Schwarzschild case at the workshop “Mathematical Aspects of General Relativity” in Oberwolfach, Germany, in August 2012 [Holzegel and Smulevici 2012b]. We thank Matthieu Léautaud for showing us the Agmon estimates and Maciej Zworski for pointing out the work of Gannot [2014]. Holzegel acknowledges support through NSF grant DMS-1161607 and thanks the Department of Mathematics at Orsay for its hospitality.

References

- [Anderson 2006] M. T. Anderson, “On the uniqueness and global dynamics of AdS spacetimes”, *Classical Quantum Gravity* **23**:23 (2006), 6935–6953. MR 2008c:83010 Zbl 1107.83015
- [Andersson and Blue 2009] L. Andersson and P. Blue, “Hidden symmetries and decay for the wave equation on the Kerr spacetime”, preprint, 2009. arXiv 0908.2265
- [Aretakis 2012] S. Aretakis, “Decay of axisymmetric solutions of the wave equation on extreme Kerr backgrounds”, *J. Funct. Anal.* **263**:9 (2012), 2770–2831. MR 2967306 Zbl 1254.83021
- [Aretakis 2013] S. Aretakis, “Horizon instability of extremal black holes”, preprint, 2013. arXiv 1206.6598
- [Bachelot 2008] A. Bachelot, “The Dirac system on the Anti-de Sitter universe”, *Comm. Math. Phys.* **283**:1 (2008), 127–167. MR 2009f:83029 Zbl 1153.83013
- [Bachelot 2011] A. Bachelot, “The Klein–Gordon equation in the Anti-de Sitter cosmology”, *J. Math. Pures Appl.* (9) **96**:6 (2011), 527–554. MR 2851681 Zbl 1235.81059
- [Bizon and Rostworowski 2011] P. Bizon and A. Rostworowski, “On weakly turbulent instability of Anti-de Sitter space”, *Phys. Rev. Lett.* **107**:3 (2011), 031102.
- [Bony and Häfner 2008] J.-F. Bony and D. Häfner, “Decay and non-decay of the local energy for the wave equation on the de Sitter–Schwarzschild metric”, *Comm. Math. Phys.* **282**:3 (2008), 697–719. MR 2010h:58041 Zbl 1159.35007
- [Burq 1998] N. Burq, “Décroissance de l’énergie locale de l’équation des ondes pour le problème extérieur et absence de résonance au voisinage du réel”, *Acta Math.* **180**:1 (1998), 1–29. MR 99j:35119 Zbl 0918.35081
- [Dafermos 2006] M. Dafermos, “The black hole stability problem”, Web seminar, Newton Institute, Cambridge, 2006, Available at <http://www.newton.ac.uk/webseminars/pg+ws/2006/gmx/1010/dafermos>.
- [Dafermos and Holzegel 2006] M. Dafermos and G. Holzegel, “Dynamic instability of solitons in 4 + 1-dimensional gravity with negative cosmological constant”, 2006, Available at <https://www.dpmms.cam.ac.uk/~md384/ADSinstability.pdf>.
- [Dafermos and Rodnianski 2005] M. Dafermos and I. Rodnianski, “A proof of Price’s law for the collapse of a self-gravitating scalar field”, *Invent. Math.* **162**:2 (2005), 381–457. MR 2006i:83016 Zbl 1088.83008

- [Dafermos and Rodnianski 2009] M. Dafermos and I. Rodnianski, “The red-shift effect and radiation decay on black hole spacetimes”, *Comm. Pure Appl. Math.* **62**:7 (2009), 859–919. [MR 2011b:83059](#) [Zbl 1169.83008](#)
- [Dafermos and Rodnianski 2010a] M. Dafermos and I. Rodnianski, “Decay for solutions of the wave equation on Kerr exterior spacetimes I-II: The cases $|a| \ll M$ or axisymmetry”, preprint, 2010. [arXiv 1010.5132](#)
- [Dafermos and Rodnianski 2010b] M. Dafermos and I. Rodnianski, “The black hole stability problem for linear scalar perturbations”, preprint, 2010. To appear in *Proceedings of the 12 Marcel Grossmann Meeting*. [arXiv 1010.5137](#)
- [Dafermos and Rodnianski 2013] M. Dafermos and I. Rodnianski, “Lectures on black holes and linear waves”, pp. 97–205 in *Evolution equations* (Zürich, 2008), edited by D. Ellwood et al., Clay Math. Proc. **17**, Amer. Math. Soc., Providence, RI, 2013. [MR 3098640](#) [Zbl 1268.35002](#)
- [Dias et al. 2012] Ó. J. C. Dias, G. T. Horowitz, D. Marolf, and J. E. Santos, “On the nonlinear stability of asymptotically Anti-de Sitter solutions”, *Classical Quantum Gravity* **29**:23 (2012), 235019. [MR 3002881](#) [Zbl 1258.83022](#)
- [Dyatlov 2011] S. Dyatlov, “Quasi-normal modes and exponential energy decay for the Kerr–de Sitter black hole”, *Comm. Math. Phys.* **306**:1 (2011), 119–163. [MR 2012g:58055](#) [Zbl 1223.83029](#)
- [Dyatlov 2012] S. Dyatlov, “Asymptotic distribution of quasi-normal modes for Kerr–de Sitter black holes”, *Ann. Henri Poincaré* **13**:5 (2012), 1101–1166. [MR 2935116](#) [Zbl 1246.83111](#)
- [Festuccia and Liu 2009] G. Festuccia and H. Liu, “A Bohr–Sommerfeld quantization formula for quasinormal frequencies of AdS black holes”, *Adv. Sci. Lett.* **2**:2 (2009), 221–235.
- [Fournais and Helffer 2010] S. Fournais and B. Helffer, *Spectral methods in surface superconductivity*, Progress in Nonlinear Differential Equations and their Applications **77**, Birkhäuser, Boston, 2010. [MR 2011j:35003](#) [Zbl 1256.35001](#)
- [Friedrich 1995] H. Friedrich, “Einstein equations and conformal structure: Existence of Anti-de Sitter-type space-times”, *J. Geom. Phys.* **17**:2 (1995), 125–184. [MR 96j:83008](#) [Zbl 0840.53055](#)
- [Gannot 2014] O. Gannot, “Quasinormal modes for Schwarzschild–AdS black holes: Exponential convergence to the real axis”, *Comm. Math. Phys.* **330**:2 (2014), 771–799. [MR 3223487](#)
- [Hawking and Reall 2000] S. W. Hawking and H. S. Reall, “Charged and rotating AdS black holes and their CFT duals”, *Phys. Rev. D* (3) **61**:2 (2000), 024014. [MR 2001g:83078](#)
- [Holzegel 2010] G. Holzegel, “On the massive wave equation on slowly rotating Kerr–AdS spacetimes”, *Comm. Math. Phys.* **294**:1 (2010), 169–197. [MR 2011d:58070](#) [Zbl 1210.58023](#)
- [Holzegel 2012] G. Holzegel, “Well-posedness for the massive wave equation on asymptotically Anti-de Sitter spacetimes”, *J. Hyperbolic Differ. Equ.* **9**:2 (2012), 239–261. [MR 2928108](#) [Zbl 1250.83025](#)
- [Holzegel and Smulevici 2012a] G. Holzegel and J. Smulevici, “Self-gravitating Klein–Gordon fields in asymptotically Anti-de Sitter spacetimes”, *Ann. Henri Poincaré* **13**:4 (2012), 991–1038. [MR 2913628](#) [Zbl 1253.81065](#)
- [Holzegel and Smulevici 2012b] G. Holzegel and J. Smulevici, “Waves, modes and quasimodes on asymptotically Anti-de Sitter spacetimes”, pp. 2304–2306 in *Mathematical aspects of general relativity*, edited by M. Dafermos et al., Oberwolfach Report **37**, Eur. Math. Soc., Zürich, 2012.
- [Holzegel and Smulevici 2013a] G. Holzegel and J. Smulevici, “Decay properties of Klein–Gordon fields on Kerr–AdS spacetimes”, *Comm. Pure Appl. Math.* **66**:11 (2013), 1751–1802. [MR 3103910](#) [Zbl 1277.83023](#)
- [Holzegel and Smulevici 2013b] G. Holzegel and J. Smulevici, “Stability of Schwarzschild–AdS for the spherically symmetric Einstein–Klein–Gordon system”, *Comm. Math. Phys.* **317**:1 (2013), 205–251. [MR 3010373](#) [Zbl 1259.83012](#)
- [Holzegel and Warnick 2014] G. Holzegel and C. Warnick, “Boundedness of the wave equation for asymptotically Anti-de Sitter black holes”, *J. Funct. Anal.* **266**:4 (2014), 2436–2485. [MR 3150167](#) [Zbl pre06299505](#)
- [Ishibashi and Wald 2004] A. Ishibashi and R. M. Wald, “Dynamics in non-globally-hyperbolic static spacetimes, III: Anti-de Sitter spacetime”, *Classical Quantum Gravity* **21**:12 (2004), 2981–3013. [MR 2005j:83015](#) [Zbl 1064.83006](#)
- [Marzuola et al. 2010] J. Marzuola, J. Metcalfe, D. Tataru, and M. Tohaneanu, “Strichartz estimates on Schwarzschild black hole backgrounds”, *Comm. Math. Phys.* **293**:1 (2010), 37–83. [MR 2010m:58043](#) [Zbl 1202.35327](#)
- [Ralston 1969] J. V. Ralston, “Solutions of the wave equation with localized energy”, *Comm. Pure Appl. Math.* **22** (1969), 807–823. [MR 40 #7642](#) [Zbl 0209.40402](#)

- [Reed and Simon 1978] M. Reed and B. Simon, *Methods of modern mathematical physics, IV: Analysis of operators*, Academic Press, New York, 1978. [MR 58 #12429c](#) [Zbl 0401.47001](#)
- [Sbierski 2013] J. Sbierski, “Characterisation of the energy of Gaussian beams on Lorentzian manifolds - with applications to black hole spacetimes”, preprint, 2013. [arXiv 1311.2477](#)
- [Shlapentokh-Rothman 2014] Y. Shlapentokh-Rothman, “Exponentially growing finite energy solutions for the Klein–Gordon equation on sub-extremal Kerr spacetimes”, *Comm. Math. Phys.* **329**:3 (2014), 859–891. [MR 3212872](#)
- [Tang and Zworski 1998] S.-H. Tang and M. Zworski, “From quasimodes to resonances”, *Math. Res. Lett.* **5**:3 (1998), 261–272. [MR 99i:47088](#) [Zbl 0913.35101](#)
- [Tataru and Tohaneanu 2011] D. Tataru and M. Tohaneanu, “A local energy estimate on Kerr black hole backgrounds”, *Int. Math. Res. Not.* **2011**:2 (2011), 248–292. [MR 2012a:58050](#) [Zbl 1209.83028](#)
- [Vasy 2012] A. Vasy, “The wave equation on asymptotically Anti-de Sitter spaces”, *Anal. PDE* **5**:1 (2012), 81–144. [MR 2957552](#) [Zbl 1277.35014](#)
- [Warnick 2013] C. M. Warnick, “The massive wave equation in asymptotically AdS spacetimes”, *Comm. Math. Phys.* **321**:1 (2013), 85–111. [MR 3089665](#) [Zbl 1271.83021](#)

Received 3 Jun 2013. Revised 24 Jan 2014. Accepted 20 Apr 2014.

GUSTAV HOLZEGEL: g.holzegel@imperial.ac.uk

Department of Mathematics, Imperial College London, South Kensington Campus, London SW7 2AZ, United Kingdom

JACQUES SMULEVICI: jacques.smulevici@math.u-psud.fr

Laboratoire de Mathématiques, Université Paris Sud 11, bât. 425, 91405 Orsay, France

Analysis & PDE

msp.org/apde

EDITORS

EDITOR-IN-CHIEF

Maciej Zworski
zworski@math.berkeley.edu
University of California
Berkeley, USA

BOARD OF EDITORS

Nicolas Burq	Université Paris-Sud 11, France nicolas.burq@math.u-psud.fr	Yuval Peres	University of California, Berkeley, USA peres@stat.berkeley.edu
Sun-Yung Alice Chang	Princeton University, USA chang@math.princeton.edu	Gilles Pisier	Texas A&M University, and Paris 6 pisier@math.tamu.edu
Michael Christ	University of California, Berkeley, USA mchrist@math.berkeley.edu	Tristan Rivière	ETH, Switzerland riviere@math.ethz.ch
Charles Fefferman	Princeton University, USA cf@math.princeton.edu	Igor Rodnianski	Princeton University, USA irod@math.princeton.edu
Ursula Hamenstaedt	Universität Bonn, Germany ursula@math.uni-bonn.de	Wilhelm Schlag	University of Chicago, USA schlag@math.uchicago.edu
Vaughan Jones	U.C. Berkeley & Vanderbilt University vaughan.f.jones@vanderbilt.edu	Sylvia Serfaty	New York University, USA serfaty@cims.nyu.edu
Herbert Koch	Universität Bonn, Germany koch@math.uni-bonn.de	Yum-Tong Siu	Harvard University, USA siu@math.harvard.edu
Izabella Laba	University of British Columbia, Canada ilaba@math.ubc.ca	Terence Tao	University of California, Los Angeles, USA tao@math.ucla.edu
Gilles Lebeau	Université de Nice Sophia Antipolis, France lebeau@unice.fr	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu
László Lempert	Purdue University, USA lempert@math.purdue.edu	Gunther Uhlmann	University of Washington, USA gunther@math.washington.edu
Richard B. Melrose	Massachusetts Institute of Technology, USA rbm@math.mit.edu	András Vasy	Stanford University, USA andras@math.stanford.edu
Frank Merle	Université de Cergy-Pontoise, France Frank.Merle@u-cergy.fr	Dan Virgil Voiculescu	University of California, Berkeley, USA dvv@math.berkeley.edu
William Minicozzi II	Johns Hopkins University, USA minicozz@math.jhu.edu	Steven Zelditch	Northwestern University, USA zelditch@math.northwestern.edu
Werner Müller	Universität Bonn, Germany mueller@math.uni-bonn.de		

PRODUCTION

production@msp.org

Silvio Levy, Scientific Editor

See inside back cover or msp.org/apde for submission instructions.

The subscription price for 2014 is US \$180/year for the electronic version, and \$355/year (+\$50, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to MSP.

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

APDE peer review and production are managed by EditFLOW[®] from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2014 Mathematical Sciences Publishers

ANALYSIS & PDE

Volume 7 No. 5 2014

Resonance widths for the molecular predissociation	1027
ALAIN GRIGIS and ANDRÉ MARTINEZ	
Quasimodes and a lower bound on the uniform energy decay rate for Kerr–AdS spacetimes	1057
GUSTAV HOLZEGEL and JACQUES SMULEVICI	
Cylindrical estimates for hypersurfaces moving by convex curvature functions	1091
BEN ANDREWS and MAT LANGFORD	
Uniform L^p -improving for weighted averages on curves	1109
BETSY STOVALL	
Propagation of singularities for rough metrics	1137
HART F. SMITH	
Well-posedness of Lagrangian flows and continuity equations in metric measure spaces	1179
LUIGI AMBROSIO and DARIO TREVISAN	
Erratum to “Poisson statistics for eigenvalues of continuum random Schrödinger operators”	1235
JEAN-MICHEL COMBES, FRANÇOIS GERMINET and ABEL KLEIN	