

A DIRECT PROOF OF THE BICHTELER–DELLACHERIE THEOREM AND CONNECTIONS TO ARBITRAGE

BY MATHIAS BEIGLBÖCK¹, WALTER SCHACHERMAYER²
AND BEZIRGEN VELIYEV³

University of Vienna

We give an elementary proof of the celebrated Bichteler–Dellacherie theorem which states that the class of stochastic processes S allowing for a useful integration theory consists precisely of those processes which can be written in the form $S = M + A$, where M is a local martingale and A is a finite variation process. In other words, S is a good integrator if and only if it is a semi-martingale.

We obtain this decomposition rather directly from an elementary discrete-time Doob–Meyer decomposition. By passing to convex combinations, we obtain a direct construction of the continuous time decomposition, which then yields the desired decomposition.

As a by-product of our proof, we obtain a characterization of semi-martingales in terms of a variant of *no free lunch*, thus extending a result from [Math. Ann. **300** (1994) 463–520].

1. Introduction. We fix filtered probability space $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{0 \leq t \leq T}, \mathbb{P})$ satisfying the usual conditions. A *simple integrand* is a stochastic process $H = (H_t)_{0 \leq t \leq T}$ of the form

$$(1) \quad H_t = \sum_{j=1}^n f_j \mathbb{1}_{[\tau_{j-1}, \tau_j]}(t), \quad 0 \leq t \leq T,$$

where n is a finite number, $0 = \tau_0 \leq \tau_1 \leq \dots \leq \tau_n = T$ is an increasing sequence of stopping times, and $f_j \in L^\infty(\Omega, \mathcal{F}_{\tau_{j-1}}, \mathbb{P})$.

Denote by $\mathcal{SI} = \mathcal{SI}(\Omega, \mathcal{F}, (\mathcal{F}_t)_{0 \leq t \leq T}, \mathbb{P})$ the vector space (of equivalence classes of) simple integrands.

Received April 2010.

¹Supported by the Austrian Science Fund (FWF) under Grant P21209.

²Supported by the Austrian Science Fund (FWF) under Grant P19456, the Vienna Science and Technology Fund (WWTF) under Grant MA13, the Christian Doppler Research Association (CDG) and the ERC Advanced grant.

³Supported by Austrian Science Fund (FWF) under Grant P19456.

MSC2010 subject classifications. Primary 60G05, 60H05; secondary 91G99.

Key words and phrases. Bichteler–Dellacherie theorem, Doob–Meyer decomposition, arbitrage, Komlós’ lemma.

For every bounded, $\mathcal{F} \otimes \mathfrak{B}_{[0,T]}$ -measurable function $G : \Omega \times [0, T] \rightarrow \mathbb{R}$ we define

$$\|G\|_\infty = \sup_{0 \leq t \leq T} \|G_t\|_{L^\infty(\mathbb{P})}$$

so that $\|\cdot\|_\infty$ is a norm on \mathcal{ST} . Given a (càdlàg, adapted) stochastic process $S = (S_t)_{0 \leq t \leq T}$ we may well define the integration operator $I_S : \mathcal{ST} \rightarrow L^0(\Omega, \mathcal{F}, \mathbb{P})$,

$$(2) \quad I_S \left(\sum_{j=1}^n f_j \mathbb{1}_{\llbracket \tau_{j-1}, \tau_j \rrbracket} \right) = \sum_{j=1}^n f_j (S_{\tau_j} - S_{\tau_{j-1}}) =: (H \cdot S)_T.$$

Note that only a finite Riemann sum is involved in this definition of an integral, so that we do not (yet) encounter any subtleties of limiting procedures.

However, if we seek to extend this operator to a larger class of integrands by approximation with simple integrands, we have to demand that the operator I_S enjoys some minimal continuity properties. A particularly weak requirement is that uniform convergence of a sequence of simple integrands H^n should imply convergence of the integrals $I_S(H^n)$ in probability.

DEFINITION 1.1 (see, e.g., [13], page 52, [16], page 24). A real-valued, càdlàg, adapted process $S = (S_t)_{0 \leq t \leq T}$ is called a *good integrator* if the integration operator

$$I_S : \mathcal{ST} \rightarrow L^0(\Omega, \mathcal{F}, \mathbb{P})$$

is continuous, if we equip \mathcal{ST} with the norm topology induced by $\|\cdot\|_\infty$ and $L^0(\Omega, \mathcal{F}, \mathbb{P})$ with the topology of convergence in probability, respectively.

If S is a good integrator, it is possible to extend the operator I_S to the space of all bounded adapted càglàd processes without major technical difficulties ([13], Chapter 2).

In other words, the above notion ensures, *essentially by definition*, that the procedures involved in extending the integration (2) from finite Riemann sums to their appropriate limits, work out properly for a good integrator S . But of course, the above definition of a good integrator is purely formal, and simply translates the delicacy of the well-definedness of an integral (which involves a limiting procedure) into an equivalent condition.

The achievements of the Strasbourg school of P. A. Meyer and the work of G. Mokobodzki culminated in the theorem of Bichteler–Dellacherie ([13], Theorem 43, Chapter 3, [16], Theorem 16.4), which provides an explicit and practically useful characterization of the set of processes allowing for a powerful stochastic integration theory.

THEOREM 1.2 [3, 4, 6]. *For a real-valued, càdlàg, adapted process $S = (S_t)_{0 \leq t \leq T}$ the following are equivalent:*

1. S is a good integrator.
2. S may be decomposed as $S = M + A$, where $M = (M_t)_{0 \leq t \leq T}$ is a local martingale and $A = (A_t)_{0 \leq t \leq T}$ an adapted process of finite variation.

We then say S is a semi-martingale.

The implication (2) \Rightarrow (1) is a straightforward verification. Indeed it is rather trivial that a càdlàg, adapted process A with a.s. paths of finite variation is a good integrator, where the integral may be defined pathwise. As regards the local martingale part M , the assertion that M is a good integrator, is an extension of Itô's fundamental insight [8, 12] that an L^2 -bounded martingale defines an integration operator which is continuous from $(\mathcal{ST}, \|\cdot\|_\infty)$ to $L^2(\mathbb{P})$.

The remarkable implication is (1) \Rightarrow (2) which provides an explicit characterization of good integrators.

The main aim of this paper is to give a proof of this implication which is inspired by (no) arbitrage-arguments. We note that our argument does not rely on the continuous time Doob–Meyer decomposition nor any change of measure techniques. Instead, we shall construct the desired representation rather directly from a discrete time Doob–Meyer decomposition.⁴ As an important by-product, we also obtain a direct proof of the decomposition of a locally bounded semi-martingale (see Theorem 1.6 below).

Let us now enter the realm of Mathematical Finance.

Here S models the (discounted) price process of some “stock” S , say, a share of company XY. People may trade the stock S : at time t they can hold H_t units of the stock S . Following a trading strategy $H = (H_t)_{0 \leq t \leq T}$, which is assumed to be a predictable process, the accumulated gains or losses up to time t then are given by the random variable

$$(3) \quad (H \cdot S)_t = \int_0^t H_u dS_u, \quad 0 \leq t \leq T.$$

The intuition is that during an infinitesimal interval $[u, u + du]$ the strategy H leads to a random gain/loss $H_u dS_u$. In the case when the predictable process H is a step function, that is, if H is a simple integrand, the stochastic integral (3) becomes a finite Riemann sum. Hence, in this case it is straightforward to justify this infinitesimal reasoning.

The dream of an investor is the possibility of an *arbitrage*.⁵ Roughly speaking, this means the existence of a trading strategy, where you are sure not to lose, but where you possibly may win. Mathematically speaking—and leaving

⁴Thus, our proof is—in spirit—closely related to the proofs of the continuous time Doob–Meyer theorem for super-martingales given by [14] (see also [7, 9]) and [2] (see also [13]).

⁵The basic axiom of mathematical finance is that arbitrages do not exist: there is no such thing as a free lunch!

aside technicalities—this translates into the existence of a predictable process $H = (H_t)_{0 \leq t \leq T}$ such that the negative part $(H \cdot S)_T^-$ of the gains/losses accumulated up to the terminal date T is zero, while the positive part $(H \cdot S)_T^+$ is not. We now give a technical variant of this intuitive idea of an arbitrage.

DEFINITION 1.3 ([5], Section 7). A real-valued, càdlàg, adapted process $S = (S_t)_{0 \leq t \leq T}$ allows for a *free lunch with vanishing risk for simple integrands* if there is a sequence $(H^n)_{n=1}^\infty$ of simple integrands such that, for $n \rightarrow \infty$,

$$(FL) \quad (H^n \cdot S)_T^+ \not\rightarrow 0 \quad \text{in probability,}$$

$$(VR) \quad \sup_{0 \leq t \leq T} \|(H^n \cdot S)_t^-\|_\infty = \|(H^n \cdot S)^-\|_\infty \rightarrow 0,$$

Rephrasing the converse, S therefore admits *no free lunch with vanishing risk* (NFLVR) for simple integrands if for every sequence $(H^n)_{n=1}^\infty \in \mathcal{ST}$ satisfying (VR) we have

$$(NFL) \quad (H^n \cdot S)_T \rightarrow 0 \quad \text{in probability.}$$

The Mathematical Finance context allows for the following interpretation: a *free lunch with vanishing risk for simple integrands* indicates that S allows for a sequence of trading schemes $(H^n)_{n=1}^\infty$, each H^n involving only finitely many rebalancings of the portfolio, such that the losses tend to zero in the sense of (VR), while the terminal gains (FL) remain substantial as n goes to infinity.⁶

It is important to note that the condition (VR) of *vanishing risk* pertains to the maximal losses of the trading strategy H^n during the entire interval $[0, T]$: if the left-hand side of (VR) equals ε_n this implies that, with probability one, the strategy H^n never, that is, for no $t \in [0, T]$, causes an accumulated loss of more than ε_n .

Here is the mathematical theorem which gives the precise relation to the notion of semi-martingales.

THEOREM 1.4 ([5], Theorem 7.2). *Let $(S_t)_{0 \leq t \leq T}$ be a real-valued, càdlàg, locally bounded process based on and adapted to a filtered probability space $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{0 \leq t \leq T}, \mathbb{P})$. If S satisfies the condition of no free lunch with vanishing risk for simple integrands, then S is a semi-martingale.*

In this theorem, we only get one implication, as opposed to the characterization of a semi-martingale in the Bichteler–Dellacherie Theorem 1.2. Indeed, trivial examples show that a semi-martingale $S = (S_t)_{0 \leq t \leq T}$ does not need to satisfy the condition of “no free lunch with vanishing risk for simple integrands.” For example, consider $S_t = t$ and $H_t^n \equiv H_t \equiv 1$, for $0 \leq t \leq T$. Then, for each $n \in \mathbb{N}$, we

⁶The sequence of random variables $((H^n \cdot S)_T^+)_{n=1}^\infty$ does not converge to zero in probability iff there is some $\alpha > 0$ such that $\mathbb{P}[(H^n \cdot S)_T^+ \geq \alpha] \geq \alpha$, for infinitely many $n \in \mathbb{N}$.

have that $(H^n \cdot S)_T = T$ which certainly provides a “free lunch with vanishing risk.”

The proof of Theorem 1.4 which is given in [5], Theorem 7.2, relies on the Bichteler–Dellacherie theorem. The starting point of the present paper was the aim to find a proof of Theorem 1.4 which does not rely on this theorem. Rather we wanted to use Komlos’ lemma and its ramifications which allows in rather general situations to pass to limits of sequences of functions and/or processes by forming convex combinations.

It came as a pleasant surprise that not only it is possible to prove Theorem 1.4 in this way, but that these arguments also yield a constructive proof of the Bichteler–Dellacherie theorem which is based on an intuitive and seemingly naive idea.

To relate the themes of Theorem 1.4 and the Bichteler–Dellacherie Theorem 1.2, we introduce for the context of this paper the following definition which combines the two theorems.

DEFINITION 1.5. Given a process $S = (S_t)_{0 \leq t \leq T}$, we say that S allows for a *free lunch with vanishing risk and little investment*, if there is a sequence $(H^n)_{n=1}^\infty$ of simple integrands as in Definition 1.3 above, satisfying (FL), (VR), and in addition

$$(LI) \quad \lim_{n \rightarrow \infty} \|H^n\|_\infty = 0.$$

The finance interpretation of (LI) above is that, on top of the requirements of *free lunch with vanishing risk*, the holdings H_t^n in the stock S is small when n tends to infinity, a.s. for all $0 \leq t \leq T$. Speaking loosely in more economic terms: S allows for a *free lunch with vanishing risk and little investment* if there are strategies which are almost an arbitrage and which are only involve the holding (or short-selling) of a few stocks.

We may resume our findings in the following theorem which combines and strengthens the content of Theorem 1.4 and the Bichteler–Dellacherie Theorem 1.2.

THEOREM 1.6. For a locally bounded, real valued, càdlàg, adapted process $S = (S_t)_{0 \leq t \leq T}$ the following are equivalent.

1. S admits no free lunch with vanishing risk and little investment, that is, for any sequence $H^n \in \mathcal{SI}$ with $\lim_n \|(H^n \cdot S)^-\|_\infty = \lim_n \|H^n\|_\infty = 0$ we find that $\lim_n (H^n \cdot S)_T^+ = 0$ in probability.
2. S is a semi-martingale in the sense of Theorem 1.2(2).

In the case of general processes S , which need not necessarily be locally bounded, Theorem 1.6 does not hold true any more. Indeed, [5], Example 7.5,

provides an adapted càdlàg process $S = (S_t)_{0 \leq t \leq 1}$ which is not a semi-martingale and for which every simple process $H \in \mathcal{ST}$ satisfying

$$\|(H^n \cdot S)^-\|_\infty \leq 1$$

is constant. Therefore, S trivially verifies the condition of *no free lunch with vanishing risk* (and in particular *no free lunch with vanishing risk and little investment*).

But by appropriately altering the condition (VR) above, we can also formulate a theorem which is analogous to Theorem 1.6 and which implies, in particular, the classical theorem of Bichteler–Dellacherie in its general setting.

THEOREM 1.7. *For a real valued, càdlàg, adapted process $S = (S_t)_{0 \leq t \leq T}$ the following are equivalent.*

1. *For any sequence of simple integrands H^n with $\lim_n \|H^n\|_\infty = 0$ and*

$$\lim_n \sup_{0 \leq t \leq T} ((H^n \cdot S)_t)^- = 0 \quad \text{in probability}$$

we find that $\lim_n (H^n \cdot S)_T^+ = 0$ in probability too.

2. *S is a semi-martingale in the sense of Theorem 1.2(2).*

REMARK 1.8. We also mention the interesting paper [10]. In the setting of a nonnegative process S , it is shown that S is a semi-martingale if and only if it satisfies a weakened NFLVR-condition. Moreover, it is pointed out in [10] what has to be altered to include the case where S is just locally bounded from below. The authors also succeed to avoid the Bichteler–Dellacherie theorem, which is replaced by the continuous time Doob–Meyer theorem.

Finally, we thank Dirk Becherer and Johannes Muhle-Karbe for useful remarks.

2. The idea of the proof. We consider a càdlàg, real-valued, adapted process $S = (S_t)_{0 \leq t \leq T}$. We want to decide whether S is a semi-martingale, and whether S allows for a “free lunch with vanishing risk and little investment.” We only consider the finite horizon case, where from now on we normalize to $T = 1$; the extension to the infinite horizon case is straightforward (see [13, 16]). We also assume that $S_0 = 0$.

We start with the situation when S is locally bounded and shall discuss the general case later.

Noting the fact that being a semi-martingale is a local property, we may and do assume by stopping that S is uniformly bounded, say $\|S\|_\infty \leq 1$ (compare [5]). For $n \in \mathbb{N}$, consider the discrete process $S^n = (S_{j/2^n})_{j=0}^{2^n}$ obtained by sampling the process S at the n th dyadic points. The process S^n may be uniquely decomposed into its Doob–Meyer components

$$S^n = M^n + A^n,$$

where $(M^n_{j/2^n})_{j=0}^{2^n}$ is a martingale and $(A^n_{j/2^n})_{j=0}^{2^n}$ a predictable process with respect to the filtration $(\mathcal{F}_{j/2^n})_{j=0}^{2^n}$: indeed, letting $A^n_0 = 0$ it suffices to define

$$(4) \quad A^n_{j/2^n} - A^n_{(j-1)/2^n} = \mathbb{E}[S_{j/2^n} - S_{(j-1)/2^n} | \mathcal{F}_{(j-1)/2^n}], \quad j = 1, \dots, 2^n,$$

$$(5) \quad M^n_j = S^n_j - A^n_j, \quad j = 0, \dots, 2^n.$$

Observe that we do not have any integrability problems in (4) as S is bounded.

The idea of our proof is—speaking very roughly and somewhat oversimplifying—to distinguish two cases.

Case 1: The processes $(M^n)_{n=1}^\infty$ and $(A^n)_{n=1}^\infty$ remain bounded (in a sense to be clarified below). In this case, we shall apply Komlos type arguments to pass to limiting processes $M = \lim_{n \rightarrow \infty} M^n$ and $A = \lim_{n \rightarrow \infty} A^n$ which then will turn out to be a local martingale and a finite variation process (in continuous time) with respect to the filtration $(\mathcal{F}_t)_{0 \leq t \leq T}$. Hence, in this case, we find that S is a semi-martingale in the sense of Theorem 1.2(2).

Case 2: The processes $(M^n)_{n=1}^\infty$ and/or $(A^n)_{n=1}^\infty$ do not remain bounded. In this case, we shall construct a sequence of simple integrands $(H^k)_{k=1}^\infty = ((H^k_t)_{0 \leq t \leq 1})_{k=1}^\infty$ for the process $(S_t)_{0 \leq t \leq 1}$ which yield a *free lunch with vanishing risk and little investment*. Here is some finance intuition why such a construction should be possible: under the assumption of Case 2 we may find a sequence $\varepsilon_n > 0$ tending to zero such that $(\varepsilon_n M^n)_{n=1}^\infty$ and $(\varepsilon_n A^n)_{n=1}^\infty$ still do not “remain bounded.” Noting that $\|\varepsilon_n M^n + \varepsilon_n A^n\|_\infty = \|\varepsilon_n S\|_\infty \leq \varepsilon_n$ we get an unbounded sequence $(\varepsilon_n M^n)_{n=1}^\infty$ of local martingales and/or an unbounded sequence $(-\varepsilon_n A^n)_{n=1}^\infty$ of predictable processes which are close to each other in the uniform topology. Oversimplifying things slightly, this may be interpreted that *the predictable process $-A^n$ traces closely the martingale M^n* . This ability of nearly reproducing the random movements of the martingale M^n by the predictable movements of the process A^n should enable a smart investor to achieve a free lunch by forming simple integrands $(H^k)_{k=1}^\infty$ which can be chosen such that $\lim_{k \rightarrow \infty} \|H^k\|_\infty = 0$.

Of course, this is only a very crude motivation for the arguments in the next section, where we have to be more precise what we mean by “to remain bounded” (in the sense of quadratic variation or total variation, in the sense of L^∞ , L^2 or L^0 , etc.) and where we have to do a lot of stopping and passing to convex combinations to make the above ideas mathematically rigorous. The crucial issue is that a successful completion of the above program will simultaneously yield proofs for the Bichteler–Dellacherie theorem (Theorem 1.2) as well as for Theorem 1.4. Indeed, it will prove Theorems 1.6 and 1.7 which contain these theorems.

3. Two preliminary decomposition results. In this section, we give two auxiliary results which are somewhat technical but already establish the major portion of our proof of the Bichteler–Dellacherie theorem.

PROPOSITION 3.1. *Let $S = (S_t)_{0 \leq t \leq 1}$ be a càdlàg, adapted process satisfying $S_0 = 0$, $\|S\|_\infty \leq 1$ and no free lunch with vanishing risk and little investment. Denote by A^n and M^n the discrete time Doob decompositions as in (4), respectively, (5).*

For $\varepsilon > 0$, there exist a constant $C > 0$ and a sequence of $\{\frac{j}{2^n}\}_{j=1}^{2^n} \cup \{\infty\}$ -valued stopping times $(\varrho_n)_{n=1}^\infty$ such that $\mathbb{P}(\varrho_n < \infty) < \varepsilon$, and the stopped processes $A^{n,\varrho_n} = (A_{j/2^n \wedge \varrho_n}^n)_{j=1}^{2^n}$, $M^{n,\varrho_n} = (M_{j/2^n \wedge \varrho_n}^n)_{j=1}^{2^n}$ satisfy

$$(6) \quad \text{TV}(A^{n,\varrho_n}) = \sum_{j=1}^{2^n(\varrho_n \wedge 1)} |A_{j/2^n}^n - A_{(j-1)/2^n}^n| \leq C,$$

$$(7) \quad \|M_1^{n,\varrho_n}\|_{L^2(\mathbb{P})}^2 = \|M_{\varrho_n \wedge 1}^n\|_{L^2(\mathbb{P})}^2 \leq C.$$

The proof of Theorem 3.1 will be obtained as a consequence of three lemmas, the first of which is a slightly altered version of [5], Lemma 7.4:

LEMMA 3.2. *Under the assumptions of Proposition 3.1, the sequence of random variables $(QV^n)_{n=1}^\infty$ is bounded in $L^0(\mathbb{P})$, where*

$$QV^n = \sum_{j=1}^{2^n} (S_{j/2^n} - S_{(j-1)/2^n})^2.$$

PROOF. Set $h_t^n = -\sum_{j=1}^{2^n} S_{(j-1)/2^n} \mathbb{1}_{(j-1)/2^n, j/2^n}(t)$. Then $\|h^n\|_\infty \leq 1$ since $\|S\|_\infty \leq 1$. Moreover,

$$(h^n \cdot S)_t = \frac{1}{2} \sum_{j=1}^{2^n} (S_{j/2^n \wedge t} - S_{(j-1)/2^n \wedge t})^2 + \frac{1}{2} (S_0^2 - S_t^2) \geq -\frac{1}{2}.$$

For $t = 1$, we find

$$(8) \quad (h^n \cdot S)_1 = \frac{1}{2} QV^n + \frac{1}{2} (S_0^2 - S_1^2).$$

Since S satisfies no free lunch with vanishing risk and small investments the family $\{(h^n \cdot S)_1 : n \geq 1\}$ is bounded in $L^0(\mathbb{P})$, hence (8) proves the lemma. \square

For $c > 0$ we define, for each $n \geq 1$,

$$\sigma_n(c) = \inf \left\{ \frac{k}{2^n} : \sum_{j=1}^k (S_{j/2^n} - S_{(j-1)/2^n})^2 \geq c - 4 \right\}.$$

The $\{\frac{1}{2^n}, \dots, \frac{2^n-1}{2^n}, 1\} \cup \{+\infty\}$ -valued functions $\sigma_n(c)$ are stopping times with respect to the filtration $(\mathcal{F}_t)_{0 \leq t \leq 1}$. By the preceding lemma, there is a constant $c_1 > 0$

such that, for all $n \geq 1$,

$$(9) \quad \mathbb{P}[\sigma_n(c_1) < \infty] < \frac{\varepsilon}{2}.$$

LEMMA 3.3. *Under the assumptions of Proposition 3.1 and assuming that c_1 satisfies (9) the stopped martingales $M^{n,\sigma_n(c_1)} = (M_{j/2^n}^n)_{j=0}^{2^n(\sigma_n(c_1) \wedge 1)}$ are bounded in $L^2(\Omega, \mathcal{F}, \mathbb{P})$ by*

$$\|M^{n,\sigma_n(c_1)}\|_{L^2(\Omega, \mathcal{F}, \mathbb{P})}^2 \leq c_1.$$

PROOF. Fix $n \in \mathbb{N}$. For any $k \in \{0, 1, \dots, 2^n - 1\}$

$$\begin{aligned} & \mathbb{E}[(S_{k/2^n}^{\sigma_n(c_1)} - S_{(k-1)/2^n}^{\sigma_n(c_1)})^2] \\ &= \mathbb{E}[(M_{k/2^n}^{n,\sigma_n(c_1)} - M_{(k-1)/2^n}^{n,\sigma_n(c_1)} + A_{k/2^n}^{n,\sigma_n(c_1)} - A_{(k-1)/2^n}^{n,\sigma_n(c_1)})^2] \\ &= \mathbb{E}[(M_{k/2^n}^{n,\sigma_n(c_1)} - M_{(k-1)/2^n}^{n,\sigma_n(c_1)})^2] + \mathbb{E}[(A_{k/2^n}^{n,\sigma_n(c_1)} - A_{(k-1)/2^n}^{n,\sigma_n(c_1)})^2] \\ &\geq \mathbb{E}[(M_{k/2^n}^{n,\sigma_n(c_1)} - M_{(k-1)/2^n}^{n,\sigma_n(c_1)})^2]. \end{aligned}$$

Hence, we obtain

$$\begin{aligned} \mathbb{E}[(M_1^{n,\sigma_n(c_1)})^2] &= \mathbb{E}[(M_1^{n,\sigma_n(c_1)})^2] - \mathbb{E}[(M_0^{n,\sigma_n(c_1)})^2] \\ &= \mathbb{E}\left[\sum_{k=1}^{2^n} (M_{k/2^n}^{n,\sigma_n(c_1)} - M_{(k-1)/2^n}^{n,\sigma_n(c_1)})^2\right] \leq c_1. \end{aligned} \quad \square$$

We write $A^{n,\sigma_n(c_1)}$ for the stopped process $(A_{j/2^n}^n)_{j=0}^{2^n(\sigma_n(c_1) \wedge 1)}$ and abbreviate

$$\text{TV}^n = \text{TV}(A^{n,\sigma_n(c_1)}) = \sum_{j=1}^{2^n(\sigma_n(c_1) \wedge 1)} |A_{j/2^n}^n - A_{(j-1)/2^n}^n|.$$

LEMMA 3.4. *Under the assumptions of Proposition 3.1, the sequence $(\text{TV}^n)_{n=1}^\infty$ is bounded in $L^0(\mathbb{P})$.*

PROOF. Suppose not. Then there is some $\alpha > 0$ and for each k some n such that

$$(10) \quad \mathbb{P}[\text{TV}^n \geq k] \geq \alpha.$$

For $n \geq 1$ let $b_{j-1}^n = \text{sign}(A_{j/2^n}^{n,\sigma_n(c_1)} - A_{(j-1)/2^n}^{n,\sigma_n(c_1)})$ and define

$$h^n(t) = \sum_{j=1}^{2^n} b_{j-1}^n \mathbb{1}_{[(j-1)/2^n, j/2^n]}(t).$$

Note that $\|h^n(t)\|_\infty \leq 1$. Also, $(h^n, \sigma_n(c_1) \cdot S)_t = (h^n \cdot S^{\sigma_n(c_1)})_t$ can be estimated from below by

$$\begin{aligned}
 & (h^n \cdot S^{\sigma_n(c_1)})_t \\
 (11) \quad &= \sum_{j \leq t2^n} b_{j-1}^n [S_{j/2^n}^{n, \sigma_n(c_1)} - S_{(j-1)/2^n}^{n, \sigma_n(c_1)}] + b_{\lfloor t2^n \rfloor}^n [S_t^{\sigma_n(c_1)} - S_{\lfloor t2^n \rfloor / 2^n}^{\sigma_n(c_1)}] \\
 &\geq (h^{n, \sigma_n(c_1)} \cdot A)_{\lfloor t2^n \rfloor / 2^n} + (h^{n, \sigma_n(c_1)} \cdot M)_{\lfloor t2^n \rfloor / 2^n} - 2.
 \end{aligned}$$

As $\|(h^{n, \sigma_n(c_1)} \cdot M)\|_{L^2(\mathbb{P})}^2 \leq \|M^{n, \sigma_n(c_1)}\|_{L^2(\mathbb{P})}^2 \leq c_1$ we obtain in particular that

$$(h^{n, \sigma_n(c_1)} \cdot S)_1 = TV^n + (h^{n, \sigma_n(c_1)} \cdot M^n)_1$$

does not remain bounded in $L^0(\mathbb{P})$.

We would like to assure that (11) is uniformly bounded from below, but since we do not have a proper control on the martingale part, we need to perform some further stopping. By Doob’s maximal inequality,

$$\mathbb{E} \left[\sup_{1 \leq j \leq 2^n} ((h^{n, \sigma_n(c_1)} \cdot M)_j)^2 \right] \leq 4c_1.$$

Hence, for c_2 sufficiently large

$$\tau_n = \inf \left\{ \frac{j}{2^n} : |(h^{n, \sigma_n(c_1)} \cdot M)_{j/2^n}| \geq c_2 \right\}$$

satisfies $\mathbb{P}[\tau_n < \infty] = \frac{\alpha}{2}$. We thus obtain that $(h^{n, \tau_n \wedge \sigma_n(c_1)} \cdot S)_t, n \geq 1$ is uniformly bounded from below, whereas $(h^{n, \tau_n \wedge \sigma_n(c_1)} \cdot S)_1 \geq 1$ is still unbounded in $L^0(\mathbb{P})$, hence we obtain a free lunch with vanishing risk and small investments. \square

PROOF OF PROPOSITION 3.1. We define $\tau_n(c) = \inf \{ \frac{k}{2^n} : \sum_{j=1}^k |A_{j/2^n}^n - A_{(j-1)/2^n}^n| \geq c - 2 \}$, so that the stopped processes $A^{n, \tau_n(c)}, n \geq 1$ are bounded in total variation by c . By the preceding lemma, there is a constant $c_2 > 0$ such that, for all $n \geq 1$,

$$\mathbb{P}[\tau_n(c_2) < \infty] < \frac{\varepsilon}{2}.$$

Finally, set $\varrho_n = \sigma_n(c_1) \wedge \tau_n(c_2)$ and $C = c_1 \vee c_2$. \square

In the next step, we extend the decompositions obtained in Proposition 3.1 to continuous time. In the course of the proof, we will use the following technical but elementary lemma.

LEMMA 3.5. *Let $f, g : [0, 1] \rightarrow \mathbb{R}$ be deterministic functions such that f takes only finitely many values and is left continuous, that is, f can be written as*

$$(12) \quad f = \sum_{k=1}^N f(s_k) \mathbb{1}_{]s_{k-1}, s_k]}$$

for appropriate $0 \leq s_0 \leq \dots \leq s_N \leq 1$. For $t \in [0, 1]$ define [in analogy to (2)]

$$(13) \quad (f \cdot g)_t = \sum_{k=1}^{n(t)} f(s_k)(g(s_k) - g(s_{k-1})) + f(t)(g(t) - g(s_{n(t)})),$$

where $n(t) \leq N$ is the maximal number subject to the condition $s_{n(t)} < t$. For any increasing finite sequence $0 \leq t_0 \leq \dots \leq t_m \leq 1$, we then have

$$\sum_{i=1}^m |(f \cdot g)(t_i) - (f \cdot g)(t_{i-1})| \leq 2\text{TV}(f)\|g\|_\infty + \|f\|_\infty \sum_{i=1}^m |g(t_i) - g(t_{i-1})|.$$

PROOF. Define for $i \in \{0, \dots, m-1\}$ numbers $t_{i,0} \leq t_{i,1} \leq \dots \leq t_{i,n}$ satisfying $t_i = t_{i,0}$ and $t_{i+1} = t_{i,n}$ and so that all jumps of f on the interval $[t_0, t_m]$ occur at some $t_{i,j}$. Then we obtain

$$\begin{aligned} & \sum_{i=1}^m |(f \cdot g)(t_i) - (f \cdot g)(t_{i-1})| \\ &= \sum_{i=1}^m \left| \sum_{j=1}^n f(t_{i,j-1})(g(t_{i,j}) - g(t_{i,j-1})) \right| \\ &= \sum_{i=1}^m \left| \left(\sum_{j=1}^n (f(t_{i,j}) - f(t_{i,j-1}))(g(t_{i,n}) - g(t_{i,j-1})) \right) \right. \\ & \qquad \qquad \qquad \left. + f(t_{i,0})(g(t_{i,n}) - g(t_{i,0})) \right| \\ &\leq 2\text{TV}(f) \cdot \|g\|_\infty + \|f\|_\infty \cdot \sum_{i=1}^m |g(t_i) - g(t_{i-1})|. \quad \square \end{aligned}$$

PROPOSITION 3.6. Let $S = (S_t)_{0 \leq t \leq 1}$ be a càdlàg, adapted process satisfying $S_0 = 0$, $\|S\|_\infty \leq 1$ and the condition of no free lunch with vanishing risk and little investment (Definition 1.5).

For $\varepsilon > 0$, there exist a constant $C > 0$, a $[0, 1] \cup \{\infty\}$ -valued stopping time α such that $\mathbb{P}[\alpha < \infty] < \varepsilon$ and a sequence of continuous time càdlàg, adapted processes $\mathcal{A}^n, \mathcal{M}^n$ such that $\mathcal{A}(0) = \mathcal{M}(0) = 0$, (\mathcal{M}^n) is a martingale and

$$(14) \quad \mathcal{A}^{n,\alpha} + \mathcal{M}^{n,\alpha} = S^\alpha,$$

$$(15) \quad \|\mathcal{M}^{n,\alpha}\|_{L^2(\mathbb{P})}^2 \leq C,$$

$$(16) \quad \sum_{j=1}^{2^n} |\mathcal{A}_{j/2^n}^{n,\alpha} - \mathcal{A}_{(j-1)/2^n}^{n,\alpha}| \leq C.$$

PROOF. Fix $\varepsilon > 0$ and let C, M^n, A^n, ϱ_n be as in Proposition 3.1. As a first step, we extend the discrete time martingales to martingales in continuous time [which, by slight abuse of notation, we still denote by $(M_t^n)_{0 \leq t \leq 1}$] by letting

$$M_t^n = \mathbb{E}[M_1^n | \mathcal{F}_t], \quad 0 \leq t \leq 1.$$

We also extend the discrete processes $(A_{j/2^n}^n)_{j=0}^{2^n}$ to processes $(A_t^n)_{0 \leq t \leq 1}$ by letting

$$A_t^n = S_t - M_t^n, \quad 0 \leq t \leq 1.$$

We note that these extended processes $(A_t^n)_{0 \leq t \leq 1}$ need neither be predictable nor do we have a control on their total variation. But we do have a control on the total variation of the restriction of the stopped process $(A_t^{n, \varrho_n})_{0 \leq t \leq 1}$ to the grid $\{0, \frac{1}{2^n}, \dots, \frac{2^n-1}{2^n}, 1\}$, that is,

$$(17) \quad \sum_{j=1}^{2^n(\varrho_n \wedge 1)} |A_{j/2^n}^n - A_{(j-1)/2^n}^n| \leq C.$$

We also note for further use that for $j \in \{0, \frac{1}{2^n}, \dots, \frac{2^n-1}{2^n}, 1\}$ and $t \in]\frac{j-1}{2^n}, \frac{j}{2^n}]$

$$(18) \quad \|A_t^n - A_{j/2^n}^n\|_{L^\infty(\mathbb{P})} \leq 2,$$

which readily follows from the representation

$$\begin{aligned} A_t^n &= S_t - M_t^n = S_t - \mathbb{E}[M_{j/2^n}^n | \mathcal{F}_t] \\ &= S_t - \mathbb{E}[S_{j/2^n} - A_{j/2^n}^n | \mathcal{F}_t] \\ &= A_{j/2^n}^n - (S_t - \mathbb{E}[S_{j/2^n} | \mathcal{F}_t]). \end{aligned}$$

Combining (17) and (18) (and using that A^n is càdlàg), we find that

$$(19) \quad \|A^{n, \varrho_n}\|_\infty \leq C + 2.$$

The most delicate issue in the present proof is to pass—in some sense—to a limit of the stopping times ϱ_n in order to find the desired stopping time α . To this end, we define the left-continuous process $R^n = \mathbb{1}_{[0, \varrho_n \wedge 1]}$ as

$$R_t^n = \begin{cases} 1, & \text{for } 0 \leq t \leq \varrho_n, \\ 0, & \text{for } \varrho_n < t \leq 1, \end{cases}$$

which is a decreasing, simple predictable integrand starting at $R_0^n = 1$ and satisfying $\mathbb{E}[R_1^n] \geq 1 - \varepsilon$. Also note that $A^{n, \varrho_n} = (R^n \cdot A^n)$ and $M^{n, \varrho_n} = (R^n \cdot M^n)$.

We now apply Komlós’ Lemma A.2 to the sequence $(R_1^n)_{n=1}^\infty$ of random variables in $L^\infty(\mathbb{P})$ to pick, for each $n \geq 1$, convex combinations $\mathcal{R}_1^n = \sum_{i=n}^{N_n} \mu_i^n R_1^i$ and a random variable $\mathcal{R}_1 \in L^\infty(\mathbb{P})$ such that $\lim_{n \rightarrow \infty} \mathcal{R}_1^n = \mathcal{R}_1$, convergence taking place almost surely. Note that by dominated convergence $\mathbb{E}[\mathcal{R}_1] \geq 1 - \varepsilon$. Subsequently, we also consider the convex combinations $\mathcal{R}^n = \sum_{i=n}^{N_n} \mu_i^n R^i$ of the

processes $(R^n)_{n=1}^\infty$, and note that convergences of \mathcal{R}_t^n is of course only granted if $t = 1$.

In order to analyze the sequence $(\mathcal{R}^n \cdot S)_{n=1}^\infty$ of simple integrals, we write

$$\begin{aligned}
 \mathcal{R}^n \cdot S &= \left(\sum_{i=n}^{N_n} \mu_i^n R^i \right) \cdot S \\
 (20) \quad &= \sum_{i=n}^{N_n} \mu_i^n (R^i \cdot (M^i + A^i)) \\
 &= \sum_{i=n}^{N_n} \mu_i^n (R^i \cdot M^i) + \sum_{i=n}^{N_n} \mu_i^n (R^i \cdot A^i).
 \end{aligned}$$

Note that for each $n \in \mathbb{N}$, the first term is a martingale bounded in $\|\cdot\|_2$ by $C^{1/2}$, while the total variation of the second term on the grid $\{0, \frac{1}{2^n}, \dots, 1\}$ is bounded by C .

Define the stopping time α_n by

$$\alpha_n = \inf\{t : \mathcal{R}_t^n < \frac{1}{2}\}.$$

As $\mathbb{E}[\mathcal{R}_1^n] \geq 1 - \varepsilon$, we deduce from $\varepsilon \geq \mathbb{E}[1 - \mathcal{R}_1^n] \geq \frac{1}{2}\mathbb{P}[\alpha_n < \infty]$ that $\mathbb{P}[\alpha_n < \infty] \leq 2\varepsilon$. Define \mathcal{T}^n by

$$\mathcal{T}^n = (\mathcal{R}^n)^{-1} \mathbb{1}_{[0, \alpha_n]},$$

so that \mathcal{T}^n is a simple predictable integrand satisfying $\|\mathcal{T}^n\|_\infty \leq 2$. Note that $\mathcal{T}^n \cdot (\mathcal{R}^n \cdot S) = (\mathcal{T}^n \mathcal{R}^n) \cdot S = \mathbb{1}_{[0, \alpha_n]} \cdot S$ equals the stopped process S^{α_n} defined by

$$S_t^{\alpha_n} = \begin{cases} S_t, & \text{for } 0 \leq t \leq \alpha_n \wedge 1, \\ S_{\alpha_n}, & \text{for } \alpha_n \leq t \leq 1. \end{cases}$$

We therefore end up with the following extension of (20):

$$(21) \quad S^{\alpha_n} = \mathcal{T}^n \cdot (\mathcal{R}^n \cdot S) = \underbrace{\mathcal{T}^n \cdot \left(\sum_{i=n}^{N_n} \mu_i (R^i \cdot M^i) \right)}_{=: \mathcal{M}^n} + \underbrace{\mathcal{T}^n \cdot \left(\sum_{i=n}^{N_n} \mu_i (R^i \cdot A^i) \right)}_{=: \mathcal{A}^n}.$$

Next, we establish that \mathcal{M}^n and \mathcal{A}^n are bounded as required. As $\|\mathcal{T}^n\|_\infty \leq 2$,

$$(22) \quad \|\mathcal{M}^n\|_{L^2(\mathbb{P})} \leq 2C^{1/2}.$$

Applying Lemma 3.5 to the functions \mathcal{T}_t^n [which a.s. satisfy $\text{TV}(\mathcal{T}_t^n) \leq 3$, $\|\mathcal{T}_t^n\|_\infty \leq 2$] and $\sum_{i=n}^{N_n} \mu_i (R^i \cdot A^i)_t$ [whose total variation on $\{0, \frac{1}{2^n}, \dots, 1\}$ is bounded by C and which are uniformly bounded by (19)], we obtain

$$(23) \quad \sum_{j=1}^{2^n} |\mathcal{A}_{j/2^n}^n - \mathcal{A}_{(j-1)/2^n}^n| \leq 6 \cdot (C + 2) + 2 \cdot C.$$

We thus have established the boundedness results (15) and (16) claimed in Proposition 3.6, except for the fact that the stopping times α_n still depend on n .

We claim that there exists an increasing sequence $(n_k)_{k=1}^\infty$ such that the stopping time $\alpha = \inf_{k \geq 1} \alpha_{n_k}$ satisfies

$$(24) \quad \mathbb{P}[\alpha < \infty] \leq 4\varepsilon.$$

Combining $\mathbb{E}[\mathcal{R}_1] \geq 1 - \varepsilon$ with the inequality $(1 - a)\mathbb{P}[\mathcal{R}_1 \leq a] \leq \varepsilon$, we have $\mathbb{P}[\mathcal{R}_1 \leq \frac{2}{3}] \leq 3\varepsilon$ and we know that the sequence of random variables $(\mathcal{R}_1^n)_{n=1}^\infty$ converges a.s. to \mathcal{R}_1 . Hence, there is an increasing sequence $(n_k)_{k=1}^\infty$ such that for all $k \geq 1$

$$\mathbb{P}(|\mathcal{R}_1^{n_k} - \mathcal{R}_1| \geq \frac{1}{15}) \leq \varepsilon 2^{-k}.$$

It follows that $\mathbb{P}[\inf_{k \geq 1} \mathcal{R}_1^{n_k} \leq \frac{3}{5}] \leq 4\varepsilon$, which implies (24).

Summing up, we obtain that the sequences $(\mathcal{M}^{n_k})_{k=1}^\infty$ and $(\mathcal{A}^{n_k})_{k=1}^\infty$ satisfy Proposition 3.6. \square

4. Proof of the main theorems. The major work has been done in Proposition 3.6; it is now sufficient to “pass to a limit” to establish Theorem 1.6.

PROOF OF THEOREM 1.6. By stopping S , if necessary, we may assume that $|S|$ is uniformly bounded by 1. We fix $\varepsilon > 0$ and pick C, α and, for each $n \geq 1$, \mathcal{M}^n and \mathcal{A}^n , according to Proposition 3.6. Denote by \mathcal{D} the dyadic numbers in the interval $[0, 1]$. We now apply Komlós lemma (cf. the discussion in the Appendix) to the sequence of $L^2(\mathbb{P})$ -martingales $(\mathcal{M}^{n,\alpha})_{n=1}^\infty$ and, for each $t \in \mathcal{D}$, to the sequence of bounded random variables $(\mathcal{A}_t^{n,\alpha})_{n=1}^\infty$ to find a càdlàg martingale \mathcal{M} , a process $(\mathcal{A}_t)_{t \in \mathcal{D}}$ and for each k some convex weights $\lambda_n^n, \dots, \lambda_{N_n}^n$ such that

$$(25) \quad \lambda_n^n \mathcal{M}_1^{n,\alpha} + \dots + \lambda_{N_n}^n \mathcal{M}_1^{N_n,\alpha} \rightarrow \mathcal{M}$$

and

$$(26) \quad \lambda_n^n \mathcal{A}_t^{n,\alpha} + \dots + \lambda_{N_n}^n \mathcal{A}_t^{N_n,\alpha} \rightarrow \mathcal{A}_t \quad \text{for each } t \in \mathcal{D},$$

where the convergence in (25) and (26) is a.s. as well as in $L^2(\mathbb{P})$.

For the process $(\mathcal{A}_t)_{t \in \mathcal{D}}$, indexed by the dyadic numbers $\mathcal{D} \subseteq [0, 1]$, we then have

$$(27) \quad \sum_{j=1}^N |\mathcal{A}_{t_j} - \mathcal{A}_{t_{j-1}}| \leq C \quad \text{a.s.}$$

for any collection $t_0 \leq t_1 \leq \dots \leq t_N$ in \mathcal{D} . Also, for every $t \in \mathcal{D}$ we have

$$S_t^\alpha = \mathcal{M}_t + \mathcal{A}_t,$$

so that $(\mathcal{A}_t)_{t \in \mathcal{D}}$ is càdlàg on \mathcal{D} . Using (27), we conclude that we may extend $(\mathcal{A}_t)_{t \in \mathcal{D}}$ by right continuity to a process $(\mathcal{A}_t)_{0 \leq t \leq 1}$ via

$$\mathcal{A}_t = \lim_{s \downarrow t, s \in \mathcal{D}} \mathcal{A}_s, \quad 0 \leq t \leq 1,$$

where a.s. the above limit exists for all $t \in [0, 1]$. Using again right continuity, we conclude that

$$S_t^\alpha = \mathcal{M}_t + \mathcal{A}_t \quad \text{for } 0 \leq t \leq 1,$$

hence we obtain desired decomposition on $[[0, \alpha \wedge 1]]$. As $\mathbb{P}(\alpha < \infty) < \varepsilon$ and $\varepsilon > 0$ was chosen arbitrarily, it follows that S is a semi-martingale. \square

The unbounded case can be reduced from Theorem 1.6 rather directly.

PROOF OF THEOREM 1.7. It is sufficient to establish that (1) \Rightarrow (2). We collect the big jumps of S in the process

$$J_t = \sum_{0 < s \leq t} \Delta S_s \mathbb{1}_{\{|\Delta S_s| \geq 1\}},$$

where $\Delta S_t = S_t - S_{t-}$. As S_t is càdlàg, J is of finite total variation. It remains to prove that the locally bounded, càdlàg process $X = S - J$ is a semi-martingale. We want to apply Theorem 1.6. Let (H^n) be a sequence of simple integrands satisfying $\lim_n \|(H^n \cdot X)^-\|_\infty = \lim_n \|H^n\|_\infty = 0$. We claim that (H^n) satisfies

$$(28) \quad \lim_n \sup_{0 \leq t \leq T} ((H^n \cdot S)_t)^- = 0 \quad \text{in probability.}$$

Indeed

$$(29) \quad \sup_{0 \leq t \leq T} ((H^n \cdot S)_t)^- \leq \sup_{0 \leq t \leq T} ((H^n \cdot X)_t)^- + \sup_{0 \leq t \leq T} |(H^n \cdot J)_t|$$

$$(30) \quad \leq \sup_{0 \leq t \leq T} ((H^n \cdot X)_t)^- + (|H^n| \cdot \text{TV}(J))_T.$$

In (30), the first term tends to 0 in probability since it tends to 0 in $\|\cdot\|_{L^\infty(\mathbb{P})}$ and the second term tends to 0 since J is a finite variation process so that $\text{TV}(J)_T = \sum_{0 < s \leq t} |\Delta S_s| \mathbb{1}_{\{|\Delta S_s| \geq 1\}} < \infty$ almost surely.

Having (28) established, assumption (1) of Theorem 1.7 implies that $(H^n \cdot S)_T \rightarrow 0$ in probability. Since we have $(H^n \cdot J)_T \rightarrow 0$ this yields that also $(H^n \cdot X)_T = (H^n \cdot S)_T - (H^n \cdot J)_T$ converges to 0 in probability.

Thus, X satisfies *no free lunch with vanishing risk and little investment* and hence is a semi-martingale by Theorem 1.6. \square

APPENDIX: KOMLÓS’ LEMMA

Komlós’ original result reads as follows.

LEMMA A.1 [11]. *Let $(f_n)_{n \geq 1}$ be a sequence of measurable functions on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$ such that $\sup_{n \geq 1} \|f_n\|_1 < \infty$. Then there exists a subsequence $(\tilde{f}_n)_{n \geq 1}$ such that the functions $\frac{1}{n}(\tilde{f}_1 + \dots + \tilde{f}_n)$ converge almost surely.*

For our purposes, a (much simpler) L^2 -version is sufficient. For the convenience of the reader, we state it together with the short proof.

LEMMA A.2. *Let $(f_n)_{n \geq 1}$ be a sequence of measurable functions on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$ such that $\sup_{n \geq 1} \|f_n\|_2 < \infty$. Then there exist functions $g_n \in \text{conv}(f_n, f_{n+1}, \dots)$ such that $(g_n)_{n \geq 1}$ converges in $\|\cdot\|_{L^2(\mathbb{P})}$ and almost surely.⁷*

PROOF. Let \mathbb{H} be the Hilbert space generated by $(f_n)_{n \geq 1}$. For $n \geq 1$, denote by K_n the (strong) closure of $\text{conv}(f_n, f_{n+1}, \dots)$ which of course coincides with the weak closure by convexity. As the K_n are weakly compact, we may pick $g \in \bigcap_{n=1}^\infty K_n$ and for each n some $g_n \in \text{conv}(f_n, f_{n+1}, \dots)$ such that $g_n \rightarrow g$ in $L^2(\mathbb{P})$. By passing to a subsequence if necessary, we may assume that g_n converges also almost surely. \square

In the course of the paper, we need to apply Lemma A.2 to countably many sequences simultaneously. Just as we may extract a diagonal subsequence of a sequence of refining subsequences, we may do so analogously in the case of convex combinations. Assume that for each $m \geq 1$, $(f_n^m)_{n \geq 1}$ is a sequence of functions bounded in $L^2(\mathbb{P})$. Then we may choose for each n some convex weights $\lambda_n^n, \dots, \lambda_{N_n}^n$ (independent of m) such that

$$(\lambda_n^n f_n^m + \dots + \lambda_{N_n}^n f_{N_n}^m)_{n \geq 1}$$

converges for every $m \in \mathbb{N}$, in $L^2(\mathbb{P})$ and almost surely.

To see this, one first uses Lemma A.2 to find convex weights $\lambda_n^n, \dots, \lambda_{N_n}^n$ such that $(\lambda_n^n f_n^1 + \dots + \lambda_{N_n}^n f_{N_n}^1)_{n \geq 1}$ converges. In the second step, one applies Lemma A.2 to the sequence $(\lambda_n^n f_n^2 + \dots + \lambda_{N_n}^n f_{N_n}^2)_{n \geq 1}$, to obtain convex weights which work for the first two sequences. Repeating this procedure inductively, we obtain sequences of convex weights which work for the first m sequences. Then a standard diagonalization argument proves the assertion.

⁷We may also formulate this result in terms of Cesaro means. (This is due to [1]; see also [15], page 80.) But we do not need this.

REFERENCES

- [1] BANACH, S. and SAKS, S. (1930). Sur la convergence forte dans les champs l^p . *Studia Math.* **2** 51–57.
- [2] BASS, R. F. (1996). The Doob–Meyer decomposition revisited. *Canad. Math. Bull.* **39** 138–150. [MR1390349](#)
- [3] BICHTELER, K. (1979). Stochastic integrators. *Bull. Amer. Math. Soc. (N.S.)* **1** 761–765. [MR0537627](#)
- [4] BICHTELER, K. (1981). Stochastic integration and L^p -theory of semimartingales. *Ann. Probab.* **9** 49–89. [MR0606798](#)
- [5] DELBAEN, F. and SCHACHERMAYER, W. (1994). A general version of the fundamental theorem of asset pricing. *Math. Ann.* **300** 463–520. [MR1304434](#)
- [6] DELLACHERIE, C. (1980). Un survol de la théorie de l’intégrale stochastique. *Stochastic Process. Appl.* **10** 115–144. [MR0587420](#)
- [7] IKEDA, N. and WATANABE, S. (1981). *Stochastic Differential Equations and Diffusion Processes*. North-Holland Mathematical Library **24**. North-Holland, Amsterdam. [MR0637061](#)
- [8] ITÔ, K. (1944). Stochastic integral. *Proc. Imp. Acad. Tokyo* **20** 519–524. [MR0014633](#)
- [9] KARATZAS, I. and SHREVE, S. E. (1991). *Brownian Motion and Stochastic Calculus*, 2nd ed. *Graduate Texts in Mathematics* **113**. Springer, New York. [MR1121940](#)
- [10] KARDARAS, C. and PLATEN, E. (2009). On the semimartingale property of discounted asset-price processes. Boston Univ. and Univ. Technology, Sydney. Available at <http://Arxiv.org/abs/0803.1890>.
- [11] KOMLÓS, J. (1967). A generalization of a problem of Steinhaus. *Acta Math. Acad. Sci. Hungar.* **18** 217–229. [MR0210177](#)
- [12] KUNITA, H. and WATANABE, S. (1967). On square integrable martingales. *Nagoya Math. J.* **30** 209–245. [MR0217856](#)
- [13] PROTTER, P. E. (2004). *Stochastic Integration and Differential Equations: Stochastic Modelling and Applied Probability*, 2nd ed. *Applications of Mathematics (New York)* **21**. Springer, Berlin. [MR2020294](#)
- [14] RAO, K. M. (1969). On decomposition theorems of Meyer. *Math. Scand.* **24** 66–78. [MR0275510](#)
- [15] RIESZ, F. and SZ.-NAGY, B. (1990). *Functional Analysis*. Dover, New York. [MR1068530](#)
- [16] ROGERS, L. C. G. and WILLIAMS, D. (2000). *Diffusions, Markov Processes, and Martingales. Vol. 2. Itô Calculus*. Cambridge Univ. Press, Cambridge. [MR1780932](#)

FAKULTÄT FÜR MATHEMATIK
UNIVERSITÄT WIEN
NORDBERGSTRASSE 15
1090 WIEN
AUSTRIA
E-MAIL: mathias.beiglboeck@univie.ac.at
walter.schachermayer@univie.ac.at
bezirgen.veliyev@univie.ac.at