

## THE SLOW REGIME OF RANDOMLY BIASED WALKS ON TREES<sup>1</sup>

BY YUEYUN HU AND ZHAN SHI

*Université Paris XIII and Université Paris VI*

We are interested in the randomly biased random walk on the supercritical Galton–Watson tree. Our attention is focused on a slow regime when the biased random walk  $(X_n)$  is null recurrent, making a maximal displacement of order of magnitude  $(\log n)^3$  in the first  $n$  steps. We study the localization problem of  $X_n$  and prove that the quenched law of  $X_n$  can be approximated by a certain invariant probability depending on  $n$  and the random environment. As a consequence, we establish that upon the survival of the system,  $\frac{|X_n|}{(\log n)^2}$  converges in law to some non-degenerate limit on  $(0, \infty)$  whose law is explicitly computed.

**1. Introduction.** Let  $\mathbb{T}$  be a supercritical Galton–Watson tree rooted at  $\emptyset$ , so it survives with positive probability. For any pair of vertices  $x$  and  $y$  of  $\mathbb{T}$ , we say  $x \sim y$  if  $x$  is either a child, or the parent, of  $y$ . Let  $\omega := (\omega(x), x \in \mathbb{T})$  be a sequence of vectors; for each vertex  $x \in \mathbb{T}$ ,  $\omega(x) := (\omega(x, y), y \in \mathbb{T})$  is such that  $\omega(x, y) \geq 0$  for all  $y \in \mathbb{T}$  and that  $\sum_{y \in \mathbb{T}} \omega(x, y) = 1$ . We assume that for each pair of vertices  $x$  and  $y$ ,  $\omega(x, y) > 0$  if and only if  $y \sim x$ .

For given  $\omega$ , let  $(X_n, n \geq 0)$  be a random walk on  $\mathbb{T}$  with transition probabilities  $\omega$ , that is, a  $\mathbb{T}$ -valued Markov chain, started at  $X_0 = \emptyset$ , such that

$$P_\omega\{X_{n+1} = y | X_n = x\} = \omega(x, y).$$

For any vertex  $x \in \mathbb{T} \setminus \{\emptyset\}$ , let  $\overleftarrow{x}$  be its parent, and let  $(x^{(1)}, \dots, x^{(N(x))})$  be its children, where  $N(x) \geq 0$  is the number of children of  $x$ . Define  $A(x) := (A_i(x), 1 \leq i \leq N(x))$  by

$$(1.1) \quad A_i(x) := \frac{\omega(x, x^{(i)})}{\omega(x, \overleftarrow{x})}, \quad 1 \leq i \leq N(x).$$

A special example is when  $A_i(x) = \lambda$  for all  $x \in \mathbb{T} \setminus \{\emptyset\}$  and all  $1 \leq i \leq N(x)$ , where  $\lambda$  is a finite and positive constant: the random walk  $(X_n)$  is then the  $\lambda$ -biased random walk on  $\mathbb{T}$  introduced and studied in depth by Lyons [27, 28], Lyons,

---

Received October 2014; revised September 2015.

<sup>1</sup>Supported in part by ANR project MEMEMO2 (2010-BLAN-0125).

*MSC2010 subject classifications.* 60J80, 60G50, 60K37.

*Key words and phrases.* Biased random walk on the Galton–Watson tree, branching random walk, slow movement, local time, convergence in law.

Pemantle and Peres [32, 33]. In particular, if  $A_i(x) = 1, \forall x, \forall i$ , we get the simple random walk on  $\mathbb{T}$ .

It is known that when the transition probabilities are *random*—the resulting random walk  $(X_n)$  is then a random walk in random environment—the walk possesses a regime of *slow movement*. We are interested in this slow movement in this paper.

In the language of Neveu [36],  $(\mathbb{T}, \omega)$  is a marked tree. Note that  $A(x), x \in \mathbb{T} \setminus \{\emptyset\}$  depends entirely on the marked tree. We assume, from now on, that  $A(x), x \in \mathbb{T} \setminus \{\emptyset\}$ , are i.i.d., and write  $A = (A_1, \dots, A_N)$  for a generic random vector having the law of  $A(x)$  (for any  $x \in \mathbb{T} \setminus \{\emptyset\}$ ). We mention that the dimension  $N \geq 0$  of  $A$  is random, and is governed by the law of reproduction of  $\mathbb{T}$ . We use  $\mathbf{P}$  to denote the probability with respect to the environment, and  $\mathbb{P} := \mathbf{P} \otimes P_\omega$  the annealed probability, that is,  $\mathbb{P}(\cdot) := \int P_\omega(\cdot) \mathbf{P}(d\omega)$ .

Throughout the paper, we assume

$$(1.2) \quad \mathbf{E}\left(\sum_{i=1}^N A_i\right) = 1, \quad \mathbf{E}\left(\sum_{i=1}^N A_i \log A_i\right) = 0.$$

In the language of branching random walks (see Section 2), (1.2) refers to the “boundary case;” in this case, the biased walks produce some unusual phenomena that have still been beyond good understanding. We also assume the following integrability condition: there exists  $\delta > 0$  such that

$$(1.3) \quad \mathbf{E}\left(\sum_{i=1}^N A_i^{1+\delta}\right) + \mathbf{E}\left(\sum_{i=1}^N A_i^{-\delta}\right) + \mathbf{E}(N^{1+\delta}) < \infty.$$

Lyons and Pemantle [30] established a recurrence vs. transience criterion for random walks on general trees; applied to the special setting of Galton–Watson trees, it says that (1.2) ensures that the biased walk  $(X_n)$  is  $\mathbb{P}$ -a.s. recurrent. Menshikov and Petritis [35] gave another proof of the recurrence by means of Mandelbrot’s multiplicative cascades, assuming some additional integrability condition. The proofs of the recurrence in both [30] and [35] required an extra exchangeability assumption on  $(A_1, \dots, A_N)$ , which turned out to be superfluous, and was removed by Faraud [15], who furthermore proved that  $(X_n)$  is null recurrent under (1.2).

Introduced by Lyons and Pemantle [30] as an extension of deterministically biased walks studied in Lyons [27, 28], randomly biased walks on trees have received much research interest. Deep results were obtained by Lyons, Pemantle and Peres [32] and [33], who also raised further open problems. Often motivated by these results and problems, both the transient regimes [1, 2] and the recurrent regimes [6, 7, 15, 16, 18, 19] have been under intensive study for these walks. For a general account of biased walks on trees, we refer to [26, 37] and [40].

We add a special vertex, denoted by  $\overleftarrow{\emptyset}$ , which is the parent of  $\emptyset$ , and assume that  $(\omega(\overleftarrow{\emptyset}, y), |y| = 1 \text{ or } y = \overleftarrow{\emptyset})$  is independent of other random vectors

$(\omega(x, y), y \sim x)$  for  $x \in \mathbb{T} \setminus \{\emptyset\}$ , having the same distribution as any of these random vectors; whenever the biased walk  $(X_i)$  hits  $\overleftarrow{\emptyset}$ , it comes back to  $\emptyset$  in the next step. [However,  $\overleftarrow{\emptyset}$  is not considered as a vertex of  $\mathbb{T}$ ; so, for example,  $\sum_{x \in \mathbb{T}} f(x)$  does not contain the term  $f(\overleftarrow{\emptyset})$ .] This makes the presentation of our model more pleasant, since the family of i.i.d. random vectors  $A(x)$  also includes the element  $A(\emptyset)$  from now on.

It was proved in [16] that under (1.2) and (1.3), almost surely upon the survival of the system,

$$\lim_{n \rightarrow \infty} \frac{1}{(\log n)^3} \max_{0 \leq i \leq n} |X_i| = \frac{8}{3\pi^2\sigma^2},$$

where

$$(1.4) \quad \sigma^2 := \mathbf{E} \left( \sum_{i=1}^N A_i (\log A_i)^2 \right) \in (0, \infty).$$

We are interested in the typical size of  $|X_n|$ ; a natural question is to find a deterministic sequence  $a_n \rightarrow \infty$  such that  $\frac{|X_n|}{a_n}$  converges in law to some non-degenerate limit. In dimension 1 [which would be an informal analogue of the case  $N(x) = 1$  for all  $x$ ], the slow movement was discovered by Sinai [41] who showed that  $\frac{X_n}{(\log n)^2}$  converges weakly to a non-degenerate limit under the annealed measure. More precisely, Sinai [41] developed the seminal “method of valley” to localise the walk around the bottom of a certain Brownian valley with high probability. This method, however, seems hopeless to be directly adapted to the biased walk on trees. Observe that in terms of the invariant measure, we can interpret Sinai’s method of valley as the approximation of the law of the walk by a certain invariant measure whose mass is concentrated at the neighbourhood of the bottom. Our main result, stated as Theorem 2.1 below, asserts that upon the survival of the system, the (quenched) finitely-dimensional distribution of the biased walk can be approximated by the product measure of some invariant probability measures. A consequence of this result is that under (1.2) and (1.3), for all  $x > 0$ ,

$$\lim_{n \rightarrow \infty} \mathbb{P} \left( \frac{\sigma^2 |X_n|}{(\log n)^2} \leq x \mid \text{survival} \right) = \int_0^x \frac{1}{(2\pi y)^{1/2}} \mathbf{P} \left( \eta \leq \frac{1}{y^{1/2}} \right) dy,$$

where  $\sigma$  is the constant in (1.4), and  $\eta := \sup_{s \in [0, 1]} [\overline{m}(s) - m(s)]$ . Here,  $(m(s), s \in [0, 1])$  is a standard Brownian meander,<sup>2</sup> and  $\overline{m}(s) := \sup_{u \in [0, s]} m(u)$ .

We mention that  $\int_0^\infty \frac{1}{(2\pi y)^{1/2}} \mathbf{P}(\eta \leq \frac{1}{y^{1/2}}) dy = 1$  because  $\mathbf{E}(\frac{1}{\eta}) = (\frac{\pi}{2})^{1/2}$ , see [21].

In the next section, we give a precise statement of Theorem 2.1, as well as an outline of its proof.

---

<sup>2</sup>Recall that the standard Brownian meander can be realised as follows:  $m(s) := \frac{|B(g+s(1-g))|}{(1-g)^{1/2}}$ ,  $s \in [0, 1]$ , where  $(B(t), t \in [0, 1])$  is a standard Brownian motion, with  $g := \sup\{t \leq 1 : B(t) = 0\}$ .

**2. Random potential, and statement of results.** The movement of the biased random walk  $(X_n)$  depends strongly on the random environment  $\omega$ . It turns out to be more convenient to quantify the influence of the random environment via the random *potential*, which we define by  $V(\emptyset) := 0$  and

$$(2.1) \quad V(x) := - \sum_{y \in \llbracket \emptyset, x \rrbracket} \log \frac{\omega(\overleftarrow{y}, y)}{\omega(y, \overleftarrow{y})}, \quad x \in \mathbb{T} \setminus \{\emptyset\},$$

where  $\overleftarrow{y}$  is the parent of  $\overleftarrow{y}$ , and  $\llbracket \emptyset, x \rrbracket := \llbracket \emptyset, x \rrbracket \setminus \{\emptyset\}$ , with  $\llbracket \emptyset, x \rrbracket$  denoting the set of vertices (including  $x$  and  $\emptyset$ ) on the unique shortest path connecting  $\emptyset$  to  $x$ . Throughout the paper, we use  $x_i$  (for  $0 \leq i \leq |x|$ ) to denote the ancestor of  $x$  in the  $i$ th generation; in particular,  $x_0 = \emptyset$  and  $x_{|x|} = x$ . As such, the potential  $V$  in (2.1) can also be written as

$$V(x) = - \sum_{i=0}^{|x|-1} \log \frac{\omega(x_i, x_{i+1})}{\omega(x_i, x_{i-1})}, \quad x \in \mathbb{T} \setminus \{\emptyset\}, \quad (x_{-1} := \overleftarrow{\emptyset}).$$

The random potential process  $(V(x), x \in \mathbb{T})$  is a branching random walk, in the usual sense of Biggins [9]. There exists an obvious bijection between the random environment  $\omega$  and the random potential  $V$ .

In terms of the random potential, assumptions (1.2) and (1.3) become, respectively,

$$(2.2) \quad \mathbf{E} \left( \sum_{x:|x|=1} e^{-V(x)} \right) = 1, \quad \mathbf{E} \left( \sum_{x:|x|=1} V(x) e^{-V(x)} \right) = 0,$$

and

$$(2.3) \quad \mathbf{E} \left( \sum_{x:|x|=1} e^{-(1+\delta)V(x)} \right) + \mathbf{E} \left( \sum_{x:|x|=1} e^{\delta V(x)} \right) + \mathbf{E} \left[ \left( \sum_{x:|x|=1} 1 \right)^{1+\delta} \right] < \infty.$$

We refer from now on to (2.2) or (2.3) instead of to (1.2) or (1.3). In the language of branching random walks, (2.2) corresponds to the “boundary case” (Biggins and Kyprianou [12]). The branching random walk in this case is known, under some additional integrability assumptions, to have some highly non-trivial universality properties.

We are often interested in properties upon the system’s non-extinction, so let us introduce

$$\mathbf{P}^*(\cdot) := \mathbf{P}(\cdot | \text{non-extinction}),$$

$$\mathbb{P}^*(\cdot) := \mathbb{P}(\cdot | \text{non-extinction}).$$

Let us define a symmetrised version of the potential:

$$(2.4) \quad U(x) := V(x) - \log \left( \frac{1}{\omega(x, \overleftarrow{x})} \right), \quad x \in \mathbb{T}.$$

We call  $U$  the *symmetrised potential*, and use frequently the following relation between  $U$  and  $V$ :

$$(2.5) \quad e^{-U(x)} = \frac{1}{\omega(x, \overset{\leftarrow}{x})} e^{-V(x)} = e^{-V(x)} + \sum_{y \in \mathbb{T}: \overset{\leftarrow}{y}=x} e^{-V(y)}, \quad x \in \mathbb{T}.$$

We now introduce a pair of fundamental martingales associated with the potential  $V$ . Assumption (2.2) immediately implies that  $(W_n, n \geq 0)$  and  $(D_n, n \geq 0)$  are martingales under  $\mathbf{P}$ , where

$$(2.6) \quad W_n := \sum_{x:|x|=n} e^{-V(x)},$$

$$(2.7) \quad D_n := \sum_{x:|x|=n} V(x)e^{-V(x)}, \quad n \geq 0.$$

In the literature,  $(W_n)$  is referred to as an *additive martingale*, and  $(D_n)$  a *derivative martingale*. Since  $(W_n)$  is a non-negative martingale, it converges  $\mathbf{P}$ -a.s. to a finite limit; under assumption (2.2), this limit is known (Biggins [10], Lyons [29]) to be 0:

$$(2.8) \quad W_n \rightarrow 0, \quad \mathbf{P}^*\text{-a.s.}$$

[We will see in (4.2) the rate of convergence.] In view of (2.5), this yields

$$(2.9) \quad \inf_{x:|x|=n} U(x) \rightarrow \infty, \quad \mathbf{P}^*\text{-a.s.}$$

For the derivative martingale  $(D_n)$ , it is known (Biggins and Kyprianou [11], Aïdékon [4]) that (2.3) is “slightly more than” sufficient to ensure that  $D_n$  converges  $\mathbf{P}$ -a.s. to a limit, denoted by  $D_\infty$ , and that

$$D_\infty > 0, \quad \mathbf{P}^*\text{-a.s.}$$

For an optimal condition (of  $L \log L$ -type) for the positivity of  $D_\infty$ , see the recent work of Chen [14]. The two martingales  $(D_n)$  and  $(W_n)$  are asymptotically related; see Section 4.

The basic idea is to add a reflecting barrier at (notation:  $\llbracket \emptyset, x \rrbracket := \llbracket \emptyset, x \rrbracket \setminus \{x\}$ )

$$(2.10) \quad \mathcal{L}_r := \left\{ x : \sum_{z \in \llbracket \emptyset, x \rrbracket} e^{V(z)-V(x)} > r, \sum_{z \in \llbracket \emptyset, y \rrbracket} e^{V(z)-V(y)} \leq r, \forall y \in \llbracket \emptyset, x \rrbracket \right\},$$

where  $r > 1$  is a parameter.<sup>3</sup> We mention that  $\mathcal{L}_r$  does not necessarily separate  $\emptyset$  from infinity: our assumptions (2.2) and (2.3) do not exclude the existence of  $r > 1$  and a sequence of vertices  $x_0 := \emptyset < x_1 < x_2 \cdots$  with  $|x_i| = i, i \geq 0$ , such that  $\sum_{i=1}^n e^{V(x_i)-V(x_n)} \leq r$  for all  $n \geq 1$ .

---

<sup>3</sup>That is, each time the biased walk  $(X_i)$  hits any vertex  $x \in \mathcal{L}_r$ , it moves back to  $\overset{\leftarrow}{x}$  in the next step.

If  $r = r(n) := \frac{n}{(\log n)^\gamma}$  with  $\gamma < 1$ , then we will see from Lemma 5.1 that with  $\mathbb{P}^*$ -probability going to 1 (for  $n \rightarrow \infty$ ), the biased walk does not hit any vertex in  $\mathcal{L}_r$  in the first  $n$  steps.<sup>4</sup> As such, it makes no significant difference if we add a reflecting barrier at  $\mathcal{L}_r$ . An advantage, with the presence of the reflecting barrier at  $\mathcal{L}_r$ , for any  $r > 1$ , is that the biased walk becomes *positive recurrent* under the quenched probability  $P_\omega$ , and its invariant probability  $\pi_r$  is as follows:  $\pi_r(\overleftarrow{\emptyset}) := \frac{1}{Z_r}$ , and for  $x \in \mathbb{T}$ ,

$$(2.11) \quad \pi_r(x) := \begin{cases} \frac{1}{Z_r} e^{-U(x)}, & \text{if } x < \mathcal{L}_r, \\ \frac{1}{Z_r} e^{-V(x)}, & \text{if } x \in \mathcal{L}_r, \end{cases}$$

where  $Z_r$  is the normalizing factor:<sup>5</sup>

$$(2.12) \quad Z_r := 1 + \sum_{x \in \mathbb{T}: x < \mathcal{L}_r} e^{-U(x)} + \sum_{x \in \mathcal{L}_r} e^{-V(x)}.$$

We extend the definition of  $\pi_r$  to the whole tree  $\mathbb{T}$  by letting  $\pi_r(x) := 0$  if neither  $x < \mathcal{L}_r$  nor  $x \in \mathcal{L}_r$ .

Due to the periodicity of the walk  $(X_i)$ , we divide the tree  $\mathbb{T}$  into  $\mathbb{T}^{(\text{even})}$  and  $\mathbb{T}^{(\text{odd})}$  with

$$\mathbb{T}^{(\text{even})} := \{x \in \mathbb{T} : |x| \text{ is even}\}, \quad \mathbb{T}^{(\text{odd})} := \{x \in \mathbb{T} : |x| \text{ is odd}\}.$$

Depending on the parity of  $n$ , the law of  $X_n$  (starting from  $\emptyset$ ) is supported either by  $\mathbb{T}^{(\text{even})}$  or by  $\mathbb{T}^{(\text{odd})} \cup \{\overleftarrow{\emptyset}\}$ . Note that  $\pi_r(\mathbb{T}^{(\text{even})}) = \pi_r(\mathbb{T}^{(\text{odd})} \cup \{\overleftarrow{\emptyset}\}) = \frac{1}{2}$  as  $\pi_r(\cdot)$  is the invariant probability measure of a finite Markov chain of period 2. We define a new probability measure: for any  $r > 1$ ,

$$(2.13) \quad \tilde{\pi}_r(\cdot) := \begin{cases} 2\pi_r(\cdot) \mathbf{1}_{\mathbb{T}^{(\text{even})}}(\cdot), & \text{if } \lfloor r \rfloor \text{ is even,} \\ 2\pi_r(\cdot) \mathbf{1}_{\mathbb{T}^{(\text{odd})} \cup \{\overleftarrow{\emptyset}\}}(\cdot), & \text{if } \lfloor r \rfloor \text{ is odd.} \end{cases}$$

For any pair of probability measures  $\mu$  and  $\nu$  on  $\mathbb{T} \cup \{\overleftarrow{\emptyset}\}$ , we denote by  $d_{\text{tv}}(\mu, \nu)$  the distance in total variation:

$$d_{\text{tv}}(\mu, \nu) := \frac{1}{2} \sum_{x \in \mathbb{T} \cup \{\overleftarrow{\emptyset}\}} |\mu(x) - \nu(x)|.$$

The main result of the paper is as follows.

<sup>4</sup>Actually  $\gamma < 2$  will do the job (by Theorem 2.8). However, in Section 6, when we start proving our main results, only Lemma 5.1 is available, which says that  $\gamma < 1$  suffices. The proof of Theorem 2.8 comes afterwards, in Section 7.

<sup>5</sup>By  $x < \mathcal{L}_r$ , we mean  $\sum_{z \in \llbracket \emptyset, y \rrbracket} e^{V(z) - V(y)} \leq r$  for all vertex  $y \in \llbracket \emptyset, x \rrbracket$ .

**THEOREM 2.1.** *Assume (2.2) and (2.3). Then*

$$d_{\text{tv}}(P_\omega\{X_n \in \bullet\}, \tilde{\pi}_n) \rightarrow 0, \quad \text{in } \mathbf{P}^*\text{-probability.}$$

*More generally, for any  $\kappa \geq 1$  and  $0 < t_1 < t_2 < \dots < t_\kappa \leq 1$ ,*

$$d_{\text{tv}}\left(P_\omega\{(X_{\lfloor t_1 n \rfloor}, \dots, X_{\lfloor t_\kappa n \rfloor}) \in \bullet\}, \bigotimes_{i=1}^\kappa \tilde{\pi}_{t_i n}\right) \rightarrow 0, \quad \text{in } \mathbf{P}^*\text{-probability.}$$

As such,  $X_{\lfloor t_i n \rfloor}$ ,  $1 \leq i \leq \kappa$ , are asymptotically independent under  $P_\omega$ . In particular, no aging phenomenon is possible in the scale of linear time.

Let us mention that in Theorem 2.1, the dependence of  $\tilde{\pi}_{t_i n}$  on  $t_i$  is rather weak. As Lemma 2.2 below shows,  $d_{\text{tv}}(\pi_{t_i n}, \pi_n) \rightarrow 0$  in  $\mathbf{P}^*$ -probability, so asymptotically, the influence of  $t_i$  on  $\tilde{\pi}_{t_i n}$  shows up only via the parity of  $\lfloor t_i n \rfloor$ .

**LEMMA 2.2.** *For any  $a \geq 0$ , as  $r \rightarrow \infty$ ,*

$$\sup_{u \in [r/(\log r)^a, r]} d_{\text{tv}}(\pi_r, \pi_u) \rightarrow 0, \quad \text{in } \mathbf{P}^*\text{-probability.}$$

Theorem 2.1 has the following interesting consequence concerning distance between  $X_n$  and  $\emptyset$ .

**COROLLARY 2.3.** *Assume (2.2) and (2.3). Fix  $\kappa \geq 1$  and  $0 < t_1 < t_2 < \dots < t_\kappa \leq 1$ . Under  $\mathbb{P}^*$ ,  $\frac{\sigma^2}{(\log n)^2} |X_{\lfloor t_i n \rfloor}|$ ,  $1 \leq i \leq \kappa$ , are asymptotically independent and converge in law to a common non-degenerate limit on  $(0, \infty)$  whose density is given by*

$$\frac{1}{(2\pi x)^{1/2}} \mathbf{P}\left(\eta \leq \frac{1}{x^{1/2}}\right) \mathbf{1}_{\{x>0\}},$$

where  $\sigma^2 \in (0, \infty)$  is the constant in (1.4), and  $\eta := \sup_{s \in [0, 1]} [\bar{\mathbf{m}}(s) - \mathbf{m}(s)]$ . Here,  $(\mathbf{m}(s), s \in [0, 1])$  is a standard Brownian meander, and  $\bar{\mathbf{m}}(s) := \sup_{u \in [0, s]} \mathbf{m}(u)$ .

The distribution of  $\eta$  is easily seen to be absolutely continuous (Section 4), and can be characterised using a result of Lehoczký [25]. For more discussions, see [21]. Very recently, Pitman [38] has succeeded in determining the law of  $\eta$  using a relation between the Brownian meander and the Brownian bridge established by Biane and Yor [8]:  $\eta$  has the Kolmogorov–Smirnov distribution:

$$\mathbf{P}(\eta \leq x) = \sum_{k=-\infty}^{\infty} (-1)^k e^{-2k^2 x^2} = \frac{(2\pi)^{1/2}}{x} \sum_{j=0}^{\infty} \exp\left(-\frac{(2j+1)^2 \pi^2}{8x^2}\right), \quad x > 0.$$

Theorem 2.1 is proved by means of two intermediate estimates, stated below as Propositions 2.4 and 2.5. The first proposition estimates the local time at the root  $\emptyset$ , whereas the second concerns the local limit probability of the biased walk.

For any vertex  $x \in \mathbb{T}$ , let us define

$$(2.14) \quad L_n(x) := \sum_{i=1}^n \mathbf{1}_{\{X_i=x\}}, \quad n \geq 1,$$

which is the (site) local time of the biased walk at  $x$ .

PROPOSITION 2.4. *Assume (2.2) and (2.3). For any  $\varepsilon > 0$ ,*

$$(2.15) \quad P_\omega \left\{ \left| \frac{L_n(\emptyset)}{n/\log n} - \frac{\sigma^2}{4D_\infty} e^{-U(\emptyset)} \right| > \varepsilon \right\} \rightarrow 0, \quad \text{in } \mathbf{P}^*\text{-probability.}$$

Moreover,

$$(2.16) \quad E_\omega \left( \frac{L_n(\emptyset)}{n/\log n} \right) \rightarrow \frac{\sigma^2}{4D_\infty} e^{-U(\emptyset)}, \quad \text{in } \mathbf{P}^*\text{-probability,}$$

PROPOSITION 2.5. *Assume (2.2) and (2.3). As  $n \rightarrow \infty$  along even numbers,*

$$(\log n) P_\omega(X_n = \emptyset) \rightarrow \frac{\sigma^2}{2D_\infty} e^{-U(\emptyset)}, \quad \text{in } \mathbf{P}^*\text{-probability.}$$

We now say a few words about the proof. It turns out that the partition function  $Z_r$  has a simpler expression. Let  $\mathcal{L}_r$  be as in (2.10). Define

$$(2.17) \quad Y_r := \sum_{x \in \mathbb{T}: x \leq \mathcal{L}_r} e^{-V(x)},$$

with the obvious notation  $x \leq \mathcal{L}_r$  meaning  $x < \mathcal{L}_r$  or  $x \in \mathcal{L}_r$ .

LEMMA 2.6. *Let  $Y_r$  and  $Z_r$  be as in (2.17) and (2.12), respectively. Then  $Z_r = 2Y_r$ , for all  $r > 1$ .*

PROOF. If  $x \in \mathbb{T}$  is such that  $x < \mathcal{L}_r$ , we have  $Z_r \pi_r(x) = e^{-U(x)} = e^{-V(x)} + \sum_{y \in \mathbb{T}: \overleftarrow{y}=x} e^{-V(y)}$ . Therefore,

$$\begin{aligned} \sum_{x < \mathcal{L}_r} Z_r \pi_r(x) &= \sum_{x < \mathcal{L}_r} e^{-V(x)} + \sum_{x < \mathcal{L}_r} \sum_{y \in \mathbb{T}: \overleftarrow{y}=x} e^{-V(y)} \\ &= \sum_{x < \mathcal{L}_r} e^{-V(x)} + \sum_{y \in \mathbb{T}: \emptyset < y \leq \mathcal{L}_r} e^{-V(y)}, \end{aligned}$$

which is  $\sum_{x < \mathcal{L}_r} e^{-V(x)} + \sum_{y \leq \mathcal{L}_r} e^{-V(y)} - e^{-V(\emptyset)}$ . Hence

$$\sum_{x < \mathcal{L}_r} Z_r \pi_r(x) = 2 \sum_{x \leq \mathcal{L}_r} e^{-V(x)} - \sum_{x \in \mathcal{L}_r} e^{-V(x)} - 1.$$



Since  $\pi$  is a probability measure, we have  $\pi_r(\overleftarrow{\emptyset}) + \sum_{x < \mathcal{L}_r} \pi_r(x) + \sum_{x \in \mathcal{L}_r} \pi_r(x) = 1$ , so

$$\begin{aligned} Z_r &= Z_r \pi_r(\overleftarrow{\emptyset}) + \sum_{x < \mathcal{L}_r} Z_r \pi_r(x) + \sum_{x \in \mathcal{L}_r} Z_r \pi_r(x) \\ &= 1 + \left[ 2 \sum_{x \leq \mathcal{L}_r} e^{-V(x)} - \sum_{x \in \mathcal{L}_r} e^{-V(x)} - 1 \right] + \sum_{x \in \mathcal{L}_r} e^{-V(x)}, \end{aligned}$$

which is  $2 \sum_{x \leq \mathcal{L}_r} e^{-V(x)}$ . Lemma 2.6 is proved.  $\square$

So  $Y_r$  is half the partition function under the invariant measure. The following theorem, which plays an important role in the proof of Proposition 2.4 and Theorem 2.1, describes the asymptotics of  $Y_r$ .

**THEOREM 2.7.** *Assume (2.2) and (2.3). Let  $Y_r$  be as in (2.17). We have*

$$\lim_{r \rightarrow \infty} \frac{Y_r}{\log r} = \frac{2}{\sigma^2} D_\infty, \quad \text{in } \mathbf{P}^*\text{-probability,}$$

where  $\sigma^2 \in (0, \infty)$  is the constant in (1.4), and  $D_\infty$  the  $\mathbf{P}^*$ -almost sure positive limit of the derivative martingale  $(D_n)$  in (2.7). As a consequence,

$$\begin{aligned} \lim_{r \rightarrow \infty} \frac{Z_r}{\log r} &= \frac{4}{\sigma^2} D_\infty, \\ \lim_{r \rightarrow \infty} (\log r) \pi_r(\emptyset) &= \frac{\sigma^2}{4D_\infty} e^{-U(\emptyset)}, \quad \text{in } \mathbf{P}^*\text{-probability.} \end{aligned}$$

Finally, the following general estimate allows us to justify the presence of a barrier at  $\mathcal{L}_r$ .

**THEOREM 2.8.** *Assume (2.2) and (2.3). Let  $(a_n)$  be a deterministic sequence of positive real numbers such that  $\lim_{n \rightarrow \infty} \frac{a_n}{(\log n)^2} = 0$ , then*

$$\lim_{n \rightarrow \infty} \mathbb{P} \left( \bigcup_{i=1}^n \{X_i \in \mathcal{L}_{r_n}\} \right) = 0,$$

where  $r_n := \frac{n}{a_n}$ .

The rest of the paper is organised as follows:

- Section 3, environment: preliminaries on branching random walks.
- Section 4, environment: proof of Theorem 2.7.
- Section 5, biased walk: preliminaries on hitting barriers and local times.
- Section 6, biased walk: proof of Proposition 2.4.

- Section 7, biased walk: proof of Theorem 2.8.
- Section 8, biased walk: proof of Proposition 2.5.
- Section 9, biased walk: proofs of Lemma 2.2, Theorem 2.1 and Corollary 2.3.

Some comments on the organization are in order. In the next two sections, we study the behaviour of the random environment, starting in Section 3 by recalling some known results for branching random walks, and ending in Section 4 with the proof of Theorem 2.7. The biased walk  $(X_n)$  comes into picture in the last five sections. In Section 5, we collect a couple of useful results about hitting lines and local times for the biased walk. The proof of Proposition 2.4, which is the most technical part of the paper, is presented in Section 6. Once Proposition 2.4 is established, we use it to deduce Theorem 2.8 in Section 7, and Proposition 2.5 in Section 8. Finally, Theorem 2.1 and Corollary 2.3 (together with Lemma 2.2) are proved in Section 9.

Throughout the paper, for any pair of vertices  $x$  and  $y$ , we write  $x < y$  or  $y > x$  if  $y$  is a (strict) descendant of  $x$ , and  $x \leq y$  or  $y \geq x$  if either  $y > x$ , or  $y = x$ .

**3. Environment: Preliminaries on branching random walks.** We recall, in this section, some known results in the literature for branching random walks, and deduce a few useful consequences.

Under assumption (2.2), there exists a sequence of i.i.d. real-valued random variables  $(S_i - S_{i-1}, i \geq 0)$ , with  $S_0 = 0$ , such that for any  $n \geq 1$  and any Borel function  $g : \mathbb{R}^n \rightarrow \mathbb{R}_+$ ,

$$(3.1) \quad \mathbf{E} \left[ \sum_{x \in \mathbb{T}: |x|=n} g(V(x_i), 1 \leq i \leq n) \right] = \mathbf{E}[e^{S_n} g(S_i, 1 \leq i \leq n)],$$

where, for any vertex  $x \in \mathbb{T}$ ,  $x_i$  ( $0 \leq i \leq n$ ) denotes, as before, the ancestor of  $x$  in the  $i$ th generation. As such,  $V(x_0), V(x_1), \dots, V(x_n)$  (for  $|x| = n$ ) are the values of the potential  $V$  alongs the branch  $[[\emptyset, x]]$ .

Formula (3.1), often referred to as the “many-to-one formula,” is easily checked by induction on  $n$ . However, the appearance of the new, one-dimensional random walk  $(S_i, i \geq 0)$  has a deep meaning in terms of the so-called spinal decomposition via a change of probabilities. The idea of change of probabilities in the study of spatial branching processes has a long history, going back at least to Kahane and Peyrière [23] and to Bingham and Doney [13], and has led to various forms of the spinal decomposition. Since Lyons, Pemantle and Peres [31], it reaches a standard way of presentation. In our paper, we do not need any deep applications of the spinal decomposition, so we stay with the original probability  $\mathbf{P}$  without making any change of probabilities, even though we do need a “bivariate” version of (3.1):

$$(3.2) \quad \begin{aligned} & \mathbf{E} \left[ \sum_{x \in \mathbb{T}: |x|=n} g(V(x_i), \Lambda(x_{i-1}), 1 \leq i \leq n) \right] \\ & = \mathbf{E}[e^{S_n} g(S_i, \tilde{\Lambda}_{i-1}, 1 \leq i \leq n)], \end{aligned}$$

where, on the left-hand side of (3.2), we define

$$(3.3) \quad \Lambda(x) := \sum_{y: \overleftarrow{y}=x} e^{-[V(y)-V(x)]}, \quad x \in \mathbb{T},$$

and on the right-hand side of (3.2),  $(\tilde{\Lambda}_i, i \geq 0)$  is such that  $(S_i - S_{i-1}, \tilde{\Lambda}_{i-1}), i \geq 1$ , are i.i.d. random vectors whose law is characterised by

$$(3.4) \quad \mathbf{E}[h(S_1, \tilde{\Lambda}_0)] = \mathbf{E}\left[ \sum_{x \in \mathbb{T}: |x|=1} e^{-V(x)} h(V(x), \Lambda(\emptyset)) \right],$$

for any Borel function  $h : \mathbb{R}^2 \rightarrow \mathbb{R}_+$ . The last equality follows, obviously, from (3.2) by taking  $n = 1$  there. Note that by definition,  $\Lambda(\emptyset) = \sum_{x: |x|=1} e^{-V(x)}$ .

In particular, an application of the Hölder inequality, using assumption (2.3), yields the existence of  $\delta_1 > 0$  such that

$$(3.5) \quad \mathbf{E}[(\tilde{\Lambda}_0)^{\delta_1}] = \left[ \left( \sum_{x: |x|=1} e^{-V(x)} \right)^{1+\delta_1} \right] < \infty.$$

These are known facts about the spinal decomposition. For a proof of (3.2), see [20].

We now deduce several simple but useful results. The first allows us to include the random variable  $\Lambda(x)$  in the bivariate many-to-one formula (3.2). The second takes care of summation over all vertices on the stopping line  $\mathcal{L}_r$  instead of on a given generation, which leads to the third which is also the main estimate in this section.

LEMMA 3.1. *Assume (2.2). Let  $\Lambda(x)$  be as in (3.3). For any  $n \geq 1$  and any Borel function  $f : \mathbb{R}^{2n+1} \rightarrow \mathbb{R}_+$ , we have*

$$\begin{aligned} & \mathbf{E}\left[ \sum_{x \in \mathbb{T}: |x|=n} f(V(x_i), \Lambda(x_{i-1}), 1 \leq i \leq n, \Lambda(x)) \right] \\ & = \mathbf{E}[e^{S_n} F(S_i, \tilde{\Lambda}_{i-1}, 1 \leq i \leq n)], \end{aligned}$$

where  $(S_i - S_{i-1}, \tilde{\Lambda}_{i-1}), i \geq 1$ , are i.i.d. whose common distribution is given in (3.4), and

$$(3.6) \quad F(a_i, b_{i-1}, 1 \leq i \leq n) := \mathbf{E}\left[ f\left(a_i, b_{i-1}, 1 \leq i \leq n, \sum_{x \in \mathbb{T}: |x|=1} e^{-V(x)}\right) \right].$$

In particular, if  $g : \mathbb{R}^{n+1} \rightarrow \mathbb{R}_+$  is a Borel function, then

$$\mathbf{E}\left[ \sum_{x \in \mathbb{T}: |x|=n} g(V(x_1), \dots, V(x_n), \Lambda(x)) \right] = \mathbf{E}[e^{S_n} G(S_1, \dots, S_n)],$$

where  $G(a_1, \dots, a_n) := \mathbf{E}[g(a_1, \dots, a_n, \sum_{x \in \mathbb{T}: |x|=1} e^{-V(x)})]$ .

PROOF. Let  $\mathcal{F}_n := \sigma(x, V(x), x \in \mathbb{T}, |x| \leq n)$ , the  $\sigma$ -field generated by the branching random walk in the first  $n$  generations. By definition, for  $|x| = n$ ,  $\Lambda(x)$  is independent of  $\mathcal{F}_n$ , so

$$\begin{aligned} & \mathbf{E} \left[ \sum_{x \in \mathbb{T}: |x|=n} f(V(x_i), \Lambda(x_{i-1}), 1 \leq i \leq n, \Lambda(x)) \middle| \mathcal{F}_n \right] \\ &= \sum_{x \in \mathbb{T}: |x|=n} F(V(x_i), \Lambda(x_{i-1}), 1 \leq i \leq n), \end{aligned}$$

where  $F$  is given by (3.6). Taking expectation with respect to  $\mathbf{P}$  on both sides, and using the bivariate many-to-one formula (3.2), we obtain the lemma.  $\square$

LEMMA 3.2. Assume (2.2). Let  $E_1, E_2, \dots$  be Borel subsets of  $\mathbb{R}$ . Let  $r > 1$  and let  $\mathcal{L}_r$  be as in (2.10). Then

$$(3.7) \quad \mathbf{E} \left( \sum_{x \in \mathcal{L}_r} e^{-V(x)} \mathbf{1}_{\{V(x_i) \in E_i, 1 \leq i \leq |x|\}} \right) = \mathbf{P}(S_i \in E_i, 1 \leq i \leq T_r^{(S)}),$$

where  $T_r^{(S)} := \inf\{i \geq 1 : \sum_{j=1}^i e^{S_j - S_i} > r\}$ .

PROOF. We write

$$\sum_{x \in \mathcal{L}_r} e^{-V(x)} \mathbf{1}_{\{V(x_i) \in E_i, 1 \leq i \leq |x|\}} = \sum_{k=1}^{\infty} \sum_{x: |x|=k} e^{-V(x)} \mathbf{1}_{\{x \in \mathcal{L}_r\}} \mathbf{1}_{\{V(x_i) \in E_i, 1 \leq i \leq k\}}.$$

Obviously,  $\{x \in \mathcal{L}_r\} = \{\sum_{z \in \llbracket \emptyset, x \rrbracket} e^{V(z) - V(x)} > r, \sum_{z \in \llbracket \emptyset, v \rrbracket} e^{V(z) - V(v)} \leq r, \forall v \in \llbracket \emptyset, x \rrbracket\}$ . We take expectation with respect to  $\mathbf{P}$  on both sides. By the many-to-one formula (3.1),

$$\mathbf{E} \left( \sum_{x \in \mathcal{L}_r} e^{-V(x)} \mathbf{1}_{\{V(x_i) \in E_i, 1 \leq i \leq |x|\}} \right) = \sum_{k=1}^{\infty} \mathbf{P}(T_r^{(S)} = k, S_i \in E_i, 1 \leq i \leq k),$$

which is  $\mathbf{P}(S_i \in E_i, 1 \leq i \leq T_r^{(S)})$ .  $\square$

REMARK 3.3. Let  $Y_r := \sum_{x \leq \mathcal{L}_r} e^{-V(x)} = 1 + \sum_{k=1}^{\infty} \sum_{x: |x|=k} e^{-V(x)} \mathbf{1}_{\{x \leq \mathcal{L}_r\}}$  as in (2.17). Since  $x \leq \mathcal{L}_r$  means  $\sum_{z \in \llbracket \emptyset, v \rrbracket} e^{V(z) - V(v)} \leq r$  for all  $v \in \llbracket \emptyset, \overleftarrow{x} \rrbracket$  (the inequality considered as holding trivially if  $|x| = 1$ ), the proof of Lemma 3.2 yields

$$\mathbf{E}(Y_r) = 1 + \mathbf{E}(T_r^{(S)}) \leq 1 + \mathbf{E} \left[ \inf\{i \geq 1 : \max_{1 \leq j \leq i} S_j - S_i > \log r\} \right].$$

It is easy to check (for a detailed proof, see [20]) that  $\mathbf{E}[\inf\{i \geq 1 : \max_{1 \leq j \leq i} S_j - S_i > u\}]$  is bounded by  $c_1 u^2$  for some constant  $c_1 > 0$  and all  $u \geq 1$ . Hence, there exists  $c_2 > 0$  such that

$$(3.8) \quad \mathbf{E}(Y_r) \leq c_2 (\log r)^2, \quad r \geq 2.$$

We are going to use (3.8) in Section 6, in the proof of Proposition 2.4.

Although we do not need it in the present paper, an elementary argument shows that  $\frac{\mathbf{E}(Y_r)}{(\log r)^2}$  is bounded also from below.

We now present the main probabilistic estimate of the section.

LEMMA 3.4. *Assume (2.2) and (2.3). The laws of  $(\log r) \sum_{x \in \mathcal{L}_r} e^{-V(x)}$  under  $\mathbf{P}^*$ , for  $r \geq 1$ , are tight.*

*In particular, for any  $a < 1$ ,  $(\log r)^a \sum_{x \in \mathcal{L}_r} e^{-V(x)} \rightarrow 0$ ,  $r \rightarrow \infty$ , in  $\mathbf{P}^*$ -probability.*

PROOF. Let  $\varepsilon > 0$ . Our assumption ensures  $\inf_{x:|x|=n} V(x) \rightarrow \infty$  (for  $n \rightarrow \infty$ )  $\mathbf{P}^*$ -a.s. [see (2.9)]; so we can choose and fix a constant  $\alpha > 0$  such that

$$(3.9) \quad \mathbf{P}^* \left( \inf_{x \in \mathbb{T}} V(x) \geq -\alpha \right) \geq 1 - \varepsilon.$$

For any  $x \in \mathbb{T}$ , write

$$\underline{V}(x) := \min_{y \in [\emptyset, x]} V(y).$$

By Lemma 3.2,

$$(3.10) \quad \mathbf{E} \left( \sum_{x \in \mathcal{L}_r} e^{-V(x)} \mathbf{1}_{\{\underline{V}(x) \geq -\alpha\}} \right) = \mathbf{P}(\underline{S}_{T_r^{(S)}} \geq -\alpha),$$

where  $T_r^{(S)} := \inf\{i \geq 1 : \sum_{j=1}^i e^{S_j - S_i} > r\}$ , and  $\underline{S}_i := \min_{0 \leq j \leq i} S_j$ .

Let  $H(\frac{1}{2} \log r) := \inf\{i \geq 1 : S_i > \frac{1}{2} \log r\}$ . We have

$$\mathbf{P}(\underline{S}_{T_r^{(S)}} \geq -\alpha) \leq \mathbf{P}(T_r^{(S)} < H(\frac{1}{2} \log r), \underline{S}_{T_r^{(S)}} \geq -\alpha) + \mathbf{P}(\underline{S}_{H(1/2 \log r)} \geq -\alpha).$$

We bound the two probability expressions on the right-hand side. For  $\mathbf{P}(\underline{S}_{H(1/2 \log r)} \geq -\alpha)$ , we write  $H_-( -\alpha) := \inf\{i \geq 1 : S_i < -\alpha\}$ , to see that for some constant  $c_3 > 0$ ,

$$\mathbf{P}(\underline{S}_{H(1/2 \log r)} \geq -\alpha) = \mathbf{P} \left\{ H \left( \frac{1}{2} \log r \right) < H_-( -\alpha) \right\} \leq \frac{c_3 \alpha}{1/2 \log r + \alpha}.$$

(For the last inequality, which is elementary, see for example Aidékon [3] under the assumption of existence of exponential moments of  $S_1$ .) Hence  $\mathbf{P}(\underline{S}_{H(1/2 \log r)} \geq -\alpha) \leq \frac{2c_3 \alpha}{\log r}$ . Accordingly,

$$(3.11) \quad \mathbf{P}(\underline{S}_{T_r^{(S)}} \geq -\alpha) \leq \mathbf{P} \left( T_r^{(S)} < H \left( \frac{1}{2} \log r \right), \underline{S}_{T_r^{(S)}} \geq -\alpha \right) + \frac{2c_3 \alpha}{\log r}.$$

To deal with  $\mathbf{P}(T_r^{(S)} < H(\frac{1}{2} \log r), \underline{S}_{T_r^{(S)}} \geq -\alpha)$ , we note that by definition of  $T_r^{(S)}$ ,  $r < \sum_{j=1}^{T_r^{(S)}} e^{S_j - S_{T_r^{(S)}}}$ , which, on the event  $\{T_r^{(S)} < H(\frac{1}{2} \log r), \underline{S}_{T_r^{(S)}} \geq -\alpha\}$ , is

bounded by  $\sum_{j=1}^{T_r^{(S)}} e^{S_j+\alpha} \leq \sum_{j=1}^{H(1/2 \log r)-1} e^{(1/2 \log r)+\alpha} \leq r^{1/2} e^\alpha H(\frac{1}{2} \log r)$ . Consequently,

$$\mathbf{P}(T_r^{(S)} < H(\frac{1}{2} \log r), \underline{S}_{T_r^{(S)}} \geq -\alpha) \leq \mathbf{P}(H(\frac{1}{2} \log r) > r^{1/2} e^{-\alpha}).$$

By Kozlov [24],  $\mathbf{P}\{H(\frac{1}{2} \log r) > r^{1/2} e^{-\alpha}\} \leq c_4 \frac{e^{\alpha/2} \log r}{r^{1/4}}$  for some constant  $c_4 > 0$  and all  $n \geq 2$ . Going back to (3.11) and having (3.10) in mind, we obtain

$$\mathbf{E}\left(\sum_{x \in \mathcal{L}_r} e^{-V(x)} \mathbf{1}_{\{\underline{V}(x) \geq -\alpha\}}\right) \leq c_4 \frac{e^{\alpha/2} \log r}{r^{1/4}} + \frac{2c_3 \alpha}{\log r}.$$

In view of (3.9), and since  $\varepsilon > 0$  can be as small as possible, Lemma 3.4 follows readily.  $\square$

**4. Environment: Proof of Theorem 2.7.** This section is mainly devoted to the proof of Theorem 2.7, but also prepares a few useful estimates for the forthcoming sections. The material in this section concerns only the environment (thus the potential  $V$  and the symmetrised potential  $U$ ); no discussion on the movement of the biased walk  $(X_i)$  is involved.

Let  $W_n := \sum_{|x|=n} e^{-V(x)}$ ,  $n \geq 0$ , be the additive martingale as in (2.6). Consider also, for  $n \geq 0$  and  $\lambda > 0$ ,

$$(4.1) \quad W_n^{(\lambda)} := \sum_{|x|=n} e^{-V(x)} \mathbf{1}_{\{\max_{y \in \llbracket \emptyset, x \rrbracket} [\bar{V}(y) - V(y)] \leq \lambda\}},$$

where

$$\bar{V}(y) := \max_{z \in \llbracket \emptyset, y \rrbracket} V(z).$$

We mentioned earlier in (2.8) that under assumption (2.2), we have  $W_n \rightarrow 0$   $\mathbf{P}^*$ -a.s. The rate of decay of  $W_n$  is known: according to [5], under (2.2) and (2.3), we have

$$(4.2) \quad \lim_{n \rightarrow \infty} n^{1/2} W_n = \left(\frac{2}{\pi \sigma^2}\right)^{1/2} D_\infty, \quad \text{in } \mathbf{P}^*\text{-probability.}$$

The asymptotics of  $W_n^{(\lambda)}$  are also studied: according to Madaule [34], for any  $a \geq 0$ ,

$$\lim_{n \rightarrow \infty} \frac{W_n^{(n^{1/2} a)}}{W_n} = \mathbf{P}\left(\eta < \frac{a}{\sigma}\right), \quad \text{in } \mathbf{P}^*\text{-probability,}$$

where  $\eta := \sup_{s \in [0, 1]} [\bar{m}(s) - m(s)]$ , with  $\bar{m}(s) := \sup_{u \in [0, s]} m(u)$ , and  $(m(s), s \in [0, 1])$  denoting as before a standard Brownian meander. In view of (4.2), this is equivalent to saying the following convergence in  $\mathbf{P}^*$ -probability:

$$(4.3) \quad \lim_{n \rightarrow \infty} n^{1/2} W_n^{(n^{1/2} a)} = \left(\frac{2}{\pi \sigma^2}\right)^{1/2} D_\infty \mathbf{P}\left(\eta < \frac{a}{\sigma}\right).$$

This holds for any given  $a \geq 0$ .

By the absolute continuation relation between the Brownian meander  $m$  and the three-dimensional Bessel process  $R$  (see [22]), we know that the law of  $\eta := \sup_{s \in [0,1]} [\bar{m}(s) - m(s)]$  is absolutely continuous with respect to the law of  $\sup_{s \in [0,1]} [\bar{R}(s) - R(s)]$ . The latter is atomless because  $R$  is an  $h$ -transform (in the sense of Doob) of Brownian motion. As a consequence,  $a \mapsto \mathbf{P}(\eta < \frac{a}{\sigma})$  is continuous on  $\mathbb{R}$ . On the other hand, both  $a \mapsto W_n^{(n^{1/2}a)}$  and  $a \mapsto \mathbf{P}(\eta < \frac{a}{\sigma})$  are non-decreasing. It follows that (4.3) holds uniformly (in  $a \geq 0$ ) in the following sense: for any  $\varepsilon > 0$ ,

$$(4.4) \quad \lim_{n \rightarrow \infty} \mathbf{P}^* \left\{ \sup_{a \geq 0} \left| n^{1/2} W_n^{(n^{1/2}a)} - \left( \frac{2}{\pi \sigma^2} \right)^{1/2} D_\infty \mathbf{P} \left( \eta < \frac{a}{\sigma} \right) \right| \geq \varepsilon \right\} = 0.$$

We now state a lemma.

LEMMA 4.1. *We have*

$$(4.5) \quad \lim_{\lambda \rightarrow \infty} \frac{1}{\lambda} \sum_{k=1}^\infty W_k^{(\lambda)} = \frac{2}{\sigma^2} D_\infty, \quad \text{in } \mathbf{P}^* \text{-probability.}$$

PROOF. We first argue that in  $\sum_{k=1}^\infty W_k^{(\lambda)}$ , only those  $k$  that are comparable to  $\lambda^2$  make a significant contribution to the sum. More precisely, we claim that for any  $\varepsilon_1 > 0$ ,

$$(4.6) \quad \lim_{b \rightarrow 0} \limsup_{\lambda \rightarrow \infty} \mathbf{P}^* \left\{ \frac{1}{\lambda} \sum_{k=1}^{\lfloor b\lambda^2 \rfloor} W_k^{(\lambda)} \geq \varepsilon_1 \right\} = 0,$$

$$(4.7) \quad \lim_{B \rightarrow \infty} \limsup_{\lambda \rightarrow \infty} \mathbf{P}^* \left\{ \frac{1}{\lambda} \sum_{k=\lfloor B\lambda^2 \rfloor}^\infty W_k^{(\lambda)} \geq \varepsilon_1 \right\} = 0.$$

To prove (4.6) and (4.7), let  $\varepsilon > 0$  and fix  $\alpha \geq 0$  as in (3.9), that is, such that

$$(3.9) \quad \mathbf{P}^* \left\{ \inf_{x \in \mathbb{T}} V(x) \geq -\alpha \right\} \geq 1 - \varepsilon.$$

Consider the truncated version of  $W_k^{(\lambda)}$  defined by

$$W_k^{(\lambda, \alpha)} := \sum_{|x|=k} e^{-V(x)} \mathbf{1}_{\{\max_{y \in \llbracket \emptyset, x \rrbracket} [\bar{V}(y) - V(y)] \leq \lambda\}} \mathbf{1}_{\{\underline{V}(x) \geq -\alpha\}},$$

where  $\underline{V}(x) := \min_{z \in \llbracket \emptyset, x \rrbracket} V(z)$  as before. Clearly, on the set  $\{\inf_{x \in \mathbb{T}} V(x) \geq -\alpha\}$ ,  $W_k^{(\lambda, \alpha)} = W_k^{(\lambda)}$  for all  $k \geq 1$ .

By the many-to-one formula in (3.1),

$$(4.8) \quad \mathbf{E}(W_k^{(\lambda, \alpha)}) = \mathbf{P} \left\{ \max_{0 \leq j \leq k} (\bar{S}_j - S_j) \leq \lambda, \underline{S}_k \geq -\alpha \right\},$$

where  $\bar{S}_j := \max_{0 \leq i \leq j} S_i$  and  $\underline{S}_j := \min_{0 \leq i \leq j} S_i$ .

The proof of (4.6) is easy: we have  $\mathbf{E}(W_k^{(\lambda, \alpha)}) \leq \mathbf{P}\{\underline{S}_k \geq -\alpha\}$ , which is bounded by  $\frac{c_5}{k^{1/2}}$  for some constant  $c_5$  (depending on  $\alpha$ ) and all  $k \geq 1$  (see Kozlov [24]); hence

$$\frac{1}{\lambda} \mathbf{E} \left( \sum_{k=1}^{\lfloor b\lambda^2 \rfloor} W_k^{(\lambda, \alpha)} \right) \leq \frac{1}{\lambda} \sum_{k=1}^{\lfloor b\lambda^2 \rfloor} \frac{c_5}{k^{1/2}},$$

which goes to 0 when  $\lambda \rightarrow \infty$  and then  $b \rightarrow 0$ . This readily yields (4.6).

To prove (4.7), we use (4.8), and apply the Markov property at time  $\frac{k}{2}$  (treating it as an integer by dropping the symbol of the integer part), to see that

$$\begin{aligned} \mathbf{E}(W_k^{(\lambda, \alpha)}) &\leq \mathbf{P}\left\{ \underline{S}_{k/2} \geq -\alpha, \max_{k/2 \leq j \leq k} (\bar{S}_j - S_j) \leq \lambda \right\} \\ &\leq \mathbf{P}\{ \underline{S}_{k/2} \geq -\alpha \} \times \mathbf{P}\left\{ \max_{0 \leq j \leq k/2} (\bar{S}_j - S_j) \leq \lambda \right\}. \end{aligned}$$

Again,  $\mathbf{P}\{ \underline{S}_{k/2} \geq -\alpha \} \leq \frac{c_5}{(k/2)^{1/2}}$ , whereas  $\mathbf{P}\{ \max_{0 \leq j \leq k/2} (\bar{S}_j - S_j) \leq \lambda \}$  can be estimated as follows: by the Markov property,  $\mathbf{P}\{ \max_{0 \leq j \leq k/2} (\bar{S}_j - S_j) \leq \lambda \} \leq [\mathbf{P}\{ \max_{0 \leq j \leq \lfloor \lambda^2 \rfloor} (\bar{S}_j - S_j) \leq \lambda \}]^{\lfloor k/(2\lfloor \lambda^2 \rfloor) \rfloor}$ . By Donsker's theorem, there exists a constant  $0 < c_6 < 1$  such that  $\mathbf{P}\{ \max_{0 \leq j \leq \lfloor \lambda^2 \rfloor} (\bar{S}_j - S_j) \leq \lambda \} \leq 1 - c_6$  for all sufficiently large  $\lambda$  (say  $\lambda \geq \lambda_0$ ) which yields  $\mathbf{P}\{ \max_{0 \leq j \leq k/2} (\bar{S}_j - S_j) \leq \lambda \} \leq (1 - c_6)^{\lfloor k/(2\lfloor \lambda^2 \rfloor) \rfloor}$ ,  $\forall \lambda \geq \lambda_0$ . Hence, for  $\lambda \geq \lambda_0$  and  $k \geq 1$ ,

$$\mathbf{E}(W_k^{(\lambda, \alpha)}) \leq \frac{c_5}{(k/2)^{1/2}} (1 - c_6)^{\lfloor k/(2\lfloor \lambda^2 \rfloor) \rfloor}$$

from which it follows that

$$\lim_{B \rightarrow \infty} \limsup_{\lambda \rightarrow \infty} \frac{1}{\lambda} \mathbf{E} \left\{ \sum_{k=\lfloor B\lambda^2 \rfloor}^{\infty} W_k^{(\lambda, \alpha)} \right\} = 0.$$

Since  $\varepsilon > 0$  in (3.9) can be as small as possible, this implies (4.7).

Now that (4.6) and (4.7) are justified, we are ready for the proof of Lemma 4.1. Fix  $B > b > 0$ . By (4.4), for  $\lambda \rightarrow \infty$ ,

$$\frac{1}{\lambda} \sum_{k=\lfloor b\lambda^2 \rfloor}^{\lfloor B\lambda^2 \rfloor} W_k^{(\lambda)} = \frac{1}{\lambda} \left( \frac{2}{\pi \sigma^2} \right)^{1/2} D_\infty \sum_{k=\lfloor b\lambda^2 \rfloor}^{\lfloor B\lambda^2 \rfloor} \frac{1}{k^{1/2}} \mathbf{P} \left( \eta < \frac{\lambda}{\sigma k^{1/2}} \right) + o_{\mathbf{P}^*}(1),$$

where  $o_{\mathbf{P}^*}(1)$  denotes a term satisfying  $\lim_{\lambda \rightarrow \infty} o_{\mathbf{P}^*}(1) = 0$  in  $\mathbf{P}^*$ -probability (whose value may vary from line to line). On the other hand, by Fubini's theorem,

$$\begin{aligned} \frac{1}{\lambda} \int_{b\lambda^2}^{B\lambda^2} \frac{1}{u^{1/2}} \mathbf{P} \left( \eta < \frac{\lambda}{\sigma u^{1/2}} \right) du &= \frac{1}{\lambda} \mathbf{E} \left[ \int_{b\lambda^2}^{(B\lambda^2) \wedge \lambda^2 / (\sigma^2 \eta^2)} \frac{du}{u^{1/2}} \mathbf{1}_{\{\eta < 1/(\sigma b^{1/2})\}} \right] \\ &= 2\mathbf{E} \left[ \left( \left( B^{1/2} \wedge \frac{1}{\sigma \eta} \right) - b^{1/2} \right) \mathbf{1}_{\{\eta < 1/(\sigma b^{1/2})\}} \right]. \end{aligned}$$



Since  $\eta$  is atomless, this yields

$$\begin{aligned}
 & \frac{1}{\lambda} \sum_{k=\lfloor b\lambda^2 \rfloor}^{\lfloor B\lambda^2 \rfloor} W_k^{(\lambda)} \\
 (4.9) \quad &= \left( \frac{8}{\pi\sigma^2} \right)^{1/2} D_\infty \mathbf{E} \left[ \left( \left( B^{1/2} \wedge \frac{1}{\sigma\eta} \right) - b^{1/2} \right) \mathbf{1}_{\{\eta < 1/(\sigma b^{1/2})\}} \right] \\
 & \quad + o_{\mathbf{P}^*}(1).
 \end{aligned}$$

Note that  $\mathbf{E}[\left( (B^{1/2} \wedge \frac{1}{\sigma\eta}) - b^{1/2} \right) \mathbf{1}_{\{\eta < 1/(\sigma b^{1/2})\}}] \rightarrow \mathbf{E}[\frac{1}{\sigma\eta}]$  when  $B \rightarrow \infty$  and  $b \rightarrow 0$ . In view of (4.6) and (4.7), we see that when  $\lambda \rightarrow \infty$ ,

$$\frac{1}{\lambda} \sum_{k=1}^{\infty} W_k^{(\lambda)} \rightarrow \left( \frac{8}{\pi\sigma^2} \right)^{1/2} D_\infty \mathbf{E} \left[ \frac{1}{\sigma\eta} \right], \quad \text{in } \mathbf{P}^*\text{-probability.}$$

By [21],  $\mathbf{E}(\frac{1}{\eta}) = (\frac{\pi}{2})^{1/2}$ , which yields Lemma 4.1.  $\square$

We now have all the ingredients for the proof of Theorem 2.7.

**PROOF OF THEOREM 2.7.** By definition,

$$Y_r = \sum_{x \in \mathbb{T}} e^{-V(x)} \mathbf{1}_{\{x < \mathcal{L}_r\}} + \sum_{x \in \mathcal{L}_r} e^{-V(x)}.$$

We already know (Lemma 3.4) that  $\sum_{x \in \mathcal{L}_r} e^{-V(x)} \rightarrow 0$  in  $\mathbf{P}^*$ -probability. So it remains to check that

$$(4.10) \quad \lim_{r \rightarrow \infty} \frac{1}{\log r} \sum_{x \in \mathbb{T}} e^{-V(x)} \mathbf{1}_{\{x < \mathcal{L}_r\}} = \frac{2}{\sigma^2} D_\infty, \quad \text{in } \mathbf{P}^*\text{-probability.}$$

By definition,  $\{x < \mathcal{L}_r\}$  means  $\sum_{z \in \llbracket \emptyset, y \rrbracket} e^{V(z) - V(y)} \leq r, \forall y \in \llbracket \emptyset, x \rrbracket$ . So

$$\begin{aligned}
 & \sum_{x \in \mathbb{T}} e^{-V(x)} \mathbf{1}_{\{x < \mathcal{L}_r\}} \\
 (4.11) \quad &= \sum_{k=0}^{\infty} \sum_{x: |x|=k} e^{-V(x)} \mathbf{1}_{\{\sum_{z \in \llbracket \emptyset, y \rrbracket} e^{V(z) - V(y)} \leq r, \forall y \in \llbracket \emptyset, x \rrbracket\}} \\
 & \leq \sum_{k=0}^{\infty} \sum_{x: |x|=k} e^{-V(x)} \mathbf{1}_{\{\max_{y \in \llbracket \emptyset, x \rrbracket} [\bar{V}(y) - V(y)] \leq \log r\}} \\
 & = \sum_{k=0}^{\infty} W_k^{(\log r)}.
 \end{aligned}$$

A similar lower bound holds as well: we fix an arbitrary positive real number  $B > 0$ ,

$$\begin{aligned}
 & \sum_{x \in \mathbb{T}} e^{-V(x)} \mathbf{1}_{\{x < \mathcal{L}_r\}} \\
 (4.12) \quad & \geq \sum_{k=0}^{\lfloor B(\log r)^2 \rfloor} \sum_{x:|x|=k} e^{-V(x)} \mathbf{1}_{\{\max_{y \in \|\emptyset, x\|} [\bar{V}(y) - V(y)] \leq \log(r/B(\log r)^2)\}} \\
 & = \sum_{k=0}^{\lfloor B(\log r)^2 \rfloor} W_k^{(\log(r/B(\log r)^2))}.
 \end{aligned}$$

Applying Lemma 4.1 and (4.7), and noting that  $\lim_{r \rightarrow \infty} \frac{\log(r/B(\log r)^2)}{\log r} = 1$ , we obtain that under  $\mathbf{P}^*$ ,

$$\lim_{r \rightarrow \infty} \frac{1}{\log r} \sum_{x \in \mathbb{T}} e^{-V(x)} \mathbf{1}_{\{x < \mathcal{L}_r\}} = \frac{2}{\sigma^2} D_\infty, \quad \text{in probability.}$$

Theorem 2.7 is proved.  $\square$

REMARK 4.2. The proof of the upper bound in Theorem 2.7, combined with (4.7), tells us that for any  $\varepsilon > 0$ ,

$$(4.13) \quad \lim_{B \rightarrow \infty} \limsup_{r \rightarrow \infty} \mathbf{P}^* \left\{ \frac{1}{\log r} \sum_{x \in \mathbb{T}: |x| \geq B(\log r)^2, x \leq \mathcal{L}_r} e^{-V(x)} \geq \varepsilon \right\} = 0.$$

We are entitled to sum over  $x \leq \mathcal{L}_r$  instead of over  $x < \mathcal{L}_r$  because  $\sum_{x \in \mathcal{L}_r} e^{-V(x)} \rightarrow 0$  in  $\mathbf{P}^*$ -probability (Lemma 3.4); (4.13) will be useful in Section 6.

**5. Biased walks: Preliminaries on hitting barriers and local times.** In this section, we collect two preliminary results for the biased walk  $(X_i)$ . The first is a weaker version of Theorem 2.8, and the second concerns the covariance of edge local times. For the sake of clarity, we present them in two distinct subsections.

5.1. *Hitting reflecting barriers.* This subsection is devoted to a weaker version of Theorem 2.8, stated as follows. The proof of Theorem 2.8 comes much later, in Section 7.

LEMMA 5.1. Assume (2.2) and (2.3). If  $r = r(n) := \frac{n}{(\log n)^\gamma}$  with  $\gamma < 1$ , then

$$\lim_{n \rightarrow \infty} \mathbb{P} \left( \bigcup_{i=1}^n \{X_i \in \mathcal{L}_r\} \right) = 0,$$

where  $\mathcal{L}_r$  is as in (2.10).

PROOF. Define

$$(5.1) \quad T_x := \inf\{i \geq 0 : X_i = x\}, \quad x \in \mathbb{T},$$

$$(5.2) \quad T_\emptyset^+ := \inf\{i \geq 1 : X_i = \emptyset\}.$$

In words,  $T_x$  is the first hitting time at  $x$  by the biased walk, whereas  $T_\emptyset^+$  is the first return time to the root  $\emptyset$ .

Let  $x \in \mathbb{T} \setminus \{\emptyset\}$ . The probability  $P_\omega(T_x < T_\emptyset^+)$  only involves a one-dimensional random walk in random environment [namely, the restriction at  $\llbracket \emptyset, x \rrbracket$  of the biased walk  $(X_i)$ ], so a standard result for one-dimensional random walks in random environment (Golosoov [17]) tells us that

$$(5.3) \quad P_\omega(T_x < T_\emptyset^+) = \frac{\omega(\emptyset, x_1)e^{V(x_1)}}{\sum_{z \in \llbracket \emptyset, x \rrbracket} e^{V(z)}} = \frac{\omega(\emptyset, \overleftarrow{\emptyset})}{\sum_{z \in \llbracket \emptyset, x \rrbracket} e^{V(z)}},$$

where  $x_1$  is the ancestor of  $x$  in the first generation.

Define  $T_\emptyset^{(0)} := 0$  and inductively  $T_\emptyset^{(k)} := \inf\{i > T_\emptyset^{(k-1)} : X_i = \emptyset\}$ ,  $k \geq 1$ . In words,  $T_\emptyset^{(k)}$  is the  $k$ th return time of the biased walk  $(X_i)$  to the root  $\emptyset$ . (In particular,  $T_\emptyset^{(1)} = T_\emptyset^+$ .) For  $n \geq 1$ , we have

$$\begin{aligned} P_\omega(T_x \leq n) &= \sum_{k=0}^\infty P_\omega[T_x \leq n, T_\emptyset^{(k)} \leq T_x < T_\emptyset^{(k+1)}] \\ &= \sum_{k=0}^\infty E_\omega[\mathbf{1}_{\{T_\emptyset^{(k)} \leq n\}} P_\omega(T_x < T_\emptyset, T_x \leq n - j) |_{j:=T_\emptyset^{(k)}}] \\ &\leq \sum_{k=0}^\infty E_\omega[\mathbf{1}_{\{T_\emptyset^{(k)} \leq n\}} P_\omega(T_x < T_\emptyset^+)] \\ &= P_\omega(T_x < T_\emptyset^+) E_\omega(L_n(\emptyset) + 1), \end{aligned}$$

where  $L_n(\emptyset) := \sum_{i=1}^n \mathbf{1}_{\{X_i = \emptyset\}}$  is the local time at  $\emptyset$ . By (5.3), we get

$$P_\omega(T_x \leq n) \leq \frac{E_\omega(L_n(\emptyset) + 1)}{\sum_{z \in \llbracket \emptyset, x \rrbracket} e^{V(z)}}.$$

Let  $r > 1$ , and let  $\mathcal{L}_r$  be as in (2.10). We have

$$P_\omega\left(\bigcup_{i=1}^n \{X_i \in \mathcal{L}_r\}\right) \leq \sum_{x \in \mathcal{L}_r} P_\omega(T_x \leq n) \leq E_\omega(L_n(\emptyset) + 1) \sum_{x \in \mathcal{L}_r} \frac{1}{\sum_{z \in \llbracket \emptyset, x \rrbracket} e^{V(z)}}.$$

By definition of  $\mathcal{L}_r$ ,  $\frac{1}{\sum_{z \in \llbracket \emptyset, x \rrbracket} e^{V(z)}} \leq \frac{1}{r} e^{-V(x)}$  for  $x \in \mathcal{L}_r$ ; hence

$$(5.4) \quad P_\omega\left(\bigcup_{i=1}^n \{X_i \in \mathcal{L}_r\}\right) \leq \frac{E_\omega(L_n(\emptyset) + 1)}{r} \sum_{x \in \mathcal{L}_r} e^{-V(x)}.$$

We use the trivial inequality  $L_n(\emptyset) \leq n$ , so  $E_\omega(L_n(\emptyset)) \leq n$ . We now take  $r = r(n) := \frac{n}{(\log n)^\gamma}$ . With this choice of  $r$ , Lemma 3.4 tells us that if  $\gamma < 1$ , then  $(\log n)^\gamma \sum_{x \in \mathcal{L}_r} e^{-V(x)} \rightarrow 0$  in  $\mathbf{P}^*$ -probability. This yields Lemma 5.1.  $\square$

5.2. *Covariance for edge local times of biased walks.* In the proof of Proposition 2.4 in Section 6, we are going to estimate the covariance of local time of the biased walk  $(X_i)$ . It turns out to be more convenient to deal with covariance of edge local time instead of site local time. More precisely, for any  $k \geq 1$  and any vertex  $x \in \mathbb{T} \setminus \{\emptyset\}$ , let us define the edge local time

$$(5.5) \quad \bar{L}_k(x) := \sum_{i=1}^k \mathbf{1}_{\{X_{i-1}=\overleftarrow{x}, X_i=x\}},$$

which is the number of passages of the walk  $(X_i)$ , in the first  $k$  steps, on the oriented edge from  $\overleftarrow{x}$  to  $x$ . We are interested in the (edge) local time during an excursion away from  $\emptyset$ .

Notation:  $x \wedge y$  is the youngest common ancestor of  $x$  and  $y$  (or, equivalently, the unique vertex satisfying  $[\emptyset, x \wedge y] = [\emptyset, x] \cap [\emptyset, y]$ ).

LEMMA 5.2. *Let  $T_\emptyset^+ := \inf\{i \geq 1 : X_i = \emptyset\}$  denote the first return to the root  $\emptyset$  as in (5.2).*

(i) *We have, for  $x \neq y \in \mathbb{T}$ ,*

$$(5.6) \quad \begin{aligned} & \text{Cov}_\omega[\bar{L}_{T_\emptyset^+}(x), \bar{L}_{T_\emptyset^+}(y)] \\ & \leq 2e^{-[V(x)-V(x \wedge y)]-[V(y)-V(x \wedge y)]} E_\omega[\bar{L}_{T_\emptyset^+}(x \wedge y)^2], \end{aligned}$$

where  $\text{Cov}_\omega$  stands for covariance under the quenched probability  $P_\omega$ .

(ii) *We have, for  $x \in \mathbb{T} \setminus \{\emptyset\}$ ,*

$$(5.7) \quad E_\omega[\bar{L}_{T_\emptyset^+}(x)] = \omega(\emptyset, \overleftarrow{\emptyset})e^{-V(x)}.$$

$$(5.8) \quad E_\omega[\bar{L}_{T_\emptyset^+}(x)^2] = \omega(\emptyset, \overleftarrow{\emptyset})e^{-V(x)} \left( 2 \sum_{y \in [\emptyset, x]} e^{V(y)-V(x)} - 1 \right).$$

PROOF. (i) We use the following elementary identity: for any pairs of random variables  $\xi_1$  and  $\xi_2$  defined on a probability space  $(\Omega, \mathcal{F}, \mathbb{P})$ , having finite second moments, and any  $\sigma$ -field  $\mathcal{G} \subset \mathcal{F}$ , we have

$$(5.9) \quad \text{Cov}(\xi_1, \xi_2) = \mathbb{E}[\text{Cov}_{\mathcal{G}}(\xi_1, \xi_2)] + \text{Cov}[\mathbb{E}(\xi_1|\mathcal{G}), \mathbb{E}(\xi_2|\mathcal{G})],$$

where  $\text{Cov}_{\mathcal{G}}(\xi_1, \xi_2) := \mathbb{E}(\xi_1 \xi_2 | \mathcal{G}) - \mathbb{E}(\xi_1 | \mathcal{G}) \mathbb{E}(\xi_2 | \mathcal{G})$  is the conditional covariance of  $\xi_1$  and  $\xi_2$  given  $\mathcal{G}$ .

We first treat the case that neither of  $x$  and  $y$  is an ancestor of the other.

We write  $u = u(x, y) := x \wedge y$  for brevity, and let  $x_{|u|+1}$  and  $y_{|u|+1}$  be the ancestor, at generation  $|u| + 1$ , of  $x$  and  $y$  respectively. By definition of  $x \wedge y$ , the vertices  $x_{|u|+1}$  and  $y_{|u|+1}$  are distinct children of  $u$ . Conditionally on  $\bar{L}_{T_\emptyset^+}(x_{|u|+1})$  and  $\bar{L}_{T_\emptyset^+}(y_{|u|+1})$ , the edge local times  $\bar{L}_{T_\emptyset^+}(x)$  and  $\bar{L}_{T_\emptyset^+}(y)$  are independent. We apply (5.9) to  $\xi_1 := \bar{L}_{T_\emptyset^+}(x)$ ,  $\xi_2 := \bar{L}_{T_\emptyset^+}(y)$  and  $\mathcal{G} := \sigma(\bar{L}_{T_\emptyset^+}(x_{|u|+1}), \bar{L}_{T_\emptyset^+}(y_{|u|+1}))$ , the  $\sigma$ -field generated by the edge local times  $\bar{L}_{T_\emptyset^+}(x_{|u|+1})$  and  $\bar{L}_{T_\emptyset^+}(y_{|u|+1})$ . Since the conditional covariance vanishes, (5.9) gives that

$$(5.10) \quad \text{Cov}_\omega[\bar{L}_{T_\emptyset^+}(x), \bar{L}_{T_\emptyset^+}(y)] = \text{Cov}_\omega[E_\omega(\bar{L}_{T_\emptyset^+}(x)|\mathcal{G}), E_\omega(\bar{L}_{T_\emptyset^+}(y)|\mathcal{G})],$$

with  $\mathcal{G} := \sigma(\bar{L}_{T_\emptyset^+}(x_{|u|+1}), \bar{L}_{T_\emptyset^+}(y_{|u|+1}))$ . Let us compute  $E_\omega(\bar{L}_{T_\emptyset^+}(x)|\mathcal{G})$ , which is nothing else but  $E_\omega(\bar{L}_{T_\emptyset^+}(x)|\bar{L}_{T_\emptyset^+}(x_{|u|+1}))$ . Write  $|x| =: j > i := |u|$ . Then for any  $k \in (i, j) \cap \mathbb{Z}$ , and given  $\bar{L}_{T_\emptyset^+}(x_k) = \ell \geq 1$ ,  $\bar{L}_{T_\emptyset^+}(x_{k+1})$  has the law of  $\sum_{m=1}^\ell G_m$ , where  $G_m, m \geq 1$ , are i.i.d. geometric random variables with parameter  $p_k := \frac{\omega(x_k, x_{k-1})}{\omega(x_k, x_{k+1}) + \omega(x_k, x_{k-1})}$  [i.e.,  $G_m$  takes value  $r$  with probability  $(1 - p_k)^r p_k$  for all non-negative integer  $r$ ]. Since  $G_m$  has mean  $\frac{1-p_k}{p_k}$ , we have  $E_\omega(\bar{L}_{T_\emptyset^+}(x_{k+1})|\bar{L}_{T_\emptyset^+}(x_k)) = \bar{L}_{T_\emptyset^+}(x_k) \frac{1-p_k}{p_k} = \bar{L}_{T_\emptyset^+}(x_k) e^{-[V(x_{k+1})-V(x_k)]}$ . As a consequence, we deduce from the Markov property of  $k \rightarrow \bar{L}_{T_\emptyset^+}(x_k)$  (under  $P_\omega$ ) that

$$\begin{aligned} E_\omega(\bar{L}_{T_\emptyset^+}(x)|\bar{L}_{T_\emptyset^+}(x_{|u|+1})) &= \bar{L}_{T_\emptyset^+}(x_{|u|+1}) \prod_{k=i+1}^{j-1} e^{-[V(x_{k+1})-V(x_k)]} \\ &= \bar{L}_{T_\emptyset^+}(x_{|u|+1}) e^{-[V(x)-V(x_{|u|+1})]}. \end{aligned}$$

Similarly,  $E_\omega(\bar{L}_{T_\emptyset^+}(y)|\bar{L}_{T_\emptyset^+}(y_{|u|+1})) = \bar{L}_{T_\emptyset^+}(y_{|u|+1}) e^{-[V(y)-V(y_{|u|+1})]}$ . Going back to (5.10), this leads to

$$(5.11) \quad \begin{aligned} &\text{Cov}_\omega[\bar{L}_{T_\emptyset^+}(x), \bar{L}_{T_\emptyset^+}(y)] \\ &= e^{-[V(x)-V(x_{|u|+1})]-[V(y)-V(y_{|u|+1})]} \text{Cov}_\omega[\bar{L}_{T_\emptyset^+}(x_{|u|+1}), \bar{L}_{T_\emptyset^+}(y_{|u|+1})]. \end{aligned}$$

To compute the covariance on the right-hand side, we write  $(u^{(1)}, \dots, u^{(N(u))})$  for the children of  $u$  [among which are  $x_{|u|+1}$  and  $y_{|u|+1}$ ; so  $N(u) \geq 2$ ], and observe that conditionally on  $\bar{L}_{T_\emptyset^+}(u) = \ell \geq 1$ , the law of the random vector  $(\bar{L}_{T_\emptyset^+}(u^{(k)}), 1 \leq k \leq N(u))$  under  $P_\omega$  is multinomial with parameter  $(\sum_{k=1}^\ell \mathfrak{G}_k, (p^{(k)}(u) := \frac{\omega(u, u^{(k)})}{1-\omega(u, \bar{u})}, 1 \leq k \leq N(u)))$ , where  $\mathfrak{G}_k, k \geq 1$ , are i.i.d. random variables having the geometric distribution of parameter  $\omega(u, \bar{u})$ .<sup>6</sup> Accord-

<sup>6</sup>A random vector  $(\xi_1, \dots, \xi_N)$  defined on  $(\Omega, \mathcal{F}, \mathbb{P})$  has the multinomial distribution with parameter  $(m, (p^{(1)}, \dots, p^{(N)}))$  if  $\mathbb{P}(\xi_1 = m_1, \dots, \xi_N = m_N) = \frac{m!}{m_1! \dots m_N!} \prod_{k=1}^N (p^{(k)})^{m_k}$  for all non-

ingly, for all  $\ell \geq 1$ ,

$$E_\omega \left[ \prod_{k=1}^{N(u)} (s_k)^{\bar{L}_{T_\emptyset^+}(u^{(k)})} \mid \bar{L}_{T_\emptyset^+}(u) = \ell \right] = E_\omega \left[ \left( \sum_{k=1}^{N(u)} \frac{s_k \omega(u, u^{(k)})}{1 - \omega(u, \bar{u})} \right)^{\sum_{k=1}^\ell \mathfrak{G}_k} \right] \\ = \left\{ E_\omega \left[ \left( \sum_{k=1}^{N(u)} \frac{s_k \omega(u, u^{(k)})}{1 - \omega(u, \bar{u})} \right)^{\mathfrak{G}_1} \right] \right\}^\ell.$$

Since  $E_\omega(s^{\mathfrak{G}_1}) = \frac{\omega(u, \bar{u})}{1 - s(1 - \omega(u, \bar{u}))}$ , this yields

$$(5.12) \quad E_\omega \left[ \prod_{k=1}^{N(u)} (s_k)^{\bar{L}_{T_\emptyset^+}(u^{(k)})} \mid \bar{L}_{T_\emptyset^+}(u) \right] = \left\{ \frac{\omega(u, \bar{u})}{1 - \sum_{k=1}^{N(u)} s_k \omega(u, u^{(k)})} \right\}^{\bar{L}_{T_\emptyset^+}(u)}.$$

[We proved it assuming that  $\bar{L}_{T_\emptyset^+}(u) \geq 1$ , but it is trivially true on the set  $\{\bar{L}_{T_\emptyset^+}(u) = 0\}$ .] In particular, for  $1 \leq k \neq m \leq N(u)$ ,

$$E_\omega[\bar{L}_{T_\emptyset^+}(u^{(k)}) \mid \bar{L}_{T_\emptyset^+}(u)] = e^{-[V(u^{(k)}) - V(u)]} \bar{L}_{T_\emptyset^+}(u), \\ E_\omega[\bar{L}_{T_\emptyset^+}(u^{(k)}) \bar{L}_{T_\emptyset^+}(u^{(m)}) \mid \bar{L}_{T_\emptyset^+}(u)] \\ = e^{-[V(u^{(k)}) - V(u)] - [V(u^{(m)}) - V(u)]} \bar{L}_{T_\emptyset^+}(u) (\bar{L}_{T_\emptyset^+}(u) + 1).$$

Applying again (5.9), this time to  $\xi_1 := \bar{L}_{T_\emptyset^+}(u^{(k)})$ ,  $\xi_2 := \bar{L}_{T_\emptyset^+}(u^{(m)})$  and  $\mathcal{G} := \sigma(\bar{L}_{T_\emptyset^+}(u))$ , we obtain (Var $_\omega$  denoting variance under  $P_\omega$ )

$$\text{Cov}_\omega[\bar{L}_{T_\emptyset^+}(u^{(k)}), \bar{L}_{T_\emptyset^+}(u^{(m)})] = e^{-[V(u^{(k)}) - V(u)] - [V(u^{(m)}) - V(u)]} E_\omega[\bar{L}_{T_\emptyset^+}(u)] \\ + e^{-[V(u^{(k)}) - V(u)] - [V(u^{(m)}) - V(u)]} \text{Var}_\omega[\bar{L}_{T_\emptyset^+}(u)] \\ \leq 2e^{-[V(u^{(k)}) - V(u)] - [V(u^{(m)}) - V(u)]} E_\omega[\bar{L}_{T_\emptyset^+}(u)^2],$$

the last inequality following from the fact that  $\bar{L}_{T_\emptyset^+}(u) \leq \bar{L}_{T_\emptyset^+}(u)^2$  [recalling that  $\bar{L}_{T_\emptyset^+}(u)$  is integer-valued]. We take  $k$  and  $m$  be such that  $u^{(k)} = x_{|u|+1}$  and  $u^{(m)} = y_{|u|+1}$ . In view of (5.11), this yields the desired inequality (5.6) in Lemma 5.2.

It remains to deal with the special case that either  $x$  is an ancestor of  $y$ , or  $y$  is an ancestor of  $x$ .

This, however, is easy. Without loss of generality, let us assume that  $y$  is an ancestor of  $x$ , in which case we have seen that  $E_\omega(\bar{L}_{T_\emptyset^+}(x) \mid \bar{L}_{T_\emptyset^+}(y)) =$

---

negative integers  $m_k$ ,  $1 \leq k \leq N$ , satisfying  $m_1 + \dots + m_N = m$ ; in particular,  $\mathbb{E}(s_1^{\xi_1} \dots s_N^{\xi_N}) = (\sum_{k=1}^N p^{(k)} s_k)^m$ , for all  $s_k \geq 0$ ,  $1 \leq k \leq N$ .

$e^{-[V(x)-V(y)]}\bar{L}_{T_{\varnothing}^+}(y)$ . So applying (5.9) to  $\xi_1 := \bar{L}_{T_{\varnothing}^+}(x)$ ,  $\xi_2 := \bar{L}_{T_{\varnothing}^+}(y)$  and  $\mathcal{G} := \sigma(\bar{L}_{T_{\varnothing}^+}(y))$  gives

$$\text{Cov}_{\omega}[\bar{L}_{T_{\varnothing}^+}(x), \bar{L}_{T_{\varnothing}^+}(y)] = 0 + e^{-[V(x)-V(y)]} \text{Var}_{\omega}[\bar{L}_{T_{\varnothing}^+}(y)],$$

yielding (5.6).

(ii) We already noted that  $E_{\omega}[\bar{L}_{T_{\varnothing}^+}(x)] = e^{-[V(x)-V(x_1)]} E_{\omega}[\bar{L}_{T_{\varnothing}^+}(x_1)]$ , where  $x_1$  denotes, as before, the ancestor of  $x$  in the first generation. Since  $E_{\omega}[\bar{L}_{T_{\varnothing}^+}(x_1)] = \omega(\varnothing, x_1)$ , and by definition,  $\omega(\varnothing, x_1) = \omega(\varnothing, \overleftarrow{\varnothing})e^{-V(x_1)}$ , this yields  $E_{\omega}[\bar{L}_{T_{\varnothing}^+}(x)] = \omega(\varnothing, \overleftarrow{\varnothing})e^{-V(x)}$ , as stated in (5.7).

It remains to compute  $E_{\omega}[\bar{L}_{T_{\varnothing}^+}(x)^2]$ . From (5.12), we get that

$$E_{\omega}[\bar{L}_{T_{\varnothing}^+}(u^{(k)})[\bar{L}_{T_{\varnothing}^+}(u^{(k)}) - 1]|\bar{L}_{T_{\varnothing}^+}(u)] = e^{-2[V(u^{(k)})-V(u)]}\bar{L}_{T_{\varnothing}^+}(u)[\bar{L}_{T_{\varnothing}^+}(u) + 1].$$

Taking expectation on both sides, and replacing the pair  $(u^{(k)}, u)$  by  $(x, \overleftarrow{x})$ , we obtain

$$\begin{aligned} E_{\omega}[\bar{L}_{T_{\varnothing}^+}(x)^2] &= e^{-2[V(x)-V(\overleftarrow{x})]} E_{\omega}[\bar{L}_{T_{\varnothing}^+}(\overleftarrow{x})^2] \\ &\quad + (e^{-2[V(x)-V(\overleftarrow{x})]} + e^{-[V(x)-V(\overleftarrow{x})]}) E_{\omega}[\bar{L}_{T_{\varnothing}^+}(\overleftarrow{x})]. \end{aligned}$$

By the already proved (5.7),  $E_{\omega}[\bar{L}_{T_{\varnothing}^+}(\overleftarrow{x})] = \omega(\varnothing, \overleftarrow{\varnothing})e^{-V(\overleftarrow{x})}$ . Solving this difference equation (with initial condition  $E_{\omega}[\bar{L}_{T_{\varnothing}^+}(x_1)^2] = \omega(\varnothing, x_1) = \omega(\varnothing, \overleftarrow{\varnothing})e^{-V(x_1)}$ ) yields (5.8). This completes the proof of the lemma.  $\square$

**6. Biased walks: Proof of Proposition 2.4.** Let  $P_{\omega}^{(r)}$  denote the quenched law of the biased walk with a reflecting barrier at  $\mathcal{L}_r$ . Under  $P_{\omega}^{(r)}$ , the biased walk  $(X_i)$  is positive recurrent taking values in  $\{x \in \mathbb{T} : x \leq \mathcal{L}_r\} \cup \{\overleftarrow{\varnothing}\}$ , with invariant probability  $\pi_r$  as in (2.11). In particular, if  $T_{\varnothing}^+$  denotes, as in (5.2), the first return time to  $\varnothing$ , and  $L_{T_{\varnothing}^+}$  (site) local time as in (2.14),

$$(6.1) \quad E_{\omega}^{(r)}(T_{\varnothing}^+) = \frac{1}{\pi_r(\varnothing)},$$

$$(6.2) \quad E_{\omega}^{(r)}[L_{T_{\varnothing}^+}(y)] = \frac{\pi_r(y)}{\pi_r(\varnothing)}, \quad y \in \{x \in \mathbb{T} : x \leq \mathcal{L}_r\} \cup \{\overleftarrow{\varnothing}\}.$$

We now proceed to study  $L_n(\varnothing)$  under  $P_{\omega}^{(r)}$ . Let  $\ell \geq 1$ , and let  $T_{\varnothing}^{(\ell)}$  denote the  $\ell$ th return time to  $\varnothing$  (so  $T_{\varnothing}^{(1)}$  is  $T_{\varnothing}^+$ , under  $P_{\omega}^{(r)}$ ). Under  $P_{\omega}^{(r)}$ ,  $T_{\varnothing}^{(\ell)}$  is the sum of  $\ell$  independent copies of  $T_{\varnothing}^+$ . In particular,  $E_{\omega}^{(r)}(T_{\varnothing}^{(\ell)}) = \ell \times E_{\omega}^{(r)}(T_{\varnothing}^+) = \frac{\ell}{\pi_r(\varnothing)}$ .

By the simple relation  $\{L_n(\emptyset) \leq \ell\} = \{T_\emptyset^{(\ell)} \geq n\}$ , we have

$$P_\omega^{(r)}\{L_n(\emptyset) \leq \ell\} = P_\omega^{(r)}\left\{T_\emptyset^{(\ell)} - \frac{\ell}{\pi_r(\emptyset)} \geq n - \frac{\ell}{\pi_r(\emptyset)}\right\},$$

which, by Chebyshev's inequality, is bounded by  $\frac{\ell}{(n-\ell/\pi_r(\emptyset))^2} \text{Var}_\omega^{(r)}(T_\emptyset^+)$  if  $n > \frac{\ell}{\pi_r(\emptyset)}$  ( $\text{Var}_\omega^{(r)}$  denoting the variance under the probability  $P_\omega^{(r)}$ ). However, it has not been clear to us whether  $\text{Var}_\omega^{(r)}(T_\emptyset^+)$  is sufficiently small. This is why some care is in order when applying the method of second moment. We are not going to estimate the variance (under  $P_\omega^{(r)}$ ) of  $T_\emptyset^+$ ; instead, we are going to decompose  $T_\emptyset^+$  into three distinct parts, in such a way that the variance of a part is sufficiently small for our needs and that the expectation of the other parts is also sufficiently small.

Recall from (2.10) that  $\mathcal{L}_r := \{x : \sum_{z \in \llbracket \emptyset, x \rrbracket} e^{V(z)-V(x)} > r, \sum_{z \in \llbracket \emptyset, y \rrbracket} e^{V(z)-V(y)} \leq r, \forall y \in \llbracket \emptyset, x \rrbracket\}$ . The reason for which we have not been able to make  $\text{Var}_\omega^{(r)}(T_\emptyset^+)$  small is that  $r$  is too large. Our solution is to consider *two* scales:  $\mathcal{L}_r$  and  $\mathcal{L}_s$  with  $s := \frac{r}{(\log r)^\theta} \leq r$ , where  $\theta \geq 0$ .

The promised decomposition for  $T_\emptyset^+$  is as follows, the constant  $\delta_1$  being defined in (3.5):

$$(6.3) \quad T_\emptyset^{(a)} := \sum_{y \in \mathbb{T}: y < \mathcal{L}_s} L_{T_\emptyset^+}(y) \mathbf{1}_{\{\min_{u \in \llbracket \emptyset, y \rrbracket} \omega(u, \hat{u}) \geq (\log r)^{-\delta/\delta_1}\}},$$

$$(6.4) \quad T_\emptyset^{(b)} := \sum_{y \in \mathbb{T}: y < \mathcal{L}_s} L_{T_\emptyset^+}(y) \mathbf{1}_{\{\min_{u \in \llbracket \emptyset, y \rrbracket} \omega(u, \hat{u}) < (\log r)^{-\delta/\delta_1}\}},$$

$$(6.5) \quad T_\emptyset^{(c)} := \sum_{y \in \mathbb{T}: \mathcal{L}_s \leq y \leq \mathcal{L}_r} L_{T_\emptyset^+}(y).$$

Then

$$(6.6) \quad T_\emptyset^+ - 1 \leq T_\emptyset^{(a)} + T_\emptyset^{(b)} + T_\emptyset^{(c)} \leq T_\emptyset^+.$$

[The quantities  $T_\emptyset^+$  and  $T_\emptyset^{(a)} + T_\emptyset^{(b)} + T_\emptyset^{(c)}$  can differ by 1 in case  $X_1 = \overleftarrow{\emptyset}$ .]

The next pair of lemmas summarise basic properties of  $T_\emptyset^{(a)}$ ,  $T_\emptyset^{(b)}$  and  $T_\emptyset^{(c)}$  that are needed in this paper: loosely speaking, we control in a satisfying way the first two moments of  $T_\emptyset^{(a)}$ , and although we have no control on the variances of  $T_\emptyset^{(b)}$  and  $T_\emptyset^{(c)}$ , we show that they both have negligible expectations compared to the expectation of  $T_\emptyset^+$ .

LEMMA 6.1. *Let  $\theta \geq 0$  and let  $s := \frac{r}{(\log r)^\theta}$ . When  $r \rightarrow \infty$ ,*

$$(6.7) \quad \frac{E_\omega^{(r)}(T_\emptyset^{(a)})}{E_\omega^{(r)}(T_\emptyset^+)} \rightarrow 1, \quad \text{in } \mathbf{P}^*\text{-probability,}$$



$$(6.8) \quad \frac{E_\omega^{(r)}(T_\emptyset^{(b)})}{E_\omega^{(r)}(T_\emptyset^+)} \rightarrow 0, \quad \text{in } \mathbf{P}^*\text{-probability,}$$

$$(6.9) \quad \frac{E_\omega^{(r)}(T_\emptyset^{(c)})}{E_\omega^{(r)}(T_\emptyset^+)} \rightarrow 0, \quad \text{in } \mathbf{P}^*\text{-probability.}$$

In particular,

$$(6.10) \quad \frac{E_\omega^{(r)}(T_\emptyset^{(b)}) + E_\omega^{(r)}(T_\emptyset^{(c)})}{\log r} \rightarrow 0, \quad \text{in } \mathbf{P}^*\text{-probability.}$$

LEMMA 6.2. Let  $\theta \geq 0$  and let  $s := \frac{r}{(\log r)^\theta}$ . There exists a constant  $c_7 > 0$  such that for all  $r \geq 2$ ,

$$(6.11) \quad \mathbf{E}[\text{Var}_\omega^{(r)}(T_\emptyset^{(a)})] \leq c_7 s (\log r)^{18/\delta_1 + 6},$$

where  $\delta_1 > 0$  is the constant in (3.5).

By admitting Lemmas 6.1 and 6.2 for the time being, we are able to prove Proposition 2.4.

PROOF OF PROPOSITION 2.4. Let  $\theta \geq 0$  and let  $s := \frac{r}{(\log r)^\theta}$ . Let

$$T_\emptyset^{(\ell),(a)} := \sum_{y \in \mathbb{T}: y < \mathcal{L}_s} L_{T_\emptyset^{(\ell)}}(y) \mathbf{1}_{\{\min_{u \in \llbracket \emptyset, y \rrbracket} \omega(u, \overleftarrow{u}) \geq (\log r)^{-6/\delta_1}\}},$$

$$T_\emptyset^{(\ell),(b)} := \sum_{y \in \mathbb{T}: y < \mathcal{L}_s} L_{T_\emptyset^{(\ell)}}(y) \mathbf{1}_{\{\min_{u \in \llbracket \emptyset, y \rrbracket} \omega(u, \overleftarrow{u}) < (\log r)^{-6/\delta_1}\}},$$

$$T_\emptyset^{(\ell),(c)} := \sum_{y \in \mathbb{T}: \mathcal{L}_s \leq y \leq \mathcal{L}_r} L_{T_\emptyset^{(\ell)}}(y).$$

Then  $T_\emptyset^{(\ell)} - \ell \leq T_\emptyset^{(\ell),(a)} + T_\emptyset^{(\ell),(b)} + T_\emptyset^{(\ell),(c)} \leq T_\emptyset^{(\ell)}$ .

For any  $n_1 \geq 1$  and  $n_2 \geq 1$  with  $n_1 + n_2 \leq n - \ell$ ,

$$\begin{aligned} P_\omega^{(r)}\{L_n(\emptyset) \leq \ell\} &= P_\omega^{(r)}\{T_\emptyset^{(\ell)} \geq n\} \\ &\leq P_\omega^{(r)}\{T_\emptyset^{(\ell),(a)} + T_\emptyset^{(\ell),(b)} + T_\emptyset^{(\ell),(c)} \geq n - \ell\} \\ &\leq P_\omega^{(r)}\{T_\emptyset^{(\ell),(a)} \geq n_1\} + P_\omega^{(r)}\{T_\emptyset^{(\ell),(b)} + T_\emptyset^{(\ell),(c)} \geq n_2\}. \end{aligned}$$

Observe that  $E_\omega^{(r)}[T_\emptyset^{(\ell),(b)} + T_\emptyset^{(\ell),(c)}] = \ell[E_\omega^{(r)}(T_\emptyset^{(b)}) + E_\omega^{(r)}(T_\emptyset^{(c)})]$ , so by Markov's inequality,

$$P_\omega^{(r)}\{T_\emptyset^{(\ell),(b)} + T_\emptyset^{(\ell),(c)} \geq n_2\} \leq \frac{\ell}{n_2} [E_\omega^{(r)}(T_\emptyset^{(b)}) + E_\omega^{(r)}(T_\emptyset^{(c)})].$$

For  $P_\omega^{(r)}\{T_\emptyset^{(\ell),(a)} \geq n_1\}$ , we note that  $E_\omega^{(r)}(T_\emptyset^{(\ell),(a)}) = \ell \times E_\omega^{(r)}(T_\emptyset^{(a)}) \leq \ell \times E_\omega^{(r)}(T_\emptyset^+) = \frac{\ell}{\pi_r(\emptyset)}$ , and that  $\text{Var}_\omega^{(r)}(T_\emptyset^{(\ell),(a)}) = \ell \text{Var}_\omega^{(r)}(T_\emptyset^{(a)})$ . If  $n_1 - \frac{\ell}{\pi_r(\emptyset)} > 0$ , then by Chebyshev's inequality,

$$\begin{aligned} P_\omega^{(r)}\{T_\emptyset^{(\ell),(a)} \geq n_1\} &\leq P_\omega^{(r)}\left\{T_\emptyset^{(\ell),(a)} - E_\omega^{(r)}(T_\emptyset^{(\ell),(a)}) \geq n_1 - \frac{\ell}{\pi_r(\emptyset)}\right\} \\ &\leq \frac{\ell \text{Var}_\omega^{(r)}(T_\emptyset^{(a)})}{[n_1 - \ell/\pi_r(\emptyset)]^2}. \end{aligned}$$

Let us now fix the choice for  $\ell$ ,  $n_1$  and  $n_2$ . Let  $0 < \varepsilon < 1$ . We take  $n_1 := \lceil (1 + \varepsilon) \frac{\ell}{\pi_r(\emptyset)} \rceil$  and  $n_2 := \lfloor \varepsilon \frac{\ell}{\pi_r(\emptyset)} \rfloor - \ell - 1$  so that  $n_1 + n_2 \leq (1 + 2\varepsilon) \frac{\ell}{\pi_r(\emptyset)} - \ell$ , which is indeed bounded by  $n - \ell$  if we take  $\ell := \lfloor \frac{1}{1+2\varepsilon} n \pi_r(\emptyset) \rfloor$ . With the choice made for  $(\ell, n_1, n_2)$ , we have

$$P_\omega^{(r)}\{L_n(\emptyset) \leq \ell\} \leq \frac{\ell \text{Var}_\omega^{(r)}(T_\emptyset^{(a)})}{[n_1 - \ell/\pi_r(\emptyset)]^2} + \frac{\ell}{n_2} [E_\omega^{(r)}(T_\emptyset^{(b)}) + E_\omega^{(r)}(T_\emptyset^{(c)})].$$

Recall from Theorem 2.7 that  $(\log r)\pi_r(\emptyset) \rightarrow \frac{\sigma^2}{4D_\infty} e^{-U(\emptyset)}$  in  $\mathbf{P}^*$ -probability (for  $r \rightarrow \infty$ ). We choose  $r = n$  so that we are entitled to apply Lemma 5.1. With the definition of  $s := \frac{r}{(\log r)^\theta}$ , we apply Lemma 6.2 (choosing  $\theta > \frac{18}{\delta_1} + 5$ ) and Lemma 6.1 [part (6.10)], to see that  $P_\omega^{(r)}\{L_n(\emptyset) \leq \ell\} \rightarrow 0$  in  $\mathbf{P}^*$ -probability. By Lemma 5.1,  $\mathbb{P}(\cup_{i=1}^n \{X_i \in \mathcal{L}_r\}) \rightarrow 0$ , so this is equivalent to saying that  $P_\omega\{L_n(\emptyset) \leq \ell\} \rightarrow 0$  in  $\mathbf{P}^*$ -probability, with the choice of  $\ell := \lfloor \frac{1}{1+2\varepsilon} n \pi_r(\emptyset) \rfloor$ . Again, since  $(\log r)\pi_r(\emptyset) \rightarrow \frac{\sigma^2}{4D_\infty} e^{-U(\emptyset)}$  in  $\mathbf{P}^*$ -probability (Theorem 2.7), this yields the lower bound in (2.15).

The proof of the upper bound is similar, with the same choice  $r := n$ , and is slightly easier because we do not need to care about  $T_\emptyset^{(\ell),(b)}$  and  $T_\emptyset^{(\ell),(c)}$  any more. Indeed, for any  $\ell \geq 1$ ,

$$\begin{aligned} P_\omega^{(r)}\{L_n(\emptyset) \geq \ell\} &= P_\omega^{(r)}\{T_\emptyset^{(\ell)} \leq n\} \\ &\leq P_\omega^{(r)}\{T_\emptyset^{(\ell),(a)} \leq n\} \\ &\leq \frac{\text{Var}_\omega^{(r)}(T_\emptyset^{(\ell),(a)})}{[E_\omega^{(r)}(T_\emptyset^{(\ell),(a)}) - n]^2}, \end{aligned}$$

as long as  $E_\omega^{(r)}(T_\emptyset^{(\ell),(a)}) > n$ . Again,  $E_\omega^{(r)}(T_\emptyset^{(\ell),(a)}) = \ell E_\omega^{(r)}(T_\emptyset^{(a)})$ , and  $\text{Var}_\omega^{(r)}(T_\emptyset^{(\ell),(a)}) = \ell \text{Var}_\omega^{(r)}(T_\emptyset^{(a)})$ . This time, with  $\varepsilon > 0$ , our choice is  $\ell := \lfloor (1 + \varepsilon) n \pi_r(\emptyset) \rfloor$ . For this new choice of  $\ell$ , part (6.7) of Lemma 6.1 ensures that  $\mathbf{P}^*\{E_\omega^{(r)}(T_\emptyset^{(\ell),(a)}) > (1 + \frac{\varepsilon}{2})n\} \rightarrow 1$  for  $n \rightarrow \infty$ . So by Lemma 6.2, if  $s := \frac{r}{(\log r)^\theta}$  with  $\theta > \frac{18}{\delta_1} + 5$ , then  $P_\omega^{(r)}\{L_n(\emptyset) \geq \ell\} \rightarrow 0$  in  $\mathbf{P}^*$ -probability, which, in view of

Lemma 5.1, is equivalent to saying that  $P_\omega\{L_n(\emptyset) \geq \ell\} \rightarrow 0$  in  $\mathbf{P}^*$ -probability. This yields (2.15).

It remains to check (2.16). In view of (2.15), it suffices to show the following:

$$(6.12) \quad \left(\frac{\log n}{n}\right)^2 E_\omega[(L_n(\emptyset))^2] \text{ is tight under } \mathbf{P}^*.$$

Clearly,  $E_\omega[(L_n(\emptyset))^2] \leq E_\omega^{(r)}[(L_n(\emptyset))^2]$ , for any  $r > 1$ . Observe that

$$E_\omega^{(r)}[(L_n(\emptyset))^2] \leq 2 \sum_{j=1}^\infty j P_\omega^{(r)}\{L_n(\emptyset) \geq j\} = 2 \sum_{j=1}^\infty j P_\omega^{(r)}\{T_\emptyset^{(j)} \leq n\}.$$

By Chebyshev’s inequality,  $P_\omega^{(r)}\{T_\emptyset^{(j)} \leq n\} \leq e \times E_\omega^{(r)}(e^{-T_\emptyset^{(j)}/n})$ , which, by the strong Markov property, is  $e \times [E_\omega^{(r)}(e^{-T_\emptyset^+/n})]^j$ . As such,

$$E_\omega[(L_n(\emptyset))^2] \leq 2e \sum_{j=1}^\infty j [E_\omega^{(r)}(e^{-T_\emptyset^+/n})]^j \leq \frac{2e}{[1 - E_\omega^{(r)}(e^{-T_\emptyset^+/n})]^2},$$

where, in the last inequality, we used the elementary fact that  $\sum_{j=1}^\infty jx^j = \frac{x}{(1-x)^2} \leq \frac{1}{(1-x)^2}$  for any  $x \in [0, 1]$ .

Note that for any nonnegative random variable  $\xi$  with  $\mathbb{E}(\xi^2) < \infty$ , we have  $\mathbb{E}(1 - e^{-\xi}) \geq \mathbb{E}(\xi - \frac{\xi^2}{2}) = \mathbb{E}(\xi) - \frac{1}{2}[\mathbb{E}(\xi)]^2 - \frac{1}{2} \text{Var}(\xi)$ . Therefore,

$$\begin{aligned} 1 - E_\omega^{(r)}(e^{-T_\emptyset^+/n}) &\geq 1 - E_\omega^{(r)}(e^{-T_\emptyset^{(a)}/n}) \\ &\geq \frac{E_\omega^{(r)}(T_\emptyset^{(a)})}{n} - \frac{[E_\omega^{(r)}(T_\emptyset^{(a)})]^2}{2n^2} - \frac{\text{Var}_\omega^{(r)}(T_\emptyset^{(a)})}{2n^2}. \end{aligned}$$

We choose again  $r := n$ . By Lemma 6.1 and Theorem 2.7,  $E_\omega^{(r)}(T_\emptyset^{(a)}) = (\frac{4}{\sigma^2} D_\infty + op^*(1)) \log r$ , where  $op^*(1)$  denotes, as before, a term converging to 0 in  $\mathbf{P}^*$ -probability and its value may vary from line to line. By Lemma 6.2, if we choose  $s := \frac{r}{(\log r)^\theta}$  with  $\theta > \frac{18}{\delta_1} + 5$ , then  $\frac{\text{Var}_\omega^{(r)}(T_\emptyset^{(a)})}{n \log n} \rightarrow 0$  in  $\mathbf{P}^*$ -probability. This yields (6.12), and thus (2.16). Proposition 2.4 is proved.  $\square$

The rest of the section is devoted to the proof of Lemmas 6.1 and 6.2.

**PROOF OF LEMMA 6.1.** Clearly, (6.10) follows from (6.8) and (6.9) (combined with Theorem 2.7). On the other hand,  $T_\emptyset^{(a)} + T_\emptyset^{(b)} + T_\emptyset^{(c)}$  and  $T_\emptyset^+$  differ by at most 1 [see (6.6)], so (6.8) and (6.9) together imply (6.7). As a consequence, we only need to prove (6.8) and (6.9).

Let us start with the *proof of (6.9)*. By definition of  $T_\emptyset^{(c)}$  [see (6.5)],  $E_\omega^{(r)}(T_\emptyset^{(c)}) = \sum_{\mathcal{L}_s \leq y \leq \mathcal{L}_r} E_\omega^{(r)}[L_{T_\emptyset^+}(y)]$ . We have seen in (6.2) that  $E_\omega^{(r)}[L_{T_\emptyset^+}(y)] = \frac{\pi_r(y)}{\pi_r(\emptyset)}$  for

$y \leq \mathcal{L}_r$ . Since  $E_\omega^{(r)}(T_\emptyset^+) = \frac{1}{\pi_r(\emptyset)}$  [see (6.1)], we have

$$\frac{E_\omega^{(r)}(T_\emptyset^{(c)})}{E_\omega^{(r)}(T_\emptyset^+)} = \sum_{\mathcal{L}_s \leq y \leq \mathcal{L}_r} \pi_r(y) = \sum_{\mathcal{L}_s \leq y < \mathcal{L}_r} \frac{e^{-U(y)}}{Z_r} + \sum_{y \in \mathcal{L}_r} \frac{e^{-V(y)}}{Z_r}.$$

Noting  $e^{-U(y)} = e^{-V(y)} + \sum_{z \in \mathbb{T}: \overleftarrow{z}=y} e^{-V(z)}$ , we arrive at

$$\begin{aligned} \frac{E_\omega^{(r)}(T_\emptyset^{(c)})}{E_\omega^{(r)}(T_\emptyset^+)} &= \sum_{\mathcal{L}_s \leq y < \mathcal{L}_r} \frac{e^{-V(y)}}{Z_r} + \sum_{\mathcal{L}_s < z \leq \mathcal{L}_r} \frac{e^{-V(z)}}{Z_r} + \sum_{y \in \mathcal{L}_r} \frac{e^{-V(y)}}{Z_r} \\ &\leq \frac{2}{Z_r} \sum_{\mathcal{L}_s \leq y \leq \mathcal{L}_r} e^{-V(y)}. \end{aligned}$$

By Theorem 2.7,  $\frac{1}{\log r} \sum_{y \leq \mathcal{L}_r} e^{-V(y)} \rightarrow \frac{2}{\sigma^2} D_\infty$  in  $\mathbf{P}^*$ -probability and  $\frac{1}{\log r} \sum_{y < \mathcal{L}_s} e^{-V(y)} \rightarrow \frac{2}{\sigma^2} D_\infty$  in  $\mathbf{P}^*$ -probability (noting that  $\frac{1}{\log r} \sum_{y \in \mathcal{L}_s} e^{-V(y)} \rightarrow 0$  in  $\mathbf{P}^*$ -probability according to Lemma 3.4). Hence  $\frac{1}{\log r} \sum_{\mathcal{L}_s \leq y \leq \mathcal{L}_r} e^{-V(y)} \rightarrow \frac{2}{\sigma^2} D_\infty - \frac{2}{\sigma^2} D_\infty = 0$  in  $\mathbf{P}^*$ -probability. Since  $\frac{Z_r}{\log r} \rightarrow \frac{4}{\sigma^2} D_\infty > 0$  in  $\mathbf{P}^*$ -probability (Theorem 2.7), we conclude that

$$\frac{2}{Z_r} \sum_{\mathcal{L}_s \leq y \leq \mathcal{L}_r} e^{-V(y)} \rightarrow 0, \quad \text{in } \mathbf{P}^*\text{-probability.}$$

This yields (6.9).

We now turn to the *proof of (6.8)*. Recall that  $E_\omega^{(r)}[L_{T_\emptyset^+}(y)] = \frac{\pi_r(y)}{\pi_r(\emptyset)} = e^{U(\emptyset) - U(y)} = e^{U(\emptyset)} [e^{-V(y)} + \sum_{z \in \mathbb{T}: \overleftarrow{z}=y} e^{-V(z)}]$  for  $y < \mathcal{L}_s$ . By definition of  $T_\emptyset^{(b)}$  in (6.4),

$$E_\omega^{(r)}(T_\emptyset^{(b)}) = e^{U(\emptyset)} \sum_{y < \mathcal{L}_s} \left( e^{-V(y)} + \sum_{z \in \mathbb{T}: \overleftarrow{z}=y} e^{-V(z)} \right) \mathbf{1}_{\{y \text{ bad}\}},$$

where, by “ $y$  bad,” we mean  $\min_{u \in \llbracket \emptyset, y \rrbracket} \omega(u, \overleftarrow{u}) < (\log r)^{-6/\delta_1}$ . So

$$E_\omega^{(r)}(T_\emptyset^{(b)}) \leq 2e^{U(\emptyset)} \sum_{y \leq \mathcal{L}_s} e^{-V(y)} \mathbf{1}_{\{y \text{ bad}\}} \leq 2e^{U(\emptyset)} \sum_{y \leq \mathcal{L}_r} e^{-V(y)} \mathbf{1}_{\{y \text{ bad}\}}.$$

Let  $B > 0$  be a constant. Then

$$E_\omega^{(r)}(T_\emptyset^{(b)}) \leq 2e^{U(\emptyset)} (\Sigma_{(6.13)} + \Sigma_{(6.14)}),$$

where<sup>7</sup>

$$(6.13) \quad \Sigma_{(6.13)} := \sum_{y \in \mathbb{T}: |y| \leq B(\log r)^2} e^{-V(y)} \mathbf{1}_{\{y \text{ bad}\}},$$

<sup>7</sup>For notational convenience, we treat  $B(\log r)^2$  as an integer.

$$(6.14) \quad \Sigma_{(6.14)} := \sum_{y \leq \mathcal{L}_r, |y| > B(\log r)^2} e^{-V(y)}.$$

In view of (4.13), we have, for any  $\varepsilon > 0$ ,

$$(6.15) \quad \lim_{B \rightarrow \infty} \limsup_{r \rightarrow \infty} \mathbf{P}^* \{ \Sigma_{(6.14)} \geq \varepsilon \log r \} = 0.$$

We now bound  $\Sigma_{(6.13)}$ . When  $y$  is bad,  $\max_{u \in \llbracket \emptyset, y \rrbracket} \frac{1}{\omega(u, \overleftarrow{u})} > (\log r)^{6/\delta_1}$ , which means  $\max_{u \in \llbracket \emptyset, y \rrbracket} \Lambda(u) > (\log r)^{6/\delta_1} - 1$ , with  $\Lambda(u) := \sum_{z \in \mathbb{T}: \overleftarrow{z} = u} e^{-[V(z) - V(u)]}$  as in (3.3). Accordingly, writing  $a(r) := (\log r)^{6/\delta_1} - 1$  for brevity,

$$\begin{aligned} \Sigma_{(6.13)} &\leq \sum_{y \in \mathbb{T}: |y| \leq B(\log r)^2} e^{-V(y)} \mathbf{1}_{\{\max_{u \in \llbracket \emptyset, y \rrbracket} \Lambda(u) > a(r)\}} \\ &\leq 1 + \sum_{k=1}^{B(\log r)^2} \sum_{y \in \mathbb{T}: |y|=k} e^{-V(y)} (\mathbf{1}_{\{\max_{u \in \llbracket \emptyset, y \rrbracket} \Lambda(u) > a(r)\}} + \mathbf{1}_{\{\Lambda(y) > a(r)\}}), \end{aligned}$$

the first term “1” on the right-hand side resulting from  $y = \emptyset$ . We take expectation with respect to  $\mathbf{P}$  on both sides. By Lemma 3.1 and in its notation,

$$\mathbf{E}(\Sigma_{(6.13)}) \leq 1 + \sum_{k=1}^{B(\log r)^2} \mathbf{E} \left[ \mathbf{1}_{\{\max_{1 \leq i \leq k} \tilde{\Lambda}_{i-1} > a(r)\}} + \mathbf{P} \left( \sum_{x: |x|=1} e^{-V(x)} > a(r) \right) \right].$$

Since  $\tilde{\Lambda}_{i-1}$  (for  $i \geq 1$ ) is distributed as  $\tilde{\Lambda}_0$ , we have, for all  $b > 0$  and all  $i \geq 1$ ,  $\mathbf{P}(\tilde{\Lambda}_{i-1} > b) \leq c_8 b^{-\delta_1}$ , where  $\delta_1 > 0$  is the constant in (3.5), and  $c_8 := \mathbf{E}[(\tilde{\Lambda}_0)^{\delta_1}] = \mathbf{E}[(\sum_{x: |x|=1} e^{-V(x)})^{1+\delta_1}]$  which is finite according to (3.5). On the other hand, (3.5) also yields  $\mathbf{P}(\sum_{x: |x|=1} e^{-V(x)} > b) \leq c_8 b^{-(1+\delta_1)}$  (for  $b > 0$ ). Hence,

$$\mathbf{E}(\Sigma_{(6.13)}) \leq 1 + B(\log r)^2 [B(\log r)^2 c_8 a(r)^{-\delta_1} + c_8 a(r)^{-(1+\delta_1)}].$$

This yields  $\frac{\Sigma_{(6.13)}}{\log r} \rightarrow 0$  in  $L^1(\mathbf{P})$  and equivalently, in  $L^1(\mathbf{P}^*)$ , and a fortiori in  $\mathbf{P}^*$ -probability. Together with (6.15), and since  $E_\omega^{(r)}(T_\emptyset^{(b)}) \leq 2e^{U(\emptyset)}(\Sigma_{(6.13)} + \Sigma_{(6.14)})$ , we obtain  $\frac{E_\omega^{(r)}(T_\emptyset^{(b)})}{\log r} \rightarrow 0$  in  $\mathbf{P}^*$ -probability. Recalling that  $E_\omega^{(r)}(T_\emptyset^+) = \frac{1}{\pi_r(\emptyset)}$  and that  $(\log r)\pi_r(\emptyset)$  converges in  $\mathbf{P}^*$ -probability to a positive limit (Theorem 2.7), we deduce that

$$\frac{E_\omega^{(r)}(T_\emptyset^{(b)})}{E_\omega^{(r)}(T_\emptyset^+)} \rightarrow 0, \quad \text{in } \mathbf{P}^*\text{-probability,}$$

which is the desired conclusion in (6.8). Lemma 6.1 is proved.  $\square$

PROOF OF LEMMA 6.2. Recall that  $T_\emptyset^{(a)} := \sum_{y < \mathcal{L}_s} L_{T_\emptyset^+}(y) \mathbf{1}_{\{y \text{ good}\}}$ , where

$$\{y \text{ good}\} := \left\{ \min_{u \in \llbracket \emptyset, y \rrbracket} \omega(u, \overleftarrow{u}) \geq (\log r)^{-6/\delta_1} \right\},$$

with  $\delta_1 > 0$  denoting the constant in (3.5). For any  $y < \mathcal{L}_s$ , we have  $y < \mathcal{L}_r$ , so  $T_\emptyset^{(a)}$  has the same distribution under  $P_\omega^{(r)}$  and under  $P_\omega$ . In particular,

$$\text{Var}_\omega^{(r)}(T_\emptyset^{(a)}) = \text{Var}_\omega(T_\emptyset^{(a)}) = \text{Var}_\omega\left(\sum_{y < \mathcal{L}_s} L_{T_\emptyset^+}(y) \mathbf{1}_{\{y \text{ good}\}}\right).$$

Let  $\bar{L}_k(x) := \sum_{i=1}^k \mathbf{1}_{\{X_{i-1}=\bar{x}, X_i=x\}}$  be edge local time as in (5.5). Then

$$\begin{aligned} \sum_{y < \mathcal{L}_s} L_{T_\emptyset^+}(y) \mathbf{1}_{\{y \text{ good}\}} &= \sum_{y < \mathcal{L}_s} \bar{L}_{T_\emptyset^+}(y) \mathbf{1}_{\{y \text{ good}\}} \\ &\quad + \sum_{y \leq \mathcal{L}_s, y \neq \emptyset} \bar{L}_{T_\emptyset^+}(y) \mathbf{1}_{\{\bar{y} \text{ good}\}}. \end{aligned}$$

By the elementary inequality  $\text{Var}(\xi_1 + \xi_2) \leq 2[\text{Var}(\xi_1) + \text{Var}(\xi_2)]$  (for random variables  $\xi_1$  and  $\xi_2$  having finite second moments), this leads to

$$\begin{aligned} \text{Var}_\omega^{(r)}(T_\emptyset^{(a)}) &\leq 2 \text{Var}_\omega\left(\sum_{y < \mathcal{L}_s} \bar{L}_{T_\emptyset^+}(y) \mathbf{1}_{\{y \text{ good}\}}\right) \\ &\quad + 2 \text{Var}_\omega\left(\sum_{y \leq \mathcal{L}_s, y \neq \emptyset} \bar{L}_{T_\emptyset^+}(y) \mathbf{1}_{\{\bar{y} \text{ good}\}}\right). \end{aligned}$$

We write

$$\begin{aligned} &\text{Var}_\omega\left(\sum_{y \leq \mathcal{L}_s, y \neq \emptyset} \bar{L}_{T_\emptyset^+}(y) \mathbf{1}_{\{\bar{y} \text{ good}\}}\right) \\ &= \sum_{y \leq \mathcal{L}_s, y \neq \emptyset} \text{Var}_\omega[\bar{L}_{T_\emptyset^+}(y)] \mathbf{1}_{\{\bar{y} \text{ good}\}} \\ &\quad + \sum_{y \neq z \leq \mathcal{L}_s, y, z \neq \emptyset} \text{Cov}_\omega[\bar{L}_{T_\emptyset^+}(y), \bar{L}_{T_\emptyset^+}(z)] \mathbf{1}_{\{\bar{y} \text{ good}\}} \mathbf{1}_{\{\bar{z} \text{ good}\}}, \end{aligned}$$

and we have a similar expression for  $\text{Var}_\omega(\sum_{y < \mathcal{L}_s} \bar{L}_{T_\emptyset^+}(y) \mathbf{1}_{\{y \text{ good}\}})$ . Lemma 6.2 will be a straightforward consequence of the following inequalities: for some constants  $c_9 > 0$  and  $c_{10} > 0$ , and all  $r \geq 2$ ,

$$(6.16) \quad \mathbf{E}\left(\sum_{y \leq \mathcal{L}_s} E_\omega[\bar{L}_{T_\emptyset^+}(y)^2] \mathbf{1}_{\{\bar{y} \text{ good}\}}\right) \leq c_9 s (\log r)^{6/\delta_1+2},$$

$$(6.17) \quad \mathbf{E}\left(\sum_{y \neq z \leq \mathcal{L}_s} (\text{Cov}_\omega[\bar{L}_{T_\emptyset^+}(y), \bar{L}_{T_\emptyset^+}(z)])^+ \mathbf{1}_{\{\bar{y} \text{ good}\}}\right) \leq c_{10} s (\log r)^{18/\delta_1+6},$$

where  $\delta_1 > 0$  is in (3.5), and  $(\text{Cov}_\omega[\dots])^+$  denotes the positive part of  $\text{Cov}_\omega[\dots]$ .

So it remains to check inequalities (6.16) and (6.17). We start with the *proof of (6.16)*. Recall from Lemma 5.2 that

$$\begin{aligned} E_\omega[\bar{L}_{T_\emptyset^+}(y)^2] &= \omega(\emptyset, \overleftarrow{\emptyset})e^{-V(y)}\left(2 \sum_{z \in \llbracket \emptyset, y \rrbracket} e^{V(z)-V(y)} - 1\right) \\ &\leq 2e^{-V(y)} \sum_{z \in \llbracket \emptyset, y \rrbracket} e^{V(z)-V(y)}. \end{aligned}$$

For the sum on the right-hand side, we write ( $\overleftarrow{y}$  denoting as before the parent of  $\overleftarrow{y}$ )

$$\begin{aligned} \sum_{z \in \llbracket \emptyset, y \rrbracket} e^{V(z)-V(y)} &= e^{-[V(y)-V(\overleftarrow{y})]} \sum_{z \in \llbracket \emptyset, \overleftarrow{y} \rrbracket} e^{V(z)-V(\overleftarrow{y})} + 1 \\ &= \frac{\omega(\overleftarrow{y}, y)}{\omega(\overleftarrow{y}, \overleftarrow{y})} \sum_{z \in \llbracket \emptyset, \overleftarrow{y} \rrbracket} e^{V(z)-V(\overleftarrow{y})} + 1 \\ &\leq \frac{1}{\omega(\overleftarrow{y}, \overleftarrow{y})} \sum_{z \in \llbracket \emptyset, \overleftarrow{y} \rrbracket} e^{V(z)-V(\overleftarrow{y})} + 1. \end{aligned}$$

If  $y \leq \mathcal{L}_s$ , then by definition,  $\overleftarrow{y} < \mathcal{L}_s$ , so that  $\sum_{z \in \llbracket \emptyset, \overleftarrow{y} \rrbracket} e^{V(z)-V(\overleftarrow{y})} \leq s$ . On the other hand, if  $\overleftarrow{y}$  is good, then by definition,  $\frac{1}{\omega(\overleftarrow{y}, \overleftarrow{y})} \leq (\log r)^{6/\delta_1}$ . Consequently,

$$(6.18) \quad E_\omega[\bar{L}_{T_\emptyset^+}(y)^2] \mathbf{1}_{\{\overleftarrow{y} \text{ good}\}} \mathbf{1}_{\{y \leq \mathcal{L}_s\}} \leq 2[s(\log r)^{6/\delta_1} + 1]e^{-V(y)}.$$

Since  $\mathbf{E}(\sum_{y \leq \mathcal{L}_s} e^{-V(y)}) \leq c_2(\log s)^2$  [see (3.8)], this yields (6.16).

We now turn to the *proof of (6.17)*. Consider a pair  $y \neq z \leq \mathcal{L}_s$ . By Lemma 5.2,

$$\text{Cov}_\omega[\bar{L}_{T_\emptyset^+}(y), \bar{L}_{T_\emptyset^+}(z)] \leq 2e^{-[V(y)-V(y \wedge z)]-[V(z)-V(y \wedge z)]} E_\omega[\bar{L}_{T_\emptyset^+}(y \wedge z)^2].$$

Hence, writing  $\text{LHS}_{(6.17)} := \sum_{y \neq z \leq \mathcal{L}_s} (\text{Cov}_\omega[\bar{L}_{T_\emptyset^+}(y), \bar{L}_{T_\emptyset^+}(z)]^+ \mathbf{1}_{\{\overleftarrow{y} \text{ good}\}})$ , we have

$$\text{LHS}_{(6.17)} \leq 2 \sum_{u < \mathcal{L}_s} \sum_{y \neq z \leq \mathcal{L}_s: y \wedge z = u} e^{-[V(y)-V(u)]-[V(z)-V(u)]} E_\omega[\bar{L}_{T_\emptyset^+}(u)^2] \mathbf{1}_{\{u \text{ good}\}}.$$

Observe that  $\mathbf{1}_{\{u \text{ good}\}} \leq \mathbf{1}_{\{\overleftarrow{u} \text{ good}\}}$ , so by (6.18),  $E_\omega[\bar{L}_{T_\emptyset^+}(u)^2] \mathbf{1}_{\{u \text{ good}\}} \leq 2[s(\log r)^{6/\delta_1} + 1]e^{-V(u)}$ . Consequently,

$$(6.19) \quad \begin{aligned} \text{LHS}_{(6.17)} &\leq 4[s(\log r)^{6/\delta_1} + 1] \sum_{u < \mathcal{L}_s} e^{-V(u)} \mathbf{1}_{\{u \text{ good}\}} \\ &\times \sum_{y \neq z \leq \mathcal{L}_s: y \wedge z = u} e^{-[V(y)-V(u)]-[V(z)-V(u)]}. \end{aligned}$$

Let us consider the double sum  $\sum_{y \neq z \leq \mathcal{L}_s: y \wedge z = u} e^{-[V(y)-V(u)]-[V(z)-V(u)]}$  on the right-hand side. Write  $k = k(u) := |u|$  for brevity. Then

$$\begin{aligned} & \sum_{y \neq z \leq \mathcal{L}_s: y \wedge z = u} e^{-[V(y)-V(u)]-[V(z)-V(u)]} \\ &= \sum_{a \neq b, \overleftarrow{a} = u = \overleftarrow{b}} e^{-[V(a)-V(u)]-[V(b)-V(u)]} \\ & \quad \times \sum_{y, z \leq \mathcal{L}_s: y_{k+1} = a, z_{k+1} = b} e^{-[V(y)-V(a)]-[V(z)-V(b)]}. \end{aligned}$$

Observe that if  $y \leq \mathcal{L}_s$  is such that  $|y| \geq k + 1$  and  $y_{k+1} = a$ , then by definition of  $\mathcal{L}_s$  in (2.10),  $\sum_{w \in \llbracket a, v \rrbracket} e^{V(w)-V(v)} \leq s, \forall v \in \llbracket a, y \rrbracket$ ; so writing  $y = a\tilde{y}$  (the concatenation of  $a$  and  $\tilde{y}$ ), then  $\tilde{y}$  as a vertex of the subtree rooted at  $a$  satisfies  $\tilde{y} \leq \mathcal{L}_s(a)$ , where  $\mathcal{L}_s(a)$  is defined as  $\mathcal{L}_s$ , but associated with the subtree rooted at vertex  $a$ . Accordingly, with  $\mathcal{F}_{k+1}$  denoting the  $\sigma$ -field generated by  $(V(x), |x| \leq k + 1)$ , we have that on the set  $\{|u| = k\}$ ,

$$\begin{aligned} & \mathbf{E} \left( \sum_{y \neq z \leq \mathcal{L}_s: y \wedge z = u} e^{-[V(y)-V(u)]-[V(z)-V(u)]} \middle| \mathcal{F}_{k+1} \right) \\ & \leq \sum_{a \neq b, \overleftarrow{a} = u = \overleftarrow{b}} e^{-[V(a)-V(u)]-[V(b)-V(u)]} [\mathbf{E}(Y_s)]^2. \end{aligned}$$

If  $u$  is good, then by definition,  $\omega(u, \overleftarrow{u}) \geq (\log r)^{-6/\delta_1}$ ; in particular,

$$\sum_{a \in \mathbb{T}: \overleftarrow{a} = u} e^{-[V(a)-V(u)]} = \frac{1}{\omega(u, \overleftarrow{u})} - 1 \leq \frac{1}{\omega(u, \overleftarrow{u})} \leq (\log r)^{6/\delta_1},$$

which implies that  $\sum_{a \neq b, \overleftarrow{a} = u = \overleftarrow{b}} e^{-[V(a)-V(u)]-[V(b)-V(u)]} \leq (\log r)^{12/\delta_1}$ . Hence on the set  $\{|u| = k\}$ ,

$$\begin{aligned} & \mathbf{E} \left( \sum_{y \neq z \leq \mathcal{L}_s: y \wedge z = u} e^{-[V(y)-V(u)]-[V(z)-V(u)]} \middle| \mathcal{F}_{k+1} \right) \mathbf{1}_{\{u \text{ good}\}} \\ & \leq (\log r)^{12/\delta_1} [\mathbf{E}(Y_s)]^2. \end{aligned}$$

Going back to (6.19), this yields

$$\mathbf{E}(\text{LHS}_{(6.17)}) \leq \mathbf{E} \left( 4[s(\log r)^{6/\delta_1} + 1](\log r)^{12/\delta_1} [\mathbf{E}(Y_s)]^2 \sum_{u < \mathcal{L}_s} e^{-V(u)} \right).$$

Since  $\sum_{u < \mathcal{L}_s} e^{-V(u)} \leq \sum_{u \leq \mathcal{L}_s} e^{-V(u)} = Y_s$ , we obtain:  $\mathbf{E}(\text{LHS}_{(6.17)}) \leq 4[s(\log r)^{6/\delta_1} + 1](\log r)^{12/\delta_1} [\mathbf{E}(Y_s)]^3$ . In view of (3.8), this yields (6.17), and completes the proof of Lemma 6.2.  $\square$



**7. Biased walks: Proof of Theorem 2.8.** Recall from (5.4) that

$$P_\omega \left( \bigcup_{i=1}^n \{X_i \in \mathcal{L}_r\} \right) \leq \frac{E_\omega(L_n(\emptyset) + 1)}{r} \sum_{x \in \mathcal{L}_r} e^{-V(x)}.$$

By Lemma 3.4,  $(\log r) \sum_{x \in \mathcal{L}_r} e^{-V(x)}$  is tight under  $\mathbf{P}^*$ . Theorem 2.8 follows from (2.16) of Proposition 2.4.

**8. Biased walks: Proof of Proposition 2.5.** We begin with a general fact for reversible Markov chains. The fact is well known. For a simple proof for finite chains, see Saloff-Coste ([39], Lemma 1.3.3 (1), page 323), applied to  $P^2$ .

FACT 8.1. Let  $P$  be the transition probability of a reversible Markov chain taking values in a countable space  $E$ . Then for any  $x \in E$ , the sequence  $k \rightarrow P^{2k}(x, x)$  is non-increasing.

We prepare for the proof of Proposition 2.5. Let  $P_\omega^{(r)}$  be, as before, the quenched probability with a reflecting barrier at  $\mathcal{L}_r$ , and  $E_\omega^{(r)}$  the corresponding expectation.

LEMMA 8.2. Let  $\gamma \in \mathbb{R}$ , and let  $r = r(n) := \frac{n}{(\log n)^\gamma}$ . Then

$$(\log n)^{2-\gamma} \sup_{B \in \sigma\{X_1, \dots, X_n\}} |P_\omega^{(r)}(B) - P_\omega(B)|$$

is tight under  $\mathbf{P}^*$ .

PROOF. For  $B \in \sigma\{X_1, \dots, X_n\}$ ,

$$|P_\omega^{(r)}(B) - P_\omega(B)| \leq P_\omega \left( \bigcup_{i=1}^n \{X_i \in \mathcal{L}_r\} \right),$$

which is bounded by  $\frac{E_\omega(L_n(\emptyset)+1)}{r} \sum_{x \in \mathcal{L}_r} e^{-V(x)}$  [see (5.4)]. We conclude by means of Lemma 3.4 and (2.16) of Proposition 2.4.  $\square$

We are now ready to prove Proposition 2.5.

PROOF OF PROPOSITION 2.5. We choose  $r := n$  so that we are entitled to apply Lemma 8.2 (with  $\gamma = 0$ ). We claim that for any  $a_n \rightarrow \infty$  satisfying  $\lim_{n \rightarrow \infty} \frac{\log a_n}{\log n} = 0$ ,

$$(8.1) \quad \max_{k \text{ even: } n/a_n \leq k \leq n} \left| (\log n) P_\omega^{(r)}(X_k = \emptyset) - \frac{\sigma^2}{2D_\infty} e^{-U(\emptyset)} \right| \rightarrow 0,$$

in  $\mathbf{P}^*$ -probability.

By Lemma 8.2, (8.1) will imply Proposition 2.5.

Let  $m := m_n$  be the smallest even number such that  $m \geq \frac{n}{a_n}$ . Clearly  $\frac{\log m}{\log n} \rightarrow 1$ . Using the trivial upper bound  $\frac{L_m(\emptyset)}{m} \leq 1$ , we deduce from part (2.16) of Proposition 2.4 and Lemma 8.2 that for  $n \rightarrow \infty$ ,

$$(8.2) \quad E_\omega^{(r)}\left(\frac{L_m(\emptyset)}{m/\log m}\right) \rightarrow \frac{\sigma^2}{4D_\infty}e^{-U(\emptyset)}, \quad \text{in } \mathbf{P}^*\text{-probability.}$$

By Fact 8.1,  $i \mapsto P_\omega^{(r)}(X_{2i} = \emptyset)$  is non-increasing, so  $E_\omega^{(r)}\left(\frac{L_m(\emptyset)}{m/\log m}\right) = \frac{\log m}{m} \sum_{i=1}^m P_\omega^{(r)}(X_i = \emptyset) \geq \frac{1}{2}(\log m)P_\omega^{(r)}(X_m = \emptyset)$ , the factor  $\frac{1}{2}$  coming from the fact we sum over even numbers  $i \in [1, m]$ . Combined with (8.2), we see that

$$(8.3) \quad \begin{aligned} \max_{k \text{ even: } n/a_n \leq k \leq n} (\log n) P_\omega^{(r)}(X_k = \emptyset) &= (\log n) P_\omega^{(r)}(X_m = \emptyset) \\ &\leq \frac{\sigma^2 + \text{op}^*(1)}{2D_\infty}e^{-U(\emptyset)}, \end{aligned}$$

where  $\text{op}^*(1)$  denotes a quantity which goes to 0 in  $\mathbf{P}^*$ -probability as  $n \rightarrow \infty$ .

To obtain the lower bound for  $P_\omega^{(r)}(X_k = \emptyset)$ , we consider the Markov chain  $(X_{2i}, i \geq 0)$  under  $P_\omega^{(r)}$ , starting from  $X_0 := \emptyset$ . This chain takes values in  $E_r := \{x \in \mathbb{T} : x \leq \mathcal{L}_r, |x| \text{ even}\}$ , with  $\pi_r(E_r) = \frac{1}{2}$  due to periodicity. In other words,  $2\pi_r(x)$  for  $x \in E_r$ , is the invariant probability measure of  $(X_{2i}, i \geq 0)$ . By Fact 8.1, we see that for integer  $i \geq 0$ ,  $P_\omega^{(r)}(X_{2i} = \emptyset) \geq 2\pi_r(\emptyset)$ . In particular, for  $k := 2\lfloor \frac{n}{2} \rfloor$ ,  $P_\omega^{(r)}(X_k = \emptyset) \geq 2\pi_r(\emptyset) = \frac{2}{Z_r}e^{-U(\emptyset)}$ . As such, (8.1) follows from Theorem 2.7 and (8.3).  $\square$

REMARK 8.3. By definition,  $\frac{1}{\pi_r(\emptyset)} = Z_r e^{U(\emptyset)}$ , which is  $\frac{4+\text{op}^*(1)}{\sigma^2} D_\infty e^{U(\emptyset)} \times \log n$  according to Theorem 2.7, where  $\text{op}^*(1) \rightarrow 0$  in  $\mathbf{P}^*$ -probability as  $n \rightarrow \infty$ . So (8.1) can also be stated as follows: For any  $a_n \rightarrow \infty$  such that  $\lim_{n \rightarrow \infty} \frac{\log a_n}{\log n} = 0$ , uniformly in even integers  $k \in [\frac{n}{a_n}, n]$ ,

$$(8.4) \quad P_\omega^{(n)}(X_k = \emptyset) = (2 + \text{op}^*(1))\pi_n(\emptyset).$$

This will be useful in the proof of Theorem 2.1 in Section 9.

**9. Biased walks: Proofs of Lemma 2.2, Theorem 2.1 and Corollary 2.3.**

PROOF OF LEMMA 2.2. For  $0 < u \leq r$  and  $x \in \mathbb{T} \cup \{\overleftarrow{\emptyset}\}$ ,

$$\begin{aligned} |\pi_r(x) - \pi_u(x)| &= \left| \frac{Z_r \pi_r(x) - Z_u \pi_u(x)}{Z_u} - Z_r \pi_r(x) \left( \frac{1}{Z_u} - \frac{1}{Z_r} \right) \right| \\ &\leq \frac{Z_r \pi_r(x) - Z_u \pi_u(x)}{Z_u} + Z_r \pi_r(x) \left( \frac{1}{Z_u} - \frac{1}{Z_r} \right), \end{aligned}$$

by using the facts that  $Z_r \pi_r(x) \geq Z_u \pi_u(x)$  and  $Z_r \geq Z_u$ . By taking the summation on  $x$ ,

$$2d_{\text{TV}}(\pi_u, \pi_r) \leq \frac{Z_r - Z_u}{Z_u} + Z_r \left( \frac{1}{Z_u} - \frac{1}{Z_r} \right) = 2 \frac{Z_r - Z_u}{Z_u},$$

which is bounded by  $2 \frac{Z_r - Z_r/(\log r)^a}{Z_r/(\log r)^a}$  if  $u \in [\frac{r}{(\log r)^a}, r]$ . By Theorem 2.7,  $\frac{1}{\log r} (Z_r - Z_r/(\log r)^a) \rightarrow 0$  in  $\mathbf{P}^*$ -probability, from which Lemma 2.2 follows.  $\square$

PROOF OF THEOREM 2.1. (i) Case  $\kappa = 1$ . We prove the following stronger statement: Fix  $0 < c < 1$ . As  $n \rightarrow \infty$ ,

$$(9.1) \quad \max_{\lfloor cn \rfloor \leq m \leq n} \sup_{A \subset \mathbb{T} \cup \{\emptyset\}} |P_\omega(X_m \in A) - \tilde{\pi}_m(A)| \rightarrow 0, \quad \text{in } \mathbf{P}^*\text{-probability.}$$

The fact that (9.1) holds uniformly in  $m$  will be useful in the proof for the case  $\kappa \geq 2$ .

In view of Lemma 8.2, it suffices to prove (9.1) for  $P_\omega^{(r)}$  in lieu of  $P_\omega$ , with  $r := n$ .

Let  $n$  be large and put  $b_n := \lfloor \frac{n}{(\log n)^2} \rfloor$ . For  $cn \leq m \leq n$  (we treat  $cn$  as an integer) and  $A \subset \mathbb{T} \cup \{\emptyset\}$ , we have

$$(9.2) \quad 0 \leq P_\omega^{(r)}(X_m \in A) - \sum_{k=b_n}^m P_\omega^{(r)}(X_m \in A, \mathfrak{g}_m = k) \leq P_\omega^{(r)}\{L_{cn}(\emptyset) < b_n\},$$

where  $\mathfrak{g}_m := \max\{i \leq m : X_i = \emptyset\}$  is the last return time to  $\emptyset$  before  $m$  and the second inequality follows from the fact that  $\{\mathfrak{g}_m \leq b_n\} \subset \{L_m(\emptyset) \leq b_n\} \subset \{L_{cn}(\emptyset) \leq b_n\}$ .

For any  $\varepsilon > 0$ , we have  $b_n < \varepsilon \frac{cn}{\log(cn)}$  for sufficiently large  $n$ ; so

$$\begin{aligned} &P_\omega\{L_{cn}(\emptyset) < b_n\} \\ &\leq P_\omega\left(\left| \frac{L_{cn}(\emptyset)}{(cn)/\log(cn)} - \frac{\sigma^2}{4D_\infty} e^{-U(\emptyset)} \right| > \varepsilon\right) + \mathbf{1}_{\{\sigma^2/(4D_\infty)e^{-U(\emptyset)} \leq 2\varepsilon\}}. \end{aligned}$$

Applying Proposition 2.4, and since  $\varepsilon > 0$  can be as small as possible, we see that  $P_\omega\{L_{cn}(\emptyset) < b_n\} \rightarrow 0$  in  $\mathbf{P}^*$ -probability. A fortiori,  $P_\omega^{(r)}\{L_{cn}(\emptyset) < b_n\} \rightarrow 0$  in  $\mathbf{P}^*$ -probability. Going back to (9.2), we obtain

$$(9.3) \quad P_\omega^{(r)}(X_m \in A) - \sum_{k=b_n}^m P_\omega^{(r)}(X_m \in A, \mathfrak{g}_m = k) \rightarrow 0, \quad \text{in } \mathbf{P}^*\text{-probability,}$$

uniformly in  $A \subset \mathbb{T} \cup \{\emptyset\}$  and in  $m \in [cn, n] \cap \mathbb{Z}$ .

Let us deal with the sum on the left-hand side of (9.3). By the Markov property at time  $k$ ,

$$P_\omega^{(r)}(X_m \in A, \mathfrak{g}_m = k) = P_\omega^{(r)}(X_k = \emptyset) P_\omega^{(r)}(X_{m-k} \in A, m - k < T_\emptyset^+),$$

where  $T_\emptyset^+$  denotes, as before, the first return time to  $\emptyset$ .

By (8.4), uniformly in even numbers  $k \in [b_n, n]$ ,  $P_\omega^{(r)}(X_k = \emptyset) = (2 + o_{\mathbf{P}^*}(1))\pi_r(\emptyset)$ . It follows that uniformly in  $A$  and in  $m$ ,

$$P_\omega^{(r)}(X_m \in A) - 2\pi_r(\emptyset) \sum_{k=b_n, k \text{ even}}^m P_\omega^{(r)}(X_{m-k} \in A, m-k < T_\emptyset^+) \rightarrow 0,$$

in  $\mathbf{P}^*$ -probability. If  $m$  is even, so is  $m - k$ , then we can restrict  $A$  to  $A \cap \mathbb{T}^{(\text{even})}$ . A similar restriction holds if  $m$  is odd. Define

$$A_m := \begin{cases} A \cap \mathbb{T}^{(\text{even})}, & \text{if } m \text{ is even,} \\ A \cap (\mathbb{T}^{(\text{odd})} \cup \{\overleftarrow{\emptyset}\}), & \text{if } m \text{ is odd.} \end{cases}$$

We have [with  $o_{\mathbf{P}^*}(1)$  denoting an expression tending to 0 in  $\mathbf{P}^*$ -probability, uniformly in  $A$  and in  $m$ ]

$$\begin{aligned} P_\omega^{(r)}(X_m \in A) &= 2\pi_r(\emptyset) \sum_{k=b_n}^m P_\omega^{(r)}(X_{m-k} \in A_m, m-k < T_\emptyset^+) + o_{\mathbf{P}^*}(1) \\ (9.4) \quad &= 2\pi_r(\emptyset) \sum_{i=0}^{m-b_n} P_\omega^{(r)}(X_i \in A_m, i < T_\emptyset^+) + o_{\mathbf{P}^*}(1), \end{aligned}$$

which implies that

$$\begin{aligned} P_\omega^{(r)}(X_m \in A) &\leq 2\pi_r(\emptyset) E_\omega^{(r)} \left[ \sum_{i=0}^{T_\emptyset^+-1} \mathbf{1}_{\{X_i \in A_m\}} \right] + o_{\mathbf{P}^*}(1) \\ (9.5) \quad &= 2\pi_r(A_m) + o_{\mathbf{P}^*}(1), \end{aligned}$$

by the fact that  $\pi_r(\emptyset) = \frac{1}{E_\omega^{(r)}(T_\emptyset^+)}$ . By Lemma 2.2,  $\pi_r(A_m) = \pi_m(A_m) + o_{\mathbf{P}^*}(1)$ .

Since  $2\pi_m(A_m) = \tilde{\pi}_m(A)$ , we obtain that  $P_\omega^{(r)}(X_m \in A) \leq \tilde{\pi}_m(A) + o_{\mathbf{P}^*}(1)$ .

To get (9.1), it remains to check that  $P_\omega^{(r)}(X_m \in A) \geq \tilde{\pi}_m(A) + o_{\mathbf{P}^*}(1)$ , which will be done if we are able to reverse the inequality in (9.5). By (9.4) and tightness of  $(\log n)\pi_r(\emptyset)$ , it suffices to prove that

$$(9.6) \quad \frac{1}{\log n} \sum_{i=m-b_n+1}^\infty P_\omega^{(r)}(i < T_\emptyset^+) \rightarrow 0, \quad \text{in } \mathbf{P}^*\text{-probability.}$$

Of course,  $\sum_{i=m-b_n+1}^\infty P_\omega^{(r)}(i < T_\emptyset^+) = E_\omega^{(r)}[(T_\emptyset^+ - (m - b_n + 1))^+]$ . By (6.6) and in its notation (with  $s := \frac{r}{(\log r)^\theta}$  and  $\theta \geq 0$ ),  $T_\emptyset^+ \leq T_\emptyset^{(a)} + T_\emptyset^{(b)} + T_\emptyset^{(c)} + 1$ ; so  $\sum_{i=m-b_n+1}^\infty P_\omega^{(r)}(i < T_\emptyset^+) \leq E_\omega^{(r)}[(T_\emptyset^{(a)} - (m - b_n))^+] + E_\omega^{(r)}[T_\emptyset^{(b)}] + E_\omega^{(r)}[T_\emptyset^{(c)}]$ . Lemma 6.1 entails that  $E_\omega^{(r)}[T_\emptyset^{(b)}] + E_\omega^{(r)}[T_\emptyset^{(c)}] = (\log n) \times o_{\mathbf{P}^*}(1)$ . On the other hand,

$$E_\omega^{(r)}[(T_\emptyset^{(a)} - (m - b_n))^+] \leq E_\omega^{(r)}[T_\emptyset^{(a)} \mathbf{1}_{\{T_\emptyset^{(a)} \geq m-b_n\}}] \leq \frac{1}{m - b_n} E_\omega^{(r)}[(T_\emptyset^{(a)})^2].$$

By Lemma 6.2 and tightness of  $\frac{1}{\log n} E_\omega^{(r)}[T_\emptyset^{(a)}]$ , we take a large parameter  $\theta$  such that  $\frac{18}{\delta_1} + 6 - \theta < 1$  and arrive at (9.6). This completes the proof of (9.1).

(ii) *Case  $\kappa \geq 2$ .* We only check the case  $\kappa = 2$  because the general case can be proved exactly in the same way. Without loss of generality, we take  $t_2 = 1$  and  $t_1 = s \in (0, 1)$ . For brevity, we treat  $sn$  as an integer. It suffices to prove that, for  $n \rightarrow \infty$ ,

$$(9.7) \quad \sup_{A_1, A_2 \subset \mathbb{T} \cup \overleftarrow{\{\emptyset\}}} |P_\omega(X_{sn} \in A_1, X_n \in A_2) - \tilde{\pi}_{sn}(A_1)\tilde{\pi}_n(A_2)| \rightarrow 0,$$

in  $\mathbf{P}^*$ -probability.

Fix  $t \in (s, 1)$ . Let  $\mathfrak{d}_{sn} := \min\{i > sn : X_i = \emptyset\}$  and  $B_n := \{\mathfrak{d}_{sn} \leq tn\} = \{L_{tn}(\emptyset) > L_{sn}(\emptyset)\}$ . By (2.15) of Proposition 2.4,

$$(9.8) \quad P_\omega(B_n^c) \rightarrow 0, \quad \text{in } \mathbf{P}^*\text{-probability.}$$

Hence,  $P_\omega(X_{sn} \in A_1, X_n \in A_2) = P_\omega(X_{sn} \in A_1, X_n \in A_2, B_n) + o_{\mathbf{P}^*}(1)$  where  $o_{\mathbf{P}^*}(1)$  denotes an expression converging to 0 in  $\mathbf{P}^*$ -probability uniformly in  $A_1, A_2 \subset \mathbb{T} \cup \overleftarrow{\{\emptyset\}}$ . Applying the strong Markov property at  $\mathfrak{d}_{sn}$ , this gives

$$\begin{aligned} &P_\omega(X_{sn} \in A_1, X_n \in A_2) \\ &= \sum_{k=sn+1}^{tn} P_\omega(X_{sn} \in A_1, \mathfrak{d}_{sn} = k)P_\omega(X_{n-k} \in A_2) + o_{\mathbf{P}^*}(1), \end{aligned}$$

which is  $\sum_{k=sn+1}^{tn} P_\omega(X_{sn} \in A_1, \mathfrak{d}_{sn} = k)\tilde{\pi}_{n-k}(A_2) + o_{\mathbf{P}^*}(1)$  by (9.1).

For even numbers  $k \in (sn, tn]$ ,  $n$  and  $n - k$  have the same parity and  $d_{\text{tv}}(\tilde{\pi}_{n-k}, \tilde{\pi}_n) \leq 2d_{\text{tv}}(\pi_{n-k}, \pi_n)$ . So by Lemma 2.2,  $d_{\text{tv}}(\tilde{\pi}_{n-k}, \tilde{\pi}_n) \rightarrow 0$  in  $\mathbf{P}^*$ -probability, uniformly in even numbers  $k \in [sn, tn]$ . As such,

$$\begin{aligned} P_\omega(X_{sn} \in A_1, X_n \in A_2) &= \sum_{k=sn+1}^{tn} P_\omega(X_{sn} \in A_1, \mathfrak{d}_{sn} = k)\tilde{\pi}_n(A_2) + o_{\mathbf{P}^*}(1) \\ &= P_\omega(X_{sn} \in A_1)\tilde{\pi}_n(A_2) + o_{\mathbf{P}^*}(1), \end{aligned}$$

by means of (9.8). Applying the already proved case  $\kappa = 1$  of Theorem 2.1 to  $sn$ , we get that  $P_\omega(X_{sn} \in A_1) = \tilde{\pi}_{sn}(A_1) + o_{\mathbf{P}^*}(1)$ , which yields (9.7) and completes the proof of Theorem 2.1.  $\square$

**PROOF OF COROLLARY 2.3.** We only prove the case  $\kappa = 1$  and  $t_\kappa = 1$ . The general case can be handled exactly in the same way.

By Lemma 2.2,  $d_{\text{tv}}(\pi_n(\cdot), \pi_{n-1}(\cdot)) \rightarrow 0$  in  $\mathbf{P}^*$ -probability, from which follows that

$$(9.9) \quad d_{\text{tv}}\left(\frac{1}{2}(\tilde{\pi}_n(\cdot) + \tilde{\pi}_{n-1}(\cdot)), \pi_n(\cdot)\right) \rightarrow 0, \quad \text{in } \mathbf{P}^*\text{-probability.}$$

Applying Theorem 2.1 (case  $\kappa = 1$ ) to  $n$  and  $n - 1$ , we get from (9.9) that

$$(9.10) \quad \sup_{A \subset \mathbb{T} \cup \{\emptyset\}} |P_\omega(X_n \in A) + P_\omega(X_{n-1} \in A) - 2\pi_n(A)| \rightarrow 0,$$

in  $\mathbf{P}^*$ -probability.

Let  $B > b > 0$  be constants and let  $n$  be large. We treat  $b(\log n)^2$  and  $B(\log n)^2$  as integers for brevity. By (9.10) [with  $o_{\mathbf{P}^*}(1)$  denoting an expression converging to 0 in  $\mathbf{P}^*$ -probability, uniformly in  $B > b > 0$ ]

$$(9.11) \quad \begin{aligned} I_{(9.11)} &:= P_\omega\left(b \leq \frac{|X_n|}{(\log n)^2} \leq B\right) + P_\omega\left(b \leq \frac{|X_{n-1}|}{(\log n)^2} \leq B\right) \\ &= 2 \sum_{b(\log n)^2 \leq |x| \leq B(\log n)^2} \pi_n(x) + o_{\mathbf{P}^*}(1). \end{aligned}$$

By definition of  $\pi_n$  in (2.11) (and the fact that  $Z_n = 2Y_n$  as in Lemma 2.6),

$$\begin{aligned} 2 \sum_{b(\log n)^2 \leq |x| \leq B(\log n)^2} \pi_n(x) &= \frac{1}{Y_n} \sum_{k=b(\log n)^2}^{B(\log n)^2} \sum_{|x|=k} (\mathbf{1}_{\{x < \mathcal{L}_n\}} e^{-U(x)} + \mathbf{1}_{\{x \in \mathcal{L}_n\}} e^{-V(x)}) \\ &= \frac{2}{Y_n} \left( \sum_{k=b(\log n)^2}^{B(\log n)^2} \sum_{|x|=k} \mathbf{1}_{\{x < \mathcal{L}_n\}} e^{-V(x)} + \Delta_n \right), \end{aligned}$$

where  $|\Delta_n| \leq \sum_{|x|=b(\log n)^2} e^{-V(x)} + \sum_{|x|=B(\log n)^2+1} e^{-V(x)} + \sum_{x \in \mathcal{L}_n} e^{-V(x)} = W_{b(\log n)^2} + W_{B(\log n)^2+1} + \sum_{x \in \mathcal{L}_n} e^{-V(x)}$ , where  $(W_i)$  is the additive martingale in (2.6). Since  $W_i \rightarrow 0$  (for  $i \rightarrow \infty$ )  $\mathbf{P}^*$ -a.s. [see (2.8)], and  $\sum_{x \in \mathcal{L}_n} e^{-V(x)} \rightarrow 0$  in  $\mathbf{P}^*$ -probability (Lemma 3.4), we have  $\Delta_n \rightarrow 0$  in  $\mathbf{P}^*$ -probability. On the other hand,  $\frac{Y_n}{\log n} \rightarrow \frac{2}{\sigma^2} D_\infty$  in  $\mathbf{P}^*$ -probability (Theorem 2.7). Consequently,

$$I_{(9.11)} = \frac{\sigma^2}{D_\infty \log n} \sum_{k=b(\log n)^2}^{B(\log n)^2} \sum_{|x|=k} \mathbf{1}_{\{x < \mathcal{L}_n\}} e^{-V(x)} + o_{\mathbf{P}^*}(1).$$

By an obvious analogue of (4.11) and (4.12),

$$\sum_{k=b(\log n)^2}^{B(\log n)^2} \sum_{|x|=k} \mathbf{1}_{\{x < \mathcal{L}_n\}} e^{-V(x)} \leq \sum_{k=b(\log n)^2}^{B(\log n)^2} W_k^{(\log n)},$$

and

$$\sum_{k=b(\log n)^2}^{B(\log n)^2} \sum_{|x|=k} \mathbf{1}_{\{x < \mathcal{L}_n\}} e^{-V(x)} \geq \sum_{k=b(\log n)^2}^{B(\log n)^2} W_k^{(\log(n/B(\log n)^2))}.$$

Applying (4.9) to  $\lambda = \log n$  and noting that  $\lim_{n \rightarrow \infty} \frac{\log(n/B(\log n)^2)}{\log n} = 1$ , this yields that for any fixed  $B > b > 0$ ,

$$(9.12) \quad I_{(9.11)} = \left(\frac{8\sigma^2}{\pi}\right)^{1/2} \mathbb{E} \left[ \left[ \left( B^{1/2} \wedge \frac{1}{\sigma\eta} \right) - b^{1/2} \right] \mathbf{1}_{(\eta \leq 1/(\sigma b^{1/2}))} \right] + o_{\mathbf{P}^*}(1).$$

Note that  $\mathbb{E}\{[(B^{1/2} \wedge \frac{1}{\sigma\eta}) - b^{1/2}] \mathbf{1}_{(\eta \leq 1/(\sigma b^{1/2}))}\}$  is continuous in  $B$  and  $b$ . Since  $|X_n|$  and  $|X_{n-1}|$  only differ 1, we get that

$$\begin{aligned} 2P_\omega \left( b + \frac{1}{(\log n)^2} \leq \frac{|X_n|}{(\log n)^2} \leq B - \frac{1}{(\log n)^2} \right) \\ \leq I_{(9.11)} \leq 2P_\omega \left( b - \frac{1}{(\log n)^2} \leq \frac{|X_n|}{(\log n)^2} \leq B + \frac{1}{(\log n)^2} \right), \end{aligned}$$

which, in view of (9.12), readily yields the case  $\kappa = 1$  of Corollary 2.3, as claimed.  $\square$

**Acknowledgements.** We are grateful to Dayue Chen, Nina Gantert and Ofer Zeitouni for stimulating discussions on biased walks, and to Laurent Saloff-Coste for enlightenment on Markov chains.

## REFERENCES

- [1] AIDÉKON, E. (2008). Transient random walks in random environment on a Galton–Watson tree. *Probab. Theory Related Fields* **142** 525–559. [MR2438700](#)
- [2] AIDÉKON, E. (2010). Large deviations for transient random walks in random environment on a Galton–Watson tree. *Ann. Inst. Henri Poincaré Probab. Stat.* **46** 159–189. [MR2641775](#)
- [3] AĪDÉKON, E. (2010). Tail asymptotics for the total progeny of the critical killed branching random walk. *Electron. Commun. Probab.* **15** 522–533. [MR2737710](#)
- [4] AĪDÉKON, E. (2013). Convergence in law of the minimum of a branching random walk. *Ann. Probab.* **41** 1362–1426. [MR3098680](#)
- [5] AIDEKON, E. and SHI, Z. (2014). The Seneta–Heyde scaling for the branching random walk. *Ann. Probab.* **42** 959–993. [MR3189063](#)
- [6] ANDROLETTI, P. and DEBS, P. (2014). The number of generations entirely visited for recurrent random walks in a random environment. *J. Theoret. Probab.* **27** 518–538. [MR3195824](#)
- [7] ANDROLETTI, P. and DEBS, P. (2014). Spread of visited sites of a random walk along the generations of a branching process. *Electron. J. Probab.* **19** no. 42, 22. [MR3210543](#)
- [8] BIANE, PH. and YOR, M. (1987). Valeurs principales associées aux temps locaux browniens. *Bull. Sci. Math. (2)* **111** 23–101. [MR0886959](#)
- [9] BIGGINS, J. D. (1977). Chernoff’s theorem in the branching random walk. *J. Appl. Probab.* **14** 630–636. [MR0464415](#)
- [10] BIGGINS, J. D. (1977). Martingale convergence in the branching random walk. *J. Appl. Probab.* **14** 25–37. [MR0433619](#)
- [11] BIGGINS, J. D. and KYPRIANOU, A. E. (2004). Measure change in multitype branching. *Adv. in Appl. Probab.* **36** 544–581. [MR2058149](#)
- [12] BIGGINS, J. D. and KYPRIANOU, A. E. (2005). Fixed points of the smoothing transform: The boundary case. *Electron. J. Probab.* **10** 609–631. [MR2147319](#)

- [13] BINGHAM, N. H. and DONEY, R. A. (1975). Asymptotic properties of supercritical branching processes. II. Crump–Mode and Jirina processes. *Adv. in Appl. Probab.* **7** 66–82. [MR0378125](#)
- [14] CHEN, X. (2015). A necessary and sufficient condition for the nontrivial limit of the derivative martingale in a branching random walk. *Adv. in Appl. Probab.* **47** 741–760. [MR3406606](#)
- [15] FARAUD, G. (2011). A central limit theorem for random walk in a random environment on marked Galton–Watson trees. *Electron. J. Probab.* **16** 174–215. [MR2754802](#)
- [16] FARAUD, G., HU, Y. and SHI, Z. (2012). Almost sure convergence for stochastically biased random walks on trees. *Probab. Theory Related Fields* **154** 621–660. [MR3000557](#)
- [17] GOLOSOV, A. O. (1984). Localization of random walks in one-dimensional random environments. *Comm. Math. Phys.* **92** 491–506. [MR0736407](#)
- [18] HU, Y. and SHI, Z. (2007). A subdiffusive behaviour of recurrent random walk in random environment on a regular tree. *Probab. Theory Related Fields* **138** 521–549. [MR2299718](#)
- [19] HU, Y. and SHI, Z. (2007). Slow movement of random walk in random environment on a regular tree. *Ann. Probab.* **35** 1978–1997. [MR2349581](#)
- [20] HU, Y. and SHI, Z. (2014). The potential energy of biased random walks on trees. Available at [arXiv:1403.6799](https://arxiv.org/abs/1403.6799).
- [21] HU, Y., SHI, Z. and YOR, M. (2015). The maximal drawdown of the Brownian meander. *Electron. Commun. Probab.* **20** no. 39, 6. [MR3352334](#)
- [22] IMHOF, J.-P. (1984). Density factorizations for Brownian motion, meander and the three-dimensional Bessel process, and applications. *J. Appl. Probab.* **21** 500–510. [MR0752015](#)
- [23] KAHANE, J.-P. and PEYRIÈRE, J. (1976). Sur certaines martingales de Benoit Mandelbrot. *Adv. Math.* **22** 131–145. [MR0431355](#)
- [24] KOZLOV, M. V. (1976). The asymptotic behavior of the probability of non-extinction of critical branching processes in a random environment. *Teor. Veroyatnost. i Primenen.* **21** 813–825. [MR0428492](#)
- [25] LEHOCZKY, J. P. (1977). Formulas for stopped diffusion processes with stopping times based on the maximum. *Ann. Probab.* **5** 601–607. [MR0458570](#)
- [26] LYONS, R. and PERES, Y. (2015). *Probability on Trees and Networks*. Cambridge Univ. Press, Cambridge. Available at <http://mypage.iu.edu/~rdlyons/prbtree/prbtree.html>.
- [27] LYONS, R. (1990). Random walks and percolation on trees. *Ann. Probab.* **18** 931–958. [MR1062053](#)
- [28] LYONS, R. (1992). Random walks, capacity and percolation on trees. *Ann. Probab.* **20** 2043–2088. [MR1188053](#)
- [29] LYONS, R. (1997). A simple path to Biggins’ martingale convergence for branching random walk. In *Classical and Modern Branching Processes (Minneapolis, MN, 1994)* (K. B. Athreya and P. Jagers, eds.). *IMA Vol. Math. Appl.* **84** 217–221. Springer, New York. [MR1601749](#)
- [30] LYONS, R. and PEMANTLE, R. (1992). Random walk in a random environment and first-passage percolation on trees. *Ann. Probab.* **20** 125–136. [MR1143414](#)
- [31] LYONS, R., PEMANTLE, R. and PERES, Y. (1995). Conceptual proofs of  $L \log L$  criteria for mean behavior of branching processes. *Ann. Probab.* **23** 1125–1138. [MR1349164](#)
- [32] LYONS, R., PEMANTLE, R. and PERES, Y. (1995). Ergodic theory on Galton–Watson trees: Speed of random walk and dimension of harmonic measure. *Ergodic Theory Dynam. Systems* **15** 593–619. [MR1336708](#)
- [33] LYONS, R., PEMANTLE, R. and PERES, Y. (1996). Biased random walks on Galton–Watson trees. *Probab. Theory Related Fields* **106** 249–264. [MR1410689](#)
- [34] MADAULE, T. (2016). First order transition for the branching random walk at the critical parameter. *Stochastic Process. Appl.* **126** 470–502. [MR3434991](#)



- [35] MENSHIKOV, M. and PETRITIS, D. (2002). On random walks in random environment on trees and their relationship with multiplicative chaos. In *Mathematics and Computer Science, II (Versailles, 2002)*. *Trends Math.* 415–422. Birkhäuser, Basel. [MR1940150](#)
- [36] NEVEU, J. (1986). Arbres et processus de Galton–Watson. *Ann. Inst. Henri Poincaré Probab. Stat.* **22** 199–207. [MR0850756](#)
- [37] PERES, Y. (1999). Probability on trees: An introductory climb. In *Lectures on Probability Theory and Statistics (Saint-Flour, 1997)*. *Lecture Notes in Math.* **1717** 193–280. Springer, Berlin. [MR1746302](#)
- [38] PITMAN, J. W. (2015). Some conditional expectations and identities for Bessel processes related to the maximal drawdown of the Brownian meander. Unpublished manuscript.
- [39] SALOFF-COSTE, L. (1997). Lectures on finite Markov chains. In *Lectures on Probability Theory and Statistics (Saint-Flour, 1996)*. *Lecture Notes in Math.* **1665** 301–413. Springer, Berlin. [MR1490046](#)
- [40] SHI, Z. (2015). *Branching Random Walks, École D’été Saint-Flour XLII (2012)*. *Lecture Notes in Math.* **2151**. Springer, Berlin.
- [41] SINAI, YA. G. (1982). The limit behavior of a one-dimensional random walk in a random environment. *Teor. Veroyatnost. i Primenen.* **27** 247–258. [MR0657919](#)

LAGA  
UNIVERSITÉ PARIS XIII  
99 AVENUE J-B CLÉMENT  
F-93430 VILLETANEUSE  
FRANCE  
E-MAIL: [yueyun@math.univ-paris13.fr](mailto:yueyun@math.univ-paris13.fr)

LPMA  
UNIVERSITÉ PARIS VI  
4 PLACE JUSSIEU  
F-75252 PARIS CEDEX 05  
FRANCE  
E-MAIL: [zhan.shi@upmc.fr](mailto:zhan.shi@upmc.fr)