

## A NOTE ON THE DE LA GARZA PHENOMENON FOR LOCALLY OPTIMAL DESIGNS

BY HOLGER DETTE<sup>1</sup> AND VIATCHESLAV B. MELAS<sup>2</sup>

*Ruhr-Universität Bochum and St. Petersburg State University*

The celebrated de la Garza phenomenon states that for a polynomial regression model of degree  $p - 1$  any optimal design can be based on at most  $p$  design points. In a remarkable paper, Yang [*Ann. Statist.* **38** (2010) 2499–2524] showed that this phenomenon exists in many locally optimal design problems for nonlinear models. In the present note, we present a different view point on these findings using results about moment theory and Chebyshev systems. In particular, we show that this phenomenon occurs in an even larger class of models than considered so far.

**1. Introduction.** Nonlinear regression models are widely used for modeling dependencies between response and explanatory variables [see [Seber and Wild \(1989\)](#) or [Ratkowsky \(1990\)](#)]. It is well known that an appropriate choice of an experimental design can improve the quality of statistical analysis substantially, and therefore the problem of constructing optimal designs for nonlinear regression models has found considerable attention in the literature. Most authors concentrate on locally optimal designs which assume that a guess for the unknown parameters of the model is available [see [Chernoff \(1953\)](#), [Ford, Torsney and Wu \(1992\)](#), [He, Studden and Sun \(1996\)](#), [Fang and Hedayat \(2008\)](#)]. These designs are usually used as benchmarks for commonly used designs. Additionally, they serve as a basis for constructing optimal designs with respect to more sophisticated optimality criteria which address for a less precise knowledge about the unknown parameters [see [Pronzato and Walter \(1985\)](#) or [Chaloner and Verdinelli \(1995\)](#), [Dette \(1997\)](#), [Müller and Pázman \(1998\)](#)]. It is a well-known fact that the numerical or analytical calculation of optimal designs simplifies substantially if it is known that the optimal design is saturated, which means that the number of different experimental conditions coincides with the number of parameters in the model [see, e.g., [He, Studden and Sun \(1996\)](#), [Dette and Wong \(1996\)](#), [Imhof and Studden \(2001\)](#), [Imhof \(2001\)](#), [Melas \(2006\)](#), [Fang and Hedayat \(2008\)](#) among many others].

---

Received July 2010; revised January 2011.

<sup>1</sup>Supported in part by the Collaborative Research Center “Statistical modeling of nonlinear dynamic processes” (SFB 823) of the German Research Foundation (DFG).

<sup>2</sup>Supported in part by Russian Foundation for Basic Research (RFBR), Project 09-01-00508. *MSC2010 subject classification.* 62K05.

*Key words and phrases.* Locally optimal designs, saturated designs, complete class theorem, moment spaces, Chebyshev systems.

So, the ideal situation appears if the optimal design is in the sub-class of all saturated designs. In a celebrated paper, [de la Garza \(1954\)](#) proved that for a  $(p - 1)$ th-degree polynomial regression model, any optimal design can be based on at most  $p$  points. [Khuri et al. \(2006\)](#) considered a nonlinear regression model and introduced the terminology of the de la Garza phenomenon, which means that for any design there exists a saturated design, such that the information matrix of the saturated design is not inferior to that of the given design under the Loewner ordering. In a remarkable paper, [Yang \(2010\)](#) derived sufficient conditions on the nonlinear regression model for the occurrence of the de la Garza phenomenon and demonstrated that this situation appears in a broad class of nonlinear regression models. These results generalize recent findings of [Yang and Stufken \(2009\)](#) for nonlinear models with two parameters.

However, some care is necessary if these results are applied as indicated in the following simple example of homoscedastic linear regression on the interval  $[0, 1]$ . Here the information matrix of the design which advises the experimenter to take all  $n$  observations at the point 0 is given by

$$X_1^T X_1 = \begin{pmatrix} n & 0 \\ 0 & 0 \end{pmatrix}$$

while any other design (using the experimental conditions  $x_1, \dots, x_n$ ) yields an information matrix

$$X_2^T X_2 = \begin{pmatrix} n & \sum_{i=1}^n x_i \\ \sum_{i=1}^n x_i & \sum_{i=1}^n x_i^2 \end{pmatrix}.$$

It is easy to see that the matrix  $X_2^T X_2 - X_1^T X_1$  is indefinite (i.e., it has positive and negative eigenvalues) whenever one of the  $x_i$  is positive. Consequently, the design corresponding to  $X_1^T X_1$  cannot be improved. On the other hand, it is also easy to see that for any  $k \in \{1, \dots, \lfloor n/2 \rfloor - 1\}$  the information matrix of the design, which takes observations at  $x_1 = \dots = x_{n-2k} = 0$  and at  $x_{n-2k+1} = \dots = x_n = 1/2$  can be improved (with respect to the Loewner ordering) by the information matrix corresponding to the design  $x_1 = \dots = x_{n-k} = 0$  and  $x_{n-k+1} = \dots = x_n = 1$ . Thus, there exist designs where a “real” improvement is possible, while other designs cannot be improved. Note that the results in [Yang \(2010\)](#) do not provide a classification of the two types of designs.

It is the purpose of the present paper to present a more detailed view point on these problems, which clarifies this—on a first glance—contradiction. In contrast to the method used by [Yang \(2010\)](#), which is mainly algebraic, our approach is analytic and based on the theory of Chebyshev systems and moment spaces [see [Karlin and Studden \(1966b\)](#)]. In particular, we will demonstrate that the de la Garza phenomenon appears in any nonlinear regression model, where the functions in the

Fisher information matrix form a Chebyshev system. Additionally, we will solve the problem described in the previous paragraph and we will identify the sufficient conditions stated in Yang (2010) as a special case of an extended Chebyshev system. Therefore, our results generalize the recent findings of Yang (2010) in a nontrivial way and, additionally, provide—in our opinion—a more transparent and more complete explanation of the de la Garza phenomenon for optimal designs in nonlinear regression models.

The remaining part of this paper is organized as follows. Section 2 provides a brief introduction in the problem, while Section 3 contains our main results. Finally, the new results are illustrated in a rational regression model, where the currently available methodology cannot be used to establish the de la Garza phenomenon.

**2. Locally optimal designs.** Consider the common nonlinear regression model

$$(2.1) \quad Y = \eta(x, \theta) + \varepsilon,$$

where  $\theta \in \Theta \subset \mathbb{R}^p$  is the vector of unknown parameters, and different observations are assumed to be independent. The errors are normally distributed with mean 0 and variance  $\sigma^2$ . The variable  $x$  denotes the explanatory variable, which varies in the design space  $[A, B] \subset \mathbb{R}$ . We assume that  $\eta$  is a continuous and real valued function of both arguments  $(x, \theta) \in [A, B] \times \Theta$  and differentiable with respect to the variable  $\theta$ . A design is defined as a probability measure  $\xi$  on the interval  $[A, B]$  with finite support [see Kiefer (1974)]. If the design  $\xi$  has masses  $w_i$  at the points  $x_i$  ( $i = 1, \dots, k$ ) and  $n$  observations can be made by the experimenter, this means that the quantities  $w_i n$  are rounded to integers, say  $n_i$ , satisfying  $\sum_{i=1}^k n_i = n$ , and the experimenter takes  $n_i$  observations at each location  $x_i$  ( $i = 1, \dots, k$ ). The information matrix of an approximate design  $\xi$  is defined by

$$(2.2) \quad M(\xi, \theta) = \int_A^B \left( \frac{\partial}{\partial \theta} \eta(x, \theta) \right) \left( \frac{\partial}{\partial \theta} \eta(x, \theta) \right)^T d\xi(x),$$

and it is well known [see Jennrich (1969)] that under appropriate assumptions of regularity the covariance matrix of the least squares estimator is approximately given by  $\sigma^2 M^{-1}(\xi, \theta)/n$ , where  $n$  denotes the total sample size and we assume that the observations are taken according to the approximate design  $\xi$ .

An optimal design maximizes an appropriate functional of the information matrix and numerous criteria have been proposed in the literature to discriminate between competing designs [see Silvey (1980), Pázman (1986) or Pukelsheim (2006) among others]. Note that in nonlinear regression models the information matrix (and as a consequence the corresponding optimal designs) depend on the unknown parameters and are therefore called locally optimal designs [see Chernoff (1953)]. These designs require an initial guess of the unknown parameters in the model and are used as benchmarks for many commonly used designs.

Most of the available optimality criteria satisfy a monotonicity property with respect to the Loewner ordering, that is

$$(2.3) \quad M(\xi_1, \theta) \leq M(\xi_2, \theta) \implies \Phi(M(\xi_1, \theta)) \leq \Phi(M(\xi_2, \theta)),$$

where the parameter  $\theta$  is fixed,  $\xi_1, \xi_2$  are two competing designs and  $\Phi$  denotes an information function in the sense of Pukelsheim (2006). For this reason, it is of interest to derive a complete class theorem in this general context which characterizes the class of designs, which cannot be improved with respect to the Loewner ordering of their information matrices. We call a design  $\xi_1$  admissible if there does not exist a design  $\xi_2$ , such that  $M(\xi_1, \theta) \neq M(\xi_2, \theta)$  and

$$(2.4) \quad M(\xi_1, \theta) \leq M(\xi_2, \theta).$$

As pointed out in Yang (2010) for many nonlinear regression models the information matrix defined in (2.2) has a representation of the form

$$(2.5) \quad M(\xi, \theta) = P(\theta)C(\xi, \theta)P^T(\theta),$$

where  $P(\theta)$  is a nonsingular  $p \times p$  matrix, which does not depend on the design  $\xi$ , the matrix  $C$  is defined by

$$(2.6) \quad C(\xi, \theta) = \begin{pmatrix} \int_A^B \Psi_{11}(x) d\xi(x) & \cdots & \int_A^B \Psi_{1p}(x) d\xi(x) \\ \vdots & \ddots & \vdots \\ \int_A^B \Psi_{p1}(x) d\xi(x) & \cdots & \int_A^B \Psi_{pp}(x) d\xi(x) \end{pmatrix}$$

and  $\Psi_{11}, \Psi_{12}, \dots, \Psi_{pp}$  are functions defined on the interval  $[A, B]$ . Note that these functions usually depend on the parameter  $\theta$ , but for the sake of simplicity we do not reflect this dependence in our notation. Obviously the inequality (2.4) is satisfied if and only if the inequality

$$(2.7) \quad C(\xi_1, \theta) \leq C(\xi_2, \theta)$$

is satisfied.

**3. Chebyshev systems and complete class theorems.** In the following discussion, we make extensive use of the property that a system of functions has the Chebyshev property. Following Karlin and Studden (1966b), a set of  $k + 1$  continuous functions  $u_0, \dots, u_k : [A, B] \rightarrow \mathbb{R}$  is called a Chebyshev system (on the interval  $[A, B]$ ) if the inequality

$$(3.1) \quad \begin{vmatrix} u_0(x_0) & u_0(x_1) & \cdots & u_0(x_k) \\ u_1(x_0) & u_1(x_1) & \cdots & u_1(x_k) \\ \vdots & \vdots & \ddots & \vdots \\ u_k(x_0) & u_k(x_1) & \cdots & u_k(x_k) \end{vmatrix} > 0$$

holds for all  $A \leq x_0 < x_1 < \dots < x_k \leq B$ . Note that if the determinant in (3.1) does not vanish then either the functions  $u_0, u_1, \dots, u_{k-1}, u_k$  or the functions  $u_0, u_1, \dots, u_{k-1}, -u_k$  form a Chebyshev system. The Chebyshev property has widely been used to determine explicitly  $c$ -optimal designs [see He, Studden and Sun (1996), Dette et al. (2003) or Dette et al. (2008) among many others]. On the other hand, its application to other optimality criteria has not been studied intensively. In the following discussion, we will demonstrate that this property will essentially be the reason for the occurrence of the de la Garza phenomenon. In particular, we will show that it is essentially sufficient to obtain a complete class theorem for the design problems associated with the nonlinear regression model (2.1).

For this purpose, we define the index  $I(\xi)$  of a design  $\xi$  on the interval  $[A, B]$  as the number of support points, where the boundary points  $A$  and  $B$  (if they occur as support points) are only counted by  $1/2$ . Recall the definition of the matrix  $C$  in (2.6) and denote by  $\Psi_1, \dots, \Psi_k$  the different elements among the functions  $\{\Psi_{ij} \mid 1 \leq j, j \leq p\}$ , which are not equal to the constant function. Throughout this paper, we assume

$$(3.2) \quad \begin{aligned} \Psi_k &= \Psi_{ll} && \text{for some } l \in \{1, \dots, p\} \quad \text{and} \\ \Psi_{ij} &\neq \Psi_k && \text{for all } (i, j) \neq (l, l) \end{aligned}$$

[see Yang (2010)]. Additionally, we put  $\Psi_0(x) = 1$  and assume either that

$$(3.3) \quad \begin{aligned} &\{\Psi_0, \Psi_1, \dots, \Psi_{k-1}\} \quad \text{and} \\ &\{\Psi_0, \Psi_1, \dots, \Psi_{k-1}, \Psi_k\} \end{aligned}$$

are Chebyshev systems or that

$$(3.4) \quad \begin{aligned} &\{\Psi_0, \Psi_1, \dots, \Psi_{k-1}\} \quad \text{and} \\ &\{\Psi_0, \Psi_1, \dots, \Psi_{k-1}, -\Psi_k\} \end{aligned}$$

are Chebyshev systems then the following result characterizes the class of admissible designs.

**THEOREM 3.1.** (1) *If the functions  $\Psi_0(x) = 1, \Psi_1, \dots, \Psi_{k-1}, \Psi_k$  satisfy (3.2) and (3.3), then for any design  $\xi$  there exists a design  $\xi^+$  with at most  $\frac{k+2}{2}$  support points, such that  $M(\xi^+, \theta) \geq M(\xi, \theta)$ . If the index of the design  $\xi$  satisfies*

$$I(\xi) < \frac{k}{2}$$

*then the design  $\xi^+$  is uniquely determined in the class of all designs  $\eta$  satisfying*

$$(3.5) \quad \int_A^B \Psi_i(x) d\eta(x) = \int_A^B \Psi_i(x) d\xi(x), \quad i = 0, \dots, k - 1,$$

*and coincides with the design  $\xi$ . Otherwise [in the case  $I(\xi) \geq \frac{k}{2}$ ], the following two assertions are valid.*

- (1a) If  $k$  is odd, then  $\xi^+$  has at most  $\frac{k+1}{2}$  support points and  $\xi^+$  can be chosen such that its support contains the point  $B$ .
- (1b) If  $k$  is even, then  $\xi^+$  has at most  $\frac{k}{2} + 1$  support points and  $\xi^+$  can be chosen such that the support of  $\xi^+$  contains the points  $A$  and  $B$ .

(2) If the functions  $\Psi_0(x) = 1, \Psi_1, \dots, \Psi_{k-1}, \Psi_k$  satisfy (3.2) and (3.4), then for any design  $\xi$  there exists a design  $\xi^-$  with at most  $\frac{k+2}{2}$  support points, such that  $M(\xi^-, \theta) \geq M(\xi, \theta)$ . If the index of the design  $\xi$  satisfies

$$I(\xi) < \frac{k}{2}$$

then the design  $\xi^-$  is uniquely determined in the class of all designs  $\eta$  satisfying (3.5) and coincides with the design  $\xi$ . Otherwise [in the case  $I(\xi) \geq \frac{k}{2}$ ], the following two assertions are valid.

- (2a) If  $k$  is odd, then  $\xi^-$  has at most  $\frac{k+1}{2}$  support points and  $\xi^-$  can be chosen such that its support contains the point  $A$ .
- (2b) If  $k$  is even, then  $\xi^-$  has at most  $\frac{k}{2}$  support points.

PROOF. We only present a proof of the first part (1) of the theorem, the second part follows by similar arguments. For  $i = 0, \dots, k$  let

$$d_i(\xi) = \int_A^B \Psi_i(x) d\xi(x)$$

denote the  $i$ th “moment” and define

$$\vec{d}_k(\xi) = (d_0(\xi), \dots, d_k(\xi))^T$$

as the vector of all “moments” up to the order  $k$ . Consider two designs  $\xi_1$  and  $\xi_2$  with

$$\vec{d}_{k-1}(\xi_1) = \vec{d}_{k-1}(\xi_2) \quad \text{and} \quad d_k(\xi_1) \leq d_k(\xi_2),$$

then for any vector  $z = (z_1, \dots, z_p)^T \in \mathbb{R}^p$  we have for some  $l \in \{1, \dots, p\}$

$$z^T (C(\xi_2, \theta) - C(\xi_1, \theta))z \geq z_l^2 (d_k(\xi_2) - d_k(\xi_1)) \geq 0,$$

which means that

$$C(\xi_2, \theta) \geq C(\xi_1, \theta).$$

Now let for a fixed vector of “moments”  $\vec{d}_{k-1}(\xi)$

$$d_k^+ = \sup\{d_k(\eta) \mid \eta \text{ design on } [A, B] \text{ with } \vec{d}_{k-1}(\eta) = \vec{d}_{k-1}(\xi)\}$$

denote the maximum of the  $k$ th “moment” over the set of all designs with fixed “moments” up to the order  $k - 1$ . Due to the compactness of the design space and the continuity of the functions  $\Psi_0, \dots, \Psi_k$ , there exists a design  $\xi^+$  such that

$$(3.6) \quad d_j(\xi^+) = d_j(\xi); \quad j = 0, \dots, k - 1,$$

$$(3.7) \quad d_k(\xi^+) = d^+ \geq d_k(\xi).$$

This shows (by the argument at the beginning of the proof and the discussion at the end of the previous section)

$$(3.8) \quad M(\xi^+, \theta) \geq M(\xi, \theta).$$

Moreover, it follows from Chapter II, Section 6 of Karlin and Studden (1966b) that the point  $\vec{d}_k(\xi^+)$  is a boundary point of the “moment space”

$$\mathcal{M}_k = \{\vec{d}_k(\eta) \mid \eta \text{ design on } [A, B]\}.$$

Consequently, we obtain from Theorem 2.1 in Karlin and Studden (1966b) that the design  $\xi^+$  is based on at most  $\frac{k+2}{2}$  support points, which proves the first part of the statement.

We now consider the cases (1a) and (1b). The vector  $\vec{d}_{k-1}(\xi)$  is either a boundary point or an interior point of the  $(k - 1)$ th moment space  $\mathcal{M}_{k-1}$ . The first case is characterized by an index satisfying  $I(\xi) < k/2$  and there exists a unique measure  $\tilde{\xi}$  with “moments” up to the order  $k$  specified by  $\vec{d}_{k-1}(\xi)$ . To prove this statement regarding uniqueness suppose that  $I(\xi) < \frac{k}{2}$  and that there exists a further design, say  $\tilde{\xi}$ , with this property. A simple counting argument shows that the total number of distinct points, say  $x_1, \dots, x_t$  among the support points of both representations is at most  $k$ . If it would be less than  $k$  we could take additional support points with corresponding vanishing weights and thus without loss of generality, we can assume that the number of distinct points is equal to  $k$ . Therefore, there would exist  $k$  different points

$$A \leq x_0 < x_1 < \dots < x_{k-1} \leq B$$

such that

$$\Psi \mu = 0,$$

where the matrix  $\Psi$  is given by

$$\Psi = \begin{pmatrix} \Psi_0(x_0) & \Psi_0(x_1) & \dots & \Psi_0(x_{k-1}) \\ \Psi_1(x_0) & \Psi_1(x_1) & \dots & \Psi_1(x_{k-1}) \\ \vdots & \vdots & \ddots & \vdots \\ \Psi_{k-1}(x_0) & \Psi_{k-1}(x_1) & \dots & \Psi_{k-1}(x_{k-1}) \end{pmatrix}$$

and the vector  $\mu \neq 0$  has components

$$\mu_i = \begin{cases} \omega_i, & x_i \in \text{supp } \xi, x_i \notin \text{supp } \tilde{\xi}, \\ -\tilde{\omega}_i, & x_i \notin \text{supp } \xi, x_i \in \text{supp } \tilde{\xi}, \\ \omega_i - \tilde{\omega}_i, & x_i \in \text{supp } \xi \cap \text{supp } \tilde{\xi}, \\ 0, & x_i \notin \text{supp } \xi, x_i \notin \text{supp } \tilde{\xi} \end{cases}$$

(here  $\omega_i$  and  $\tilde{\omega}_i$  denote the weights of the designs  $\xi$  and  $\tilde{\xi}$ , resp.). Because  $\mu \neq 0$  it follows from here that  $\det \Psi = 0$  which is impossible by the definition of Chebyshev systems. Consequently, a design with moments specified by (3.5) is

uniquely determined and therefore we take  $\xi^+ = \tilde{\xi}$ , which has at most  $\frac{k+1}{2}$  support points [see Theorem 2.1 in Karlin and Studden (1966b), page 42].

If the index of the design  $\xi$  satisfies  $I(\xi) \geq k/2$  it follows from the discussion in Chapter II, Section 6 in Karlin and Studden (1966b) that the design  $\xi^+$  defined by (3.6) and (3.7) is the upper principal representation of the vector  $\vec{d}_{k-1}(\xi)$ , which means that its index is precisely  $\frac{k}{2}$  and its support includes the point  $B$ . Note that for this argument we require condition (3.3).

Consequently, if  $k = 2m + 1$  is odd, the upper principal representation  $\xi^+$  has index  $m + \frac{1}{2}$  and precisely  $m + 1$  support points including the point  $B$ . On the other hand, if  $k = 2m$  is even,  $\xi^+$  has  $m + 1$  support points and the boundary points  $A$  and  $B$  of the design interval are support points because the index of the design  $\xi^+$  is  $m$ .

The proof of part (2) of Theorem 3.1 is similar [where the upper principal representation has to be replaced by the lower principal representation using condition (3.4)] and omitted.  $\square$

REMARK 3.2. (a) Note that Theorem 2.1 in Karlin and Studden [(1966b), Chapter II] refers to moment spaces corresponding to not necessarily bounded measures and the inclusion of the constant function in the system under consideration guarantees its application to a moment space corresponding to probability measures as required in the proof of Theorem 3.1. An alternative explanation can be given by the generalized equivalence theorem as stated in Pukelsheim (2006). It follows from this result that for an optimal design (with respect to the commonly used criteria) there exist some constants, say  $a_i \in \mathbb{R}$ ,  $i = 1, \dots, k$ , such that for all support points of the optimal design the identity

$$\sum_{i=1}^k a_i \Psi_i(x) = c$$

is satisfied, where  $c$  denotes a constant (e.g., for the  $D$ -optimality criterion  $c$  is the number of parameters). Since an optimal design is admissible, the inclusion of the constant function guarantees that the index of these designs is at most  $k/2$ . Note that this is a sufficient but, generally speaking, not necessary condition.

(b) Note that it follows from the proof of Theorem 3.1 that the conditions (3.6) and (3.7) imply (3.8), that is, the superiority of the information matrix of the design  $\xi^+$  with respect to the Loewner ordering. In many cases (e.g., polynomial regression models), the converse direction is also true and in these cases it follows from the proof of Theorem 3.1 that a design  $\xi$  with index  $I(\xi) < \frac{k}{2}$  can only be “improved” (with respect to the Loewner ordering of the corresponding information matrices) by itself. In fact we are not aware of any case where the converse direction does not hold.

(c) Note also that Theorem 3.1 provides a solution to the problem indicated in the example of the Introduction. In the linear regression model we have  $k = 2$ ,



therefore we can use the given design  $\xi_1$  (concentrating all observations at  $x = 0$ ) as an “improvement” of  $\xi_1$ . However, because the index of  $\xi_1$  is  $1/2 < 1$  the design  $\xi_1$  can only be improved by itself (see the previous remark). In particular, there does not exist a design  $\xi$  which takes observations at  $x = 1$  and improves  $\xi_1$  in the sense  $M(\xi) \geq M(\xi_1)$ .

(d) It is also worthwhile to mention that a design improving the given design  $\xi$  is not necessarily unique. Consider, for example, again the linear regression model on the interval  $[0, 1]$  and the design  $\xi$  which has equal masses at the points 0 and  $3/4$ . The information matrix of  $\xi$  is given by

$$M(\xi) = \begin{pmatrix} 1 & \frac{3}{8} \\ \frac{3}{8} & \frac{9}{32} \end{pmatrix}.$$

Now define for any  $p \in [\frac{1}{2}, \frac{5}{8}]$  a design  $\xi_p^+$  with masses  $p$  and  $1 - p$  at the points 0 and  $\frac{3}{8(1-p)}$ , respectively. Then it follows that

$$M(\xi_p^+) = \begin{pmatrix} 1 & \frac{3}{8} \\ \frac{3}{8} & \frac{9}{64(1-p)} \end{pmatrix}$$

and  $M(\xi_p^+) \geq M(\xi)$  for any  $p \in [\frac{1}{2}, \frac{5}{8}]$ . Note that the choice  $p = \frac{5}{8}$  gives the upper principal representation  $\xi^+ = \xi_{5/8}^+$  with index 1 and support points 0 and 1, while for  $p \in [\frac{1}{2}, \frac{5}{8})$  we have index  $I(\xi_p^+) = 3/2$ .

In the remaining part of this section, we will relate the result of Theorem 3.1 to the recent findings of Yang (2010). Note that—in contrast to Theorem 1 and 2 of Yang (2010)—our Theorem 3.1 does not require the differentiability of the functions  $\Psi_j$ . Moreover, in some cases it provides a better description of the admissible designs. For a more detailed explanation, we note that a Chebyshev system of functions  $\{u_0, \dots, u_k\}$  is called an extended Chebyshev system, if and only if for any  $a_0, \dots, a_k \in \mathbb{R}$  with  $\sum_{i=0}^k a_i^2 \neq 0$  the function

$$\sum_{i=0}^k a_i u_i(x)$$

has at most  $k$  zeros counted with multiplicities in the interval  $[A, B]$ . Note that this definition is equivalent to the definition given in Karlin and Studden (1966b). It is in fact proved in Karlin and Studden [(1966b), Section 1.2] for the case of system  $u_i(t) = t^i, i = 0, \dots, n$ . And the argument can be applied for general case. Moreover, by definition, an extended Chebyshev system is always a Chebyshev system.

A simple way of constructing an extended Chebyshev system is the following [see Karlin and Studden (1966b), page 19]. Let  $w_0, \dots, w_k$  be functions on the

interval  $[A, B]$  which are either positive or negative. We now consider the new functions

$$\begin{aligned}
 u_0(x) &= w_0(x), \\
 u_1(x) &= w_0(x) \int_A^x w_1(t_1) dt_1, \\
 &\vdots \\
 u_k(x) &= w_0(x) \int_A^x w_1(t_1) \int_A^{t_2} w_2(t_2) \cdots \int_A^{t_{k-1}} w_k(t_k) dt_k \cdots dt_1.
 \end{aligned}
 \tag{3.9}$$

A direct calculation shows that the Wronskian determinant of the functions  $u_0, \dots, u_k$  is given by

$$\begin{aligned}
 W_x(u_0, \dots, u_k) &= \begin{vmatrix} u_0(x) & u_0'(x) & \cdots & u_0^{(k)}(x) \\ u_1(x) & u_1'(x) & \cdots & u_1^{(k)}(x) \\ \vdots & \vdots & \ddots & \vdots \\ u_k(x) & u_k'(x) & \cdots & u_k^{(k)}(x) \end{vmatrix} \\
 &= (w_0(x))^{k+1} (w_1(x))^k \cdots (w_{k-1}(x))^2 w_k(x)
 \end{aligned}
 \tag{3.10}$$

and it is shown in Chapter XI in [Karlin and Studden \(1966b\)](#) that the set  $\{u_0, \dots, u_k\}$  of  $k$  times differentiable function is an extended Chebyshev system if and only if

$$W_x(u_0, \dots, u_k) > 0$$

for all  $x \in [A, B]$ . On the other hand, this representation provides a constructive method for checking if a given system of  $k$  times differentiable functions  $\{u_0, \dots, u_k\}$  is a Chebyshev system on the interval  $[A, B]$ . To be precise, define  $w_0(x) = u_0(x)$  and recursively differential operators

$$D_j f = \frac{d}{dx} \left( \frac{f}{w_j} \right); \quad j = 0, \dots, k,$$

$$w_{j+1} = (D_j D_{j-1} \cdots D_0) u_{j+1}; \quad j = 0, 1, \dots, k - 1.$$

Consequently, the set  $\{u_0, \dots, u_k\}$  is a Chebyshev system if the functions  $w_0, \dots, w_k$  calculated by (3.11) and (3.12) are all positive on the interval  $[A, B]$ .

**REMARK 3.3.** [Yang \(2010\)](#) constructed a triangle array of functions  $\{f_{l,t} \mid t = 1, \dots, k; t \leq l \leq k\}$  from the functions  $\Psi_1, \dots, \Psi_k$  induced by the nonlinear regression model (2.1) using the recursion

$$f_{l,t}(x) = \begin{cases} \Psi_l'(x), & t = 1, \dots, k, \\ \left( \frac{f_{l,t-1}(x)}{f_{l-1,t-1}(x)} \right)', & 2 \leq t \leq k; t \leq l \leq k. \end{cases}$$

It is now easy to see that the functions  $w_1, \dots, w_k$  obtained from (3.11) and (3.12) with  $w_0 = 1, u_j = \Psi_j (j = 1, \dots, k)$  are precisely the functions  $f_{jl}$  defined by Yang (2010). As a consequence, we will obtain the main result of Yang (2010) as a special case of our Theorem 3.1 (note that our assumptions regarding the differentiability are slightly weaker than in this reference).

**THEOREM 3.4.** *Let  $\Psi_1, \dots, \Psi_k$  denote the  $k$  different functions in the information matrix (3.1) corresponding to the nonlinear regression model which are not equal to the constant function. Assume that  $\Psi_j$  is  $(j + 1)$  times continuously differentiable, define  $w_0 = 1$  and for  $j = 0, \dots, k - 1$*

$$w_{j+1} = D_j D_{j-1} \cdots D_0 \Psi_{j+1}$$

and assume that condition (3.2) is satisfied. If

$$F(x) = w_1(x) \cdots w_k(x) \neq 0$$

for all  $x \in [A, B]$ , then for any given design  $\xi$  there exists a design  $\tilde{\xi}$ , such that  $I(\tilde{\xi}) \leq \frac{k}{2}$

$$M(\tilde{\xi}, \theta) \geq M(\xi, \theta).$$

If the index of the design  $\xi$  satisfies  $I(\xi) < \frac{k}{2}$  then  $\tilde{\xi}$  is uniquely determined in the class of all designs  $\eta$  with moments specified by (3.5) and coincides with the design  $\xi$ . Otherwise [in the case  $I(\xi) \geq \frac{k}{2}$ ] the following assertions are valid.

- (1a) *If  $k$  is odd and  $F(x) < 0$  on the interval  $[A, B]$ , then the design  $\tilde{\xi}$  has at most  $(k + 1)/2$  support points and  $\tilde{\xi}$  can be chosen such that the point  $A$  is a support point.*
- (1b) *If  $k$  is odd and  $F(x) > 0$  on the interval  $[A, B]$ , then the design  $\tilde{\xi}$  has at most  $(k + 1)/2$  support points and  $\tilde{\xi}$  can be chosen such that the point  $B$  is a support point.*
- (2a) *If  $k$  is even and  $F(x) < 0$  on the interval  $[A, B]$ , then the design  $\tilde{\xi}$  has at most  $k/2$  support points.*
- (2b) *If  $k$  is even and  $F(x) > 0$  on the interval  $[A, B]$ , then the design  $\tilde{\xi}$  has at most  $k/2 + 1$  support points and  $\tilde{\xi}$  can be chosen such that the points  $A$  and  $B$  are support points.*

**PROOF.** Let us define  $\Psi_0(x) = 1$  and note that

$$F(x) = \frac{W_x(\Psi_0, \dots, \Psi_k)}{W_x(\Psi_0, \dots, \Psi_{k-1})}.$$

Thus if  $F(x) > 0$  then condition (3.3) is fulfilled and if  $F(x) < 0$ , then condition (3.4) is fulfilled. Now Theorem 3.4 is an immediate corollary of Theorem 3.1.  $\square$

REMARK 3.5. Note that if the constant function appears among the different functions  $\{\Psi_{ij} \mid 1 \leq i \leq j \leq p\}$  in the information matrix (3.1) it is not counted in Theorem 3.4 or Theorem 2 of Yang (2010) (see the proof of Theorems 3 and 5–7 in this reference).

A number of interesting applications of Theorem 3.4 are given in Yang (2010). Note that in all examples considered there the functions under consideration generate a special type of Chebyshev systems, namely extended Chebyshev systems that can be generated by formulas (3.7). This follows from Remark 3.3 and the discussion before Theorem 3.4. Note that several other interesting examples for the case of two parameters are given in Yang and Stufken (2009). All these examples are based on Lemma 1 from that paper and the conditions of this lemma are in fact imply that the system of the three functions (corresponding to different elements of the information matrix) is an extended Chebyshev system. Thus, these examples can also be considered as particular cases of Theorem 3.1.

The main advantage of Theorem 3.1 consists in the fact that the de la Garza phenomenon can be established by proving that the system under consideration is a Chebyshev system. For this purpose, several methods are available which differ from the approach presented in Yang (2010) and in the next section we will consider an example illustrating the usefulness of Theorem 3.1.

**4. An application to rational regression models.** In this section, we present a class of nonlinear regression models where Theorem 3.4 [or Theorem 2 in Yang (2010)] is not directly applicable, but the de la Garza phenomenon can be established by an application of Theorem 3.1. For this purpose, we consider rational regression models of the form

$$(4.1) \quad \eta(x, \theta) = \frac{P(x, \theta_{(1)})}{Q(x, \theta_{(2)})},$$

where

$$P(x, \theta_{(1)}) = \theta_1 + \theta_2 x + \dots + \theta_l x^{(l-1)},$$

$$Q(x, \theta_{(2)}) = 1 + \theta_{l+1} x + \dots + \theta_{s+l} x^s$$

are polynomials of degree  $l - 1$  and  $s$ , respectively, with corresponding parameters

$$\theta_{(1)} = (\theta_1, \dots, \theta_l)^T, \quad \theta_{(2)} = (\theta_{l+1}, \dots, \theta_{l+s})^T.$$

It is shown in He, Studden and Sun (1996) that the information matrix for this model can be written in the form

$$M(\xi, \theta) = B(\theta)C(\xi, \theta)B(\theta),$$

where  $\theta = (\theta_1, \dots, \theta_{l+s})^T$ ,  $B$  denotes an appropriate matrix [see He, Studden and Sun (1996)], the matrix  $C$  is given by

$$C(\xi, \theta) = \int_A^B [1/Q^4(x)]h(x)h(x)^T d\xi(x),$$

$h(x) = (1, x, \dots, x^{p-1})^T$  denotes the vector of monomials with  $p = l + s$  and  $Q(x)$  is a polynomial of degree  $s$ . Therefore, it follows that the different functions in the information matrix are given by

$$\Psi_1(x) = 1/Q^4(x), \dots, \Psi_k(x) = x^{k-1}/Q^4(x),$$

where  $k = 2p - 1$ . Define  $\Psi_0(x) = 1$ , then it is well known [see Karlin and Studden (1966a)] that under the conditions:

- (a)  $Q(x)$  does not vanish in the interval  $[A, B]$ ;
- (b)  $[Q^4(x)]^{(2p-1)}$  does not vanish in the interval  $[A, B]$

the functions  $\Psi_0, \Psi_1, \dots, \Psi_{2p-1}$  generate a Chebyshev system on the interval  $[A, B]$  and Theorem 3.1 is applicable here.

However, we will give an alternative proof of this property which yields—as a by-product—a constructive condition under which the condition (b) is fulfilled. Assume that  $Q^4(x) > 0$  for all  $x \in [A, B]$  and note that a Chebyshev system remains a Chebyshev system after multiplication of all functions by a positive function. Thus, in order to apply Theorem 3.1 it is sufficient to prove that the functions

$$1, x, x^2, \dots, x^{2p-2}, -Q^4(x)$$

generate a Chebyshev system on the interval  $[A, B]$ . The following lemma provides a sufficient condition for this property.

LEMMA 4.1. *Assume that the polynomial  $Q(x)$  has only real roots which are either all smaller than  $A$  or larger than  $B$ . If  $s > l - 1$ , then the functions*

$$1, x, x^2, \dots, x^{2p-2}, \epsilon Q^4(x),$$

*generate a Chebyshev system on the interval  $[A, B]$ , where  $\epsilon = +1$  if the roots are smaller than  $A$  and  $\epsilon = -1$  if the roots larger than  $B$ .*

PROOF. Based on the assumptions about  $Q(x)$ , the polynomial  $Q^4(x)$  can be written as  $c \prod_{i=1}^{4s} (x - \alpha_i)$ , where  $\alpha_i$  are not necessary distinct. Clearly,  $(Q^4(x))^{(k)} = c \sum_{A_k} \prod_{j \in A_k} (x - \alpha_j)$ , where  $A_k$  is the set of all possible subsets of  $\{1, \dots, 4s\}$  with  $4s - k$  elements. Define  $x_{\min}$  and  $x_{\max}$  as the smallest and largest root of  $Q(x)$ , then all derivatives of  $Q^4(x)$  of even order less than  $4s - 1$  are positive outside of the interval  $[x_{\min}, x_{\max}]$ . Define  $u_0(x) = 1, u_1(x) = x, \dots, u_{2p-2}(x) = x^{2p-2}, u_{2p-1}(x) = Q^4(x)$ . By formulas (3.11) and (3.12), we can easily calculate that  $w_0(x) = 1, w_j(x) = j, j = 1, \dots, 2p - 2, w_{2p-1}(x) = [Q^4(x)]^{(2p-1)}$ . Thus, if  $s > l - 1$  it follows that  $w_{2p-1}(x)$  is negative for  $x < x_{\min}$  and positive for  $A > x > x_{\max}$ . Therefore (note that  $[Q^4(x)]^{(2p-1)}$  has no roots in the interval  $[A, B]$ ), we have  $w_{2p-1}(x) > 0$  for all  $x \in [A, B]$ . Now the assertion of Lemma 4.1 follows from the formula for the Wronskian determinant in (3.10) and the fact that a positive Wronskian determinant is sufficient for the Chebyshev property of the functions  $u_0, \dots, u_{2p-1}$ .  $\square$

The following result is now an immediate consequence of Lemma 4.1 and Theorem 3.1 (note that we do not repeat the statement of uniqueness of the latter result).

**THEOREM 4.2.** *Consider the rational regression model (4.1). Assume that  $s > l - 1$  and that the polynomial  $Q(x)$  has only real roots, which are either all smaller than  $A$  or larger than  $B$ . Then for any design  $\xi$  there exists a design  $\tilde{\xi}$  with at most  $p$  support points, such that  $M(\xi, \theta) \leq M(\tilde{\xi}, \theta)$ . Moreover:*

- (1) *if the index of  $\xi$  satisfies  $I(\xi) \geq p - \frac{1}{2}$  and all roots of the polynomial  $Q$  are smaller than  $A$ , then  $\tilde{\xi}$  can be chosen such that the support of  $\tilde{\xi}$  contains the point  $A$ ,*
- (2) *if the index of  $\xi$  satisfies  $I(\xi) \geq p - \frac{1}{2}$  and all roots of the polynomial  $Q$  are larger than  $B$ , then  $\tilde{\xi}$  can be chosen such that the support of  $\tilde{\xi}$  contains the point  $B$ .*

**REMARK 4.3.** (a) Theorem 4.2 is an extension of Theorem 5 in He, Studden and Sun (1996) who investigated only locally D-optimal designs.

(b) Note that Yang (2010) considered the classical weighted polynomial regression model where the different functions in the information matrix are given by  $\Psi_j(x) = \lambda(x)x^{j-1}$ ,  $j = 1, \dots, 2p - 1$ , where  $\lambda$  is a positive function on the interior of the design space, which is called efficiency function [see Dette and Trampusch (2010)]. His findings can be generalized in the following way. If there exists a function  $g(x)$  such that

$$(4.2) \quad \left(\frac{d}{dx}\right)^j \left(\frac{d}{dx}(\lambda(x)x^{j-1})/g(x)\right) = c_j, \quad g(x) > 0, x \in [A, B],$$

for some constants  $c_j \in \mathbb{R} \setminus \{0\}$ ,  $j = 1, \dots, 2p - 1$ , then one can denote

$$\hat{\Psi}_1(x) = \int_0^x g(t) dt, \quad \hat{\Psi}_j = \Psi_{j-1}, \quad j = 1, \dots, 2p - 1,$$

and obtains a system of functions satisfying the assumptions of Theorem 3.4. In particular, in Theorem 9 of Yang (2010) for the case  $\lambda(x) = \exp(x^2)$  the function  $g(x) = \lambda(x) = \exp(x^2)$  is appropriate, while the case  $\lambda(x) = (1 - x)^{\alpha+1}(1 + x)^{\beta+1}$ ,  $\alpha > -1$ ,  $\beta > -1$  requires the choice  $g(x) = (1 - x)^\alpha(1 + x)^\beta$ . Moreover, the differential equation (4.2) shows that there are many other efficiency functions for which the de la Garza phenomenon in the weighted polynomial regression model occurs. For example, if  $\lambda(x) = 1/(1 + x)^n$ ,  $A > -1$ ,  $n > 2p - 2$  one could use

$$g(x) = 1/(1 + x)^{(n+2)}$$

and it follows that for the weighted polynomial regression model with this efficiency function any optimal design can be based on at most  $p$  points. However,

for the rational model of the form (4.1) such a technique seemingly does not work. The alternative way is to prove that the functions  $1, x, \dots, x^k, \lambda(x)^{-1}$  generate a Chebyshev system and to use the new Theorem 3.1 to establish the de la Garza phenomenon. Such a method has been realized for the rational model (4.1) in the proof of Theorem 4.2.

**Acknowledgments.** The authors would like to thank Martina Stein, who typed parts of this manuscript with considerable technical expertise. We are also grateful to M. Yang and W. J. Studden for helpful discussions on an earlier version of this paper. The authors would also like to thank two anonymous referees for very constructive comments on an earlier version of this paper.

## REFERENCES

- CHALONER, K. and VERDINELLI, I. (1995). Bayesian experimental design: A review. *Statist. Sci.* **10** 273–304. [MR1390519](#)
- CHERNOFF, H. (1953). Locally optimal designs for estimating parameters. *Ann. Math. Statist.* **24** 586–602. [MR0058932](#)
- DE LA GARZA, A. (1954). Spacing of information in polynomial regression. *Ann. Math. Statist.* **25** 123–130. [MR0060777](#)
- DETTE, H. (1997). Designing experiments with respect to “standardized” optimality criteria. *J. Roy. Statist. Soc. Ser. B* **59** 97–110. [MR1436556](#)
- DETTE, H. and TRAMPISCH, M. (2010). A general approach to  $D$ -optimal designs for weighted univariate polynomial regression models. *J. Korean Statist. Soc.* **39** 1–26. [MR2655806](#)
- DETTE, H. and WONG, W. K. (1996). Optimal Bayesian designs for models with partially specified heteroscedastic structure. *Ann. Statist.* **24** 2108–2127. [MR1421164](#)
- DETTE, H., MELAS, V. B., PEPELYSHEV, A. and STRIGUL, N. (2003). Efficient design of experiments in the Monod model. *J. R. Stat. Soc. Ser. B Stat. Methodol.* **65** 725–742. [MR1998631](#)
- DETTE, H., BRETZ, F., PEPELYSHEV, A. and PINHEIRO, J. (2008). Optimal designs for dose-finding studies. *J. Amer. Statist. Assoc.* **103** 1225–1237. [MR2462895](#)
- FANG, X. and HEDAYAT, A. S. (2008). Locally  $D$ -optimal designs based on a class of composed models resulted from blending  $E_{\max}$  and one-compartment models. *Ann. Statist.* **36** 428–444. [MR2387978](#)
- FORD, I., TORSNEY, B. and WU, C. F. J. (1992). The use of a canonical form in the construction of locally optimal designs for nonlinear problems. *J. Roy. Statist. Soc. Ser. B* **54** 569–583. [MR1160483](#)
- HE, Z., STUDDEN, W. J. and SUN, D. (1996). Optimal designs for rational models. *Ann. Statist.* **24** 2128–2147. [MR1421165](#)
- IMHOF, L. A. (2001). Maximin designs for exponential growth models and heteroscedastic polynomial models. *Ann. Statist.* **29** 561–576. [MR1863970](#)
- IMHOF, L. A. and STUDDEN, W. J. (2001).  $E$ -optimal designs for rational models. *Ann. Statist.* **29** 763–783. [MR1865340](#)
- JENNRICH, R. I. (1969). Asymptotic properties of non-linear least squares estimators. *Ann. Math. Statist.* **40** 633–643. [MR0238419](#)
- KARLIN, S. and STUDDEN, W. J. (1966a). Optimal experimental designs. *Ann. Math. Statist.* **37** 783–815. [MR0196871](#)
- KARLIN, S. and STUDDEN, W. J. (1966b). *Tchebycheff Systems: With Applications in Analysis and Statistics. Pure and Applied Mathematics, Vol. XV.* Wiley, New York. [MR0204922](#)

- KHURI, A. I., MUKHERJEE, B., SINHA, B. K. and GHOSH, M. (2006). Design issues for generalized linear models: A review. *Statist. Sci.* **21** 376–399. [MR2339137](#)
- KIEFER, J. (1974). General equivalence theory for optimum designs (approximate theory). *Ann. Statist.* **2** 849–879. [MR0356386](#)
- MELAS, V. B. (2006). *Functional Approach to Optimal Experimental Design. Lecture Notes in Statistics* **184**. Springer, New York. [MR2193670](#)
- MÜLLER, C. H. and PÁZMAN, A. (1998). Applications of necessary and sufficient conditions for maximin efficient designs. *Metrika* **48** 1–19. [MR1647889](#)
- PÁZMAN, A. (1986). *Foundations of Optimum Experimental Design. Mathematics and Its Applications (East European Series)* **14**. Reidel, Dordrecht. [MR0838958](#)
- PRONZATO, L. and WALTER, E. (1985). Robust experiment design via stochastic approximation. *Math. Biosci.* **75** 103–120. [MR0800967](#)
- PUKELSHEIM, F. (2006). *Optimal Design of Experiments. Classics in Applied Mathematics* **50**. SIAM, Philadelphia, PA. [MR2224698](#)
- RATKOWSKY, D. A. (1990). *Handbook of Nonlinear Regression Models*. Dekker, New York.
- SEBER, G. A. F. and WILD, C. J. (1989). *Nonlinear Regression*. Wiley, New York. [MR0986070](#)
- SILVEY, S. D. (1980). *Optimal Design: An Introduction to the Theory for Parameter Estimation*. Chapman & Hall, London. [MR0606742](#)
- YANG, M. (2010). On the de la Garza phenomenon. *Ann. Statist.* **38** 2499–2524. [MR2676896](#)
- YANG, M. and STUFKEN, J. (2009). Support points of locally optimal designs for nonlinear models with two parameters. *Ann. Statist.* **37** 518–541. [MR2488361](#)

FAKULTÄT FÜR MATHEMATIK  
RUHR-UNIVERSITÄT BOCHUM  
44780 BOCHUM  
GERMANY  
E-MAIL: [holger.dette@rub.de](mailto:holger.dette@rub.de)

DEPARTMENT OF MATHEMATICS  
ST. PETERSBURG STATE UNIVERSITY  
ST. PETERSBURG  
RUSSIA  
E-MAIL: [vbmelas@post.ru](mailto:vbmelas@post.ru)