

ON A GENERALIZATION OF THE WILSON-GLAISHER THEOREM

BY GIOVANNI RICCI

1. *Introduction.* J. W. L. Glaisher* has shown that, if n be any number, p any odd prime not exceeding n , k the integral part of the quotient n/p , and if A_r denote the sum of the products of the first $n-1$ consecutive integers taken r together, then

$$A_{p-1} + k \equiv 0 \pmod{p}.$$

This theorem contains Wilson's theorem as the special case $n=p$, and it has been extended by R. E. Moritz† in the following form. If $n=kp+q$, p an odd prime, $0 \leq q < p$, and if ${}^m A_r$ denote the sum of the products of any $n-1$ consecutive numbers, $m+1, m+2, \dots, m+n-1$ taken r together; if $0 < q < p$, then ${}^m A_{p-1} + k \equiv 0 \pmod{p}$. If $q=0$, then ${}^m A_{p-1} + k \equiv 0$, or $\equiv 1 \pmod{p}$, according as m is, or is not, a multiple of p .

It is the purpose of the present paper to show that the Wilson-Glaisher theorem, the Moritz theorem, and other theorems are special cases of a still more general theorem relating to the symmetric functions of special systems of numbers, these systems being composed of the residues of powers, eventually repeated, for different moduli.

2. *The Generalized Theorem.* We shall prove the following general theorem.

Let $m = p^\alpha q^\beta \dots r^\gamma$, (p, q, \dots, r odd primes, $p < q < \dots < r$; $\alpha \geq 1, \beta \geq 1, \dots, \gamma \geq 1$), be an odd number and let $\rho, \sigma, \dots, \chi$ be any divisors respectively of $\phi(p^\alpha), \phi(q^\beta), \dots, \phi(r^\gamma)$, where $\phi(n)$ denotes Euler's Indicator; we shall write $\rho = p^\alpha \lambda$ (λ divisor

* J. W. L. Glaisher, *Congruences relating to the sums of products of the first n numbers and to other sums of products*, Quarterly Journal of Mathematics, vol. 31 (1900), pp. 1-35; see p. 23. See also L. E. Dickson, *History of the Theory of Numbers*, vol. I, p. 99.

† R. E. Moritz, *On an extension of Glaisher's generalization of Wilson's theorem*, Tôhoku Mathematical Journal, vol. 28 (1927), p. 198-201.

of $p-1$), $\sigma = q^b \mu$ (μ divisor of $q-1$), \dots , $\chi = r^c \nu$ (ν divisor of $r-1$). Let $(u_0, u_1, \dots, u_{\rho-1}), (v_0, v_1, \dots, v_{\rho-1}), \dots, (w_0, w_1, \dots, w_{\chi-1})$ be the complete root systems respectively of the congruences

$$u^{\rho} \equiv 1 \pmod{p^{\alpha}}; \quad v^{\sigma} \equiv 1 \pmod{q^{\beta}}; \quad \dots; \quad w^{\chi} \equiv 1 \pmod{r^{\gamma}}.$$

Consider the $\tau = \rho\sigma \dots \chi$ numbers

$$t_1, t_2, \dots, t_{\tau}$$

two by two incongruent $(\text{mod } m)$, represented by the form

$$A u_d + B v_e + \dots + C w_f,$$

$$(0 \leq d \leq \rho - 1, 0 \leq e \leq \sigma - 1, \dots, 0 \leq f \leq \chi - 1),$$

in which A, B, \dots, C denote auxiliary integers satisfying the congruences

$$A \equiv 1 \pmod{p^{\alpha}}; \quad B \equiv 1 \pmod{q^{\beta}}; \quad \dots; \quad C \equiv 1 \pmod{r^{\gamma}};$$

$$A \equiv 0 \left(\text{mod } \frac{m}{p^{\alpha}} \right); \quad B \equiv 0 \left(\text{mod } \frac{m}{q^{\beta}} \right); \quad \dots; \quad C \equiv 0 \left(\text{mod } \frac{m}{r^{\gamma}} \right).$$

Consider the $k\tau$ integers $t_{j,i}$, ($j=1, 2, \dots, \tau; i=1, 2, \dots, k$), k by k congruent $(\text{mod } m)$, precisely,

$$(1) \quad t_{j,i} \equiv t_j \pmod{m}, \quad (i = 1, 2, \dots, k), \quad j = 1, 2, \dots, \tau,$$

and h other arbitrary integers

$$(2) \quad z_1, z_2, \dots, z_h, \quad (h \geq 0).$$

If $R_{s\tau}(m; \rho, \sigma, \dots, \chi | k, h)$ denotes the sum of the products of integers (1) and (2) taken $s\tau$ together, ($R_0=1$), then from any one of the inequalities

$$h < \lambda, \quad h < \mu, \quad \dots, \quad h < \nu$$

there follows the corresponding congruence

$$(A) \ R_{s\tau}(m; \rho, \sigma, \dots, \chi \parallel k, h) \left\{ \begin{array}{l} \equiv (-1)^{s\tau(\lambda-1)/\lambda} \begin{pmatrix} \frac{k\tau}{\lambda} \\ s\tau \\ \lambda \end{pmatrix} \pmod{p^\alpha}, \\ \dots \\ \equiv (-1)^{s\tau(\mu-1)/\mu} \begin{pmatrix} \frac{k\tau}{\mu} \\ s\tau \\ \mu \end{pmatrix} \pmod{q^\beta}, \\ \dots \\ \equiv (-1)^{s\tau(\nu-1)/\nu} \begin{pmatrix} \frac{k\tau}{\nu} \\ s\tau \\ \nu \end{pmatrix} \pmod{r^\gamma}. \end{array} \right.$$

In the special case $s = k, h < \lambda, h < \mu, \dots, h < \nu$, if the integers

$$\frac{k\tau}{\lambda}(\lambda - 1), \frac{k\tau}{\mu}(\mu - 1), \dots, \frac{k\tau}{\nu}(\nu - 1)$$

are all even or all odd, we have the congruence

$$R_{k\tau}(m; \rho, \sigma, \dots, \chi \parallel k, h) \equiv \pm 1 \pmod{m}.$$

3. *Special Cases.* We observe that the $\phi(m)$ integers \pmod{m} , prime to m , are characterized by their satisfying the congruences $t^{\phi(p^\alpha)} \equiv 1 \pmod{p^\alpha}; \ t^{\phi(q^\beta)} \equiv 1 \pmod{q^\beta}; \ \dots; \ t^{\phi(r^\gamma)} \equiv 1 \pmod{r^\gamma}$.

Therefore we may express Wilson's classic theorem and its generalizations by means of the forms

$$\left. \begin{array}{l} R_{p-1}(p; p - 1 \parallel 1, 0) \equiv -1 \pmod{p} \quad \text{(WILSON)*,} \\ R_{\phi(p^\alpha)}(p^\alpha; \phi(p^\alpha) \parallel 1, 0) \equiv -1 \pmod{p^\alpha} \\ R_{\phi(m)}(m; \phi(p^\alpha), \phi(q^\beta), \dots, \phi(r^\gamma) \parallel 1, 0) \equiv 1 \pmod{m} \end{array} \right\} \text{(GAUSS) \dagger,}$$

$$R_{p-1}(p; p - 1 \parallel k, h) \equiv -k \pmod{p} \quad \text{GLAISHER-MORITZ \ddagger,}$$

* See L. E. Dickson, op. cit., p. 62.

\dagger See L. E. Dickson, op. cit., p. 65.

\ddagger See L. E. Dickson, op. cit., p. 99; and R. E. Moritz, loc. cit.

$$\left. \begin{aligned}
 R_{s\tau}(m; \phi(p^\alpha), \phi(q^\beta), \dots, \phi(r^\gamma) \parallel k, 0) &\equiv (-1)^{s\tau/(p-1)} \left(\frac{\frac{k\tau}{p-1}}{s\tau} \right) \pmod{p^\alpha}, \\
 (\tau = \phi(m) = \phi(p^\alpha) \cdots \phi(r^\gamma)) &\equiv (-1)^{s\tau/(q-1)} \left(\frac{\frac{k\tau}{q-1}}{s\tau} \right) \pmod{q^\beta}, \\
 \text{(M. BAUER)*} &\dots\dots\dots \\
 &\equiv (-1)^{s\tau/(r-1)} \left(\frac{\frac{k\tau}{r-1}}{s\tau} \right) \pmod{r^\gamma}.
 \end{aligned} \right\}$$

For the ω -ic residues $\pmod{p^\alpha}$,

$$R_\rho(p^\alpha; \rho \parallel 1, 0) \equiv (-1)^{\rho-1} \pmod{p^\alpha}; \quad \rho = \frac{\phi(p^\alpha)}{D(\omega, \phi(p^\alpha))} \dagger.$$

And also

$$(3) \quad R_{s\lambda}(m; \rho, \sigma, \dots, \chi \parallel 1, 0) \equiv (-1)^{s(\lambda-1)} \left(\frac{\tau}{\lambda} \right) \pmod{p^\alpha}, \quad \text{(RICCI) } \ddagger$$

4. PROOF. Let $R_{n_i}(m; \rho, \sigma, \dots, \chi \parallel 1, 0)$ denote the sum of the products of the τ numbers $t_{j,i}$, ($j=1, 2, \dots, \tau$), of the system (1) taken n_i together, and let R'_n be the sum of the products of the h numbers z_1, z_2, \dots, z_h taken n together ($R'_n = 1$, if $hn = 0$). Obviously we have the equality

$$\begin{aligned}
 (4) \quad R_{s\tau}(m; \rho, \sigma, \dots, \chi \parallel k, h) \\
 = \sum \left\{ \prod_{i=1}^k R_{n_i}(m; \rho, \sigma, \dots, \chi \parallel 1, 0) \right\} R'_n,
 \end{aligned}$$

* See L. E. Dickson, op. cit., p. 88. (Bauer ¹⁸⁶.)

† See P. Bachmann, *Niedere Zahlentheorie*, Teil I, Leipzig, 1902, p. 347.

‡ See G. Ricci, *Sulle funzioni simmetriche delle radici dell'unit  secondo un modulo composto*, Annali di Matematica, (4), vol. 9 (1931), p. 190, formula (B₄).

in which the sum is extended to the solutions in integers ≥ 0 of the equation

$$n_1 + n_2 + \cdots + n_k + n = s\tau,$$

$$(0 \leq n_i \leq \tau, i = 1, 2, \cdots, k; 0 \leq n \leq h).$$

If $n_i \not\equiv 0 \pmod{\lambda}$, then it is known* that

$$R_{n_i}(m; \rho, \sigma, \cdots, \chi \parallel 1, 0) \equiv 0 \pmod{p^\alpha},$$

and if $n_i = s\lambda$, then the congruence (3) stands. Therefore if one at least of the integers n_1, n_2, \cdots, n_k is $\not\equiv 0 \pmod{\lambda}$, then the corresponding term on the right of (4) is divisible by p^α ; hence, for the relation $0 \leq n \leq h < \lambda$, we obtain

$$R_{s\tau}(m; \rho, \sigma, \cdots, \chi \parallel k, h)$$

$$\equiv \sum \prod_{i=1}^k R_{s_i\lambda}(m; \rho, \sigma, \cdots, \chi \parallel 1, 0) \pmod{p^\alpha},$$

$$\left(s_1 + s_2 + \cdots + s_k = s \frac{\tau}{\lambda}; 0 \leq s_i \leq \frac{\tau}{\lambda} \right).$$

Then, by (3), we may write

$$R_{s\tau}(m; \rho, \sigma, \cdots, \chi \parallel k, h) \equiv (-1)^{s\tau(\lambda-1)/\lambda} \sum \prod_{i=1}^k \binom{\frac{\tau}{\lambda}}{s_i} \pmod{p^\alpha},$$

$$\left(s_1 + s_2 + \cdots + s_k = \frac{s\tau}{\lambda} \right);$$

and, by a well known formula on binomial coefficients, from this congruence we deduce the first formula of (A). We may deduce the other formulas in a similar manner.

R. SCUOLA NORMALE SUPERIORE
PISA, ITALY

* See G. Ricci, loc. cit., formula (B₄).