

A CLASSIFICATION OF MODULES OVER COMPLETE DISCRETE VALUATION RINGS

BY CHANG MO BANG¹

Communicated by Richard Pierce, October 17, 1969

1. Introduction. The purpose of this paper is to announce the completion of a classification (up to isomorphism) of all modules which are direct sums of countably generated modules over complete discrete valuation rings. The detailed proofs will appear elsewhere. Throughout this paper, let R denote a fixed but arbitrary complete discrete valuation ring and p a fixed but arbitrary prime element of R . For the sake of convenience, a cardinal is viewed as the first ordinal having that cardinality. Let (c, R, k) be the class of all countably generated reduced R -modules of (torsion-free) rank $\leq k$ and $D(c, R, k)$ that of all direct sums of members of (c, R, k) . Clearly

$$\begin{array}{ccccccc} (c, R, 0) & \subset & (c, R, 1) & \subset & \cdots & \subset & (c, R, \omega) \\ \cap & & \cap & & & & \cap \\ D(c, R, 0) & \subset & D(c, R, 1) & \subset & \cdots & \subset & D(c, R, \omega). \end{array}$$

Notice that a p -primary abelian group is a member of $(c, R, 0)$, particularly if R is a ring of p -adic integers. A classification (of all members) of (c, R, k) was done by Ulm (1933) when $k=0$ [8], by Kaplansky and Mackey (1951) when $k=1$ [4], by Rotman and Yen (1961) when $k < \omega$ [7], and that of $D(c, R, k)$ was done by Kolettis (1960) when $k=0$ [5]. First, we complete a classification of (c, R, ω) and then, utilizing this, we finish that of $D(c, R, \omega)$.

2. Invariants. We need two kinds of invariants, namely, the Ulm invariants and the basis types. Since the celebrated Ulm invariants are well known, a brief explanation of the basis types only is in order [2], [4], [7]. Let $R^k = \bigoplus \{R : i < k\}$ for each k . Define $f(R)$ to be the class of all ordinal (ordinal or ∞) valued functions on R^k for all cardinals k , and $m(Q)$ that of all square row-finite matrices over Q , the quotient field of R . Suppose that $f, g \in f(R)$. Define $f \sim g$ to mean

AMS Subject Classifications. Primary 2030, 2027; Secondary 1340, 1350.

Key Words and Phrases. Countably generated module, complete discrete valuation ring, classification up to isomorphism, torsion-free rank, Ulm invariant, reduced module, p -adic integers, basis, basic free submodule, basis type, summandwise basis, height-preserving isomorphism, p -height.

¹ The author was supported in part by NSF Grant GP-8725 and GP-11804.

both that $\text{Dom } f = \text{Dom } g = R^k$ for some cardinal k and that there is a matrix γ and a diagonal matrix δ , both $k \times k$ invertible integral (that is, all entries are elements of R) in $m(Q)$, such that $f(\alpha\gamma) = g(\alpha\delta)$ for all $\alpha \in R^k$. It is routine to show that \sim is an equivalence relation on $f(R)$.

Let M be an R -module of rank k . Then, every basis $\eta = \{y_i : i < k\}$ defines a function g of $f(R)$ by

$$g(\alpha) = h_p(\alpha\eta) = h_p(\sum \{a_i y_i : i < k\})$$

for all $\alpha = \{a_i : i < k\} \in R^k$. Notice that $g = \infty$ if $k = 0$ since a sum without term is 0. It is routine to show that $g \sim g'$ if g' is defined by another basis of M . Thus, M determines uniquely a class of $f(R)/\sim$, which we call the *basis type* of M . It is easy to show the following lemma.

LEMMA 1. *Two reduced R -modules M and M' have the same basis type if and only if they contain basic free submodules F and F' , respectively, with a height-preserving isomorphism from F onto F' .*

3. A classification of (c, R, ω) .

THEOREM 1. *Let M and M' be countably generated reduced R -modules. Then, $M \simeq M'$ if and only if they have the same Ulm invariant and the same basis type.*

Only the “if” part needs a proof. Let $\alpha = \{a_i : i < k\}$. Define $\alpha(r) = \{a_i : i < r\}$ for each number r . Let k be the same rank of M and M' . Then, by Lemma 1, there are ordered bases $\eta = \{y_i : i < k\}$ and $\eta' = \{y'_i : i < k\}$ of M and M' , respectively, with a height-preserving isomorphism ρ such that $\rho(\alpha\eta) = \alpha\eta'$ for all $\alpha \in R^k$. We may assume that there are countable subsets $\xi = \{x_i : i < \omega\}$ and $\xi' = \{x'_i : i < \omega\}$ of M and M' , respectively, such that

$$M = [\xi \cup \eta] \quad \text{and} \quad M' = [\xi' \cup \eta'] \quad \text{with}$$

$$px_i \in [\xi(i) \cup \eta(i)] \quad \text{and} \quad px'_i \in [\xi'(i) \cup \eta'(i)] \quad \text{for each } i < \omega.$$

The main idea of the proof is to construct a sequence of height-preserving isomorphisms $\{\phi_i : i < \omega\}$ in such a way that the following conditions are satisfied.

(a) $\phi_i : A_i \rightarrow A'_i$ where

$$A_i = [\xi(i) \cup \eta(i) \cup \phi_i^{-1}(\xi'(i) \cup \eta'(i))],$$

$$A'_i = [\xi'(i) \cup \eta'(i) \cup \phi_i(\xi(i) \cup \eta(i))].$$

(b) $\phi_0 \leq \dots \leq \phi_i \leq \phi_{i+1} \leq \dots$

(c) There exists a nonnegative integer $n(i)$ such that $p^{n(i)}A_i$

$\subseteq [\eta(i)]$ and $p^{n(i)}A'_i = [\eta'(i)]$ and $\phi_i = \rho$ as height-preserving isomorphism from $p^{n(i)}A_i$ onto $p^{n(i)}A'_i$.

The supremum of $\{\phi_i : i < \omega\}$ gives the required isomorphism from M onto M' . For more detailed proof, see [1] or [2].

4. A classification of $D(c, R, \omega)$.

THEOREM 2. *Let M and M' be direct sums of countably generated reduced R -modules. Then, $M \simeq M'$ if and only if they have the same Ulm invariant and the same basis type.*

Again, only the "if" part needs a proof. We may write as

$$M = \bigoplus \{M_i : i \in I\} \quad \text{and} \quad M' = \bigoplus \{M'_i : i \in I\}$$

where all $M_i, M'_i \in (c, R, \omega)$ and I is a cardinal. For notational convenience, define $M(T) = \bigoplus \{M_i : i \in T\}$, $T \subseteq I$. The main idea of the proof is to show that there is a partition of I into countable subsets $\{I_j : j < I\}$ such that, for each $j < I$, $M(I_j)$ and $M'(I_j)$ have the same Ulm invariant and the same basis type. Then by Theorem 1, they are isomorphic and, consequently, $M \simeq M'$. In fact, by the Kolettis theorem [3], [5], [6], we may assume that M_i and M'_i have already the same Ulm invariant for each i . The following lemmas indicate the route of the proof.

LEMMA 2. *Let $N = A \oplus B \oplus C$ be a reduced R -module such that the following conditions are satisfied.*

(a) *There are in N disjoint subsets η_A and η_B such that η_A and $\eta_A \cup \eta_B$ are bases of A and $A \oplus B$, respectively.*

(b) *If $x_A \in [\eta_A]$ and $x_B \in [\eta_B]$, then (h_N denotes the p -height in N)*

$$h_N(x_A + x_B) = \min\{h_N(x_A), h_N(x_B)\}.$$

Then, if we write $\eta_B = \{y_i : i < k\}$, $k = |\eta_B|$, there is in $m(Q)$ a $k \times k$ diagonal invertible integral matrix $\delta = \{d_i : i < k\}$ such that the following conditions are satisfied.

(c) $\tau = \Pi_B(\delta\eta_B)$ *is an ordered basis of B . (Here, Π_B is the canonical projection of N onto B .)*

(d) *There is a height-preserving isomorphism ρ from $[\delta\eta_B]$ onto $[\tau]$ such that $\rho(\alpha\delta\eta_B) = \alpha\tau$ for all $\alpha \in R^k$.*

LEMMA 3. *Let k be the rank of M and M' . Let $\eta = \{y_i : i < k\}$ and $\eta' = \{y'_i : i < k\}$ be ordered bases of M and M' , respectively, with η' summandwise (that is, each $y'_i \in M_j$ for a j). If J is a countable subset of I , then there is a set T such that the following conditions are satisfied.*

(a) *T is countable and $J \subseteq T \subseteq I$.*

(b) Define $\eta(T) = \{y_i \in \eta : y_i \in M(T)\}$. $\eta(T)$ and $\eta'(T)$ are bases of $M(T)$ and $M'(T)$, respectively.

(c) $y_i \in \eta(T)$ if and only if $y'_i \in \eta'(T)$.

LEMMA 4. M and M' have the same basis type if and only if there is a partition of I into countable subsets $\{I_j : j < I\}$ such that $M(I_j)$ and $M'(I_j)$ have the same basis type for each index $j < I$.

Using Lemma 2, 3 and a transfinite induction, we can prove Lemma 4. Theorem 2 is immediate from Lemma 4.

COROLLARY. $M \simeq M'$ if and only if they have isomorphic torsion parts and contain basic free submodules F and F' of, respectively, with a height-preserving isomorphism from F onto F' .

The author would like to acknowledge his gratitude to Professor Charles K. Megibben for suggesting the problems solved in this paper and for his invaluable guidance.

BIBLIOGRAPHY

1. C. Bang, *Countably generated modules over complete discrete valuation rings*, J-Algebra (to appear).
2. ———, *Modules over complete discrete valuation rings*, Dissertation, Vanderbilt University, Nashville, Tenn., 1969.
3. P. Hill, *Sums of countable primary groups*, Proc. Amer. Math. Soc. **17** (1966), 1469–1470. MR **33** #7408.
4. I. Kaplansky and G. Mackey, *A generalization of Ulm's theorem*, Summa Brasil. Math. **2** (1951), 195–202. MR **14**, 128.
5. G. Kolettis, Jr., *Direct sums of countable groups*, Duke Math. J. **27** (1960), 111–125.
6. F. Richman and E. Walker, *Extending Ulm's theorem without group theory*, Proc. Amer. Math. Soc. **21** (1969), 194–196.
7. J. Rotman and T. Yen, *Modules over a complete discrete valuation ring*, Trans. Amer. Math. Soc. **98** (1961), 242–254. MR **23** #A227.
8. H. Ulm, *Zur Theorie der abzählbar-unendlichen Abelschen Gruppen*, Math. Ann. **107** (1933), 774–803.

EMORY UNIVERSITY, ATLANTA, GEORGIA 30322 AND
VANDERBILT UNIVERSITY, NASHVILLE, TENNESSEE 37203.