

ON UNIQUENESS OF A SOLUTION OF $Lu = u^\alpha$ WITH GIVEN TRACE

S. E. KUZNETSOV¹

*Department of Mathematics, University of Colorado, Boulder, CO 80309-0395
USA*

email: Sergei.Kuznetsov@Colorado.edu

submitted August 12, 1998 Final version accepted May 7, 1999

AMS 2000 Subject classification: 35J67, 35J75, 60J50, 60J60, 60J85, 60H30

superdiffusion, moderate solutions, sigma-moderate solutions, stochastic boundary values, trace of a solution, explosion points

Abstract

A boundary trace (Γ, ν) of a solution of $\Delta u = u^\alpha$ in a bounded smooth domain in \mathbb{R}^d was first constructed by Le Gall [12] who described all possible traces for $\alpha = 2, d = 2$ in which case a solution is defined uniquely by its trace. In a number of publications, Marcus, Véron, Dynkin and Kuznetsov gave analytic and probabilistic generalization of the concept of trace to the case of arbitrary $\alpha > 1, d \geq 1$. However, it was shown by Le Gall [13] that the trace, in general, does not define a solution uniquely in case $d \geq (\alpha + 1)/(\alpha - 1)$. He offered a sufficient condition for the uniqueness and conjectured that a uniqueness should be valid if the singular part Γ of the trace coincides with the set of all explosion points of the measure ν . Here, we establish a necessary condition for the uniqueness which implies a negative answer to the above conjecture.

1 Introduction and Results

1.1 Moderate solutions

Let L be a second order uniformly elliptic differential operator with smooth coefficients in \mathbb{R}^d and let $E \subset \mathbb{R}^d$ be a bounded smooth domain. We consider a class \mathcal{U} of all positive solutions of the equation

$$Lu = u^\alpha \quad \text{in } E \tag{1.1}$$

where $\alpha \in (1, 2]$ is a parameter. A solution u is called *moderate* if $u \leq h$ for an L -harmonic function h . The class of all moderate solutions is denoted by \mathcal{U}_1 .

For every moderate solution u , there exists a minimal L -harmonic function that dominates u . It is called the minimal (L -harmonic) majorant of u . A solution u can be recovered from

¹PARTIALLY SUPPORTED BY NATIONAL SCIENCE FOUNDATION GRANTS DMS-9623190 AND DMS-9971009

its majorant as the maximal solution to (1.1) dominated by h . Moreover, u is related to its minimal majorant h by the integral equation

$$u(x) + \Pi_x \int_0^\zeta u^\alpha(\xi_s) ds = h(x). \quad (1.2)$$

Here (ξ_t, Π_x) is the corresponding L -diffusion in E and ζ is its life time. See [5] for more detail. Every positive L -harmonic function h has a unique representation

$$h(x) = \int_{\partial E} k(x, y) \nu(dy) \quad (1.3)$$

where $k(x, y)$ is the Poisson kernel for L in E and ν is a finite measure on ∂E . We denote by h_ν the function given by (1.3). For a moderate solution $u \in \mathcal{U}_1$, we write $u = u_\nu$ if h_ν is the minimal majorant of u .

1.2 Superdiffusions and stochastic boundary values

An (L, α) -superdiffusion is a probabilistic model for an evolution of a random cloud of branching particles. A spatial movement of particles is described by an L -diffusion, and $\alpha \in (1, 2]$ characterizes branching. See, for instance, [2]. To every open set D there corresponds a random measure (X_D, P_μ) on ∂D , called the exit measure from D . It represents the total accumulation of mass on ∂D assuming that the evolution starts from μ and particles are instantly frozen if they reach the complement of D . Relations between X_D and equation (1.1) can be described as follows. Let f be a positive continuous function on ∂E . The function

$$u(x) = -\log P_x e^{-\langle f, X_E \rangle}, \quad (1.4)$$

where P_x stands for P_{δ_x} , is the only solution of the boundary value problem

$$\begin{aligned} Lu &= u^\alpha & \text{in } E, \\ u &= f & \text{on } \partial E. \end{aligned} \quad (1.5)$$

An arbitrary solution u of (1.1) can also be represented in a form similar to (1.4) in terms of its *stochastic boundary value* Z_u (cf. [3]). It can be defined as a limit

$$Z_u = \lim \langle u, X_{D_n} \rangle \quad (1.6)$$

where D_n is an increasing sequence of bounded smooth domains approximating E . A solution u can be recovered from its stochastic boundary value by the formula

$$u(x) = -\log P_x e^{-Z_u}. \quad (1.7)$$

We write Z_ν instead of Z_{u_ν} . See [3] for more detail.

We define the range \mathcal{R} of a superdiffusion in E as the minimal closed set that supports all X_D for $D \subset E$. A set $\Gamma \subset \partial E$ is called a polar set for the superdiffusion if, for any x , $P_x\{\mathcal{R} \cap \Gamma \neq \emptyset\} = 0$. According to [6], the class of polar sets coincides with the class of all removable boundary singularities for the equation (1.1). By [4], the equation (1.2) has a solution if and only if the corresponding measure ν does not charge polar sets. Therefore the mapping $\nu \rightarrow u_\nu$ defines a 1-1 correspondence between the class \mathcal{N}_1 of all finite measures on

∂E which don't charge polar sets and the class \mathcal{U}_1 of all moderate solutions of (1.1); see [5], [4], [9], [7].

For every Borel subset $B \subset \partial E$,

$$w_B(x) = -\log P_x\{\mathcal{R} \cap B = \emptyset\} \tag{1.8}$$

is a solution of (1.1). Its stochastic boundary value is given by the formula $Z_B = Z_{w_B} = \infty 1_{\{\mathcal{R} \cap B \neq \emptyset\}}$. If B is closed, then w_B is the maximal solution of (1.1) such that $w_B = 0$ on $\partial E \setminus B$. See [3], Sect. 6.

1.3 σ -moderate solutions

A solution u of (1.1) is called σ -moderate if there exists an increasing sequence of moderate solutions u_n such that $u_n \uparrow u$ as $n \rightarrow \infty$. It follows from (1.2) that the corresponding measures ν_n also increase to some measure ν . The measure ν does not charge polar sets, but it may be not finite and not even σ -finite. However, it is always Σ -finite. We denote by \mathcal{N}_0 the class of all Σ -finite measures that don't charge polar sets. Every measure $\nu \in \mathcal{N}_0$ can be represented as a limit of an increasing sequence of finite measures ν_n and therefore defines a σ -moderate solution $u = \lim u_{\nu_n}$. We denote this solution by u_ν and we write Z_ν for its stochastic boundary value. (It follows from [9], Theorem 4.2 that u_ν and Z_ν do not depend on the choice of $\nu_n \uparrow \nu$.) Every σ -moderate solution can be represented this way. However, in contrast to moderate solutions, this representation is not unique. σ -moderate solutions have been studied in Section 4 of [9] by means of continuous linear additive functionals.

The class of all σ -moderate solutions is denoted by \mathcal{U}_0 . Existence of non- σ -moderate solutions remains an open question: all known elements of \mathcal{U} either belong to \mathcal{U}_0 or, at least, it is not proved that this is not true. See [11], [7].

1.4 Sweeping and the trace

First definition of the trace was introduced by Le Gall [12], [14], [13], who used it to describe all solutions of the equation $\Delta u = u^2$ in a smooth planar domain. In a more general setting, a definition of a trace was introduced by Marcus and Véron [15], [16], [17], [18] and, in a probabilistic way, by Dynkin and Kuznetsov [9], [8].

Let $u \in \mathcal{U}$. For a closed set $B \subset \partial E$, we define $Q_B(u)$ as the maximal element of \mathcal{U} such that $Q_B(u) \leq u$ and $Q_B(u) = 0$ on $\partial E \setminus B$. We consider the maximal open subset O of ∂E such that $Q_B(u)$ is moderate for every compact $B \subset O$ and we set $\Gamma = O^c$. It can be shown that there exists a Radon measure ν on O such that $Q_B(u) = u_{\nu_B}$ for every compact $B \subset O$ where ν_B stands for the restriction of ν to B . The pair (Γ, ν) is called the trace of u . Cf. [9].

Let ν be a measure on ∂E . A point $x \in \partial E$ is called an explosion point for ν if $\nu(O) = \infty$ for every open set O containing x . The collection of all explosion points of ν is denoted by $Ex(\nu)$. Clearly, $Ex(\nu)$ is a closed set. Let Γ be a closed subset of ∂E and ν be a Radon measure on Γ^c not charging polar sets. The pair (Γ, ν) is called *normal* if there exists no nontrivial relatively open polar subset $B \subset \Gamma \setminus Ex(\nu)$.

Proposition 1.1 (See [9]). *The trace (Γ, ν) of a solution $u \in \mathcal{U}$ is always a normal pair. Each normal pair (Γ, ν) is the trace of some solution u . The maximal solution with the given trace (Γ, ν) is given by the formula*

$$w_{\Gamma, \nu}(x) = -\log P_x\{\mathcal{R} \cap \Gamma = \emptyset, e^{-Z_\nu}\}. \tag{1.9}$$

1.5 Essential explosion points

For an arbitrary Borel set $B \subset \partial E$, put

$$\text{Cap}_R(B) = P_c\{\mathcal{R} \cap B \neq \emptyset\} \quad (1.10)$$

where c is a reference point and \mathcal{R} is the range of the (L, α) -superdiffusion in E . According to [1], Theorem III.32, $\text{Cap}_R(B)$ is a Choquet capacity. By [3], Sect. 6.2, $\text{Cap}_R(B) = 0$ if and only if B is polar.

Let $x \in Ex(\nu)$. We call x a point of *non-essential explosion* if there exists a neighborhood U of x and a sequence of open sets $O_n \subset U$ such that $\text{Cap}_R(O_n) \downarrow 0$ as $n \rightarrow \infty$ and $\nu(U \setminus O_n) < \infty$ for all n . Otherwise x is called a point of *essential explosion*. We denote the set of all essential explosion points by $Ess(\nu)$. Note that $Ex(\nu) = Ess(\nu)$ if single points on the boundary are not polar (this happens if $d < (\alpha + 1)/(\alpha - 1)$; see [10], [6]).

Properties of $Ess(\nu)$ can be summarized as follows.

Theorem 1.1. *Let ν be a Σ -finite measure that doesn't charge polar sets. Then:*

- (i) *The set $Ess(\nu)$ is always closed;*
- (ii) *There exist open sets $U_n \supset Ess(\nu)$ such that $\text{Cap}_R(U_n) \downarrow \text{Cap}_R(Ess(\nu))$ and $\nu(U_n^c) < \infty$;*
- (iii) *$Ess(\nu)$ is either empty or non-polar;*
- (iv) *ν is σ -finite on the complement of $Ess(\nu)$.*

The following result clarifies the difference between $Ex(\nu)$ and $Ess(\nu)$.

Theorem 1.2. *Let $d \geq (\alpha + 1)/(\alpha - 1)$. For every closed set $\Gamma \subset \partial E$, there exists a σ -finite measure $\nu \in \mathcal{N}_0$ such that $Ex(\nu) = \Gamma$ and $Ess(\nu)$ is empty. The measure ν can be chosen to have a density with respect to the surface measure.*

Main result of the paper is given by the following

Theorem 1.3. *Let $d \geq (\alpha + 1)/(\alpha - 1)$ and let (Γ, ν) be a normal pair. If $\Gamma \setminus Ess(\nu)$ is not polar, then there exist at least two solutions with the trace (Γ, ν) .*

Remarks. 1. For $\nu = 0$ this is, essentially, Proposition 3 in [13] (to be precise, [13] is devoted to the initial trace of the corresponding semilinear parabolic equation. However, the arguments can be easily extended to an elliptic case.)

2. Le Gall proved in [13] that the uniqueness takes place if Γ is polar and that there is no uniqueness if $\nu = 0$ and Γ is not polar. He also conjectured that the uniqueness is valid if $\Gamma = Ex(\nu)$. The following example shows that it is not true. Let Γ be a non-polar closed subset of ∂E with the surface measure 0 and let ν be the measure constructed in Theorem 1.2. Since ν has the density with respect to the surface measure, $\nu(\Gamma) = 0$. By Theorem 1.2, $\Gamma = Ex(\nu)$ and therefore (Γ, ν) is a normal pair. However, $Ess(\nu)$ is empty. By Theorem 1.3, there exist at least two solutions with the trace (Γ, ν) . This implies a negative answer to the above conjecture.

1.6 Remarks on fine trace

To overcome the difficulties related to the non-uniqueness, Dynkin and Kuznetsov have defined in [11], [7] a fine trace of a solution and they have shown that σ -moderate solutions are defined uniquely by their fine traces. The fine trace is a pair (Γ, ν) where Γ is closed in a certain topology (fine topology) related to the equation (1.1) and ν is a σ -finite measure on the complement of Γ not charging polar sets. See [11], [7] for all details.

Theorem 1.4. *Let ν be a Σ -finite measure that doesn't charge polar sets. If $\text{Ess}(\nu)$ is empty, then the fine trace of u_ν is equal to (Γ, ν) with polar Γ .*

By applying this result to measures ν constructed in Theorem 1.2, we see that the corresponding solutions u_ν are not determined by their traces and they can be recovered from their fine traces.

2 Proofs

2.1 Some Lemmas

We start with an important lemma.

Lemma 2.1. *Let $u \leq v$ be two solutions of (1.1). If $u(c) = v(c)$ at some interior point $c \in E$, then $u = v$ everywhere in E .*

Proof. It is sufficient to show that $u = v$ in any bounded smooth domain D such that $c \in D$ and $\bar{D} \subset E$. Solutions u and v are bounded and continuous in D and therefore they admit a representation

$$u(x) = -\log P_x e^{-\langle u, X_D \rangle}, \quad v(x) = -\log P_x e^{-\langle v, X_D \rangle}. \quad (2.1)$$

Since $u \leq v$, we conclude from (2.1) that $\langle u, X_D \rangle = \langle v, X_D \rangle$ P_c -a.s. Therefore

$$\Pi_c u(\xi_{\tau_D}) = P_c \langle u, X_D \rangle = P_c \langle v, X_D \rangle = \Pi_c v(\xi_{\tau_D})$$

and $u = v$ on ∂D . By (2.1), this yields $u = v$ in D . \square

As a next step, we compute the trace of w_B for Borel B .

Lemma 2.2. *Let B be a Borel subset of ∂E . The trace of w_B is equal to $(\Gamma, 0)$ where Γ is the smallest closed set such that $B \setminus \Gamma$ is polar.*

Proof. Let (Γ, ν) be the trace of w_B . Suppose $B \setminus \Gamma$ is not polar. There exists a non-polar compact $K \subset B \setminus \Gamma$. Since $K \subset B$, $w_B \geq w_K$ and therefore the sweeping $Q_K(w_B) \geq Q_K(w_K) = w_K$. But, w_K is not moderate.

Suppose now that a closed $\tilde{\Gamma}$ is such that $B \setminus \tilde{\Gamma}$ is polar. Then $w_B \leq w_{\tilde{\Gamma}}$ by (1.8) and therefore $Q_K(w_B) \leq Q_K(w_{\tilde{\Gamma}})$ for every K . This implies $Q_K(w_B) = 0$ for every compact $K \subset \tilde{\Gamma}^c$, and therefore $\Gamma \subset \tilde{\Gamma}$. Same argument applied to Γ instead of $\tilde{\Gamma}$ shows that $\mu = 0$. \square

Lemma 2.3. *Let $\nu \in \mathcal{N}_0$. The trace of u_ν is equal to (Γ, μ) where $\Gamma = \text{Ex}(\nu)$ and μ coincides with the restriction of ν to Γ^c .*

Proof. Let $B \subset \partial E$ be a compact. Clearly, $u_\nu \geq u_{\nu_B}$ where ν_B is the restriction of ν to B . Therefore

$$Q_B(u_\nu) \geq Q_B(u_{\nu_B}) \geq u_{\nu_B}. \quad (2.2)$$

Suppose now that $B \cap Ex(\nu) = \emptyset$. Show that

$$Q_B(u_\nu) = u_{\nu_B} \quad (2.3)$$

for such B . There exists a relatively open $U \subset \partial E$ such that $B \subset U$ and $\nu(U) < \infty$. Let λ and κ be the restrictions of ν to U and U^c , respectively. Let $u_1 = u_\lambda$ and $u_2 = u_\kappa$. By [3], Theorem 2.3,

$$u \leq u_1 + u_2$$

and therefore

$$Q_B(u) \leq Q_B(u_1) + Q_B(u_2). \quad (2.4)$$

However, solution u_1 is moderate and therefore $Q_B(u_1) = u_{\lambda_B} = u_{\nu_B}$. On the other hand, κ vanishes on U and therefore $Q_B(u_2) = 0$ by [9], 4.4.A and Theorem 3.1. Combining (2.2) and (2.4), we get (2.3).

The statement of the lemma follows from (2.2), (2.3) and the definition of the trace. \square

Lemma 2.4. *For every Borel set $B \subset \partial E$, and every $\nu \in \mathcal{N}_0$,*

$$w_{B,\nu}(x) = -\log P_x\{\mathcal{R} \cap B = \emptyset, e^{-Z_\nu}\}$$

is a solution of (1.1). Its trace (Γ, μ) can be characterized by the following properties. The set Γ is the smallest closed set such that $\Gamma \supset Ex(\nu)$ and $B \setminus \Gamma$ is polar, and μ is the restriction of ν to Γ^c .

Proof. Note that $w_{B,\nu}(x) = -\log P_x e^{-Z_B - Z_\nu}$ and therefore the first part follows easily from Theorems 2.3 and 6.1 in [3]. Second part follows easily from Lemmas 2.2 and 2.3 and from the inequalities

$$w_B \leq w_{B,\nu}, \quad u_\nu \leq w_{B,\nu}, \quad w_{B,\nu} \leq u_\nu + w_B$$

(for the last inequality, see, e.g., [3], Theorem 2.3). \square

Lemma 2.5. *Let Γ be a closed subset of ∂E and let $B \supset \Gamma$ be such that $B \setminus \Gamma$ is not polar. Then $\text{Cap}_R(B) > \text{Cap}_R(\Gamma)$.*

Proof. Suppose $\text{Cap}_R(B) = \text{Cap}_R(\Gamma)$. Then $w_B(c) = w_\Gamma(c)$ and therefore $w_B = w_\Gamma$ everywhere by Lemma 2.1. By assumption, there exists a compact $K \subset B \setminus \Gamma$ such that $\text{Cap}_R(K) > 0$. Clearly, $w_K \leq w_B$ and therefore $w_K \leq w_\Gamma$. However, $w_K = 0$ on $\partial E \setminus K$ and $w_\Gamma = 0$ on $\partial E \setminus \Gamma$, which implies $w_K = 0$ on ∂E and therefore $w_K = 0$ in E , that is $\text{Cap}_R(K) = 0$. \square

2.2 Proof of Theorem 1.1

1°. Let $x \notin Ess(\nu)$. If $x \notin Ex(\nu)$, then there exists an open set $U \subset \partial E$ such that $x \in U$ and $\nu(U) < \infty$. By definition of explosion points, all points of U do not belong to $Ex(\nu) \supset Ess(\nu)$. Suppose now that x is a point of non-essential explosion and U is as in definition. Clearly, any $y \in Ex(\nu) \cap U$ must also be a point of non-essential explosion. Therefore each point $x \notin Ess(\nu)$ has a neighborhood that does not intersect with $Ess(\nu)$. Hence $Ess(\nu)$ is closed.

2°. To prove (ii), it is enough to show that, for every $\varepsilon > 0$, there exists an open set $O \supset \text{Ess}(\nu)$ such that $\text{Cap}_R(O) < \text{Cap}(\text{Ess}(\nu)) + \varepsilon$ and $\nu(O^c) < \infty$.

Let $U \supset \text{Ess}(\nu)$ be an open set such that $\text{Cap}_R(U) \leq \text{Cap}_R(\text{Ess}(\nu)) + \varepsilon$. Denote $B = U^c$ and $F = B \cap \text{Ex}(\nu)$. By construction, F consists of non-essential explosion points. For $x \in F$, denote by U_x the neighborhood of x described in the definition of a non-essential explosion point. Open sets U_x cover a compact set F and therefore there exists a finite set $x_1, \dots, x_k \in F$ such that $F \subset U_{x_1} \cup \dots \cup U_{x_k}$. For each x_j , there exists an open set $O_j \subset U_{x_j}$ such that $\nu(U_{x_j} \setminus O_j)$ is finite and $\text{Cap}_R(O_j) \leq \varepsilon/k$. Put $O = U \cup O_1 \cup \dots \cup O_k$. By construction,

$$\text{Cap}_R(O) \leq \text{Cap}_R(U) + \sum_j \text{Cap}_R(O_j) \leq \text{Cap}_R(\text{Ess}(\nu)) + 2\varepsilon$$

On the other hand, the set O^c is contained in the union of the sets $K_j = U_{x_j} \setminus O_j$ and the set $K = E \setminus \{U \cup U_{x_1} \cup \dots \cup U_{x_k}\}$. Sets K_j have finite measure by construction. The set K is a compact set disjoint from $\text{Ex}(\nu)$ and therefore $\nu(K) < \infty$. (Recall that ν is a Radon measure on the complement of $\text{Ex}(\nu)$ and therefore $\nu(K) < \infty$ for every compact K that contains no explosion points.)

3°. Suppose $\text{Ess}(\nu)$ is polar. Let O_n be the sequence constructed in (ii). Put $U = E$. Since $\text{Cap}_R(O_n) \downarrow 0$ and $\nu(E \setminus O_n) < \infty$, all explosion points are non-essential.

4°. Again, let O_n be the sequence constructed in (ii). Denote $B = \bigcap_n O_n$. By construction, ν is σ -finite on B^c . Besides, $\text{Cap}_R(B) = \text{Cap}_R(\text{Ess}(\nu))$ and therefore $B \setminus \text{Ess}(\nu)$ is polar by Lemma 2.5. Hence $\nu(B \setminus \text{Ess}(\nu)) = 0$ and ν is σ -finite on the complement of $\text{Ess}(\nu)$ as well. \square

2.3 Essential explosion points and stochastic boundary values

Lemma 2.6. *Let $\nu \in \mathcal{N}_0$. Then Z_ν is finite a.s. on the set $\{\mathcal{R} \cap \text{Ess}(\nu) = \emptyset\}$.*

Proof. Let K be a compact with $\nu(K) < \infty$ and let ν_K be the restriction of ν to K . The solution u_{ν_K} is moderate, and therefore Z_{ν_K} is finite a.s. As a first step, we prove that

$$Z_\nu = Z_{\nu_K} \quad \text{a.s. on the set } \{\mathcal{R} \cap K^c = \emptyset\}. \quad (2.5)$$

Indeed, let $B \subset K^c$ be a compact. Then $u_{\nu_B} \leq w_B$ and

$$Z_{\nu_B} \leq Z_{w_B} = \infty 1_{\{\mathcal{R} \cap B \neq \emptyset\}}$$

and therefore $Z_{\nu_B} = 0$ on $\{\mathcal{R} \cap K^c = \emptyset\}$. Let now B_n be an increasing sequence of compacts with the union K^c . Since ν is the increasing limit of $\nu_n = \nu_K + \nu_{B_n}$,

$$Z_\nu = \lim Z_{\nu_n} = \lim Z_{\nu_K} + Z_{\nu_{B_n}}$$

(see [3], Sect. 3.8) and therefore

$$Z_\nu = Z_{\nu_K} \quad \text{on } \{\mathcal{R} \cap K^c = \emptyset\}.$$

By Theorem 1.1(ii), there exists a decreasing sequence of open sets O_n such that $\nu(O_n^c) < \infty$, $\text{Ess}(\nu) \subset O_n$ and

$$P_c\{\mathcal{R} \cap O_n \neq \emptyset\} \downarrow P_c\{\mathcal{R} \cap \text{Ess}(\nu) \neq \emptyset\}. \quad (2.6)$$

Denote by K_n the complement of O_n . By (2.5), Z_ν is a.s. finite on the sets $A_n = \{\mathcal{R} \cap O_n = \emptyset\}$. On the other hand, $A_n \uparrow \{\mathcal{R} \cap \text{Ess}(\nu) = \emptyset\}$ by (2.6). \square

2.4 Proof of Theorem 1.2

In order to construct the measure ν , take an arbitrary countable dense subset $\{x_k\}$ of Γ . For every k , let $\phi_k(x) = |x - x_k|^{-d}$. ϕ_k is a continuous positive function on $\partial D \setminus \{x_k\}$, which is not integrable over every neighborhood of x_k . The assumption $d \geq (\alpha + 1)/(\alpha - 1)$ implies $\text{Cap}_R(x_k) = 0$. Hence $\lim_{n \rightarrow \infty} \text{Cap}_R\{\phi_k > n\} = 0$ and therefore $\text{Cap}_R\{\phi_k > C_k\} \leq 2^{-k}$ for sufficiently large C_k .

Put $f = \sup_k \phi_k/C_k$ and $\nu(dx) = f(x)\sigma(dx)$, where $\sigma(dx)$ is the surface measure. By increasing C_k , if necessary, we can make f bounded on a positive distance from Γ . For this reason, $Ex(\nu) \subset \Gamma$. On the other hand, x_n are dense in Γ . Therefore, if $x \in \Gamma$ and U is a neighborhood of x , then U contains at least one of the points x_n and therefore f is not integrable over U . Hence $\Gamma = Ex(\nu)$.

Denote $O_n = \{f > n\} = \cup_k \{\phi_k > nC_k\}$. Since ϕ_k are continuous, O_n is open for every n . Besides, $\nu(O_n^c) \leq n\sigma(O_n^c) < \infty$ for every n . By construction, $\text{Cap}_R\{O_n\} \leq \sum \text{Cap}_R\{\phi_k > nC_k\} \rightarrow 0$ as $n \rightarrow \infty$ by the dominated convergence theorem. Hence, ν has no essential explosion points. \square

2.5 Proof of Theorem 1.3

1°. Suppose $\Gamma = Ex(\nu)$ and $Ex(\nu) \setminus Ess(\nu)$ is not polar. By Theorem 1.1(i), both $Ex(\nu)$ and $Ess(\nu)$ are closed. Therefore Lemma 2.5 implies that $\text{Cap}_R(Ex(\nu)) > \text{Cap}_R(Ess(\nu))$ and therefore

$$P_c\{\mathcal{R} \cap Ex(\nu) \neq \emptyset, \mathcal{R} \cap Ess(\nu) = \emptyset\} > 0. \quad (2.7)$$

Let $v = w_{\Gamma, \nu}$ and $u = u_\nu$. By Proposition 1.1, v is the maximal solution with the trace (Γ, ν) . By Lemma 2.3, u also has the trace (Γ, ν) . However,

$$Z_v \geq Z_\Gamma = \infty 1_{\mathcal{R} \cap \Gamma \neq \emptyset}$$

and

$$Z_u < \infty \quad \text{a.s. on } \mathcal{R} \cap Ess(\nu) = \emptyset$$

by Lemma 2.6. By (2.7), $Z_u \neq Z_v$ with positive probability.

2°. The proof in case $\Gamma \neq Ex(\nu)$ is essentially the same as the proof of Proposition 3 in [13]. Since both Γ and $Ex(\nu)$ are closed, $C = \Gamma \setminus Ex(\nu)$ is relatively open in Γ and therefore it is not polar (see Proposition 1.1 and the definition of a normal pair). For the same reason, for every $x \in C$ and every neighborhood U_ε of x , $\Gamma_\varepsilon(x) = U_\varepsilon \cap C$ is not polar.

Put

$$\text{Cap}_{R, \nu}(B) = P_c\{\mathcal{R} \cap B \neq \emptyset, e^{-Z_\nu}\}.$$

Likewise Cap_R , it is a Choquet capacity by [1], Theorem III.32. Note that

$$\text{Cap}_{R, \nu}(B) = e^{-u_\nu(c)} - e^{-w_{B, \nu}(c)}. \quad (2.8)$$

By Lemma 2.5,

$$P_c\{\mathcal{R} \cap B \neq \emptyset, \mathcal{R} \cap Ex(\nu) = \emptyset\} > 0$$

for every non-polar B disjoint from $Ex(\nu)$. In addition, $Z_\nu < \infty$ on $\{\mathcal{R} \cap Ex(\nu) = \emptyset\}$ by Lemma 2.6. Therefore $\text{Cap}_{R, \nu}(B) > 0$ for every non-polar B disjoint from $Ex(\nu)$. In particular, $\text{Cap}_{R, \nu}(\Gamma_\varepsilon(x)) > 0$ for every $x \in C$.

Let x_i be everywhere dense in C . Since $\text{Cap}_{R,\nu}(x_i) = 0$, one can choose $\varepsilon_i > 0$ to have

$$\sum \text{Cap}_{R,\nu}(\Gamma_{\varepsilon_i}(x_i)) < \text{Cap}_{R,\nu}(C) \tag{2.9}$$

Put $B = \cup_i \Gamma_{\varepsilon_i}(x_i)$. By (2.9),

$$\text{Cap}_{R,\nu}(B) < \text{Cap}_{R,\nu}(C) \leq \text{Cap}_{R,\nu}(\Gamma)$$

and, by (2.8) and Lemma 2.1,

$$w_{B,\nu} < w_{\Gamma,\nu}. \tag{2.10}$$

Since (Γ, ν) is a normal pair, the trace of $w_{\Gamma,\nu}$ coincides with (Γ, ν) by Proposition 1.1. Let now $(\tilde{\Gamma}, \mu)$ be the trace of $w_{B,\nu}$. By Lemma 2.4, $\tilde{\Gamma} \subset \Gamma$ and $\mu = \nu$.

Suppose $A = \Gamma \setminus \tilde{\Gamma}$ is not empty. Since A is relatively open in Γ , it contains at least one of x_i . The set $A \cap \Gamma_{\varepsilon_i}(x_i)$ is also relatively open in Γ , and therefore it is not polar by the definition of normal pair. But,

$$A \cap \Gamma_{\varepsilon_i}(x_i) \subset A \cap B,$$

and therefore $B \setminus \tilde{\Gamma}$ is not polar, in contradiction with Lemma 2.4. Therefore $\tilde{\Gamma} = \Gamma$ and the two solutions $w_{B,\nu}$ and $w_{\Gamma,\nu}$ do not coincide and have the trace (Γ, ν) . \square

2.6 Proof of Theorem 1.4

1°. Recall some notation from [7] and [3]. A point $y \in \partial E$ is a singular point of a solution u if

$$\int_0^\zeta u^{\alpha-1}(\xi_s) ds = \infty \quad \Pi_x^y\text{-a.s.}$$

Here (ξ_t, Π_x^y) is the L -diffusion in E conditioned to exit from E at the point y , and ζ is its life time. The set of all singular points of u is denoted by $\text{SG}(u)$.

Let $\Gamma = \text{SG}(u_\nu)$. By [3], Theorem 1.1

$$P_x Z_\eta e^{-Z_\nu} = 0$$

for every measure $\eta \in \mathcal{N}_1$ concentrated on Γ . Since $Z_\nu < \infty$ a.s. by Lemma 2.6, this is possible only if $Z_\eta = 0$ a.s. and therefore $\eta = 0$. By [6], Theorem 1.2, this is equivalent to the polarity of Γ .

2°. Let B be a Borel subset of ∂E . As in [7], denote by u_B the supremum of all moderate solutions u_μ such that $\mu(B^c) = 0$. For two solutions u_1, u_2 , we define $u_1 \oplus u_2$ as the maximal solution dominated by $u_1 + u_2$. See [7] for more detail.

Since Γ is polar, $u_\Gamma = 0$ and $u_\Gamma \oplus u_\nu = u_\nu$.

3°. The fine trace of a solution u is defined as a pair (Γ, μ) where $\Gamma = \text{SG}(u)$ and μ is the maximal measure such that $\mu(\Gamma) = 0$, μ does not charge polar sets and $u_\mu \leq u$. According to [7], Theorem 1.3, the fine trace of any solution has the following properties:

- (A) (See [7],1.10.A.) The set Γ is finely closed (that is, closed in fine topology introduced in [7]).
- (B) (See [7],1.10.B.) The measure μ is a σ -finite measure on Γ^c not charging polar sets and such that $\text{SG}(u_\mu) \subset \Gamma$.

Moreover (see [7], Theorem 1.4), if (Γ, μ) is any pair satisfying (A) and (B), then $\nu = u_\Gamma \oplus u_\mu$ has the fine trace (Γ', μ) where $\Gamma' = \text{SG}(\nu)$ differs from Γ by a polar set.

The set $\Gamma = \text{SG}(u_\nu)$ is finely closed by [7], Theorem 1.3. Since Γ is polar, ν does not charge Γ . By assumption, $\text{Ess}(\nu)$ is empty and therefore ν is σ -finite by Theorem 1.1(iv). Hence the pair (Γ, ν) satisfies (A) and (B) and $u_\nu = u_\Gamma \oplus u_\nu$ has the fine trace $(\text{SG}(u_\nu), \nu) = (\Gamma, \nu)$. Since Γ is polar, the statement follows. \square

Acknowledgments. I'd like to thank E. Dynkin for numerous stimulating discussions and the anonymous referee for very helpful comments.

References

- [1] C. Dellacherie and P.-A. Meyer, *Probabilités et potentiel*, Hermann, Paris, 1975, 1980, 1983, 1987.
- [2] E. B. Dynkin, *Superprocesses and partial differential equations*, Ann. Probab. **21** (1993), 1185–1262.
- [3] ———, *Stochastic boundary values and boundary singularities for solutions of the equation $Lu = u^\alpha$* , J. Functional Analysis **153** (1998), 147–186.
- [4] E. B. Dynkin and S. E. Kuznetsov, *Linear additive functionals of superdiffusions and related nonlinear p.d.e.*, Trans. Amer. Math. Soc. **348** (1996), 1959–1987.
- [5] ———, *Solutions of $Lu = u^\alpha$ dominated by L -harmonic functions*, Journale d'Analyse Mathématique **68** (1996), 15–37.
- [6] ———, *Superdiffusions and removable singularities for quasilinear partial differential equations*, Comm. Pure & Appl. Math **49** (1996), 125–176.
- [7] ———, *Fine topology and fine trace on the boundary associated with a class of quasilinear differential equations*, Comm. Pure Appl. Math. **51** (1998), 897–936.
- [8] ———, *Solutions of nonlinear differential equations on a Riemannian manifold and their trace on the Martin boundary*, Transact. Amer. Math. Soc. **350** (1998), 4521–4552.
- [9] ———, *Trace on the boundary for solutions of nonlinear differential equations*, Transact. Amer. Math. Soc. **350** (1998), 4499–4519.
- [10] A. Gmira and L. Véron, *Boundary singularities of solutions of some nonlinear elliptic equations*, Duke Math.J. **64** (1991), 271–324.
- [11] S. E. Kuznetsov, *σ -moderate solutions of $Lu = u^\alpha$ and fine trace on the boundary*, C. R. Acad. Sci. Paris, Série I **326** (1998), 1189–1194.
- [12] J.-F. Le Gall, *Solutions positives de $\Delta u = u^2$ dans le disque unité*, C.R. Acad. Sci. Paris, Série I **317** (1993), 873–878.
- [13] ———, *A probabilistic approach to the trace at the boundary for solutions of a semilinear parabolic differential equation*, J. Appl.Math. Stochast. Analysis **9** (1996), 399–414.
- [14] ———, *A probabilistic Poisson representation for positive solutions of $\Delta u = u^2$ in a planar domain*, Comm. Pure & Appl Math. (1997), 69–103.

-
- [15] M. Marcus and L. Véron, *Trace au bord des solutions positives d'équations elliptiques non linéaires*, C.R. Acad.Sci Paris **321**, ser I (1995), 179–184.
- [16] ———, *Trace au bord des solutions positives d'équations elliptiques et paraboliques non linéaires. Résultats d'existence and d'unicité*, C.R. Acad. Sci. Paris **323** (1996), 603–608.
- [17] ———, *The boundary trace of positive solutions of semilinear elliptic equations, I: The subcritical case*, Arch. Rat. Mech. Anal. **144** (1998), 201–231.
- [18] ———, *The boundary trace of positive solutions of semilinear elliptic equations: The supercritical case*, J. Math. Pures Appl. **77** (1998), 481–524.