

***abc* TRIPLES**

GREG MARTIN, WINNIE MIAO

Abstract: The *abc* conjecture, one of the most famous open problems in number theory, claims that three relatively prime positive integers a, b, c satisfying $a + b = c$ cannot simultaneously have significant repetition among their prime factors; in particular, the product of the distinct primes dividing the three integers should never be much less than c . Triples of relatively prime numbers satisfying $a + b = c$ are called *abc triples* if the product of their distinct prime divisors is strictly less than c . We catalog what is known about *abc* triples, both numerical examples found through computation and infinite families of examples established theoretically. In addition, we collect motivations and heuristics supporting the *abc* conjecture, as well as some of its refinements and generalizations, and we describe the state-of-the-art progress towards establishing the conjecture.

Keywords: *abc* conjecture, number theory, factorization.

1. Introduction

$a, b, c \dots$ only in mathematics could such a trite trio of letters signify a major outstanding open problem with significant connections to multiple topics. The *abc conjecture* is a simple-to-state yet challenging problem in number theory that has stumped mathematicians for the past 30 years. It has become known for its large number of profound implications in number theory and particularly in Diophantine equations; among these myriad consequences are Fermat’s last theorem (up to finitely many counterexamples), Mordell’s conjecture [12], and Roth’s theorem [5] (see [31] for a more comprehensive list). The *abc* conjecture is deeply intriguing because it unveils some delicate tension between the additive and multiplicative properties of integers, the bread and butter of number theorists.

The purpose of this article is to discuss examples and constructions of *abc* triples—trios of integers demonstrating that the *abc* conjecture, if true, must be only barely true. To do so we must first describe the *abc* conjecture itself, starting with a preliminary definition: the *radical* of an integer n , denoted by $R(n)$, is the product of all the distinct prime factors of n . For example, $600 = 2^4 \cdot 3 \cdot 5^2$ and so $R(600) = 2 \cdot 3 \cdot 5 = 30$. In other words, $R(n)$ is the largest squarefree divisor

2010 Mathematics Subject Classification: primary: 11N32, 11N25, 11N05; secondary: 11R29

of n . The radical is a multiplicative function; in particular, for pairwise relatively prime integers a , b and c , we have $R(abc) = R(a)R(b)R(c)$. We may now state (the first version of) the *abc* conjecture, which postulates that the radical of three additively-related numbers normally cannot be much smaller than the numbers themselves.

***abc* Conjecture, Version 1.** *For every $\varepsilon > 0$, there exist only finitely many triples (a, b, c) of relatively prime positive integers satisfying $a + b = c$ for which*

$$R(abc) < c^{1-\varepsilon}.$$

A typical integer's radical is not too much smaller than the integer itself, and so $R(abc)$ is often about as large as abc —that is, much larger than c . Yet there are rare occurrences of integer triples (a, b, c) where c is in fact greater than $R(abc)$; these special cases are referred to as *abc triples*. The smallest such example is $(a, b, c) = (1, 8, 9)$ for which $R(abc) = R(72) = 6 < 9$.

Indeed, one can even construct an infinite sequence of *abc* triples. One such example is $(a, b, c) = (1, 9^n - 1, 9^n)$: since $9^n - 1 \equiv 1^n - 1 \equiv 0 \pmod{8}$, we see that 8 divides $9^n - 1$ for every positive integer n . Writing $b = 2^3k$ for some positive integer k , we calculate that $R(abc) = R(a)R(b)R(c) = 1 \cdot R(2^3k) \cdot 3$ is at most $2k \cdot 3 = 6k$, which is less than $c = 8k + 1$ for every n . We will see many more infinite families of *abc* triples in Section 3.

As is often the case, the literature contains various equivalent formulations of the *abc* conjecture; a few of these we will list now, and the others will appear as we proceed through the survey. For starters, the *abc* conjecture is commonly stated with the epsilon on the opposite side:

***abc* Conjecture, Version 2.** *For every $\varepsilon > 0$, there exist only finitely many triples (a, b, c) of relatively prime positive integers satisfying $a + b = c$ for which*

$$c > R(abc)^{1+\varepsilon}.$$

Version 1 and Version 2 can be effortlessly obtained from each other, although we need to remember that both statements are “for every $\varepsilon > 0$ ” statements: for example, the inequality in Version 1 with a given ε implies the inequality in Version 2 with its ε replaced by $\frac{\varepsilon}{1-\varepsilon}$. This second version of the conjecture is closely connected to the “quality” of an *abc* triple, a quantity we will define in Section 2.

For a given ε , if there are only finitely many *abc* triples for which $R(abc) < c^{1-\varepsilon}$, then there are only finitely many values of $R(abc)/c^{1-\varepsilon}$ that are less than 1. We can select the minimum such value to be $K(\varepsilon)$ and in turn transform Version 1 into:

***abc* Conjecture, Version 3.** *For every $\varepsilon > 0$ there exists a positive constant $K(\varepsilon)$ such that all triples (a, b, c) of relatively prime positive integers with $a + b = c$ satisfy*

$$R(abc) \geq K(\varepsilon)c^{1-\varepsilon}.$$

In fact, it is easily shown that Version 3 with a given positive ε implies Version 1 for any larger ε . A parallel reformulation from Version 2 is as follows:

***abc* Conjecture, Version 4.** *For every $\varepsilon > 0$ there exists a positive constant $K'(\varepsilon)$ such that all triples (a, b, c) of relatively prime positive integers with $a + b = c$ satisfy*

$$c \leq K'(\varepsilon)R(abc)^{1+\varepsilon}.$$

It is worthwhile to point out that for trios of integers related by the equation $a + b = c$, being relatively prime automatically implies being pairwise relatively prime. While it might be nice to be able to drop the coprimality condition of the hypothesis, without it nothing would stop us from multiplying any given triple by a huge power of a prime p , which would undesirably inflate c while only raising $R(abc)$ by a factor of p . The most extreme example is the triple $(a, b, c) = (2^n, 2^n, 2^{n+1})$, for which $c = 2^{n+1}$ can be made as much larger than $R(abc) = 2$ as we wish.

Likewise, the epsilon in the conjecture might seem like a nuisance but is an absolute necessity. Section 3 is devoted to recording several examples that refute similar epsilon-less statements; many of these examples are “well-known to the experts” yet decidedly hard to find in the literature. We hope amassing them here is a helpful service to those studying this topic.

First, however, we take time in Section 2 to look at some numerical examples of *abc* triples that have been garnered over the years, and examine various computational techniques of obtaining such triples. After presenting the aforementioned infinite families of *abc* triples in Section 3, we then delve into the motivation behind this deep conjecture in Section 4. Lastly, in Section 5 we present refinements and generalizations of the *abc* conjecture, and discuss progress towards and the current status of the conjecture. Although some of these later results and extensions are a bit technical, the large majority of the material we present is pleasantly elementary and accessible.

2. Numerical examples of *abc* triples

Because the *abc* conjecture has become so prominent in the last thirty years, corresponding roughly to the era of widespread and easily accessible computation, it is no surprise that people have developed a sustained interest in compiling numerical examples of *abc* triples. In fact, one can go to an online *abc* triples database [18] and list all *abc* triples of positive integers up to a bound of 10^8 , or input any integer in that range to search for *abc* triples containing it! For instance, there are exactly seven *abc* triples with $c = 100,000,000 = 2^8 \cdot 5^8$: the one with the largest value of b has $a = 351,297 = 3^4 \cdot 4,337$ and $b = 99,648,703 = 7^7 \cdot 11^2$, which gives $R(abc) = 2 \cdot 3 \cdot 5 \cdot 7 \cdot 11 \cdot 4,337 = 10,018,470$. Computations of *abc* triples have been carried out for much larger ranges. Typically such computations record the found triples according to their “quality”:

Definition. Given a triple (a, b, c) of relatively prime positive integers such that $a + b = c$, the *quality* $q(a, b, c)$ of the triple is defined to be

$$q(a, b, c) = \frac{\log c}{\log R(abc)}.$$

For example, the quality of the smallest abc triple is $q(1, 8, 9) = \frac{\log 9}{\log 6} = 1.22629\dots$. By this definition, a triple will be an abc triple only if $q(a, b, c) > 1$. We can reformulate the abc conjecture yet again by solving the inequality in Version 2 for $1 + \varepsilon$:

abc Conjecture, Version 5. *For every $\varepsilon > 0$, there exist only finitely many triples (a, b, c) of relatively prime positive integers satisfying $a + b = c$ for which $q(a, b, c) > 1 + \varepsilon$.*

By looking at de Smit’s website [10], we see for example that among numbers with at most twenty digits, there are exactly 236 abc triples of quality at least 1.4. Atop that list is the triple

$$a = 2, \quad b = 6,436,341 = 3^{10} \cdot 109, \quad c = 6,436,343 = 23^5, \quad (1)$$

for which $q(a, b, c) = 1.62991\dots$; this is the highest quality of any known abc triple (and possibly the highest quality of any abc triple in the universe!). This triple was discovered in 1987 by the French mathematician E. Reyssat (apparently by “brute force”, according to [21, page 137]). de Smit’s list also includes an abc triple, discovered by I. J. Calvo, where c has a whopping 2,958 digits: the triple

$$(a, b, c) = \left(3^3 \cdot 31^3 \cdot A, 5^{362} \cdot 7^{109} \cdot 11^7 \cdot 17^{326} \cdot 37^{11} \cdot 53^{33} \cdot 59^{179} \cdot 67^{137} \cdot 79^{76} \cdot 103^{348} \right. \\ \cdot 109^{12} \cdot 113^{103} \cdot 131^{42} \cdot 151^{12} \cdot 163^{166}, 2^{465} \cdot 13^{76} \cdot 19^{57} \cdot 23^{611} \cdot 29^{19} \cdot 41^{11} \cdot 43^{98} \\ \left. \cdot 61^{84} \cdot 71^{13} \cdot 73^{250} \cdot 83^{30} \cdot 89^{10} \cdot 97^{80} \cdot 101^{45} \cdot 127^7 \cdot 137^8 \cdot 139^3 \cdot 167^{253} \cdot 173^{25} \right) \quad (2)$$

has quality at least $1.01522\dots$, where $A = (c - b)/93^3$ is a number with 2,854 digits. (Interestingly, as is often the case with large numbers, A is easily shown to be composite—by calculating that $2^{A-1} \not\equiv 1 \pmod{A}$ and invoking Fermat’s little theorem, for example—but its factorization is unknown.)

Reken mee met ABC, hosted by the Mathematical Institute of Leiden University [35], is a distributive computing program aiming to collect experimental data on the abc conjecture. The project is based on the BOINC platform [2], and any individual with a computer can download the software and join in the hunt for abc triples. The project currently has over 150,000 users and has tested nearly three quintillion triples—not too much less than the number of insects on Earth!

People have developed many different techniques for finding abc triples, using tools from all parts of number theory and neighboring fields. To give a flavor of the wide variety of techniques, we describe six of them now.

2.1. ABC@home algorithm

The ABC@home project, which supports the *Reken mee met ABC* distributed computation described above, uses the following algorithm [1] to search exhaustively for *abc* triples.

Suppose that (a, b, c) is an *abc* triple of numbers all less than N . Rename the integers $\{a, b, c\}$ as $\{x, y, z\}$ so that x, y , and z have the smallest, middle, and largest radical, respectively. Since (a, b, c) is an *abc* triple, we have $R(a)R(b)R(c) < c < N$, and so $R(x)R(y)R(z) < N$. From this inequality and the inequalities $R(x) < R(y) < R(z)$, it is easy to deduce that $R(y) < \sqrt{N}$ and $R(x) < N/R(y)^2$.

We may therefore search for *abc* triples up to N by sorting them according to their smallest two radicals $R(x)$ and $R(y)$, both of which are at most \sqrt{N} . First, we make a list of all of the squarefree numbers less than \sqrt{N} (by a variant of the sieve of Eratosthenes, say). For every pair of relatively prime numbers (r, s) from this list that satisfy $r < \min\{s, N/s^2\}$, we calculate all pairs of numbers (x, y) for which $R(x) = r$ and $R(y) = s$. There are two ways of completing the pair (x, y) to a triple where two numbers sum to the third: we can set either $z = x + y$ or $z = |x - y|$. If $s < R(z) < N/rs$, then we have discovered a new *abc* triple, namely the sorted ordering of (x, y, z) .

2.2. Continued fractions

The (*simple*) *continued fraction* of an irrational number θ is an expression of the form

$$\theta = a_0 + \frac{1}{a_1 + \frac{1}{a_2 + \frac{1}{a_3 + \dots}}}, \tag{3}$$

where a_0 is an integer and a_j is a positive integer for each $j \geq 1$. Calculating the partial quotients a_0, a_1, \dots of a given irrational number θ turns out to be a simple variant of the Euclidean algorithm (which at its heart is simply division with remainder). If we replace the infinite tail $a_j + \frac{1}{a_{j+1} + \dots}$ of the continued fraction with a_j itself, we obtain a rational number called the *j*th *convergent* to θ . The theory of these convergents, and how they comprise the best rational approximations to θ in a suitable sense, is extremely interesting [32, Chapter 7].

For example, we calculate the continued fraction of the irrational number $\sqrt[5]{109}$, which has been cunningly chosen for its relationship to Reyssat's example (1):

$$\sqrt[5]{109} = 2 + \frac{1}{1 + \frac{1}{1 + \frac{1}{4 + \frac{1}{77,733 + \frac{1}{2 + \dots}}}}}$$

Noting that $4 + \frac{1}{77,733 + \frac{1}{2 + \dots}}$ is extremely close to 4, we form the approximation

$$\sqrt[5]{109} \approx 2 + \frac{1}{1 + \frac{1}{1 + \frac{1}{4}}} = \frac{23}{9},$$

which is the third convergent to $\sqrt[5]{109}$. (In this particular case, we might have found this approximation just by noting the decimal expansion $\sqrt[5]{109} = 2.5555553967\dots$!) This approximation tells us that $9^5 \cdot 109 \approx 23^5$, and in fact their difference is exactly 2, yielding Reyssat's triple $(2, 9^5 \cdot 109, 23^5)$.

In general, we begin with an irrational root $\theta = \sqrt[k]{m}$ of an integer m and compute its continued fraction. At any point, when we see a relatively large partial quotient a_{j+1} , we truncate the infinite continued fraction (3) after a_j to obtain the j th convergent, which we write as p/q . We have thus found integers p and q such that $p/q \approx \sqrt[k]{m}$, or equivalently $mq^k \approx p^k$. We then examine the triple candidate $(|mq^k - p^k|, mq^k, p^k)$ to see whether its quality exceeds 1. For the curious reader, [7] contains a list of ninety *abc* triples, all with quality exceeding 1.4, that can be found via this continued fraction method.

2.3. The LLL method

Another interesting method, proposed by Dokchitser [11], employs a famous “lattice basis reduction” algorithm by Lenstra, Lenstra, and Lovász [25]. The usual integer lattice \mathbb{Z}^3 is the set of all integer linear combinations of the vectors $(1, 0, 0)$, $(0, 1, 0)$, $(0, 0, 1)$ inside \mathbb{R}^3 . These three vectors form a basis for this lattice, but so do, say, $(12, 34, 39)$, $(20, 57, 65)$, $(95, 269, 309)$. Like vector spaces, a general lattice can have many bases; when given a complicated and suboptimal basis like this latter one, the *LLL algorithm* converts it into a much nicer basis—one with smaller entries, and for which the basis elements are nearly orthogonal.

To apply this tool to the construction of *abc* triples, we select large relatively prime integers r, s, t that are comparable in size and have very small radicals themselves. If we can find small integers u, v, w such that

$$ur + vs + wt = 0, \tag{4}$$

then $(|u|r, |v|s, |w|t)$ has a good chance of being an *abc* triple: the radicals of r, s, t were all chosen to be small, and the small integers $|u|, |v|, |w|$ can only contribute so much to the radical of the product.

The set of all integer vectors (u, v, w) satisfying equation (4) is a two-dimensional sublattice of \mathbb{Z}^3 , and the usual methods for finding its basis results in quite sizeable vectors. However, after running the LLL algorithm to solve for a reduced basis $\{\mathbf{b}_1, \mathbf{b}_2\}$, taking any small linear combination of \mathbf{b}_1 and \mathbf{b}_2 and setting the entries of the resultant vector as our (u, v, w) is fair game. We can then test the trio of integers $(|u|r, |v|s, |w|t)$ to see if it is an *abc* triple.

In this fashion, Dokchitser was able to obtain 41 new *abc* triples, including $a = 13^{10} \cdot 37^2$, $b = 3^7 \cdot 19^5 \cdot 71^4 \cdot 223$, $c = 2^{26} \cdot 5^{12} \cdot 1,873$ which has a quality of 1.5094, the 11th highest quality known.

2.4. Transfer method

Yet another approach to finding new *abc* triples is to take existing triples and “transfer” them, using certain polynomial identities, to create new triples.

For example, note that if $a + b = c$, then $a^2 + c(b - a) = b^2$, since $c(b - a) = (b + a)(b - a) = b^2 - a^2$. Note also that if $R(abc) < c$, then

$$\begin{aligned} R(a^2 \cdot c(b - a) \cdot b^2) &\leq R(a)R(b)R(c)R(b - a) \\ &= \frac{R(abc)}{c} cR(b - a) < c(b - a) < b^2 \end{aligned} \tag{5}$$

as well. In other words, if (a, b, c) is an *abc* triple with $a < b$, then $(a^2, c(b - a), b^2)$ is also an *abc* triple. Indeed, if the quality $q(a, b, c)$ is larger than 1, then a quick calculation [47, page 16] shows that

$$q(a^2, c(b - a), b^2) > \frac{2q(a, b, c)}{q(a, b, c) + 1} > 1.$$

For future reference, we also note a slight improvement: if (a, b, c) is an *abc* triple where a and b are both odd (which forces both c and $b - a$ to be even), then

$$\begin{aligned} R(a^2 \cdot c(b - a) \cdot b^2) &< \frac{c}{R(abc)} R(a^2 \cdot c(b - a) \cdot b^2) \\ &\leq \frac{c}{R(abc)} R(a)R(b)R(c)R\left(\frac{b - a}{2}\right) < c\left(\frac{b - a}{2}\right) < \frac{b^2}{2}. \end{aligned} \tag{6}$$

When we are looking for good numerical examples, moreover, we can try this transfer method on many known *abc* triples and hope for some extra repeated factors in $b - a$. For example, we can start with the small *abc*-triple $(7, 243, 250)$, whose radical is 210 and whose quality is about 1.03261. Using the above transfer identity leads to the triple $a = 7^2 = 49$, $b = 250(243 - 7) = 59,000$, $c = 243^2 = 59,049$. We know from the bound (5) that the radical of this new triple is at most $210 \cdot (243 - 7)$. However, $243 - 7 = 2^2 \cdot 59$, and the factors of 2 are dropped from the radical since 250 is already even. Consequently, the radical of this new triple is only $210 \cdot 59 = 12,390$, and the quality of $(49, 59,000, 59,049)$ is about 1.16568, which is quite a bit better than the original triple.

The transfer method, then, is to start with existing *abc* triples, apply a polynomial identity to obtain a new triple, and then check for fortunate coincidences that make the new triple even better than we already knew it would be. It is an experimentation game, where different starting triples can yield results from mediocre to extremely good. In fact, we can experiment not only with the starting triple but with the polynomial identity as well! Some other examples of such polynomial transfers, which are all easily seen to be valid when $c = a + b$, include:

$$\begin{aligned} (b - a)^2 + 4ab &= c^2 \\ a^3 + b^3 &= c(b^2 - ab + a^2) \\ a^2(a + 3b) + b^2(3a + b) &= c^3 \\ a^3(a + 2b) + c^3(b - a) &= b^3(2a + b) \\ 27c^5(b - a) + a^3(3a + 5b)^2(3a + 2b) &= b^3(5a + 3b)^2(2a + 3b). \end{aligned}$$

(Sometimes the resulting triples have small common factors, which we should cancel out.) Moreover, there is even a whole family of such identities

$$a^{n-k} \left(\sum_{j=0}^k \binom{n}{j} a^{k-j} b^j \right) + b^{k+1} \left(\sum_{j=0}^{n-k-1} \binom{n}{j} a^j b^{n-k-1-j} \right) = c^n$$

which comes from splitting the binomial formula for $(a+b)^n$ at some term with index $1 \leq k \leq n-1$. (Note that the third identity on the above list is the $n=3$, $k=1$ case of this general family.)

The interested reader can refer to [47, Section 2.3] for a detailed examination of these polynomial transfers as a way of generating triples.

2.5. An elliptic curve method

The study of elliptic curves is fascinating in its own right (see [51] or [37], for example, where one can find all the terminology about elliptic curves that we use in this survey), but here we are particularly interested in the overlap of elliptic curves and the *abc* conjecture. We describe (a slight variant of) a method of van der Horst for searching for *abc* triples, one which uses an idea that was anticipated earlier by Carella [8].

For any fixed integers $x_0 < y_0$, set $k = y_0^3 - x_0^3$ and consider the elliptic curve given by the equation

$$y^3 = x^3 + k, \tag{7}$$

where the variables x and y are allowed to be rational numbers. Whenever $(x, y) = (\frac{p}{d}, \frac{q}{d})$ is a point on this elliptic curve (for simplicity we assume that p, q , and d are positive), we have $q^3 = p^3 + kd^3$. Clearly $R(p^3, kd^3, q^3) \leq kdpq < kdq^2$, and so this triple is an *abc* triple whenever $q > kd$, or equivalently when $y > k$; when y is large and has small denominator, the quality of the triple will be large.

It is probably not the case that y_0 itself is larger than k ; however, we can use the group operation on the elliptic curve to search for rational solutions to equation (7) other than (x_0, y_0) . Simply adding (using the group law on the elliptic curve) the point (x_0, y_0) to itself repeatedly yields a sequence of points on the elliptic curve that is typically infinite. van der Horst even develops a way of predicting which elements of this sequence will have large y -values: he writes down a group homomorphism from the elliptic curve to the unit circle in the complex plane that takes points with large coordinates to complex numbers near 1. Since it is easy to calculate which powers of a complex number are close to 1, one can take the corresponding multiples of (x_0, y_0) back on the elliptic curve and check how good the corresponding triples' qualities are. One feature of this method is that all three numbers in the *abc* triples it generates have small radicals, not just one or two of them.

The exact algorithm and variants used by van der Horst [47, Sections 4.2–4.3] discovered some notable *abc* triples. The point $(x, y) = (\frac{19}{93}, \frac{289}{93})$ on the elliptic curve $y^3 = x^3 + 30$ does not have $y > 30$, but fortunately the numerator of y

happens to be a square, and so we get to divide the radical by an extra factor of 17. The resulting *abc* triple $a = 19^3 = 6,859$, $b = 30 \cdot 93^3 = 24,130,710$, $c = 289^3 = 24,137,569$ has radical 300,390 and quality about 1.34778. Moreover, the algorithm often finds rational solutions with huge numerators and denominators, and is thus suited for finding enormous *abc* triples; van der Horst reports [47, Chapter 5] finding a point on the elliptic curve $y^3 = x^3 + 854$ that yields an *abc* triple with quality about 1.01635, where the largest integer in the triple has 340 digits.

2.6. Differences of *j*-invariants

We conclude this section with some exotic *abc* triples that are found unexpectedly when discussing factorizations of “*j*-invariants”.

There is a beautiful link between lattices and elliptic curves: through two “elliptic functions” studied by Weierstrass, it is known that every elliptic curve can be represented as $y^2 = 4x^3 - g_2(\tau)x - g_3(\tau)$, where $g_2(\tau)$ and $g_3(\tau)$ are invariants that correspond to a fixed lattice. More specifically, they are the modular forms

$$g_2(\tau) = 60 \sum_{\substack{m,n \in \mathbb{Z} \\ (m,n) \neq (0,0)}} \frac{1}{(m\tau + n)^4}$$

$$g_3(\tau) = 140 \sum_{\substack{m,n \in \mathbb{Z} \\ (m,n) \neq (0,0)}} \frac{1}{(m\tau + n)^6}$$

where τ , a complex number with positive imaginary part, determines the relevant lattice as the set of all numbers of the form $m\tau + n$ with m, n integral.

The *j*-invariant $j(\tau)$ of an elliptic curve can now be defined by the formula

$$j(\tau) = \frac{1728g_2^3(\tau)}{g_2^3(\tau) - 27g_3^2(\tau)}.$$

This *j*-invariant is a modular function with ubiquitous remarkable properties and applications in complex analysis, algebraic number theory, transcendence theory, and so on. When the argument τ lies in an imaginary quadratic field $\mathbb{Q}(\sqrt{-d})$ for some positive integer d , the values $j(\tau)$ are called *singular moduli*, and the associated elliptic curves possess extra endomorphisms and are said to have *complex multiplication*. This singular modulus is an algebraic integer lying in some abelian extension of $\mathbb{Q}(\sqrt{-d})$; remarkably, the degree of its minimal polynomial is exactly the class number $h(-d)$. In particular, by the Stark–Heegner theorem [37, Appendix C, Section 11], there are only thirteen negative discriminants $-d$ that have class number equal to 1, namely $-3, -4, -7, -8, -11, -12, -16, -19, -27, -28, -43, -67$, and -163 ; the corresponding *j*-invariants are thus actual integers.

As it happens, these thirteen special *j*-invariants are all forced to be perfect cubes of integers. Equally marvelously, the difference of two of these special *j*-invariants is very nearly a perfect square [16, 24]. The corresponding triple of

integers is therefore a prime candidate for an abc triple (at least, once the three integers are divided by their greatest common divisor). Gross and Zagier [16] cite an example with $\tau = (-1 + i\sqrt{163})/2$, where the three integers

$$\begin{aligned}\frac{j(i)}{1728} &= 1 \\ \frac{-j(\tau)}{1728} &= 151,931,373,056,000 = 2^{12} \cdot 5^3 \cdot 23^3 \cdot 29^3 \\ \frac{j(i) - j(\tau)}{1728} &= 151,931,373,056,001 = 3^3 \cdot 7^2 \cdot 11^2 \cdot 19^2 \cdot 127^2 \cdot 163\end{aligned}$$

form an abc triple with quality about 1.20362. Going through all $\binom{13}{2} = 78$ possible pairs of special j -invariants, we find that the best resulting abc triple comes from both $j(\tau_4) - j(\tau_{43})$ and $j(\tau_{16}) - j(\tau_{67})$, where $\tau(d) = \frac{1}{2}(d + \sqrt{-d})$: the triple is $a = 1$, $b = 512,000 = 2^{12} \cdot 5^3$, $c = 512,001 = 3^5 \cdot 7^2 \cdot 43$ and has quality about 1.44331.

3. Infinite families of abc triples

All of the numerical examples from Section 2, however interesting, cannot shed any light on whether the abc conjecture is true or false: the “only finitely many” or “there exists a constant” clauses in its various versions preclude us from drawing conclusions from any finite number of examples. For that matter, any finite number of examples cannot rule out even more ambitious possible versions of the abc conjecture. For instance, could there be an absolute constant $S > 0$ such that $c < S \cdot R(abc)$ always? This statement, similar to the abc conjecture but without the messy epsilons, might be called the “simplistic abc conjecture”. Again, no finite collection of examples can resolve this question.

What we need, to help us decide whether these statements are true or false, are constructions of infinite families of abc triples. And it turns out that several such constructions exist; any one of these constructions shows that the simplistic abc conjecture is false. In other words, the constructions in this section demonstrate that the epsilons in the abc conjecture are necessary if we are to hope that the assertion is true.

3.1. The transfer method again

Recall from Section 2.4 that if (a, b, c) is an abc triple, then so is $(a^2, c(b-a), b^2)$. In particular, if $(1, c-1, c)$ is an abc triple, then so is $(1, c^2 - 2c, (c-1)^2)$. Of course, we can iterate this transfer multiple times in a row: for example, $(1, (c^2 - 2c)^2 - 1, (c^2 - 2c)^2) = (1, c^4 - 4c^3 + 4c^2 - 1, c^4 - 4c^3 + 4c^2)$ will also be an abc triple. As it happens, doing this double transfer always allow us to remove an extra factor of 2 from the radical. For example, suppose that c is odd. Then, by setting $a = 1$ and $b = c - 1$ in the third and last terms of the chain of

inequalities (5), we know that

$$\frac{(c-1)^2}{R((c^2-2c)(c-1)^2)} \geq \frac{c}{R((c-1)c)}.$$

But now $(c-1)^2$ is even, so replacing a , b , and c in the second and last terms of the chain of inequalities (6) with 1 , c^2-2c , and $(c-1)^2$, we find that

$$\frac{(c^2-2c)^2}{R(((c^2-2c)^2-1)(c^2-2c)^2)} \geq \frac{2(c-1)^2}{R((c^2-2c)(c-1)^2)} \geq \frac{2c}{R((c-1)c)}. \tag{8}$$

We can iterate this double transfer endlessly to create an infinite sequence. Let us set $c_0 = 9$, corresponding to the *abc* triple $(1, 8, 9)$, and for every $n \geq 0$ define $c_{n+1} = c_n^4 - 4c_n^3 + 4c_n^2$. For example, $c_1 = 3,969$, corresponding to the double transfer $(1, 8, 9) \rightarrow (1, 63, 64) \rightarrow (1, 3,968, 3,969)$. Equation (8) tells us that

$$\frac{c_{n+1}}{R((c_{n+1}-1)c_{n+1})} \geq 2 \frac{c_n}{R((c_n-1)c_n)}$$

for every $n \geq 0$. Since $c_0/R((c_0-1)c_0) = \frac{3}{2}$, this implies that

$$\frac{c_n}{R((c_n-1)c_n)} \geq 2^{n-1} \cdot 3 \tag{9}$$

for every $n \geq 0$. And since $2^{n-1} \cdot 3$ exceeds any constant we might care to name in advance, this infinite sequence of *abc* triples $(1, c_n-1, c_n)$ repudiates the “simplistic *abc* conjecture”.

We can convert the inequality (9) into a quantitative measure of how much smaller than c_n this radical is. Note that $c_n \leq c_{n-1}^4$ for every $n \geq 1$, and so $c_n \leq c_0^{4^n} = 9^{4^n}$. In particular, $\log c_n \leq 4^n \log 9$, and so $2^n \geq \sqrt{\log c_n} / \sqrt{\log 9}$. It now follows from (9), when $a = 1$, $b = c_n - 1$, and $c = c_n$, that

$$R(abc) \leq \frac{c}{2^{n-1} \cdot 3} \leq \frac{2\sqrt{\log 9}}{3} \frac{c}{\sqrt{\log c}}. \tag{10}$$

To this point, it hasn’t mattered which logarithm we’ve been using, but now we clarify that we are using $\log x$ to denote the natural logarithm. With that admission out of the way, we remark that the constant $\frac{2\sqrt{\log 9}}{3}$ is approximately 0.988203.

This bound for the radical of these triples can be re-expressed as an inequality about their quality: the lower bound

$$\begin{aligned} q(a, b, c) &= \frac{\log c}{\log R(abc)} \geq \frac{\log c}{\log c - \log \sqrt{\log c} + \log(\frac{2}{3}\sqrt{\log 9})} \\ &\geq \frac{\log c}{\log c - \frac{1}{2} \log \log c} > \frac{\log c + \frac{1}{2} \log \log c}{\log c} = 1 + \frac{\log \log c}{2 \log c} \end{aligned} \tag{11}$$

holds when $(a, b, c) = (1, c_n - 1, c_n)$. Notice that these qualities are all greater than 1, but the lower bound does tend to 1 as c becomes larger and larger. If the lower bound tended to a constant larger than 1, this sequence would disprove the actual abc conjecture (specifically Version 5) and this whole survey would need to be rewritten!

3.2. Folklore examples

There are several known constructions of infinite sequences of abc triples, each of which provides a counterexample to the “simplistic abc conjecture”. We present a few of these constructions in this section. Unlike the recursive construction from the previous section, these constructions have very simple closed forms which make it obvious that the smallest and largest numbers in the triples have extremely small radicals. In each case, a quick number theory lemma is required to show that the radical of the middle number is somewhat smaller than the number itself. These constructions are simple enough (to those well-versed in the field) that it is nearly impossible to determine who first came up with them; indeed, some cannot even be found explicitly in any publication, despite the fact that they are “well known”! Part of the motivation for this survey was to ensure that these families of abc triples are explicitly recorded in the literature; we have included earlier citations whenever we could locate them.

Lemma 1. *If p is an odd prime, then p^2 divides $2^{p(p-1)} - 1$.*

Proof. Euler’s theorem [17, page 63] says that if a and m are relatively prime positive integers, then $a^{\phi(m)} \equiv 1 \pmod{m}$, where $\phi(m)$ is the Euler phi-function. Applied with $a = 2$ and $m = p^2$, for which $\phi(m) = p(p-1)$, Euler’s theorem yields $2^{p(p-1)} \equiv 1 \pmod{p^2}$, which is exactly the conclusion of the lemma. ■

The following construction was recorded by Granville and Tucker [15].

Example 1. For any odd prime p , set $(a, b, c) = (1, 2^{p(p-1)} - 1, 2^{p(p-1)})$. We know by Lemma 1 that p^2 divides b , and so $R(b) \leq b/p$. It follows that

$$R(abc) = R(a)R(b)R(c) \leq 1 \cdot \frac{b}{p} \cdot 2 < \frac{2c}{p}. \quad (12)$$

Since the sequence of primes p becomes larger than any constant we want, this family of triples does contradict the “simplistic abc conjecture”.

For easier comparison to other examples, we can rewrite the right-hand side in a form involving only c . Since $c < 2^{p^2}$, we have $\log c < p^2 \log 2$ and so $p > \frac{\sqrt{\log c}}{\sqrt{\log 2}}$. Combining this with (12) yields

$$R(abc) < 2\sqrt{\log 2} \frac{c}{\sqrt{\log c}}.$$

This upper bound for the radical has the same shape as the bound in equation (10) for our first example, but with the slightly worse constant $2\sqrt{\log 2} \approx 1.66511$.

Our next infinite family involves a lemma providing divisibility by high powers of a prime, rather than just its square:

$$\text{If } n \text{ is a nonnegative integer, then } 7^{n+1} \text{ divides } 8^{7^n} - 1. \tag{13}$$

In fact, we can establish a much more general lemma, of which this divisibility is a special case (taking $u = 2, k = 3, v = 7,$ and $m = r = t = 1$). When u is a prime and v is a prime dividing $u^k - 1$, the following construction was given by Stewart [40, Theorem 1, (3)].

Lemma 2. *Fix positive integers k, m, r, t, u, v . If the divisibility*

$$t^r v^{n+m} \mid (u^{kv^n} - 1) \tag{14}$$

holds for some $n = n_0$, then it holds for all $n \geq n_0$.

Proof. Since the base case $n = n_0$ is assumed as a hypothesis, we proceed by induction. Suppose that the divisibility (14) holds for a particular n ; we write

$$\begin{aligned} u^{kv^{n+1}} - 1 &= (u^{kv^n})^v - 1 \\ &= (u^{kv^n} - 1)((u^{kv^n})^{v-1} + (u^{kv^n})^{v-2} + \dots + (u^{kv^n})^2 + u^{kv^n} + 1). \end{aligned}$$

On the right-hand side, the first factor is divisible by $t^r v^{n+m}$ by the induction hypothesis. In particular, it is divisible by v , and so $u^{kv^n} \equiv 1 \pmod{v}$. Thus each of the v terms in the second factor is also congruent to 1 modulo v , and so that second factor is actually a multiple of v . Therefore $t^r v^{n+m} \cdot v$ divides the left-hand side, which is the statement of the lemma for $n + 1$ as required. ■

Example 2. For any nonnegative integer n , set $(a, b, c) = (1, 8^{7^n} - 1, 8^{7^n})$. Equipped with the divisibility (14), we deduce that $R(b) \leq b/7^n$ and thus

$$R(abc) = R(a)R(b)R(c) \leq 1 \cdot \frac{b}{7^n} \cdot 2 < \frac{2c}{7^n}.$$

Again we have disproved the “simplistic abc conjecture”, and again we can write the right-hand side as an expression in c alone, since $\log c = 7^n \log 8$:

$$R(abc) < 2 \log 8 \frac{c}{\log c} = 6 \log 2 \frac{c}{\log c}.$$

Note that we have improved the order of magnitude of the upper bound on the radical, from the previous examples’ $c/\sqrt{\log c}$ to $c/\log c$.

We remark that the same sort of argument that led to equation (11) shows that the qualities of the abc triples arising from this example are essentially as large as $q(a, b, c) > 1 + \frac{\log \log c}{\log c}$, without the factor of 2 in the denominator. (The same bound will hold, at least asymptotically, for the rest of the examples in Section 3.)

Variants of this construction abound. In general, if we have integers k, m, r, t, u, v satisfying the divisibility (14) for some n , then the family of triples $(a, b, c) = (1, u^{kv^n} - 1, u^{kv^n})$ will satisfy

$$R(abc) = R(a)R(b)R(c) \leq 1 \cdot \frac{b}{t^{r-1}v^{n+m-1}} \cdot u < \frac{u}{t^{r-1}v^{m-1}} \frac{c}{v^n} = \frac{ku \log u}{t^{r-1}v^{m-1}} \frac{c}{\log c}. \tag{15}$$

For example, taking $u = 3, v = 2, m = 2$, and $k = t = r = 1$, we see from Lemma 2 that 2^{n+2} divides $3^{2^n} - 1$ for any $n \geq 1$; this construction employing the triple $(a, b, c) = (1, 3^{2^n} - 1, 3^{2^n})$ (attributed in [22, pages 40–41] to Jastrzebowski and Spielman) results in the upper bound $R(abc) < \frac{1 \cdot 3 \log 3}{1^{1 \cdot 2^{2^n - 1}}} \frac{c}{\log c}$. Here, the leading constant $\frac{3 \log 3}{2} \approx 1.64792$ is even better than $6 \log 2 \approx 4.15888$. The observation that $2^5 5^2$ divides $2400 = 7^4 - 1$ (in which we set $u = 7$ and $k = 4$) can be generalized in two different ways—either by taking $v = 2, m = 5$, and $t^r = 5^2$, or by taking $v = 5, m = 2$, and $t^r = 2^5$; both choices lead to families of abc triples satisfying $R(abc) < \frac{7 \log 7}{20} \frac{c}{\log c}$, where the leading constant is about 0.68107.

Connected to examples of this sort is the notion of a *Wieferich prime*, which is a prime p that satisfies $2^{p-1} \equiv 1 \pmod{p^2}$ (a stronger congruence than the $2^{p-1} \equiv 1 \pmod{p}$ guaranteed by Fermat’s little theorem). Analogously, a *base- u Wieferich prime* is a prime p that satisfies $u^{p-1} \equiv 1 \pmod{p^2}$. For example, 5 is a base-7 Wieferich prime: the divisibility $5^2 \mid (7^4 - 1)$ from the previous paragraph is equivalent to the defining criterion $7^{5-1} \equiv 1 \pmod{5^2}$. With $u = 2$, there are only two known Wieferich primes, namely $v = 1093$ and $v = 3511$, which lead to further instances of Lemma 2: taking $k = 364$ and $k = 1755$, respectively, and $(m, t, r) = (2, 1, 1)$, we obtain families of abc triples satisfying $R(abc) < \frac{728 \log 2}{1093} \frac{c}{\log c}$ and $R(abc) < \frac{3510 \log 2}{3511} \frac{c}{\log c}$, respectively. Here the two leading constants are about 0.46168 and 0.69295.

The most spectacular example of this type that we know uses $u = 239$ and $v = 13$: not only is 13 a base-239 Wieferich prime, but in fact the much stronger congruence $239^4 \equiv 1 \pmod{13^4}$ holds. (Indeed, the abc triple $(1, 239^4 - 1, 239^4)$ has a very respectable quality, exceeding 1.35.) Taking $(k, m, t, r) = (4, 4, 2, 6)$, we obtain the triples $(a, b, c) = (1, 239^{4 \cdot 13^n} - 1, 239^{4 \cdot 13^n})$ in which the middle term is divisible by $2^6 13^{n+4}$; the resulting bound on the radical is

$$R(a, b, c) < \frac{239 \log 239}{8 \cdot 13^3} \frac{c}{\log c} \approx 0.15197 \frac{c}{\log c}.$$

Our theoretical knowledge of Wieferich primes (referring now to the classical ones with $u = 2$) is quite poor. We expect that there are infinitely many Wieferich primes, but that they are quite rare; however, we cannot even prove that there are infinitely many primes that are *not* Wieferich primes. Perhaps fittingly, it has been proved [38] that there are infinitely many non-Wieferich primes—but only under the assumption of the abc conjecture!

Our last “folklore” example differs from the previous ones: the radical of the middle number of the triple is small because it is divisible by high powers of several primes, not just a high power of a single prime.

Lemma 3. *For any positive integer n , define $L = \text{lcm}[1, 2, \dots, n]$ and $t = \lfloor \frac{\log n}{\log 2} \rfloor$, and let $P = \prod_{3 \leq p \leq n} p$ be the product of all the odd primes up to n . Then $PL/2^t$ divides $2^L - 1$. In particular,*

$$R(2^L - 1) \leq \frac{2^t(2^L - 1)}{L}.$$

Proof. Given an odd prime $p \leq n$, let $r = \lfloor \frac{\log n}{\log p} \rfloor$, so that p^r is the largest power of p not exceeding n . Clearly both p^r and $p-1$, being at most n in size, divide L ; since they are relatively prime, their product $p^r(p-1)$ also divides L . As 2 is relatively prime to p^{r+1} , Euler’s theorem tells us that $2^{\phi(p^{r+1})} = 2^{p^r(p-1)} \equiv 1 \pmod{p^{r+1}}$, and therefore $2^L \equiv 1 \pmod{p^{r+1}}$ since L is a multiple of $p^r(p-1)$. Therefore p^{r+1} divides $2^L - 1$ for every odd prime $p \leq n$. All of these prime powers are relatively prime to one another, and hence their product

$$\prod_{3 \leq p \leq n} p^{r+1} = \prod_{3 \leq p \leq n} p \prod_{3 \leq p \leq n} p^r = P \frac{L}{2^t}$$

also divides $2^L - 1$, as claimed. In this last equality, we used the fact that $\text{lcm}[1, 2, \dots, n]$ is composed exactly from the highest power of each distinct prime factor found amongst the factorizations of the numbers 1 through n . In other words $L = 2^{\lfloor \frac{\log n}{\log 2} \rfloor} \prod_{3 \leq p \leq n} p^{\lfloor \frac{\log n}{\log p} \rfloor} = 2^t \prod_{3 \leq p \leq n} p^r$.

Note also that every prime dividing $L/2^t$ is an odd prime not exceeding n , hence divides P as well. The above argument shows that P divides the quotient $(2^L - 1)/(L/2^t)$, and so the primes dividing $L/2^t$ are already represented in this quotient; consequently, the radical of $2^L - 1$ is no larger than $(2^L - 1)/(L/2^t)$. ■

Example 3. For any positive integer n , define $L = \text{lcm}[1, 2, \dots, n]$, and set $(a, b, c) = (1, 2^L - 1, 2^L)$. Using the notation $t = \lfloor \frac{\log n}{\log 2} \rfloor$ and $P = \prod_{3 \leq p \leq n} p$ from Lemma 3, we have $2^t < n$ and $\log c = L \log 2$ and thus

$$R(abc) \leq 1 \cdot \frac{2^t(2^L - 1)}{L} \cdot 2 < 2 \log 2 \frac{nc}{\log c}.$$

It is a bit harder than in the previous examples to write the right-hand side solely in terms of c , since the relationship between n and c is more complicated. The Chebyshev function $\psi(n) = \log \text{lcm}[1, \dots, n]$ (often written in terms of the “von Mangoldt function” $\Lambda(n)$) satisfies $\psi(n) \sim n$ by the famous prime number theorem [3, pages 74–75]. Therefore $\log \log c = \log L + \log \log 2 = \psi(n) + \log \log 2 \sim n$, and hence we have the asymptotic inequality

$$R(abc) \lesssim 2 \log 2 \frac{c \log \log c}{\log c},$$

which has a slightly worse order of magnitude than the last two examples. For what it's worth, we can remove a factor of 2 from the right-hand side by restricting n to be just less than a power of 2.

There is another, related folklore example, involving the ‘‘Pell equation’’ $x^2 - 8y^2 = 1$. Like all Pell equations, this one has an infinite sequence (x_n, y_n) of nonnegative solutions, which can be described in various ways. For example, the pair of sequences satisfies the recursion $(x_n, y_n) = 6(x_{n-1}, y_{n-1}) - (x_{n-2}, y_{n-2})$, together with the initial values $(x_0, y_0) = (1, 0)$ and $(x_1, y_1) = (3, 1)$. Alternatively, x_n and y_n are the integers that satisfy $x_n + y_n\sqrt{8} = (3 + \sqrt{8})^n$. In particular, the sequence y_n is a ‘‘Lucas sequence’’ and has many nice divisibility properties— for example, y_m divides y_n whenever m divides n . The following more detailed divisibility fact can be established: if, for any prime p , we define

$$h(p) = \begin{cases} \frac{p-1}{2}, & \text{if } p \equiv \pm 1 \pmod{8}, \\ \frac{p+1}{2}, & \text{if } p \equiv \pm 3 \pmod{8}, \\ 2, & \text{if } p = 2, \end{cases}$$

then p^k divides $y_{p^{k-1}h(p)}$.

Consequently, one can prove an analogue of Lemma 3 for the sequence y_n : if $L = \text{lcm}[1, \dots, n]$ and $P = \prod_{3 \leq p \leq n} p$ as before, then PL divides y_L . If we then set $a = 1$, $b = y_L^2$, and $c = x_L^2$, we obtain

$$R(abc) = R(1)R(y_L)R(x_L) \leq 1 \cdot \frac{y_L}{L/2} \cdot x_L < \frac{2}{L} \frac{x_L^2}{\sqrt{8}} = \frac{c}{L\sqrt{2}}.$$

Since x_L is approximately $\frac{1}{2}(3 + \sqrt{8})^L$, this bound becomes

$$R(abc) \lesssim \sqrt{2} \log(3 + \sqrt{8}) \frac{c}{\log c},$$

where the leading constant is about 2.4929. The difference between this order of magnitude and the one in Example 3 is that here we obtain high powers of all small primes, including $p = 2$, dividing y_L , while no power of 2 divided $2^L - 1$.

3.3. A curious divisibility

All of the abc triples constructed in this section so far share the property that their smallest number equals 1. However, we have a final construction to describe, one that was discovered only recently [6], which has the feature that all three numbers in the constructed abc triples are nearly the same size. This construction relies on the following quite strange divisibility relationship.

Lemma 4. *For any positive integer n satisfying $n \equiv 2 \pmod{6}$,*

$$\left(\frac{n^2 - n + 1}{3} \right)^2 \text{ divides } n^n - (n - 1)^{n-1}.$$

Setting $n = 6k + 2$ for a nonnegative integer k reveals that the lemma is equivalent to the curious statement:

$$(12k^2 + 6k + 1)^2 \text{ divides } (6k + 2)^{6k+2} - (6k + 1)^{6k+1}. \tag{16}$$

Proof. Given a nonnegative integer k , set $Q = 12k^2 + 6k + 1$. To establish the divisibility (16), we need to show that $(6k + 2)^{6k+2} \equiv (6k + 1)^{6k+1} \pmod{Q^2}$. Our main tool will be the following observation: if $a \equiv bQ + 1 \pmod{Q^2}$, then $a^j \equiv jbQ + 1 \pmod{Q^2}$ for any positive integer j . This observation follows from the binomial expansion

$$a^j \equiv (1 + bQ)^j = \sum_{i=0}^j \binom{j}{i} (bQ)^i \equiv 1 + j \cdot bQ + \sum_{i=2}^j 0 \pmod{Q^2}.$$

Since $(6k + 1)^3 = 18kQ + 1$ and $-(6k + 2)^3 = -(18k + 9)Q + 1$, we can certainly say that

$$\begin{aligned} (6k + 1)^3 &\equiv 18kQ + 1 \pmod{Q^2} \\ -(6k + 2)^3 &\equiv -(18k + 9)Q + 1 \pmod{Q^2}. \end{aligned}$$

Raising both congruences to the $2k$ th power using our observation, we see that

$$\begin{aligned} (6k + 1)^{6k} &\equiv 2k \cdot 18kQ + 1 = (3Q - (18k + 3))Q + 1 \equiv -(18k + 3)Q + 1 \pmod{Q^2} \\ (6k + 2)^{6k} &\equiv -2k(18k + 9)Q + 1 = (-3Q + 3)Q + 1 \equiv 3Q + 1 \pmod{Q^2}. \end{aligned}$$

We now calculate that

$$\begin{aligned} (6k + 2)^{6k+2} - (6k + 1)^{6k+1} &\equiv (3Q + 1)(6k + 2)^2 - (-(18k + 3)Q + 1)(6k + 1) \\ &= (3Q + 1)(3Q + 6k + 1) + (9Q - 18k - 6)Q - (6k + 1) \\ &= 18Q^2 \equiv 0 \pmod{Q^2}, \end{aligned}$$

which is what we needed to show. ■

Remark. Although Lemma 4 has the elementary (if unilluminating) proof just given, there is in fact a deeper explanation [6, Proposition 4.3] behind this interesting divisibility. It is related to the trinomial $x^n + x + 1$, which is reducible when $n \equiv 2 \pmod{6}$, and the relationship between its discriminant $n^n - (n - 1)^{n-1}$ and the resultant of its pair of irreducible factors.

Example 4. For any odd integer $k \geq 7$, set $n = 2^k$ and

$$(a, b, c) = ((n - 1)^{n-1}, n^n - (n - 1)^{n-1}, n^n).$$

Since n is congruent to 2 (mod 6), Lemma 4 tells us that $(\frac{n^2-n+1}{3})^2$ divides b . Thus

$$R(abc) = R(a)R(b)R(c) \leq (n - 1) \cdot \frac{b}{(n^2 - n + 1)/3} \cdot 2 < \frac{6b}{n} < \frac{6c}{n}. \tag{17}$$

Seeking a lower bound on n , we write $\log c = n \log n$ and $\log \log c = \log n + \log \log n < \frac{4}{3} \log n$ when $n > 100$, hence $n = \log c / \log n > \log c / (\frac{3}{4} \log \log c)$ and so

$$R(abc) < \frac{6b}{n} < \frac{6c}{\log c / (\frac{3}{4} \log \log c)} = \frac{8c \log \log c}{\log c}$$

when $k \geq 7$.

As stated so far, this construction yields a bound on the radical comparable to the bound from Example 3, but with a worse constant (although for large n , the 8 can essentially be replaced by a 6). However, if we choose specific values for n in the previous example in a manner suggested by Carl Pomerance, we can further decrease the radicals of the corresponding abc triples to be on par with the bound from Example 2.

Example 5. For any positive integer j , set $k = 3 \cdot 2^j$ in the triple of Example 4, so that $n = 8^{7^j}$. Using the divisibility (14), we see that 7^{j+1} divides $n - 1$ and thus $R(a) = R(n - 1) \leq (n - 1) / 7^j$. Therefore for the abc triples

$$(a, b, c) = \left((8^{7^j} - 1)^{8^{7^j} - 1}, 8^{7^j 8^{7^j}} - (8^{7^j} - 1)^{8^{7^j} - 1}, 8^{7^j 8^{7^j}} \right),$$

we may improve the bound (17) to

$$\begin{aligned} R(abc) = R(a)R(b)R(c) &\leq \frac{n-1}{7^j} \cdot \frac{b}{(n^2 - n + 1)/3} \cdot 2 \\ &< \frac{6b}{7^j n} < \frac{6c}{7^j n} = 6 \log 8 \frac{c}{\log c}. \end{aligned}$$

We end this section with a question: we have seen several elementary constructions of infinite families of abc triples, all of which yield an upper bound on $R(abc)$ somewhere between $c/\sqrt{\log c}$ and $c/\log c$ in magnitude. Is there an elementary construction of a sequence of abc triples satisfying $R(abc) < c/(\log c)^\lambda$ for some $\lambda > 1$, or equivalently, satisfying $q(a, b, c) \gtrsim 1 + \lambda \log \log c / \log c$? (We will see in Section 5.1 that such sequences exist, but the proof does not supply a formula for them, merely a proof of their existence.)

4. Background, motivation, and support for the abc conjecture

The abc conjecture was proposed in 1985 by Masser and Oesterlé [28, 33], who were motivated by two analogous problems concerning polynomial rings and elliptic curves. In addition, after the abc conjecture's appearance, number theorists found a probabilistic heuristic that also supports its statement. In this section we describe these links between the abc conjecture and other branches of mathematics.

4.1. The Mason–Stothers theorem

Despite their very different appearances, the integers \mathbb{Z} and the ring of polynomials with complex coefficients $\mathbb{C}[x]$ have a lot in common. In both settings, all nonzero elements enjoy unique factorization into irreducible elements: every integer can be written uniquely as a product of primes (and possibly -1), while every polynomial can be written uniquely as a product of monic linear factors $x - \rho$ (and possibly a nonzero leading coefficient in \mathbb{C}). Indeed, each ring is a principal ideal domain (PID), which is even stronger than being a unique factorization domain (UFD). In particular, one can define the radical $R(a)$ of a polynomial $a(x) \in \mathbb{C}[x]$ to simply be the product of all distinct monic linear factors that divide it, in perfect analogy with the radical of an integer. Similarly, one can define the greatest common divisor of two polynomials and hence decide whether two polynomials are relatively prime. (For these definitions, we ignore the leading coefficients, which are “units” in $\mathbb{C}[x]$, just as we take absolute values of integers to ignore their sign for the purposes of examining their factors.) It follows that the degree of the radical of a polynomial in $\mathbb{C}[x]$ is the same as the number of distinct complex roots of the polynomial.

The integers generate the rational numbers \mathbb{Q} , which are quotients of one integer by a second nonzero integer; the polynomials generate the aptly named rational functions $\mathbb{C}(x)$, which are quotients of one polynomial by a second polynomial that is not identically zero. The rational numbers form the simplest example of a *number field* (we will say more about number fields in Section 5.3), while the field of rational functions over \mathbb{C} form a *function field*; and it is a robust phenomenon in number theory (see for example [30, Chapter 1, Section 14]) that most results in number fields have analogous formulations in function fields. We have seen that irreducible polynomials correspond to prime numbers; another entry in the “dictionary” between the two rings is that the degree of a polynomial corresponds to the logarithm of a positive integer.

Masser’s description of the *abc* conjecture was motivated by the following theorem in the “function field case”, independently discovered by Stothers and Mason [44, 27] in the 1980s:

Theorem (Mason–Stothers). *Let $a(x), b(x), c(x) \in \mathbb{C}[x]$ be relatively prime polynomials satisfying $a(x) + b(x) = c(x)$. Then*

$$\max \{ \deg(a), \deg(b), \deg(c) \} \leq \deg(R(abc)) - 1.$$

As it happens, the proof of the Mason–Stothers theorem is actually quite elementary. Some versions of the proof (see for example [23, Chapter IV, Sections 3 and 9]) rely on one important feature of polynomials that is completely absent from the integers: the ability to take derivatives. For example, it is not hard to show that a polynomial is squarefree (that is, has no repeated factors in its factorization into linear polynomials) if and only if it is relatively prime to its derivative. Number theorists would love to be able to detect squarefree integers so easily!

What would happen if, in the Mason–Stothers theorem, we translated from the function field setting to the number field setting by replacing degree with logarithm

everywhere? We would obtain the statement $\max\{\log(a), \log(b), \log(c)\} + 1 \leq \log(R(abc))$, which, after exponentiating, becomes $e \max\{a, b, c\} \leq R(abc)$, or simply $R(abc) \geq ec$ if we order the three positive integers so that $a + b = c$. This is an instance of the “simplistic abc conjecture” we disproved thoroughly in Section 3. So the analogy between function fields and number fields, while fruitful, should always be taken with an epsilon grain of salt.

4.2. The Szpiro conjecture

In addition to the analogy with triples of polynomials, Oesterlé’s motivation for formulating the abc conjecture had an additional source: the subject of elliptic curves. In the early 1980s, L. Szpiro formulated the following conjecture relating the minimal discriminant of an elliptic curve to its conductor.

Szpiro Conjecture. *For every $\varepsilon > 0$, there exists a positive constant $S(\varepsilon)$ such that for any elliptic curve E defined by an equation with rational coefficients,*

$$|\Delta| \leq S(\varepsilon)N^{6+\varepsilon},$$

where Δ is the minimal discriminant of E and N is the conductor of E .

Oesterlé observed that the newly formulated abc conjecture is stronger than Szpiro’s conjecture: one can deduce Szpiro’s conjecture from the abc conjecture, but knowing Szpiro’s conjecture for all $\varepsilon > 0$, one can deduce the abc conjecture only when the ε in Version 4 is greater than $\frac{1}{5}$ (see [37, Chapter VIII, Exercise 8.20] and [49, Chapter 5, Appendix ABC]).

In fact, Oesterlé demonstrated [33, pages 169–170] that the abc conjecture is actually equivalent to the following modification of the Szpiro conjecture:

Modified Szpiro Conjecture. *For every $\varepsilon > 0$, there exists a positive constant $S'(\varepsilon)$ such that for any elliptic curve E whose minimal model is $y^2 = x^3 + a_4x + a_6$,*

$$\max\{|a_4|^3, a_6^2\} \leq S'(\varepsilon)N^{6+\varepsilon},$$

where N is the conductor of E .

Since $\Delta = -16(4a_4^3 + 27a_6^2)$, the modified Szpiro conjecture clearly implies the original; indeed, one can take $S(\varepsilon) = 16(4 + 27)S'(\varepsilon)$ and prove the original conjecture from the modified one. But it is possible, in theory, for Δ to be small only because of extreme cancellation when the hypothetically enormous numbers $4a_4^3$ and $27a_6^2$ are added together (note that a_4 can be negative).

The modified Szpiro conjecture is usually stated in terms of two particular invariants c_4 and c_6 of an elliptic curve rather than the coefficients a_4 and a_6 ; these invariants and the conductor N can be associated with any elliptic curve, no matter what equation originally defines it. The invariants c_4 and c_6 are special in the sense that they suffice to determine any elliptic curve E up to isomorphism (indeed, E can be defined by the equation $y^2 = x^3 - 27c_4x - 54c_6$). There are algorithms for computing these invariants from various defining equations. For

example (see [9, Section 3.2]), an algorithm of Laska–Kraus–Connell takes c_4 and c_6 , computed from any model of E , and outputs the minimal model of E ; while Tate’s algorithm computes, among other things, the conductor of E .

4.3. Heuristic based on a probabilistic model

While relating the *abc* conjecture to other mathematical statements is valuable, we might be comforted by having a more intrinsic reason to believe in its truth. One way that analytic number theorists hone their beliefs about how the integers work is by creating a random-variable situation that models the integer phenomenon. By rigorously showing that something happens with probability 1 in the random model, we can gain some confidence that the analogous statement really is true in the integers. The “Cramér model” of the distribution of primes (see for example [39]) is probably the most well-known example of this paradigm.

Here we give a probabilistic argument, adapted from Tao [45], for why one should expect the *abc* conjecture to hold. Broadly speaking, the argument asserts that if the radicals of three integers are too small, then the “probability” that two of the integers sum to the third is vanishingly small; this is assuming that these numbers with small radicals are “distributed randomly”. More specifically, we will argue that the following equivalent version of the *abc* conjecture should be true:

***abc* Conjecture, Version 6.** *Suppose α, β, γ are positive real numbers satisfying $\alpha + \beta + \gamma < 1$. When M is sufficiently large (in terms of α, β, γ), there are no solutions to the equation $a + b = c$ with $c \geq M$, where a, b, c are relatively prime positive integers satisfying $R(a) \leq M^\alpha, R(b) \leq M^\beta, R(c) \leq M^\gamma$.*

We see rather easily that Version 1 of the *abc* conjecture implies this new version: given M and α, β, γ with $\alpha + \beta + \gamma < 1$, we must have

$$R(abc) = R(a)R(b)R(c) \leq M^\alpha M^\beta M^\gamma = M^{\alpha+\beta+\gamma} < c^{\alpha+\beta+\gamma},$$

for any triple satisfying the hypotheses of Version 6. But by Version 1 with $\varepsilon = 1 - (\alpha + \beta + \gamma) > 0$, there can be only finitely many such triples (a, b, c) ; simply choose M larger than the largest c that occurs in any of them.

It is a little more difficult to see that Version 6 of the *abc* conjecture implies Version 1: on the face of it, the loophole phrase “sufficiently large (in terms of α, β, γ)” might allow an infinite sequence of counterexamples to Version 1 corresponding to a sequence of distinct triples α, β, γ . However, if there were in fact an infinite sequence of counterexamples to Version 1 for some fixed ε , then the corresponding sequence $(\alpha, \beta, \gamma) = \left(\frac{\log R(a)}{\log c}, \frac{\log R(b)}{\log c}, \frac{\log R(c)}{\log c}\right)$ would all lie in a compact region of \mathbb{R}^3 (namely the simplex in the positive orthant defined by $x + y + z \leq 1 - \varepsilon$), and hence some subsequence of them would converge to a fixed point $(\alpha_0, \beta_0, \gamma_0)$ satisfying $\alpha_0 + \beta_0 + \gamma_0 \leq 1 - \varepsilon$. Consequently, Version 6 of the *abc* conjecture could be applied to the slightly larger point $(\alpha, \beta, \gamma) = ((1 + \varepsilon)\alpha_0, (1 + \varepsilon)\beta_0, (1 + \varepsilon)\gamma_0)$ to derive a contradiction.

Now let's use a probabilistic model to probe Version 6 itself. We will need the following lemma, standard in analytic number theory (see [45]), saying that the numbers with a given radical are quite sparse:

Proposition. *For every $\varepsilon > 0$, there exists a constant $T(\varepsilon)$ such that for every positive squarefree integer r and every $M > 0$, there are at most $T(\varepsilon)M^\varepsilon$ integers less than $2M$ whose radical equals r .*

(This upper bound for the number of such integers will feature again, in a more precise form, in a lower bound given in Section 5.1.) Using this lemma we can estimate, given α, β, γ , how many triples of integers (a, b, c) there are with $1 \leq a, b \leq 2M$, $M < c \leq 2M$ and $R(a) \leq M^\alpha, R(b) \leq M^\beta, R(c) \leq M^\gamma$. (Note that for the moment we are not paying attention to whether two of these numbers sum to the third number.) There are at most $M^\alpha \cdot M^\beta \cdot M^\gamma$ possibilities for the three radicals, even if we forget that they have to be squarefree and relatively prime. Choosing $\varepsilon = \frac{1}{4}(1 - (\alpha + \beta + \gamma)) > 0$, the above proposition tells us that there are $T(\varepsilon)M^\varepsilon$ possibilities for a for any given $R(a)$, and similarly for b and c . Thus the number of such triples is at most

$$M^\alpha \cdot M^\beta \cdot M^\gamma \cdot (T(\varepsilon)M^\varepsilon)^3 = T(\varepsilon)^3 M^{\alpha+\beta+\gamma+3\varepsilon} = T(\varepsilon)^3 M^{1-\varepsilon}.$$

Now, instead of looking at the specific collection of triples described in the previous paragraph, let us suppose that *we choose the same number of triples completely at random* from the set of triples (a, b, c) with $1 \leq a, b \leq 2M$, $M < c \leq 2M$; what is the probability that at least one of the chosen triples satisfies $a + b = c$? The probability of a single chosen triple satisfying $a + b = c$ is at most $1/M$, since there is at most one correct choice out of M for c , no matter what a and b are. Therefore the probability that we obtain such a chosen triple is at most $T(\varepsilon)^3 M^{1-\varepsilon} \cdot M^{-1} = T(\varepsilon)^3 M^{-\varepsilon}$. Because we have no reason to think that the actual triples described in the previous paragraph are any more or less likely to satisfy $a + b = c$ than randomly chosen triples, we are persuaded of the following heuristic: the “probability” is at most $T(\varepsilon)^3 M^{-\varepsilon}$ that there exists a triple (a, b, c) with $1 \leq a, b \leq 2M$, $M < c \leq 2M$ that successfully satisfies the conditions in the second sentence of Version 6 of the *abc* conjecture.

Every triple with $a + b = c$ satisfies $1 \leq a, b \leq 2^{k+1}$, $2^k < c \leq 2^{k+1}$ for a unique positive integer k . Let the “ k th event” be the assertion that there exists a successful triple for $M = 2^k$ as described above. The “probability” of the k th event, by the above heuristic, is at most $T(\varepsilon)^3 (2^k)^{-\varepsilon}$. Notice that the series

$$\sum_{k=0}^{\infty} T(\varepsilon)^3 (2^k)^{-\varepsilon} = \frac{T(\varepsilon)^3}{1 - 2^{-\varepsilon}}$$

converges to a finite number. Therefore, by the Borel–Cantelli lemma [20, pages 51–52], with probability 1 only finitely many of the events occur. We conclude that heuristically, only finitely many triples should successfully satisfy the conditions in the second sentence of Version 6 of the *abc* conjecture; so if we choose M large enough, we believe that there will be no counterexamples remaining.

In Section 5.2, we discuss a refinement of this heuristic that enabled the authors of [36] to propose a stronger, even more precise version of the *abc* conjecture.

It is tempting to think of the *abc* conjecture as a vast conspiracy that arranges the numbers with small radicals so precisely that no two of them ever add to a third. This temptation is even stronger when we see examples like the infinite families from Section 3: we forced the smallest and largest numbers to have tiny radicals, but the *abc* conjecture seems, like some mystical force, to keep the middle number from being divisible by too large a perfect square. However, the above heuristic (with, say, $\alpha = \gamma = \varepsilon$ and $\beta = 1 - 3\varepsilon$) offers an explanation: these families do not contain all that many triples, and probability simply dictates that it is overwhelmingly unlikely for any of the middle numbers in such a sparse set to have a small radical.

We conclude this section by remarking that the same heuristic suggests, when $\alpha + \beta + \gamma > 1$, that there do exist infinitely many triples (a, b, c) with $a + b = c$ and $c \geq M$ that satisfy $R(a) \leq M^\alpha$, $R(b) \leq M^\beta$, and $R(c) \leq M^\gamma$. In other words, we expect the set of ordered triple $(\frac{\log R(a)}{\log c}, \frac{\log R(b)}{\log c}, \frac{\log R(c)}{\log c})$ to have the entire triangular boundary $T = \{x, y, z \geq 0 : x + y + z = 1\}$ in \mathbb{R}^3 in its set of limit points.

5. Generalizations, refinements, and the state of the art

In this last section, we explore our most state-of-the-art knowledge about the *abc* conjecture. We describe some (less elementary) constructions of infinite families of *abc* triples with much better asymptotic quality than the ones we have seen so far; we present several refinements of the conjecture, which attempt to decide where in the space between the “simplistic” and actual *abc* conjectures the exact boundary lies; we generalize the *abc* conjecture to other settings and to more variables; and finally we discuss how close we are to actually proving the inequalities that the *abc* conjecture asserts.

5.1. Best known *abc* triples

We already saw in Section 3 that having examples of families of *abc* triples helps us probe how sharp the *abc* conjecture is. Presumably, thinking more deeply about how to find good *abc* triples would yield even smaller radicals than the ones we have seen thus far. Stewart and Tijdeman [41, Theorem 2] did exactly this in 1986: they came up with a construction of infinitely many *abc* triples of higher quality than those in Section 3.

They proved that for any $\delta > 0$, there exist infinitely many triples (a, b, c) of relatively prime positive integers with $a + b = c$ that satisfy

$$c > R(abc) \exp \left((4 - \delta) \frac{\sqrt{\log R(abc)}}{\log \log R(abc)} \right). \tag{18}$$

It can be shown that for any $B > 1$, every sufficiently large *abc* triple satisfying the bound (18) also satisfies $R(abc) < c/(\log c)^B$. In fact, (18) is equivalent to

a lower bound for the quality of the form

$$q(a, b, c) > 1 + \frac{4 - \delta}{\sqrt{\log c} \cdot \log \log c}. \quad (19)$$

Both of these observations show that these abc triples are far better than the ones in Section 3. Certainly they amply refute the “simplistic abc conjecture” and show that the epsilons in the statement of the real abc conjecture must be present. But again, the lower bound for the quality tends to 1 as c grows large, and so even this better construction does not disprove the actual conjecture.

Stewart and Tijdeman’s construction is quite illuminating. First, given a positive integer r and a parameter X that is far larger than r , they consider the set of integers up to X whose factorizations contain only the first r odd primes p_1, \dots, p_r ; such integers are called “ p_r -friable” (or “ p_r -smooth”). They obtain good estimates for the number of such integers by noting that the number of solutions to $p_1^{n_1} \cdots p_r^{n_r} \leq X$ is the same as the number of lattice points (n_1, \dots, n_r) in the r -dimensional simplex

$$\{x_1, \dots, x_r \geq 0 : x_1 \log p_1 + \cdots + x_r \log p_r \leq \log X\}.$$

When X is large, this number of lattice points is essentially the r -dimensional volume of the simplex, which is easy to calculate. Their resulting lower bound for the number of p_r -friable integers up to X is a more precise version of the Proposition from Section 4.3.

Then, they point out that two of these p_r -friable integers must be congruent to each other modulo a high power of 2; indeed, if the number of such integers exceeds 2^k , then the pigeonhole principle forces two of them to lie in the same residue class modulo 2^k . (And if those two happen to have common factors, their quotients by their greatest common divisor will also be congruent to each other.) These two integers and their difference form a triple satisfying $a + b = c$; the product of the radicals of the first two integers is at most $p_1 \cdots p_r$; and the radical of their difference is at most $X/2^{k-1}$.

Of course, it is necessary to write down explicitly the relationships among all of these functions and parameters; analytic number theorists learn tools that are precisely suited for converting from the above sketch to a full quantitative proof. When the dust settles, the inequality (18) is the payoff.

Later, van Frankenhuysen [48] added an improvement to the argument of Stewart and Tijdeman: instead of using the full lattice of integer points (n_1, \dots, n_r) , he chose a sublattice sitting askew inside the full integer lattice in such a way that the points in the sublattice were relatively more tightly packed together. In this way, and using high-dimensional sphere-packing bounds already in the literature, he showed that one could improve the constant $4 - \delta$ in the above inequalities to 6.068.

5.2. Refinements of the abc conjecture

In a sense, the loophole phrases “only finitely many” and “there exists a positive constant” make it hard to actually determine from data whether the abc conjecture

is acting like the truth. In 1996, Baker [4] refined the conjecture to provide some insight into how the constant $K(\varepsilon)$ in Version 4 of the *abc* conjecture should depend on ε . Let the function $\omega(n)$ denote the number of distinct primes (that is, ignoring repetitions) dividing n . Baker proposed the following refinement: there exists an absolute constant K_1 such that, if a, b, c are relatively prime positive integers satisfying $a + b = c$, then

$$c < \varepsilon^{-K_1 \min\{\omega(ab), \omega(ac), \omega(bc)\}} R(abc)^{1+\varepsilon}$$

for any $\varepsilon > 0$. (The minimum in the exponent is there to help us: we can keep whichever two of the three numbers have the fewest prime factors between them.) Although this bound has an extra dependence on a, b, c , this dependence turns out to be smaller than $R(abc)^\varepsilon$, and so we have not significantly altered the shape of the conjecture. It is true that there is still an unknown absolute constant $K_1 > 0$ in this formulation; but at least now this constant is independent of ε , which is often quite important when making deductions.

Baker did demonstrate that there exist infinitely many *abc* triples satisfying the related inequality

$$c > K_2 \varepsilon^{1-\min\{\omega(ab), \omega(ac), \omega(bc)\}} R(abc)^{1+\varepsilon}$$

for some absolute constant $K_2 > 0$. In fact, his proof relies upon estimates for *linear forms in logarithms*, a profound technical tool in Diophantine approximation for which Baker was awarded the Fields Medal in 1970. He also mentions that Granville conjectured that there exists an absolute constant $K_3 > 1$ such that $c < K_3^{-\Omega(abc)} R(abc)$ for all *abc* triples, where $\Omega(n)$ counts the number of prime factors of n with multiplicity (so for example, $\omega(72) = 2$ but $\Omega(72) = 5$). Notice that there is no exponent $1 + \varepsilon$ on the right-hand side of this conjecture! But of course the slack has to go somewhere: $\Omega(abc)$ can be far larger than $\omega(abc)$.

Most recently, another refinement has been put forward by Robert, Stewart, and Tenenbaum [36], following up on a heuristic proposed by van Frankenhuysen in his PhD thesis. In Section 4.3 we described the heuristic assumption that statistically, $R(c)$ is distributed independently from $R(a)$ and $R(b)$ when a, b, c are relatively prime and $a + b = c$; we then estimated how many integers up to M have radicals bounded by M^α and so on. Through an extremely careful study of the function that counts how many integers up to M have their radical bounded by a second parameter Y , Robert, Stewart, and Tenenbaum proposed the following two-part conjecture: First, there exists a real number K_4 such that all *abc* triples satisfy

$$c < R(abc) \exp \left(4 \sqrt{\frac{3 \log R(abc)}{\log \log R(abc)}} \left(1 + \frac{\log \log \log R(abc)}{2 \log \log R(abc)} + \frac{K_4}{\log \log R(abc)} \right) \right); \quad (20)$$

second, there exists a real number K_5 such that infinitely many *abc* triples satisfy

$$c > R(abc) \exp \left(4 \sqrt{\frac{3 \log R(abc)}{\log \log R(abc)}} \left(1 + \frac{\log \log \log R(abc)}{2 \log \log R(abc)} + \frac{K_5}{\log \log R(abc)} \right) \right). \quad (21)$$

(Indeed, they included more detailed versions of this refinement with even more secondary main terms inside the exponential.) Note that the right-hand side of their second conjecture (21) is a little bit larger than the lower bound (18) coming from the construction of Stewart and Tijdeman, since we are dividing only by $\sqrt{\log \log R(abc)}$ in the main term inside the exponential instead of $\log \log R(abc)$. It can be shown that their first conjecture (20) really does imply Version 2 of the *abc* conjecture.

Interestingly, a special quantity arises from these conjectures of Robert, Stewart, and Tenenbaum. Define the *merit* of an *abc* triple to be

$$m(a, b, c) = (q(a, b, c) - 1)^2 \log R(abc) \log \log R(abc).$$

Every infinite family of *abc* triples ever established has the property that their merit tends to 0 (it is not hard to verify this for the Stewart–Tijdeman examples, for instance, assuming that the right-hand side of the inequality (19) is in fact the correct size of the quality). If the *abc* conjecture were false, it is an easy deduction from Version 5 that the merit would be unbounded above. But it would actually follow from the conjectures (20) and (21) that the lim sup of all merits of all *abc* triples equals 48 exactly! So the merit is somehow an incredibly fine-scale measurement of an *abc* triple, one that looks at the boundary between possible and impossible through a powerful microscope.

For the record, the largest merit found to date is approximately 38.67, which comes from the *abc* triple $a = 2,543^4 \cdot 182,587 \cdot 2,802,983 \cdot 85,813,163$, $b = 2^{15} \cdot 3^{77} \cdot 11 \cdot 173$, $c = 5^{56} \cdot 245,983$ discovered by Ralf Bonse in 2011. de Smit’s web site [10] lists the 131 known *abc* triples with merit greater than 24.

5.3. Other alterations of the conjecture

Beyond these results, some interesting generalizations and refinements of the *abc* conjecture have appeared in various contexts.

Congruence *abc* conjecture

First we state yet another version of the conjecture, which looks like it concerns only a small subset of *abc* triples but is actually equivalent to the other versions we have seen so far.

***abc* Conjecture, Version 7.** *For every positive integer N and every $\varepsilon > 0$, there exists a positive constant $E(N, \varepsilon)$ such that if $N \mid abc$, where a, b, c are relatively prime positive integers with $a + b = c$, then*

$$c \leq E(N, \varepsilon) R(abc)^{1+\varepsilon}.$$

The special case $N = 1$ is of course the familiar *abc* conjecture (specifically, Version 4). Motivated by Oesterlé’s observation [33] that the special case $N = 16$ of this new version implies the full *abc* conjecture, Ellenberg [13] demonstrated that in fact 16 can be replaced by any integer N , thus showing that Version 7 of the *abc* conjecture really is equivalent to the others.

Uniform *abc* conjecture for number fields

An important focus in algebraic number theory is the study of *number fields*, which are finite field extensions of the field \mathbb{Q} of rational numbers (equivalently, a field of the form $\mathbb{Q}(\alpha)$, where α is a root of a polynomial with integer coefficients). Vojta [49, page 84] formulated a generalization of the *abc* conjecture for arbitrary number fields. A version of this generalization is given by Granville and Stark [14], which is adapted from Elkies’ reformulation [12] of Vojta’s work.

Uniform *abc* Conjecture. *For every $\varepsilon > 0$, there exists a constant $U(\varepsilon) > 0$ with the following property: for every number field K of degree n and discriminant D_K over \mathbb{Q} , and every triple (a, b, c) of elements of K satisfying $a + b + c = 0$,*

$$H(a, b, c) \leq U(\varepsilon) (D_K^{1/n} N(a, b, c))^{1+\varepsilon},$$

where $H(a, b, c)$ and $N(a, b, c)$ are the height and conductor of the triple (a, b, c) .

This generalization would imply many additional consequences, even more than the traditional *abc* conjecture. For example, Granville and Stark showed that the uniform *abc* conjecture implies a strong lower bound on the class numbers of imaginary quadratic fields (equivalently, on the number of inequivalent binary quadratic forms of a given negative discriminant) and also rules out “exceptional zeros” of certain Dirichlet L -functions, which is a major barrier to approaching the generalized Riemann hypothesis.

Additional integers

Our last generalization, stated by Browkin and Brzezinski [7], incorporates more variables than the three we have been working with so far.

***abc* Conjecture, n -Variable Version.** *For every integer $n \geq 3$ and every $\varepsilon > 0$, there exists a positive constant $B(n, \varepsilon)$ such that all relatively prime n -tuples (a_1, \dots, a_n) of nonzero integers with $a_1 + \dots + a_n = 0$ and no vanishing subsums satisfy*

$$\max\{|a_1|, \dots, |a_n|\} \leq B(n, \varepsilon) R(|a_1 \cdots a_n|)^{2n-5+\varepsilon}.$$

Here, “no vanishing subsums” means that it is not possible to reorder a_1, \dots, a_n so that $a_1 + \dots + a_k = 0 = a_{k+1} + \dots + a_n$ for some $1 \leq k \leq n - 1$; this hypothesis is necessary because of trivial examples such as $(a_1, a_2, a_3, a_4) = (2^n, -2^n, 3^n, -3^n)$, which is a relatively prime quadruple even though some pairs of terms have huge common factors. This n -variable version is our familiar friend when $n = 3$: given such a triple a_1, a_2, a_3 , one of them has a different sign than the other two, and we recover Version 4 of the *abc* conjecture by letting c be the absolute value of the one with a different sign and a, b the absolute values of the other two.

Browkin and Brzezinski constructed examples showing that the exponent $2n - 5 + \varepsilon$ on the right-hand side cannot be reduced; their constructions are rather similar to the transfer method described in Section 2.4. Taking $n = 4$ for example, if (a, b, c) is any *abc* triple, we may set

$$(a_1, a_2, a_3, a_4) = (a^3, 3abc, b^3, -c^3), \tag{22}$$

which one can check does satisfy $a_1 + a_2 + a_3 + a_4 = 0$; for these quadruples,

$$\begin{aligned} \max\{|a_1|, \dots, |a_n|\} = c^3 &\geq (R(abc))^3 \\ &\geq \left(\frac{1}{3}R(|a_1a_2a_3a_4|)\right)^3 = \frac{1}{27}(R(|a_1a_2a_3a_4|))^{2 \cdot 4 - 5}. \end{aligned}$$

Equation (22) is the $n = 4$ case of a sequence of impressive identities: when $n \geq 3$,

$$a_j = \frac{2n-5}{2j-1} \binom{n+j-4}{2j-2} a^{2j-1} (bc)^{n-j-2} \quad \text{for } 1 \leq j \leq n-2, \quad a_{n-1} = b^{2n-5}, \quad a_n = -c^{2n-5} \quad (23)$$

is an n -tuple satisfying the hypotheses of the n -variable version of the abc conjecture. For these n -tuples, the maximum absolute value is c^{2n-5} , while $R(|a_1 \cdots a_n|)$ is at most a constant (the product of all the primes up to $2n-5$, say) times $R(abc)$, and $R(abc)$ is in turn less than c when (a, b, c) is an abc triple. Not only does this construction show that the exponent $2n-5+\varepsilon$ would be best possible, it also shows that the n -variable version for any $n \geq 4$ implies the usual three-variable abc conjecture. (It seems less clear whether, for example, the 5-variable version of the abc conjecture implies the 4-variable version.)

Interestingly, the statement of the n -variable version of the abc conjecture is not what one would predict from a probabilistic heuristic like the one described in Section 4.3: the analogous argument would lead again to a conjecture with exponent $1+\varepsilon$ on the right-hand side. In this case, probability would lead us astray—but presumably because the set of counterexamples is extremely thin, coming only from constructions like equation (22). In fact, it follows from a sufficiently strong version of Vojta’s conjecture [50, Conjecture 2.3] that the exponent $2n-5+\varepsilon$ can be reduced to $1+\varepsilon$ if we exclude a finite number of constructions like equation (23) for each n . (Those constructions have the property that the constructed n -tuples are not pairwise relatively prime. However, similar constructions [19] can be found where the entries are indeed pairwise coprime; see also the last section of [34] for an example due to Granville.)

5.4. Progress towards the abc conjecture

Baker’s theory of linear forms in logarithms has been a useful aid for mathematicians to obtain effective results toward the conjecture. The first players of this game were Stewart and Tijdeman [41]: in 1986 they proved that

$$c < \exp(K_6 R(abc)^{15}) \quad (24)$$

for some constant $K_6 > 0$. Their proof used bounds on linear forms in logarithms similar to those mentioned in Section 5.2, in particular a p -adic version due to van der Poorten. Subsequently, Stewart and Yu [42, 43] improved the bound (24) to

$$c < \exp(K_7 R(abc)^{1/3} (\log R(abc))^3)$$

for some constant $K_7 > 0$. Also noteworthy is a paper by Murty and Pasten in [29], in which they obtained similar bounds via a fresh approach based on modular forms. Yet, despite the fact that all the above results were hard-earned and brought the problem of bounding c into the realm of the finite, none of these results is as good as $c < R(abc)^B$ for any fixed B .

The number theory community has been abuzz with the topic of the *abc* conjecture the past few years. In August 2012, Shinichi Mochizuki released the final installment of his series of four papers on “inter-universal Teichmüller theory”, in which he claimed to have proven the *abc* conjecture as a consequence of his work. His proof, with its incredible length and heavy dependence on a new and untested field with a limited number of practitioners, is still under verification by the mathematical community. Moreover, due to the introduction of several arcane objects such as “Frobenioids,” “log-theta-lattice”, and “alien arithmetic holomorphic structures,” a cautious response from the mathematical community was inevitable.

Mochizuki published progress reports in December 2013 and December 2014, informing the community of the advancement that had been made towards verifying his results. (See the Polymath page [26] for useful links to Mochizuki’s papers, progress reports, announcements, and other related topics.) Members of his home university have studied his preparatory papers and waded through his manuscripts on inter-universal Teichmüller theory, communicating with Mochizuki on suggested improvements and adjustments to be made; they plan to give seminars on the material and to publicly post more accessible write-ups of the work in early 2015. On the other hand, due to the esoteric nature of Mochizuki’s work (and the presence of some at least superficial mistakes in the deduction of the *abc* conjecture from his theory), it has been hard for others to attest to the validity of his results. While a wave of colleagues around the world was drawn to the task of understanding his exotic, potentially revolutionary work, the reality is that it is difficult for most academics to pause their own research to invest the necessary energy. Several skilled mathematicians spent a good deal of time trying to understand how the arguments were structured but, after making little headway in being able to verify Mochizuki’s claims, eventually abandoned the project.

This unsettled state of affairs begs the question: when does one say that a problem in mathematics has been solved? Many of us would like to think we have an absolute standard, where proofs are accepted if and only if they are completely rigorous and complete, line by line, like a successfully compiling computer program. But in practice, we tolerate typos, allusions to proofs of similar cases, sketches of arguments, and occasional exercises for the reader as acceptable parts of research papers; our standard of proof in mathematics is a social construct [46, Section 4]. Researchers in specialized fields form their own epistemic communities and move forward in clusters, building around one another’s work, and sharing their knowledge with researchers in neighboring areas as they can.

A mathematician’s results, then, are accepted only when her primary audience — the cohort of experts occupying the same niche—has validated their accuracy. In this case, with Mochizuki’s original and complex work, it will take some time for more mathematicians to surmount the barrier and begin exporting the ideas

to the wider community. In the best possible world, experts will come to agree that the papers contain a proof to one of the most significant problems in number theory, as well as the foundations of new areas of research. But until and unless that happens, we must be content with the *abc* conjecture remaining a mystery, at least for now.

Acknowledgments. The authors are happy to thank Tim Dokchitser for indicating where to locate the material in Section 2.6 in the literature; Carlo Beenakker and Cam Stewart for their help in finding attributions for some of the examples in Section 3.2; joro, Felipe Voloch, and Trevor Wooley for helpful comments on the n -variable version of the *abc* conjecture mentioned in Section 5.3 and the connection to Vojta’s work; and the referees for many helpful observations and suggestions.

References

- [1] ABC@Home, The algorithm, Mathematisch Instituut Universiteit Leiden. <http://abcathome.com/algorithm.php> (accessed August 3, 2014)
- [2] D. P. Anderson, R. Walton, C. Fenton, et al., Berkeley Open Infrastructure for Network Computing (BOINC), April 10, 2002. <http://boinc.berkeley.edu> (accessed August 3, 2014)
- [3] T.M. Apostol, *Introduction to Analytic Number Theory*, Springer, 1976.
- [4] A. Baker, *Logarithmic forms and the abc-conjecture*, in Number Theory: Diophantine, Computational and Algebraic Aspects, Eger–1996, K. Györy, A. Pethö, and V.T. Sós, Proc. Intl. Conf., de Gruyter, Berlin, 1998, 37–44.
- [5] E. Bombieri, *Roth’s theorem and the abc conjecture*, preprint (1994).
- [6] D. Boyd, G. Martin, and M. Thom, *Squarefree values of trinomial discriminants*, LMS J. Comput. Math. **18** (2015), no. 1, 148–169.
- [7] J. Browkin and J. Brzezinski, *Some remarks on the abc-conjecture*, Math. Comp. **62** (1994), no. 206, 931–939.
- [8] N.A. Carella, *Probable counterexamples of the ABC conjecture*, 2003. <http://arxiv.org/pdf/math/0503401v1.pdf>
- [9] J. E. Cremona, *Algorithms for Modular Elliptic Curves*, Cambridge University Press, 1997.
- [10] B. de Smit, *Update on abc triples*, March 2007. <http://pub.math.leidenuniv.nl/~smitbde/abc> (accessed August 3, 2014)
- [11] T. Dokchitser, *LLL and ABC*, J. Number Theory **107** **1** (2004), 161–167.
- [12] N. Elkies, *ABC implies Mordell*, Int. Math. Res. Notices (1991), no. 7, 99–109.
- [13] J.S. Ellenberg, *Congruence ABC implies ABC*, Indag. Math. N.S. **11** (2), 197–200 (2000).
- [14] A. Granville and H.M. Stark, *ABC implies no “Siegel zeros” for L-functions of characters with negative discriminant*, Invent. Math. **139**, no.3, 509–523.
- [15] A. Granville and T.J. Tucker, *It’s as easy as abc*, Notices of the A.M.S. **49** (2002), 1224–1231.

- [16] B. Gross and D. Zagier, *On singular moduli*, J. Reine Angew. Math. **355** (1985), 191–220.
- [17] G.H. Hardy and E.M. Wright, *An Introduction to the Theory of Numbers*, Oxford University Press, 1979.
- [18] C. Hurlburt, *Good abc triples*. <http://www.phfactor.net/abc> (accessed August 3, 2014)
- [19] joro, *Argument againts the abcd conjecture with extra gcd conditions*, MathOverflow. <https://mathoverflow.net/questions/185857/argument-againts-the-abcd-conjecture-with-extra-gcd-conditions> (accessed January 10, 2015)
- [20] A. Klenke, *Probability Theory*, Springer, 2014.
- [21] S.K. Lando and A.K. Zvonkin, *Graphs on Surfaces and Their Applications*, Encyclopaedia of Mathematical Sciences: Lower-Dimensional Topology II 141 (Springer-Verlag), 2004.
- [22] S. Lang, *Old and new conjectured Diophantine inequalities*, Bull. Amer. Math. Soc. (N.S.) **23** (1990), no. 1, 37–75.
- [23] S. Lang, *Undergraduate Algebra*, Springer, New York, 2005.
- [24] K. Lauter and B. Viray, *On singular moduli for arbitrary discriminants*, Int. Math. Res. Not. IMRN, to appear.
- [25] A.K. Lenstra, H.W. Lenstra, Jr., and L. Lovász, *Factoring polynomials with rational coefficients*, Mathematische Annalen **261** (1982), no. 4, 515–534.
- [26] Mabcp, MarioKappa, and T. Tao, *ABC conjecture*, Polymath, August 8, 2009. http://michaelnielsen.org/polymath1/index.php?title=ABC_conjecture (accessed August 11, 2014)
- [27] R.C. Mason, *Diophantine equations over function fields*, London Mathematical Society Lecture Note Series, 96, Cambridge University Press, Cambridge, 1984.
- [28] D.W. Masser, *Open problems*, Proc. Symp. Analytic Number Theory, (W.W.L. Chen, ed.), Imperial Coll. London, 1985.
- [29] M.R. Murty and H. Pasten, *Modular forms and effective Diophantine approximation*, Journal of Number Theory **133** (2013), no. 11, 3739–3754.
- [30] J. Neukirch, *Algebraic Number Theory* (translated by N. Schappacher), Springer-Verlag, Berlin, 1999.
- [31] A. Nitaj, <http://www.math.unicaen.fr/~nitaj/abc.html#Consequences>, 2013.
- [32] I. Niven, H.S. Zuckerman, and H.L. Montgomery, *An Introduction to the Theory of Numbers*, 5th edition, John Wiley & Sons, Inc., New York (1991).
- [33] J. Oesterlé, *Nouvelles approches du théorème de Fermat*, Sémin. Bourbaki 1987-1988, Vol.694, Astérisque **161-162** (1988), 165–186.
- [34] C. Pomerance, *Computational number theory*, in Princeton Companion to Mathematics (ed. W.T. Gowers), Princeton U. Press, Princeton, New Jersey, 2008, 348–362.
- [35] Reken mee met ABC, Mathematisch Instituut Universiteit Leiden. <http://www.rekenmeemetabc.nl> (accessed August 3, 2014)
- [36] O. Robert, C.L. Stewart, and G. Tenenbaum, *A refinement of the abc conjecture*, Bull. London Math. Soc. **46** (2014), no. 6, 1156–1166.

- [37] J.H. Silverman, *Arithmetic of Elliptic Curves*, Springer, New York, 2009.
- [38] J.H. Silverman, *Wieferich's criterion and the abc-conjecture*, J. Number Theory **30** (1988), 226–237.
- [39] K. Soundararajan, *The distribution of prime numbers*, in Equidistribution in number theory, an introduction, NATO Sci. Ser. II Math. Phys. Chem. **237**, Springer (Dordrecht), 2007, 59–83.
- [40] C.L. Stewart, *A note on the product of consecutive integers*, in Topics in classical number theory, Vol. I, II (Budapest, 1981), Colloq. Math. Soc. János Bolyai **34**, North-Holland, Amsterdam, 1984, 1523–1537.
- [41] C.L. Stewart and R. Tijdeman, *On the Oesterlé-Masser conjecture*, Monatsh. Math. **102** (1986), no.3, 251–257.
- [42] C.L. Stewart and K. Yu, *On the abc conjecture*, Math. Ann. **291** (1991), 225–230.
- [43] C.L. Stewart and K. Yu, *On the abc conjecture, II*, Duke Math. Journal **108** (2001), 169–181.
- [44] W.W. Stothers, *Polynomial identities and Hauptmoduln*, Q.J. Math. Oxford **32** (1981), 349–370.
- [45] T. Tao, *The probabilistic heuristic justification of the ABC conjecture*, What's new, September 18, 2012. <http://terrytao.wordpress.com/2012/09/18/the-probabilistic-heuristic-justification-of-the-abc-conjecture/> (accessed August 11, 2014)
- [46] W.P. Thurston, *On proof and progress in mathematics*, Bulletin of the AMS **30** (1994), no. 2, 161–177.
- [47] J. van der Horst, *Finding ABC-triples using elliptic curves*, Masters thesis, Universiteit Leiden, 2010.
- [48] M. van Frankenhuysen, *A lower bound in the abc conjecture*, J. Number Theory **82** (2000), 91–95.
- [49] P. Vojta, *Diophantine Approximations and Value Distribution Theory*, Springer, New York, 1987.
- [50] P. Vojta, *A more general ABC conjecture*, Internat. Math. Res. Notices **21** (1998), 1103–1116.
- [51] L.C. Washington, *Elliptic Curves: Number theory and cryptography*, 2nd ed., Chapman & Hall/CRC, Boca Raton, 2008.

Address: Greg Martin and Winnie Miao: Department of Mathematics, University of British Columbia, Room 121, 1984 Mathematics Road, Vancouver, BC, Canada V6T 1Z2.

E-mail: gerg@math.ubc.ca, miao.winnie@gmail.com

Received: 4 September 2015; **revised:** 16 February 2016