

Affine representability results in \mathbb{A}^1 –homotopy theory

II: Principal bundles and homogeneous spaces

ARAVIND ASOK
 MARC HOYOIS
 MATTHIAS WENDT

We establish a relative version of the abstract “affine representability” theorem in \mathbb{A}^1 –homotopy theory from part I of this paper. We then prove some \mathbb{A}^1 –invariance statements for generically trivial torsors under isotropic reductive groups over infinite fields analogous to the Bass–Quillen conjecture for vector bundles. Putting these ingredients together, we deduce representability theorems for generically trivial torsors under isotropic reductive groups and for associated homogeneous spaces in \mathbb{A}^1 –homotopy theory.

14F42, 14L10, 20G15, 55R15

1 Introduction

Suppose k is a fixed commutative unital base ring, and write $\mathcal{H}(k)$ for the Morel–Voevodsky \mathbb{A}^1 –homotopy category over k [45]. The category $\mathcal{H}(k)$ is constructed as a certain localization of the category of simplicial presheaves on Sm_k , the category of smooth k –schemes. Write $\mathrm{Sm}_k^{\mathrm{aff}}$ for the subcategory of Sm_k consisting of affine schemes. If \mathcal{X} is a simplicial presheaf on Sm_k , by an “affine representability” result for \mathcal{X} , we will mean, roughly, a description of the presheaf on $\mathrm{Sm}_k^{\mathrm{aff}}$ defined by $U \mapsto [U, \mathcal{X}]_{\mathbb{A}^1} := \mathrm{Hom}_{\mathcal{H}(k)}(U, X)$.

Here is a flavor of the description we provide: if \mathcal{X} is a simplicial presheaf on Sm_k , then for any $U \in \mathrm{Sm}_k^{\mathrm{aff}}$ one can consider the simplicial set $\mathrm{Sing}^{\mathbb{A}^1} \mathcal{X}(U)$ [45, page 87]. The 0–simplices of this simplicial set are morphisms $U \rightarrow \mathcal{X}$ and the 1–simplices are “naive” or “elementary” \mathbb{A}^1 –homotopies $U \times \mathbb{A}^1 \rightarrow \mathcal{X}$. The assignment $U \mapsto \pi_0(\mathrm{Sing}^{\mathbb{A}^1} \mathcal{X}(U))$ defines a presheaf $\pi_0(\mathrm{Sing}^{\mathbb{A}^1} \mathcal{X})$ of “naive” \mathbb{A}^1 –homotopy classes of maps $U \rightarrow \mathcal{X}$. In [9], we gave conditions that allowed us to identify $\pi_0(\mathrm{Sing}^{\mathbb{A}^1} \mathcal{X})(U) \cong [U, \mathcal{X}]_{\mathbb{A}^1}$, ie under which “naive” \mathbb{A}^1 –homotopy classes coincide with “true” \mathbb{A}^1 –homotopy classes.

In [9, Theorem 1], building on results of M Schlichting [57, Theorems 6.15 and 6.22], we simplified and generalized F Morel’s affine representability result for vector bundles; we encourage the reader to consult the introduction of [9] for a more detailed discussion of these points. Our goal in this paper is to further extend the scope of these affine representability results in \mathbb{A}^1 –homotopy theory. For example, the following result provides a generalization of the representability result from vector bundles to torsors under suitable reductive group schemes (the description in terms of naive homotopy classes is hidden here).

Theorem 1 (see Theorem 4.1.3) *Suppose k is an infinite field, and G is an isotropic reductive k –group (see Definition 3.3.5). For every smooth affine k –scheme X , there is a bijection*

$$H_{\text{Nis}}^1(X, G) \cong [X, BG]_{\mathbb{A}^1}$$

that is functorial in X .

Remark 2 Theorem 1 is essentially the strongest possible representability statement for which one could hope. First, one cannot expect the functor “isomorphism classes of Nisnevich locally trivial G –torsors” to be representable on $\mathcal{H}(k)$ in general. Indeed, if we do not restrict attention to the category Sm_k^{aff} , then this functor need not even be \mathbb{A}^1 –invariant (see eg Ramanathan [56] for a study of failure of homotopy invariance in case $X = \mathbb{P}^1$ or the introduction to [9] for other ways in which \mathbb{A}^1 –invariance can fail). Second, at least if k is infinite and perfect, the hypothesis that G is isotropic cannot be weakened. Indeed, if G is not an isotropic reductive k –group in the sense mentioned above, then even affine representability for G –torsors fails in general; see Remark 3.3.8 and Balwe and Sawant [14, Theorem 1] for more details. We do not know if Theorem 1 holds if k is finite.

Remark 3 It has been known for some time that an analog of Morel’s theorem should hold for torsors under groups like SL_n and Sp_{2n} (for SL_n this is mentioned eg in Asok and Fasel [5, Theorem 4.2]). Schlichting observed [57, Remark 6.23] that his techniques also apply to torsors under groups like SL_n or Sp_{2n} . Combined with the results of [9], one therefore expects affine representability results to hold for torsors under such groups in the same generality as for vector bundles. For completeness, we include such results here as Theorems 4.1.1 and 4.1.2.

We also establish affine representability results for various homogeneous spaces under reductive groups.

Theorem 4 (see [Theorem 4.2.10](#)) *Suppose k is an infinite field, and G is an isotropic reductive k -group. If $P \subset G$ is a parabolic k -subgroup possessing an isotropic Levi k -subgroup, then, for any smooth affine k -scheme X , there is a bijection*

$$\pi_0(\mathrm{Sing}^{\mathbb{A}^1} G/P)(X) \xrightarrow{\sim} [X, G/P]_{\mathbb{A}^1}$$

that is functorial in X .

Remark 5 As suggested prior to the statement, we actually establish representability results with targets that are more general homogeneous spaces. In this direction, observe that it is often possible to “explicitly” identify sets of naive homotopy classes and thus, via [Theorem 4](#), true \mathbb{A}^1 -homotopy classes. Barge and Lannes [[15](#), Chapter 4] provide such identifications in the case where the target is related to symmetric bilinear forms. Cazanave [[20](#)] provides such identifications in the case where the target is \mathbb{P}^n . In addition, Fasel [[27](#), Theorem 2.1] gives such an identification in the case where the target is a Stiefel variety (various homogeneous spaces of GL_n).

Building on the ideas of Schlichting and Morel, the proofs of the results above are established using the framework developed in [[9](#)]: affine representability follows from affine Nisnevich excision and affine homotopy invariance. The restrictions on k that appear in our results are imposed to guarantee that affine homotopy invariance holds for Nisnevich locally trivial torsors under G .

While affine homotopy invariance for vector bundles is precisely the Bass–Quillen conjecture (about which much is known), precise statements regarding affine homotopy invariance for torsors under other groups are harder to find in the literature (in part because such results are typically false for étale locally trivial torsors), but see Wendt [[65](#), Section 3]. The entirety of [Section 3](#) is devoted to studying affine homotopy invariance for torsors under reductive group schemes over a rather general base.

[Theorem 1](#) is a straightforward consequence of our general representability result (see [Theorem 2.3.5](#)) combined with affine homotopy invariance (see [Theorem 3.3.7](#) for a precise statement of what we mean by this term). [Theorem 4](#) follows from [Theorem 2.4.2](#) and affine homotopy invariance for isotropic reductive k -groups by a reduction from P to a Levi factor of P (which by assumption is also an isotropic reductive k -group). Again, for certain groups, significantly more general statements can be made; see [Theorem 4.2.12](#).

Our techniques also allow us to establish significant generalizations (with simpler proofs) of some results of Morel regarding when classifying spaces for split groups are \mathbb{A}^1 -local

[43, Theorems 1.3, 1.5 and A.2]. While Morel deduces these results from strong \mathbb{A}^1 -invariance of nonstable K_1 -functors, which he establishes by appeal to classical results regarding elementary matrices, we are, in sharp contrast, able to deduce such strong \mathbb{A}^1 -invariance statements as a direct consequence of our general representability result (see [Theorem 4.3.3](#) for more details). As another sample application of these results, we adapt some classical ideas of G W Whitehead [66] to establish nilpotence results for nonstable K_1 -functors (see [Theorem 4.4.3](#)), along the lines of the results of Bak [11] and Bak, Hazrat and Vavilov [12]. In particular, we are able to resolve Problem 6 of [12] in a number of new situations (see [Remark 4.4.4](#) for more details).

The representability results for homogeneous spaces are relevant when applying the methods of obstruction theory to analyze algebraic classification problems. For example, if the base k is a perfect field, the \mathbb{A}^1 -fibration sequence

$$\mathbb{A}^n \setminus \{0\} \rightarrow BGL_{n-1} \rightarrow BGL_n$$

was used by Morel [44, Chapter 8] to develop an obstruction-theoretic approach to answering the question of when a vector bundle over a smooth affine variety splits off a trivial rank 1 summand; this approach was further developed by the first author and Fasel in [6; 7], to which we refer the interested reader for a more detailed discussion. The results of this paper (specifically [Theorem 2.2.5](#)) open the possibility of studying such questions over more general base rings, eg \mathbb{Z} .

Our representability results also broaden the scope of geometric and algebraic applications of \mathbb{A}^1 -homotopy theory. We mention a few such directions here (though we do not develop the applications). First, [Theorem 1](#) allows one to give explicit classifications of principal G -bundles on certain quadric hypersurfaces; see Asok and Fasel [5] and Asok, Doran and Fasel [4]. [Theorems 4.2.1](#) and [4.2.2](#) establish affine representability results for “split” quadric hypersurfaces. The former result has relevance to questions regarding unimodular rows (see [5]). Building on the ideas of Fasel [28], affine representability results for even dimensional quadrics are a key tool in Asok and Fasel [8] to interpret Euler class groups à la Bhatwadekar and Sridharan in terms of \mathbb{A}^1 -homotopy theory. In another direction, since the homogeneous space G_2/SL_3 is a 6-dimensional “split” smooth affine quadric, we use our results in [10] to study questions regarding reductions of structure group for “generically trivial” octonion algebras. In algebraic terms, this can be rephrased as follows: when is an octonion algebra a Zorn (“vector-matrix”) algebra (see eg Springer and Veldkamp [58, page 19])?

Dependency of sections/prerequisites Section 2 is devoted to extending our results from [9]; the proofs rely on ideas from loc. cit., which we will use rather freely together with some basic properties of torsors and homogeneous spaces collected in Sections 2.3 and 2.4. Section 3 is devoted to establishing affine homotopy invariance results for torsors under reductive groups. The results of this section rely on the basic properties of torsors and homogeneous spaces recalled in Section 2 as well as the theory of (reductive) group schemes over a base; regarding the latter: we review some of the main definitions and basic properties, but we mainly provide pointers to the literature. At the very end of Section 3.3 we also rely on the representability results from Section 2. Section 4 contains applications of our main results and thus relies on all of the preceding sections. We refer the reader to the beginning of each section for a more detailed description of its contents.

Acknowledgements The authors would like to thank Brian Conrad for extremely helpful correspondence regarding [22; 23]. In particular, the proof of Lemma 3.1.5 in its current form was a product of these discussions. The authors would also like to thank Chetan Balwe and Anand Sawant for helpful discussions of [13], Philippe Gille and Anastasia Stavrova for helpful comments and corrections on a previous version of this paper, and Marco Schlichting and Jean Fasel for mentioning at some point the Zariski local triviality of the torsor appearing in the proof of Lemma 3.1.7. Finally, this paper owes an intellectual debt to Marco Schlichting: even if it is obscured in references to [9], the ideas of [57, Section 6] served to focus our attention (for example, he established a representability result for special linear groups or symplectic groups [57, Remark 6.23]).

Asok was partially supported by National Science Foundation Award DMS-1254892. Hoyois was partially supported by National Science Foundation Award DMS-1508096. Wendt was partially supported by EPSRC grant EP/M001113/1.

Preliminaries/notation All rings considered in this paper will be assumed unital. We use the symbol S for a quasicompact, quasiseparated base scheme, Sm_S for the category of finitely presented smooth S -schemes, and $\mathrm{Sm}_S^{\mathrm{aff}} \subset \mathrm{Sm}_S$ for the full subcategory of affine schemes (in the absolute sense). We also reuse some terminology and notation introduced in [9], eg the notion of affine Nisnevich excision [9, Example 2.1.2 and Definition 3.2.1], the t -localization functor R_t [9, Section 3.1], the singular construction Sing^1 [9, Section 4.1], etc.

2 Some general representability results

The goal of this section is to extend the affine representability results of [9]. In particular, Theorem 2.2.4 provides a relative version of [9, Theorem 5.1.3]. We then specialize this result to two cases of particular interest in Theorems 2.3.5 and 2.4.2.

2.1 Naive \mathbb{A}^1 -homotopy classes

Let \mathcal{F} be a simplicial presheaf on Sm_S . Given $X \in \mathrm{Sm}_S$, there is a canonical map

$$(2-1) \quad \pi_0(\mathrm{Sing}^{\mathbb{A}^1} \mathcal{F})(X) \rightarrow [X, \mathcal{F}]_{\mathbb{A}^1},$$

where the right-hand side is the set of maps in the \mathbb{A}^1 -homotopy category $\mathcal{H}(S)$. The left-hand side is the set of naive \mathbb{A}^1 -homotopy classes of maps from X to \mathcal{F} : it is the quotient of the set of maps $X \rightarrow \mathcal{F}$ by the equivalence relation generated by \mathbb{A}^1 -homotopies. For presheaves \mathcal{F} of “geometric origin”, such as representable presheaves, it is rare that (2-1) is a bijection for all $X \in \mathrm{Sm}_S$ (this happens for example when \mathcal{F} is represented by an \mathbb{A}^1 -rigid smooth scheme in the sense of Morel and Voevodsky [45, Section 3 Example 2.4], eg a smooth curve of genus $g > 0$ or an abelian variety). However, one of the main themes of this paper is that there are many examples of presheaves \mathcal{F} such that (2-1) is a bijection for every affine X . We formalize this idea in the following definition:

Definition 2.1.1 Let \mathcal{F} be a simplicial presheaf on Sm_S and let $\widetilde{\mathcal{F}}$ be a Nisnevich-local \mathbb{A}^1 -invariant fibrant replacement of \mathcal{F} . Then there is a canonical map $\mathrm{Sing}^{\mathbb{A}^1} \mathcal{F} \rightarrow \widetilde{\mathcal{F}}$ that is well-defined up to simplicial homotopy. We will say that \mathcal{F} is \mathbb{A}^1 -naive if the map $\mathrm{Sing}^{\mathbb{A}^1} \mathcal{F}(X) \rightarrow \widetilde{\mathcal{F}}(X)$ is a weak equivalence for every $X \in \mathrm{Sm}_S^{\mathrm{aff}}$.

Remark 2.1.2 If \mathcal{F} is \mathbb{A}^1 -naive, then in particular (2-1) is a bijection for every $X \in \mathrm{Sm}_S^{\mathrm{aff}}$. More generally, if \mathcal{F} is \mathbb{A}^1 -naive and pointed, then

$$\pi_n(\mathrm{Sing}^{\mathbb{A}^1} \mathcal{F})(X) \cong [S^n \wedge X_+, \mathcal{F}]_{\mathbb{A}^1, *}$$

for every $X \in \mathrm{Sm}_S^{\mathrm{aff}}$ and $n \geq 0$.

Proposition 2.1.3 If \mathcal{F} is a simplicial presheaf on Sm_S , then \mathcal{F} is \mathbb{A}^1 -naive if and only if $\mathrm{Sing}^{\mathbb{A}^1} \mathcal{F}$ satisfies affine Nisnevich excision (see [9, Section 2.1]). In that case, $R_{\mathrm{Zar}} \mathrm{Sing}^{\mathbb{A}^1} \mathcal{F}$ is Nisnevich-local and \mathbb{A}^1 -invariant.

Proof Let $\widetilde{\mathcal{F}}$ be a Nisnevich-local \mathbb{A}^1 -invariant replacement of \mathcal{F} . Suppose that \mathcal{F} is \mathbb{A}^1 -naive. Then the restriction of $\text{Sing}^{\mathbb{A}^1} \mathcal{F}$ to Sm_S^{aff} is (objectwise) weakly equivalent to $\widetilde{\mathcal{F}}$, and hence it is Nisnevich-local. But this implies that $\text{Sing}^{\mathbb{A}^1} \mathcal{F}$ satisfies affine Nisnevich excision, by [9, Theorem 3.2.5].

Conversely, suppose that $\text{Sing}^{\mathbb{A}^1} \mathcal{F}$ satisfies affine Nisnevich excision. By [9, Theorem 3.3.4], the canonical map

$$\text{Sing}^{\mathbb{A}^1} \mathcal{F}(X) \rightarrow R_{\text{Zar}} \text{Sing}^{\mathbb{A}^1} \mathcal{F}(X)$$

is a weak equivalence for every $X \in \text{Sm}_S^{\text{aff}}$, and $R_{\text{Zar}} \text{Sing}^{\mathbb{A}^1} \mathcal{F}$ is Nisnevich-local. By [9, Lemma 5.1.2], $R_{\text{Zar}} \text{Sing}^{\mathbb{A}^1} \mathcal{F}$ is also \mathbb{A}^1 -invariant. Hence, $R_{\text{Zar}} \text{Sing}^{\mathbb{A}^1} \mathcal{F} \simeq \widetilde{\mathcal{F}}$ and \mathcal{F} is \mathbb{A}^1 -naive. □

2.2 The singular construction and homotopy fiber sequences

The notion of representable interval object was formulated in [9, Definition 4.1.1]. By a homotopy fiber sequence of pointed simplicial presheaves, we mean a homotopy Cartesian square in which either the top-right or bottom-left corner is a point.

Proposition 2.2.1 *Let \mathcal{C} be a small category and I a representable interval object in \mathcal{C} . Let*

$$\mathcal{F} \rightarrow \mathcal{G} \rightarrow \mathcal{H}$$

be a homotopy fiber sequence of pointed simplicial presheaves on \mathcal{C} . If $\pi_0(\mathcal{H})$ is I -invariant, then

$$\text{Sing}^I \mathcal{F} \rightarrow \text{Sing}^I \mathcal{G} \rightarrow \text{Sing}^I \mathcal{H}$$

is a homotopy fiber sequence.

Proof For $X \in \mathcal{C}$, consider the square of bisimplicial sets

$$\begin{array}{ccc} \mathcal{F}(X \times I^\bullet) & \longrightarrow & \mathcal{G}(X \times I^\bullet) \\ \downarrow & & \downarrow \\ * & \longrightarrow & \mathcal{H}(X \times I^\bullet) \end{array}$$

which is degreewise homotopy Cartesian. Since $\pi_0(\mathcal{H})$ is I -invariant, the simplicial set $\pi_0 \mathcal{H}(X \times I^\bullet)$ is constant. By [9, Lemma 4.2.1], the diagonal of this square is homotopy Cartesian, ie

$$\text{Sing}^I \mathcal{F}(X) \rightarrow \text{Sing}^I \mathcal{G}(X) \rightarrow \text{Sing}^I \mathcal{H}(X)$$

is a homotopy fiber sequence. □

Corollary 2.2.2 *Let C be a small category and I a representable interval object in C . If \mathcal{F} is a pointed simplicial presheaf on C such that $\pi_0(\mathcal{F})$ is I -invariant, then the canonical map*

$$\mathrm{Sing}^1 R\Omega\mathcal{F} \rightarrow R\Omega \mathrm{Sing}^1 \mathcal{F}$$

is a weak equivalence.

Proof This follows from [Proposition 2.2.1](#) applied to the homotopy fiber sequence $R\Omega(\mathcal{F}) \rightarrow * \rightarrow \mathcal{F}$. □

Lemma 2.2.3 *Suppose C is a small category with an initial object and let P be a cd structure on C . If J is a small diagram and $F: J \rightarrow \mathrm{sPre}(C)$ is a functor such that $F(j)$ satisfies P -excision for every $j \in J$, then $\mathrm{holim}_J F$ satisfies P -excision as well.*

Proof This is a straightforward consequence of commutation of homotopy limits. □

Theorem 2.2.4 *Suppose*

$$\mathcal{F} \rightarrow \mathcal{G} \rightarrow \mathcal{H}$$

is a homotopy fiber sequence of pointed simplicial presheaves on Sm_S . If

- (i) \mathcal{G} and \mathcal{H} satisfy affine Nisnevich excision, and
- (ii) $\pi_0(\mathcal{G})$ and $\pi_0(\mathcal{H})$ are \mathbb{A}^1 -invariant on affine schemes,

then \mathcal{F} is \mathbb{A}^1 -naive.

Proof By [Proposition 2.2.1](#), for every $U \in \mathrm{Sm}_S^{\mathrm{aff}}$, the sequence

$$(2-2) \quad \mathrm{Sing}^{\mathbb{A}^1} \mathcal{F}(U) \rightarrow \mathrm{Sing}^{\mathbb{A}^1} \mathcal{G}(U) \rightarrow \mathrm{Sing}^{\mathbb{A}^1} \mathcal{H}(U)$$

is a homotopy fiber sequence. By [\[9, Corollary 4.2.4\]](#), both $\mathrm{Sing}^{\mathbb{A}^1} \mathcal{G}$ and $\mathrm{Sing}^{\mathbb{A}^1} \mathcal{H}$ satisfy affine Nisnevich excision. Hence, by [Lemma 2.2.3](#), $\mathrm{Sing}^{\mathbb{A}^1} \mathcal{F}$ also satisfies affine Nisnevich excision. In other words, by [Proposition 2.1.3](#), \mathcal{F} is \mathbb{A}^1 -naive. □

The following result is not used in the sequel, but it fits the theme of this section. It is a variant of a result of Morel [\[44, Theorem 6.53\]](#) that holds over arbitrary base schemes.

Theorem 2.2.5 *Let $\mathcal{F} \rightarrow \mathcal{G} \rightarrow \mathcal{H}$ be a homotopy fiber sequence of pointed simplicial presheaves on Sm_S . Assume that:*

- (i) \mathcal{H} satisfies affine Nisnevich excision.
- (ii) $\pi_0(\mathcal{H})$ is \mathbb{A}^1 -invariant on affine schemes.

Then $\mathcal{F} \rightarrow \mathcal{G} \rightarrow \mathcal{H}$ is an \mathbb{A}^1 -fiber sequence, ie it remains a homotopy fiber sequence after taking Nisnevich-local \mathbb{A}^1 -invariant replacements.

Proof As in [Theorem 2.2.4](#), the sequence (2-2) is a homotopy fiber sequence for every $U \in \text{Sm}_S^{\text{aff}}$. Let i^* be the restriction functor from Sm_S to Sm_S^{aff} and $\mathbf{R}i_*$ its derived right adjoint. By [\[9, Lemma 3.3.2\]](#), there is a natural equivalence of functors $R_{\text{Zar}} \simeq \mathbf{R}i_* R_{\text{Zar}} i^*$. Since $\mathbf{R}i_*$ and R_{Zar} preserve homotopy fiber sequences, we deduce that

$$R_{\text{Zar}} \text{Sing}^{\mathbb{A}^1} \mathcal{F} \rightarrow R_{\text{Zar}} \text{Sing}^{\mathbb{A}^1} \mathcal{G} \rightarrow R_{\text{Zar}} \text{Sing}^{\mathbb{A}^1} \mathcal{H}$$

is a homotopy fiber sequence. By [\[9, Theorem 5.1.3\]](#), $R_{\text{Zar}} \text{Sing}^{\mathbb{A}^1} \mathcal{H}$ is Nisnevich-local and \mathbb{A}^1 -invariant. But it follows from the right properness of the Morel–Voevodsky model structure [\[45, Section 2, Theorem 2.7\]](#) that every homotopy fiber sequence whose base is Nisnevich-local and \mathbb{A}^1 -invariant is an \mathbb{A}^1 -fiber sequence. \square

2.3 Application to torsors

In this subsection we specialize the general representability result of [\[9, Section 5.1\]](#) to simplicial presheaves classifying G -torsors for some group G . We start by recalling some general facts about torsors.

Definition 2.3.1 Let \mathcal{C} be a small category equipped with a Grothendieck topology t , let G be a t -sheaf of groups on \mathcal{C} , and let $X \in \mathcal{C}$. A G -torsor over X is a triple (\mathcal{P}, π, a) , where \mathcal{P} is a t -sheaf on \mathcal{C} , $a: \mathcal{P} \times G \rightarrow \mathcal{P}$ is a right action of G on \mathcal{P} and $\pi: \mathcal{P} \rightarrow X$ is a morphism that is G -equivariant for the trivial G -action on X such that:

- (i) The morphism $\mathcal{P} \times G \rightarrow \mathcal{P} \times_X \mathcal{P}$ of components π_1 and a is an isomorphism.
- (ii) π is t -locally split, ie the collection of morphisms $U \rightarrow X$ in \mathcal{C} such that $\mathcal{P} \times_X U \rightarrow U$ has a section is a t -covering sieve of X .

The collection of G -torsors over various $X \in \mathcal{C}$ can be assembled into a category $\mathbf{Tors}_t(G)$ fibered in groupoids over \mathcal{C} . We write $\mathbf{BTors}_t(G)$ for the simplicial presheaf whose value on $U \in \mathcal{C}$ is the nerve of the groupoid of sections of $\mathbf{Tors}_t(G)$ over \mathcal{C}/U (this groupoid is canonically equivalent to the groupoid of G -torsors over U , but is

strictly functorial in U ; see Hollander [34, Section 3.3]). It is well known that $\mathbf{Tors}_t(G)$ is a stack for the topology t . As shown in [34, Theorem 3.9], this is equivalent to the statement that $B\mathbf{Tors}_t(G)$ satisfies t -descent.

We denote by BG the pointed simplicial presheaf with n -simplices G^n and with the usual face and degeneracy maps, and we let

$$B_tG := R_tBG$$

be its t -local replacement (see [9, Section 3]). There is a morphism $BG \rightarrow B\mathbf{Tors}_t(G)$ sending the unique vertex of $BG(U)$ to the trivial G -torsor over U . Since $B\mathbf{Tors}_t(G)$ is t -local, we obtain a morphism of simplicial presheaves

$$(2-3) \quad B_tG \rightarrow B\mathbf{Tors}_t(G).$$

Lemma 2.3.2 *Let \mathcal{C} be a small category, t a Grothendieck topology on \mathcal{C} , and G a t -sheaf of groups on \mathcal{C} . Then:*

- (i) *The map (2-3) is a weak equivalence of simplicial presheaves.*
- (ii) *There is a natural isomorphism*

$$\pi_0(B_tG)(-) \cong H_t^1(-, G).$$

- (iii) *There is a canonical weak equivalence $R\Omega B_tG \simeq G$.*

Proof It is clear that the map (2-3) induces an isomorphism on t -sheaves of homotopy groups, so that it is a weak equivalence in the Jardine model structure. To deduce that it is a weak equivalence, it therefore suffices to show that the source and target are fibrant in the Jardine model structure. By Dugger, Hollander and Isaksen [26, Corollary A.8], it suffices to show that, for every $U \in \mathcal{C}$, the simplicial sets $B_tG(U)$ and $B\mathbf{Tors}_t(G)(U)$ have no homotopy in dimensions ≥ 2 . This statement is clear for the latter as it is the nerve of a groupoid. To treat the former case, we recall a fact from simplicial homotopy theory: if X is a simplicial set, then X has no homotopy in dimensions $\geq k$ if and only if the homotopy fibers of the diagonal map $X \rightarrow X \times^h X$ have no homotopy in dimensions $\geq k - 1$; this can be checked by assuming X is a Kan complex and studying homotopy groups. Thus, a simplicial set X has no homotopy in dimensions ≥ 2 if and only if its 3-fold diagonal

$$X \rightarrow X \times_{X \times_{X \times^h X}^h X}^h X$$

is a weak equivalence. Since R_t preserves homotopy pullbacks, it also preserves the property of having no homotopy in dimensions ≥ 2 . This proves (i). Assertions (ii) and (iii) are true essentially by definition if we replace $B_t G$ by $B\text{Tors}_t(G)$, so they both follow from (i). \square

Torsors under S -group schemes Our main interest is to representability results for torsors under group schemes, so we now discuss that situation in greater detail. Let G be an S -group scheme and let X be an S -scheme. By a G -torsor over X we will mean a G -torsor in the sense of Definition 2.3.1, for \mathcal{C} the category of S -schemes and t the fppf topology. In the sequel, G will always be affine over S , and in that case a G -torsor over X is automatically representable by an S -scheme, by Milne [41, Theorem III.4.3(a)] (note the implicit Noetherian hypothesis in Milne’s argument is unnecessary).

If moreover X and G belong to Sm_S , then taking \mathcal{C} to be the category Sm_S with t the étale topology one obtains an equivalent notion of torsor. Indeed, if $\pi: \mathcal{P} \rightarrow X$ is a G -torsor over X , then π is finitely presented and smooth by the following lemma. Since smooth morphisms admit sections étale locally, π itself is a cover of X in the étale topology which trivializes the torsor.

Lemma 2.3.3 *Suppose G is an affine S -group scheme, X is an S -scheme and $\pi: \mathcal{P} \rightarrow X$ is a G -torsor over X . If $G \rightarrow S$ is finitely presented, flat or smooth, then so is $\pi: \mathcal{P} \rightarrow X$.*

Proof By definition, there exists an fppf cover $\{U_i \rightarrow X\}_{i \in I}$ such that $\mathcal{P} \times_X U_i \rightarrow U_i$ is isomorphic to $G \times_S U_i \rightarrow U_i$, which is finitely presented, flat or smooth. We conclude using the fact that each of these properties of a morphism is fppf-local on the target, by [17, Tags 02L0, 02L2 and 02VL]. \square

Example 2.3.4 Let t be a topology on Sm_S in between the Zariski topology and the étale topology and let $n \geq 1$. The groupoid of GL_n -torsors over a scheme is canonically equivalent to the groupoid of rank n vector bundles. Since GL_n is a smooth special group, any GL_n -torsor is t -locally trivial. In particular, by Lemma 2.3.2(ii), we have

$$\pi_0(B_t \text{GL}_n)(X) \cong \mathcal{V}_n(X)$$

for any $X \in \text{Sm}_S$, where $\mathcal{V}_n(X)$ denotes the set of isomorphism classes of rank n vector bundles on X . Similarly, we have

$$\pi_0(B_t \text{SL}_n)(X) \cong \mathcal{V}_n^o(X) \quad \text{and} \quad \pi_0(B_t \text{Sp}_{2n})(X) \cong \mathcal{H}\mathcal{V}_{2n}(X),$$

where $\mathcal{V}_n^o(X)$ (resp. $\mathcal{H}\mathcal{V}_{2n}(X)$) is the set of isomorphism classes of rank n oriented (resp. rank $2n$ symplectic) vector bundles (see the beginning of Section 3.3 for reminders about oriented and symplectic vector bundles).

Affine representability for Nisnevich locally trivial G -torsors

Theorem 2.3.5 *Suppose G is a finitely presented, smooth S -group scheme. If $H_{\text{Nis}}^1(-, G)$ is \mathbb{A}^1 -invariant on Sm_S^{aff} , then:*

- (i) *The simplicial presheaf $R_{\text{Zar}} \text{Sing}^{\mathbb{A}^1} B_{\text{Nis}} G$ is Nisnevich-local and \mathbb{A}^1 -invariant.*
- (ii) *For every affine $X \in \text{Sm}_S^{\text{aff}}$, the canonical map*

$$H_{\text{Nis}}^1(X, G) \rightarrow [X, BG]_{\mathbb{A}^1}$$

is a bijection that is functorial with respect to X .

Proof Since $B_{\text{Nis}} G$ is Nisnevich-local by definition, it satisfies Nisnevich excision by [9, Theorem 3.2.5]. Taking into account the identification $\pi_0(B_{\text{Nis}} G) \cong H_{\text{Nis}}^1(-, G)$ from Lemma 2.3.2(ii), we can apply [9, Theorem 5.1.3] to $B_{\text{Nis}} G$, which implies (i) and (ii) (note also that $[X, B_{\text{Nis}} G]_{\mathbb{A}^1} \cong [X, BG]_{\mathbb{A}^1}$ since $BG \rightarrow B_{\text{Nis}} G$ is a Nisnevich-local equivalence). □

2.4 Application to homogeneous spaces

Let \mathcal{C} be a small category equipped with a Grothendieck topology t . Let G and H be t -sheaves of groups on \mathcal{C} with $H \subset G$. We then have a homotopy fiber sequence of simplicial presheaves

$$G/H \rightarrow BH \rightarrow BG,$$

where G/H denotes the presheaf $U \mapsto G(U)/H(U)$. Applying the t -localization functor R_t , we obtain a homotopy fiber sequence of t -local simplicial presheaves

(2-4)
$$a_t(G/H) \rightarrow B_t H \rightarrow B_t G.$$

We now restrict attention to $\mathcal{C} = \text{Sm}_S$ with the goal of applying Theorem 2.2.4. For geometric applications, we need to better understand the sheaf $a_t(G/H)$.

Homogeneous spaces: topologies and quotient sheaves Write rX for the presheaf on the category of S -schemes represented by an S -scheme X and $r'X$ for the restriction of the presheaf rX to Sm_S . Suppose that G and H are finitely presented,

smooth S -group schemes, and that H is a closed subgroup of G . The right translation action of H on G is scheme-theoretically free and it follows from a result of Artin [2, Corollary 6.3] that the sheaf $a_{\text{fppf}}(rG/rH)$ is representable by an S -algebraic space. Two questions naturally present themselves: first, when does the fppf sheaf quotient coincide with the Zariski or Nisnevich sheaf quotient, and second, is the fppf sheaf $a_{\text{fppf}}(rG/rH)$ representable by a smooth scheme? We address the first question here; we answer the second question in various cases in Section 3.1.

Lemma 2.4.1 *Suppose G is a finitely presented S -group scheme and $H \subset G$ is a finitely presented, closed S -subgroup scheme. Assume that H is flat over S and that the quotient G/H exists as an S -scheme. Then $G \rightarrow G/H$ is an H -torsor, and the following statements hold:*

- (i) *If t is a subcanonical topology on S -schemes such that the map $G \rightarrow G/H$ is t -locally split, then $r(G/H) \cong a_t(rG/rH)$.*
- (ii) *If G is smooth over S , then G/H is smooth over S . Moreover, if t is a subcanonical topology on Sm_S such that the map $G \rightarrow G/H$ is t -locally split, then $r'(G/H) \cong a_t(r'G/r'H)$.*

Proof By a theorem of Anantharaman [1, Appendice I, Théorème 6], we have $r(G/H) \cong a_{\text{fppf}}(rG/rH)$. In particular, $G \rightarrow G/H$ is an H -torsor, and hence it is flat by Lemma 2.3.3. If G is smooth, it follows from [33, Proposition 17.7.7] that G/H is also smooth. If $G \rightarrow G/H$ is t -locally split, then $rG \rightarrow r(G/H)$ is an epimorphism of t -sheaves. By [3, Proposition 4.3(2)], this implies that $r(G/H)$ is the coequalizer of the equivalence relation $rG \times_{r(G/H)} rG \cong rG \times rH \rightrightarrows rG$ in the category of t -sheaves, which exactly means that $r(G/H) \cong a_t(rG/rH)$. The second statement is proved in the same way. \square

Affine representability for homogeneous spaces

Theorem 2.4.2 *Suppose G is a finitely presented, smooth S -group scheme and $H \subset G$ is a finitely presented, smooth, closed S -subgroup scheme such that the quotient G/H exists as an S -scheme. Suppose that $G \rightarrow G/H$ is Nisnevich locally split and that $H_{\text{Nis}}^1(-, G)$ and $H_{\text{Nis}}^1(-, H)$ are \mathbb{A}^1 -invariant on Sm_S^{aff} . Then G/H is \mathbb{A}^1 -naive. In particular, for every $X \in \text{Sm}_S^{\text{aff}}$, there is a bijection*

$$\pi_0(\text{Sing}^{\mathbb{A}^1} G/H)(X) \cong [X, G/H]_{\mathbb{A}^1}.$$

Proof The assumption on $G \rightarrow G/H$ combined with [Lemma 2.4.1](#) allow us to conclude that $r'(G/H) \cong a_{\text{Nis}}(r'G/r'H)$ and thus the homotopy fiber sequence (2-4) has the form $r'(G/H) \rightarrow B_{\text{Nis}}H \rightarrow B_{\text{Nis}}G$. The simplicial presheaves $B_{\text{Nis}}G$ and $B_{\text{Nis}}H$ are Nisnevich-local and hence satisfy Nisnevich excision by [[9](#), Theorem 3.2.5]. The result is now a direct application of [Theorem 2.2.4](#), taking into account [Lemma 2.3.2\(ii\)](#). \square

3 Homotopy invariance for torsors under group schemes

The main goal of this section is to study \mathbb{A}^1 -invariance of the functors $H_{\text{Nis}}^1(-, G)$ for G a linear group. [Section 3.1](#) reviews basic definitions about group schemes, torsors and homogeneous spaces; it also collects a number of results that will be used later in the text. [Section 3.2](#) establishes an analog of the local-to-global principle (also called “Quillen patching”) for torsors under linear group schemes under rather general hypotheses; the main result is [Theorem 3.2.5](#). Finally, [Section 3.3](#) proves general homotopy invariance results; the main results are [Theorems 3.3.3](#) and [3.3.7](#). For simplicity, we assume throughout this section that the base scheme S is the spectrum of a commutative ring R . In general there is a tradeoff between generality of the group G under consideration and the base ring R .

3.1 Reductive group schemes and homogeneous spaces: recollections

The goal of this section is to recall some basic definitions and properties of group schemes, torsors and homogeneous spaces over rather general bases. Rather than attempting to be exhaustive, we only aim to point the reader to places in the literature where they can find the required results. The grouping of these results is slightly eclectic: only a very small portion of the definitions and results established here will be used in the remainder of [Section 3](#). Many of the results we state here are significantly easier to establish (or even unnecessary) if the base ring R is a field.

Linear and reductive group schemes We write $\text{GL}_{n,R}$ for the general linear group scheme over R and $\mathbf{G}_{m,R}$ for $\text{GL}_{1,R}$. If R is clear from context, we will drop it from the notation.

Definition 3.1.1 By a *linear R -group scheme*, we mean a group scheme G over R admitting a finitely presented closed immersion group homomorphism $G \rightarrow \text{GL}_{n,R}$.

Later, the homotopy invariance results we establish will require much more stringent hypotheses on G . We use the definition of *reductive* (resp. *semisimple*) R -group scheme of Demazure and Grothendieck [25, Exposé XIX, Définition 2.7]: a reductive (resp. *semisimple*) R -group scheme is a smooth, affine R -group scheme with geometric fibers that are connected reductive (resp. *semisimple*) groups in the usual sense [25, Exposé XIX, 1.6], ie have trivial unipotent radical (resp. radical). Recall that a reductive R -group scheme G is called *split* if it contains a split maximal torus [25, Exposé XXII, Définition 1.13]. Any split reductive group scheme is pulled back from a unique “Chevalley” group scheme over $\text{Spec } \mathbb{Z}$.

If R is a field, it is a well-known consequence of the classification of reductive groups that reductive R -group schemes are linear R -group schemes. If R is no longer a field, the connection between “reductive” and “linear” becomes more complicated, as the following example demonstrates.

Example 3.1.2 Groups of multiplicative type need not be linear in general [24, Exposé IX, Définition 1.1]. Indeed, [24, Exposé XI, Remarque 4.6] explains that if R is a Noetherian and connected ring, then a group G of multiplicative type admits an embedding in GL_n if and only if it is isotrivial.

Nevertheless, the following result shows that, assuming suitable hypotheses on the base, reductive R -group schemes are always linear.

Proposition 3.1.3 (Thomason) *Suppose G is a reductive R -group scheme. Assume, in addition, either*

- (i) R is regular and Noetherian, or
- (ii) G is split.

Then G is a linear R -group scheme.

Proof If G is split, we can assume that $R = \mathbb{Z}$ and in particular that R is regular Noetherian. In that case, the result follows from Thomason [61, Corollary 3.2(3)]. \square

Remark 3.1.4 Thomason actually gives a sufficient condition for a group scheme to admit a closed immersion group homomorphism into the automorphism group scheme of a vector bundle over an arbitrary base S [61, Theorem 3.1]. Since we have in mind applications to homotopy invariance, we have restricted attention to spectra of regular rings.

Homogeneous spaces for reductive groups Suppose G is a reductive R -group scheme and $\lambda: \mathbf{G}_m \rightarrow G$ is a homomorphism of R -group schemes. Via λ , we may consider the \mathbf{G}_m -action $\lambda: \mathbf{G}_m \times G \rightarrow G$ defined pointwise by the formula $\lambda(t, g) := \lambda(t)g\lambda(t)^{-1}$. We can define a subfunctor $P_G(\lambda) \subset G$ consisting of those points $g \in G$ such that $\lim_{t \rightarrow 0} \lambda(t, g)$ exists and a subfunctor $U_G(\lambda) \subset G$ consisting of those points $g \in G$ such that $\lim_{t \rightarrow 0} \lambda(t, g) = 1$ (see Conrad [22, Theorem 4.1.7] for precise definitions). By [22, Theorem 4.1.7], both of these functors are representable by R -subgroup schemes of G ; since we assumed G reductive, it follows also that $P_G(\lambda)$ and $U_G(\lambda)$ are smooth and connected. By [22, Example 5.2.2], $P_G(\lambda)$ is parabolic, and $U_G(\lambda)$ is a closed normal R -subgroup scheme whose geometric fibers correspond to unipotent radicals of the geometric fibers of $P_G(\lambda)$ [22, Corollary 5.2.5]; we will abuse terminology and refer to $U_G(\lambda)$ as the unipotent radical of $P_G(\lambda)$.

If $Z_G(\lambda)$ is the centralizer of λ , then, by [22, Definition 5.4.2] and the subsequent discussion, $Z_G(\lambda)$ is a Levi factor of $P_G(\lambda)$, ie $Z_G(\lambda)$ is a smooth reductive R -group scheme, and there is a semidirect product decomposition of the form $Z_G(\lambda) \ltimes U_G(\lambda) \cong P_G(\lambda)$. This description of parabolics, their unipotent radicals and Levi factors is called a “dynamic” description in [23; 22] (since it arises from a study of “flows” under an action of \mathbf{G}_m). We use these ideas to establish the following result:

Lemma 3.1.5 *Suppose R is a connected ring, G is a reductive R -group scheme, $P \subset G$ is a parabolic R -subgroup scheme and L is a Levi factor of P . Then:*

- (i) *The quotients G/L and G/P exist as smooth R -schemes.*
- (ii) *The morphism $G \rightarrow G/L$ is a generically trivial L -torsor.*
- (iii) *The morphism $G/L \rightarrow G/P$ is a composition of torsors under vector bundles.*

Proof For later use, we observe that since R is assumed connected and L is presumed to exist, by P Gille [31, Théorème 9.3.1] there is a cocharacter $\lambda: \mathbf{G}_m \rightarrow G$ such that $P = P_G(\lambda)$ and $L = Z_G(\lambda)$. If S is the spectrum of a field, which is the case we will use later, the fact that all pairs (P, L) consisting of a parabolic together with a Levi factor are of the form $(P_G(\lambda), Z_G(\lambda))$ for a suitable cocharacter λ is contained in [23, Proposition 2.2.9].

For (i), begin by observing that, since P is a parabolic subgroup of G , it is a self-normalizing subgroup [22, Corollary 5.2.8]. The quotients G/L and G/P exist as smooth R -schemes by [22, Theorems 2.3.1 and 2.3.6] (and, by Lemma 2.4.1, the morphisms $G \rightarrow G/L$ and $G \rightarrow G/P$ are an L -torsor and a P -torsor, respectively).

For (ii), set $U^- = U_G(-\lambda)$, ie the “unipotent radical” of an opposite parabolic. We know that there is a dense open subscheme of G isomorphic to $U^- \times P$ [22, Theorem 4.1.7] (here and below, we will refer to this as the “big cell”). The image of this open subscheme in G/L , which is isomorphic to $U^- \times P/L$, is again open and dense since $G \rightarrow G/L$ is smooth and surjective. The Levi decomposition yields an isomorphism of schemes $P \cong L \times U$, and thus an identification $P/L \cong U$. Under these identifications, the unit map $U \rightarrow P$ provides a morphism $U^- \times U \rightarrow U^- \times L \times U$, which yields the required generic trivialization.

For (iii), let U be the unique smooth closed normal R -subgroup scheme of P whose geometric fibers coincide with the unipotent radicals of the geometric fibers of P , which is guaranteed to exist by [22, Corollary 5.2.5]. By the uniqueness assertion, $U \cong U_G(\lambda)$ for the character whose existence we observed in the first paragraph. By [22, Theorem 5.4.3], U admits a finite descending filtration by $\text{Aut}_{P/R}$ -stable, closed, normal, smooth R -subgroup schemes U_i with successive subquotients U_i/U_{i+1} isomorphic to P -equivariant vector bundles over R . Moreover, the isomorphism $P/L \cong U$ described in (ii) is actually P -equivariant.

Now, the morphism $G/L \rightarrow G/P$ is G -equivariant by definition. The scheme-theoretic fiber over the identity coset in G/P is isomorphic to the quotient P/L and there is an induced G -equivariant isomorphism $G \times^P P/L \xrightarrow{\sim} G/L$ under which the morphism $G/L \rightarrow G/P$ is sent to the projection onto the first factor. In particular, since $P/L \cong U$ is smooth, $G \times^P P/L \rightarrow G$ is smooth and since smoothness is fppf local on the base [17, Tag 02VL], we conclude that $G/L \rightarrow G/P$ is also smooth. By the discussion of the previous paragraph, the morphism $G/L \rightarrow G/P$ thus factors successively through morphisms of the form

$$(3-1) \quad G \times^P U/U_{i+1} \rightarrow G \times^P U/U_i.$$

To finish the proof, it suffices to inductively establish that each morphism in (3-1) is a torsor under a vector bundle.

Each morphism $U/U_{i+1} \rightarrow U/U_i$ is, by construction, a torsor under the vector bundle U_i/U_{i+1} and, as we observed above, provided with a P -equivariant structure. If \mathcal{E} is a quasicoherent sheaf on a scheme X , then $H_{\text{fppf}}^1(X, \mathcal{E}) = H_{\text{Zar}}^1(X, \mathcal{E})$ by [17, Tag 03DR, Proposition 34.8.10]. Since $H_{\text{fppf}}^1(X, \mathcal{E})$ parametrizes fppf torsors under the quasicoherent sheaf \mathcal{E} , the P -equivariant structure on U_i/U_{i+1} allows us to conclude, by fppf descent, that $G \times^P U_i/U_{i+1}$ is a torsor under a vector bundle on G/P . In other words, each morphism in (3-1) is again a torsor under the vector bundle U_i/U_{i+1} . \square

Remarks 3.1.6 (1) Since R a connected ring, it is not necessary to assume in the statement above that L exists; this follows from Conrad [22, Corollary 5.4.8]. If we were to work over a nonaffine base scheme, parabolics need not have Levi factors (see [22, Example 5.4.9] for more details). By reorganizing the proof, the argument presented in (iii) actually shows that the quotient G/L exists assuming we know G/P to exist and the relevant results on the structure of U .

(2) By Lemma 2.3.3, since L is a smooth R -group scheme by assumption, $G \rightarrow G/L$ is étale locally trivial. If R is Noetherian and regular, then the morphism $G \rightarrow G/L$ being generically trivial is tantamount to $G \rightarrow G/L$ being Nisnevich locally trivial. To prove this, it suffices to show that generically trivial L -torsors over Henselian local rings are trivial. If G is split reductive, then L is as well, and the asserted triviality follows from Białynicki-Birula [18, Proposition 2]. If G is not necessarily split, then L can be an arbitrary reductive group and one can appeal to Nisnevich [48, Théorème 4.5] to deduce the required triviality result (Nisnevich makes a statement for semisimple group schemes, but it is true more generally; see Fedorov and Panin [29, Section 1.1]).

(3) If G is split, it is possible to use translation of the big cell by elements of the Weyl group to produce an explicit Zariski local trivialization of $G \rightarrow G/L$. In fact, even if G is not split, to establish Zariski local triviality of $G \rightarrow G/L$ (or, equivalently, $G \rightarrow G/P$), it suffices to know that the $G(R)$ -translates of the big cell form an open cover of G/L (or G/P). If R is an infinite field, this kind of result follows from the fact that the image of $G(R)$ in $G/P(R)$ is Zariski dense (via the unirationality of G).

(4) In contrast, if R is a finite field (and G is nonsplit), it is a priori not obvious that $G(R)$ translates of the big cell cover G/L (or G/P). Nevertheless, assuming the Grothendieck–Serre conjecture, one knows that $G \rightarrow G/L$ is Zariski locally trivial. If R is the spectrum of a finite field, the Grothendieck–Serre conjecture was established by Gabber for reductive groups coming from the ground field (unpublished), but another proof of a more general case was recently given by Panin [49] (see also [29]).

Write SO_n for the split special orthogonal group over R . We restrict attention to the case where 2 is a unit in R , so we can view SO_n as the R -subgroup scheme of GL_n consisting of automorphisms of the standard hyperbolic form q_n with trivial determinant (see eg Conrad [22, Definition C.1.2]); for more details on special orthogonal groups, see [22, Appendix C].

Lemma 3.1.7 *If R is a ring in which 2 is invertible, then:*

- (i) If $n \geq 3$, the quotient $\mathrm{SO}_n/\mathrm{SO}_{n-1}$ exists and is isomorphic to a quadric hypersurface in \mathbb{A}_R^n defined by the equation $q_n = 1$.
- (ii) If $n \geq 3$, the projection morphism $\mathrm{SO}_n \rightarrow \mathrm{SO}_n/\mathrm{SO}_{n-1}$ makes SO_n into a Zariski locally trivial SO_{n-1} -torsor over the quotient.

Proof Without loss of generality, we can take $R = \mathbb{Z}[\frac{1}{2}]$, which is Noetherian of dimension ≤ 1 . Since SO_{n-1} is a closed R -subgroup scheme of SO_n , the quotient $\mathrm{SO}_n/\mathrm{SO}_{n-1}$ exists as a scheme [1, Théorème 4.C].

To identify this quotient with the quadric in the statement, we proceed as follows: Since $\mathrm{SO}_{n-1} = \mathrm{SO}_n \cap \mathrm{SL}_{n-1}$ inside of SL_n , the inclusion $\mathrm{SO}_n \subset \mathrm{SL}_n$ induces a monomorphism $\mathrm{SO}_n/\mathrm{SO}_{n-1} \hookrightarrow \mathrm{SL}_n/\mathrm{SL}_{n-1}$. Note that if A is an R -algebra, the map sending $X \in \mathrm{SL}_n(A)$ to its first row and the first column of its inverse determines an isomorphism $\mathrm{SL}_n/\mathrm{SL}_{n-1} \cong \mathrm{Spec} R[x_1, \dots, x_{2n}]/(q_{2n} - 1)$. If we restrict $X \in \mathrm{SO}_n(A)$ and if J is the symmetric matrix corresponding to the bilinear form associated with q_n , then the orthogonality condition imposes the relation $X^{-1} = JX^T$. Using this observation, it is straightforward to check that the image is isomorphic, in suitable coordinates, to a subquadric given by the equation $q_n = 1$.

For the second statement, observe that morphisms $X \rightarrow \mathrm{SO}_n/\mathrm{SO}_{n-1}$ classify SO_{n-1} -torsors which are trivial after stabilization to SO_n -torsors. The Witt cancellation theorem (see Milnor and Husemoller [42, Lemma 6.3]) implies that, over a local ring in which 2 is invertible, such an SO_{n-1} -torsor is already trivial. \square

3.2 The local-to-global principle for torsors under linear group schemes

In this section we establish a local-to-global principle or “Quillen patching” for torsors under linear R -group schemes in the sense of Definition 3.1.1. The main result of this section is Theorem 3.2.5, which is a multivariable analog of a result of Quillen [54, Theorem 1] along the lines of Lam [39, Theorem V.1.6]. As will be clear from the presentation, the argument follows quite closely that for projective modules given in [39, Chapter V.1].

That the local-to-global principle holds for torsors under linear group schemes is certainly “well known to experts”, under suitable hypotheses. For example, Raghunathan [55] states (without proof) that Quillen’s local-to-global principle holds for linear algebraic groups over a field, and Bass, Connell and Wright developed an axiomatic method to establish such results [16, Proposition 3.1]; in particular, the latter approach

applies for various classical groups [16, Remark 4.15.4] over a general base ring. Nevertheless, since we could not find a suitable published reference for precisely what we needed, in the interest of completeness, we decided to collect the necessary results here.

Modifying automorphisms We begin by generalizing [54, Lemma 1] (also Corollary V.1.2 of [39]) and [39, Corollary V.1.3] to linear R -group schemes over an arbitrary commutative ring R . The following pair of results are due to Moser [47, Lemmas 3.5.3–3.5.5] (though our hypotheses differ slightly); we include them here for the convenience of the reader.

Lemma 3.2.1 *Let R be a commutative ring, let G be a linear R -group scheme, let $f \in R$ and let $\theta(t) \in G(R_f[t])$ be such that $\theta(0) = 1 \in G(R_f)$. There exists an integer $s \geq 0$ such that, for any $a, b \in R$ with $a - b \in f^s R$, there exists $\psi \in G(R[t])$ with $\psi(0) = 1$ and such that $\psi_f(t) = \theta(at)\theta(bt)^{-1} \in G(R_f[t])$.*

Proof Since G is a linear R -group scheme, by definition there is a finitely presented closed immersion $G \rightarrow \mathrm{GL}_n$. For $s \in \mathbb{N}$, set $\psi_s(t, x, y) := \theta((x + f^s y)t)\theta(xt)^{-1} \in G(R_f[t, x, y])$. It suffices to show that there exists s such that ψ_s can be lifted to an element $\psi_s \in G(R[t, x, y])$. Indeed, in that case, by specializing with $x = b$, $a = b + f^s \alpha$, we see that $\theta(at)\theta(bt)^{-1} = \psi_s(t, b, \alpha)$ lifts as well. By the proof of [54, Lemma 1], we know that there exists s such that $\psi_s(t, x, y)$ lifts to an element of $\mathrm{GL}_n(R[t, x, y])$ and such that $\psi_s(0, x, y) = 1$ (see also [39, Theorem V.1.1]). Observe that, by definition, $\psi_s(t, x, 0) = 1$ and thus $\psi_s(t, x, 0) \in G(R[x, t])$.

It remains to show that there exists $i \geq 0$ such that $\psi_s(t, x, f^i y) \in G(R[t, x, y])$. We first recast this in ring-theoretic terms. Set $A := R[t, x]$, let B be the coordinate ring of GL_n , and let $I \subset B$ be the finitely generated ideal defining G . The lift of ψ_s is given by a homomorphism $\varphi: B \rightarrow A[y]$, and we want to show that, for some $i \geq 0$, $\varphi(-)(f^i y)$ vanishes on I . We claim that, for every $r \in I$, there exists an integer i_r such that $\varphi(r)(f^i y) = 0$ for $i \geq i_r$. If $J \subset I$ is a finite generating set and $i = \max_{r \in J} i_r$, then i will have the desired property.

Note that φ has the following properties: if $\mathrm{ev}_0: A[y] \rightarrow A$ is the evaluation homomorphism, then the composites $\mathrm{ev}_0 \circ \varphi: B \rightarrow A$ and $B \rightarrow A[y] \rightarrow A_f[y]$ both vanish on I . If $r \in I$ and $P := \varphi(r) \in A[y]$, these properties imply that $P = yQ$ for some $Q \in A[y]$ and that $f^{i_r} P = 0$ for some $i_r \geq 0$. Combining these two observations, we have $0 = f^{i_r} P = f^{i_r} yQ$. Therefore, $f^{i_r} Q = 0$ as well. Thus, $P(f^i y) = f^i yQ(f^i y) = 0$ for all $i \geq i_r$, which is what we wanted to show. \square

Lemma 3.2.2 *Let R be a commutative ring and G a linear R -group scheme. Given $f_0, f_1 \in R$ such that $f_0R + f_1R = R$, and $\theta \in G(R_{f_0f_1}[t])$ with $\theta(0) = 1$, we can find $\tau_i \in G(R_{f_i}[t])$ with $\tau_i(0) = 1$ such that $\theta = \tau_0\tau_1^{-1}$.*

Proof Let $\theta(t) \in G(R_{f_0f_1}[t])$. We can apply Lemma 3.2.1 to the localizations $R_{f_0} \rightarrow R_{f_0f_1}$ and $R_{f_1} \rightarrow R_{f_0f_1}$; pick an integer s that suffices for both localizations. For any $b \in R$, we can write

$$\theta(t) = [\theta(t)\theta(bt)^{-1}]\theta(bt).$$

If $f_0R + f_1R = R$, then the same thing is true for f_0^s and f_1^s . Thus, we can pick $b \in f_1^sR$ such that $1 - b \in f_0^sR$. In that case, $\theta(t)\theta(bt)^{-1} \in G(R_{f_1}[t])_{f_0}$ and $\theta(bt) \in G(R_{f_0}[t])_{f_1}$ lift to elements τ_1 and τ_0 with the stated properties. \square

Remark 3.2.3 Lemma 3.2.1 implies ‘‘Axiom Q’’ (in the sense of Bass, Connell and Wright [16, Section 1.1]) holds for the functor on R -algebras determined by G . Lemma 3.2.2 essentially corresponds to [16, Theorem 2.4].

The local-to-global principle Let R be a commutative ring and suppose G is a linear R -group scheme. If A is a commutative R -algebra, by a G -torsor over A we will mean a G -torsor over $\text{Spec } A$; by assumption our G -torsors are locally trivial in the fppf topology (see Definition 2.3.1 and the discussion just prior to Lemma 2.3.3 for more details). A G -torsor over $A[t_1, \dots, t_n]$ that is pulled back from a G -torsor over A will be called *extended from A* . For the remainder of this section, we will essentially confine our attention to a *fixed* G -torsor \mathcal{P} , which will be important for subsequent applications.

Proposition 3.2.4 *Let R be a commutative ring. If \mathcal{P} is a G -torsor over $R[t]$, then the set $Q(\mathcal{P})$ consisting of $g \in R$ such that $\mathcal{P}|_{\text{Spec } R_g[t]}$ is extended from R_g is an ideal in R .*

Proof It is immediate that $Q(\mathcal{P})$ is closed under multiplication by elements in R . Thus, we have to show that if $f_0, f_1 \in Q(\mathcal{P})$, then $f = f_0 + f_1$ lies in $Q(\mathcal{P})$ as well. After replacing R by R_f , we can assume that $f_0R + f_1R = R$.

Write $0: \text{Spec } R \rightarrow \mathbb{A}^1_R$, and $\text{pr}: \mathbb{A}^1_R \rightarrow \text{Spec } R$ for the zero section and the structure morphism. Thus, suppose \mathcal{P} is a G -torsor over $R[t]$ and assume that the restrictions $\mathcal{P}_i := \mathcal{P}|_{\text{Spec } R_{f_i}[t]}$ are extended. We want to show that $\mathcal{P} \cong \text{pr}^*0^*\mathcal{P}$.

By assumption, there are isomorphisms $u_i: \mathcal{P}_i \cong \text{pr}^*0^* \mathcal{P}_i$ over $R_{f_i}[t]$. By modifying u_i if necessary, we may assume that $0^*u_i = 1$. Let \mathcal{P}_{01} be the restriction of \mathcal{P} to $R_{f_0 f_1}[t]$. Then u_0 and u_1 restrict to give two isomorphisms $(u_0)_{f_1}, (u_1)_{f_0}: \mathcal{P}_{01} \cong \text{pr}^*0^* \mathcal{P}_{01}$. If we set $\theta = (u_1)_{f_0}(u_0)_{f_1}^{-1} \in G(R_{f_0 f_1}[t])$, then there is a commutative diagram of the form

$$\begin{array}{ccccc}
 \mathcal{P}_0 & \longleftarrow & \mathcal{P}_{01} & \longrightarrow & \mathcal{P}_1 \\
 \downarrow u_0 & & \swarrow (u_0)_{f_1} & & \searrow (u_1)_{f_0} \\
 \text{pr}^*0^* \mathcal{P}_0 & \longleftarrow & \text{pr}^*0^* \mathcal{P}_{01} & \xrightarrow{\theta} & \text{pr}^*0^* \mathcal{P}_{01} & \longrightarrow & \text{pr}^*0^* \mathcal{P}_1 \\
 & & & & & & \downarrow u_1
 \end{array}$$

If θ is the identity, then, by fppf descent for G -torsors, the isomorphisms u_0 and u_1 glue to give an isomorphism $\mathcal{P} \cong \text{pr}^*0^* \mathcal{P}$, as desired. If not, since $0^*u_i = 1$, we see that θ restricts along $t = 0$ to the identity. Then Lemma 3.2.2 guarantees that we can find $\tau_i \in G(R_{f_i}[t])$ such that $\tau_i(0) = 1$ and $\theta = \tau_0 \tau_1^{-1}$. Thus, $(\tau_0 u_0)_{f_1} = (\tau_1 u_1)_{f_0}$ and, replacing u_0 by $\tau_0 u_0$ and u_1 by $\tau_1 u_1$, we can glue these isomorphisms to conclude that \mathcal{P} is extended. □

Theorem 3.2.5 (local-to-global principle) *Let R be a commutative ring and suppose G is a linear R -group scheme. If \mathcal{P} is a G -torsor over $R[t_1, \dots, t_n]$, then:*

- (A_n) *The set $Q(\mathcal{P})$ consisting of $g \in R$ such that $\mathcal{P}|_{\text{Spec } R_g[t_1, \dots, t_n]}$ is extended from R_g is an ideal in R .*
- (B_n) *If $\mathcal{P}|_{\text{Spec } R_{\mathfrak{m}}[t_1, \dots, t_n]}$ is extended for every maximal ideal $\mathfrak{m} \subset R$, then \mathcal{P} is extended.*

Proof We know that (A₁) holds by Proposition 3.2.4.

(A_n) \implies (B_n) It suffices to check that, for \mathcal{P} satisfying the conditions in (B_n), the ideal $Q(\mathcal{P})$ is the unit ideal in R . To this end, let $\mathcal{P}|_0$ be the pullback of \mathcal{P} along the zero section $\text{Spec } R \rightarrow \text{Spec } R[t_1, \dots, t_n]$ and let \mathcal{P}' be the pullback of $\mathcal{P}|_0$ along the structure map $\text{Spec } R[t_1, \dots, t_n] \rightarrow \text{Spec } R$.

For any maximal ideal $\mathfrak{m} \subset R$, since $\mathcal{P}|_{\text{Spec } R_{\mathfrak{m}}[t_1, \dots, t_n]}$ is by assumption extended, we know there is an isomorphism $\varphi: \mathcal{P}|_{\text{Spec } R_{\mathfrak{m}}[t_1, \dots, t_n]} \xrightarrow{\sim} \mathcal{P}'|_{\text{Spec } R_{\mathfrak{m}}[t_1, \dots, t_n]}$. Since G -torsors over affine bases are of finite presentation under our hypotheses by Lemma 2.3.3, there exists $g \in R \setminus \mathfrak{m}$ such that φ is the localization of an isomorphism of torsors over $\text{Spec } R_g[t_1, \dots, t_n]$. It follows that $g \in Q(\mathcal{P}) \setminus \mathfrak{m}$ and therefore that $Q(\mathcal{P})$ is not contained in \mathfrak{m} , ie $Q(\mathcal{P}) = R$.

(A₁) \implies (A_n) We proceed by induction on n . Assume therefore that (A_{n-1}) holds. By the conclusion of the previous step, this means (B_{n-1}) holds as well. Form the set $Q(\mathcal{P})$ as in (A_n) . It is straightforward to check that $R \cdot Q(\mathcal{P}) \subset Q(\mathcal{P})$ and therefore it suffices to show that if $f_0, f_1 \in Q(\mathcal{P})$, then $f_0 + f_1 \in Q(\mathcal{P})$ as well.

Write $f = f_0 + f_1$. Consider the quotient map $R[t_1, \dots, t_n] \rightarrow R[t_1, \dots, t_{n-1}]$ and set $\mathcal{P}|_{t_n=0}$ to be the restriction of \mathcal{P} under the corresponding morphism of schemes. Likewise, write $\mathcal{P}|_0$ for the restriction of \mathcal{P} along the zero section as in the previous step. Applying (A_1) to the map $R[t_1, \dots, t_{n-1}] \rightarrow R[t_1, \dots, t_{n-1}][t_n]$, we conclude that \mathcal{P}_f is extended from $(\mathcal{P}|_{t_n=0})_f$.

We claim that $(\mathcal{P}|_{t_n=0})_f$ is itself extended from R_f . If that is the case, then \mathcal{P}_f is extended and so $f \in Q(\mathcal{P})$. Since (B_{n-1}) holds, it suffices to show that $(\mathcal{P}|_{t_n=0})_f$ is extended upon restriction to every maximal ideal $\mathfrak{m} \in R_f$. Write $\mathfrak{m} = \mathfrak{p}_f$, where \mathfrak{p} is the preimage of \mathfrak{m} under the localization map $R \rightarrow R_f$. Since $f \notin \mathfrak{p}$, it follows that either f_0 or f_1 is not in \mathfrak{p} ; without loss of generality, we can assume that $f_0 \notin \mathfrak{p}$. By assumption, however, \mathcal{P}_{f_0} is extended from $(\mathcal{P}_0)_{f_0}$, so we conclude that the restriction of $(\mathcal{P}|_{t_n=0})_f$ to the maximal ideal \mathfrak{m} is extended from $(\mathcal{P}_0)_{\mathfrak{p}}$, which is what we wanted to show. □

Corollary 3.2.6 *Let G be a reductive R -group scheme. If R is regular Noetherian or G is split, then the local-to-global principle holds for G -torsors, ie a G -torsor over $R[t_1, \dots, t_n]$ is extended from R if and only if, for every maximal ideal $\mathfrak{m} \subset R$, the G -torsor on $R_{\mathfrak{m}}[t_1, \dots, t_n]$ obtained by restriction is extended from $R_{\mathfrak{m}}$.*

Proof Combine [Proposition 3.1.3](#) and [Theorem 3.2.5](#). □

3.3 Affine homotopy invariance for G -torsors

Let G be a smooth linear R -group scheme. In this section, we analyze when the pullback map

$$H_{\text{Nis}}^1(X, G) \rightarrow H_{\text{Nis}}^1(X \times \mathbb{A}^1, G)$$

is a bijection for X a smooth affine R -scheme.

Special linear groups We begin by recalling some facts about oriented vector bundles over schemes. If X is a scheme, then recall that an oriented vector bundle on X is a pair (\mathcal{E}, φ) consisting of a vector bundle \mathcal{E} on X equipped with an isomorphism

$\varphi: \det \mathcal{E} \xrightarrow{\sim} \mathcal{O}_X$. There is a standard equivalence between the groupoid of oriented vector bundles on X and that of SL_n -torsors over X . Write $\mathcal{V}_n^o(X)$ for the set of isomorphism classes of rank n oriented vector bundles on X .

Theorem 3.3.1 (special linear homotopy invariance) *Fix an integer $n \geq 1$ and suppose R is a ring such that, for every maximal ideal $\mathfrak{m} \subset R$, $R_{\mathfrak{m}}$ is ind-smooth over a Dedekind ring with perfect residue fields (for example, $R_{\mathfrak{m}}$ is Noetherian and regular over such a Dedekind ring). For every integer $m \geq 0$, the map*

$$\mathcal{V}_n^o(\mathrm{Spec} R) \rightarrow \mathcal{V}_n^o(\mathrm{Spec} R[t_1, \dots, t_m])$$

is a bijection.

Proof By [9, Theorem 5.2.1], every vector bundle on $\mathrm{Spec} R[t_1, \dots, t_m]$ is pulled back from a vector bundle on $\mathrm{Spec} R$. In particular, every oriented vector bundle on $\mathrm{Spec} R[t_1, \dots, t_m]$ is pulled back from a vector bundle on $\mathrm{Spec} R$ with trivial determinant. It remains to show that every automorphism of the trivial line bundle on $\mathrm{Spec} R[t_1, \dots, t_m]$ is extended from $\mathrm{Spec} R$. In other words, we must show that the inclusion map $R \rightarrow R[t_1, \dots, t_m]$ induces an isomorphism on unit groups.

Observe that our assumptions guarantee that $R_{\mathfrak{m}}$ is reduced for every maximal ideal $\mathfrak{m} \subset R$, and therefore R must itself be reduced. Since R is reduced, the fact that $R \rightarrow R[t_1, \dots, t_m]$ induces an isomorphism on unit groups follows from a straightforward induction argument, using the elementary observation that if A is a reduced commutative ring, then the map $A \rightarrow A[t]$ induces an isomorphism $A^\times \rightarrow A[t]^\times$. \square

Remark 3.3.2 In [44, Definition 4.3], Morel defines an orientation on a vector bundle \mathcal{E} to be an isomorphism between $\det \mathcal{E}$ and the square of a line bundle. Oriented vector bundles in this sense correspond to torsors under the *metalelinear group* ML_n defined by the pullback square

$$\begin{array}{ccc} ML_n & \longrightarrow & \mathbb{G}_m \\ \downarrow & & \downarrow 2 \\ GL_n & \xrightarrow{\det} & \mathbb{G}_m \end{array}$$

This more general notion of orientation is very natural in Morel’s theory of the Euler class, since the latter only depends on an orientation in this sense. **Theorem 3.3.1** is also true for ML_n -torsors instead of SL_n -torsors, with a very similar proof.

Symplectic groups We refer the reader to Knus [37, Section I.4] for more details about symplectic spaces over rings; we briefly fix notations in the scheme-theoretic context. If X is a scheme and \mathcal{B} is a quasicoherent sheaf on X , an *alternating bilinear form on \mathcal{B}* is a morphism of quasicoherent sheaves $\varphi: \mathcal{B} \otimes_{\mathcal{O}_X} \mathcal{B} \rightarrow \mathcal{O}_X$ such that $\varphi \circ \Delta = 0$, where $\Delta: \mathcal{B} \rightarrow \mathcal{B} \otimes_{\mathcal{O}_X} \mathcal{B}$ is the (nonlinear) diagonal map. If (\mathcal{B}, φ) is a quasicoherent sheaf equipped with an alternating bilinear form, then we will say that φ is *nondegenerate* if φ induces an isomorphism $\mathcal{B} \rightarrow \mathcal{B}^\vee := \text{Hom}_{\mathcal{O}_X}(\mathcal{B}, \mathcal{O}_X)$. By a *symplectic bundle (of rank $2n$)* we will mean a pair (\mathcal{B}, φ) consisting of a (rank $2n$) vector bundle \mathcal{B} on X equipped with a nondegenerate alternating bilinear form φ . Write $\mathcal{H}\mathcal{V}_{2n}(X)$ for the set of isomorphism classes of rank $2n$ symplectic bundles on X .

We briefly recall the standard equivalence between the groupoid of symplectic vector bundles and that of Sp_{2n} -torsors on X . In one direction, send a symplectic vector bundle (\mathcal{B}, φ) to its bundle of “symplectic frames”; by [37, Proposition I.4.1.4], this construction yields an fppf torsor under Sp_{2n} . In the other direction, given an Sp_{2n} -torsor \mathcal{P} on X , consider the vector bundle associated with the standard $2n$ -dimensional representation of Sp_{2n} , which comes equipped with a reduction of structure group to Sp_{2n} , ie an alternating form on the bundle. By [37, Corollary 4.1.2], any symplectic bundle on a scheme X is Zariski locally on X isometric to the hyperbolic space of a trivial vector bundle [37, I.3.5]. Combining these observations, we see that Sp_{2n} -torsors are Zariski locally trivial and that there is an equivalence between the groupoid of symplectic vector bundles over X and that of Nisnevich locally trivial Sp_{2n} -torsors (as mentioned in Example 2.3.4).

Theorem 3.3.3 (symplectic homotopy invariance) *Fix an integer $n \geq 1$ and suppose R is a ring such that, for every maximal ideal $\mathfrak{m} \subset R$, $R_{\mathfrak{m}}$ is ind-smooth over a Dedekind ring with perfect residue fields (for example, $R_{\mathfrak{m}}$ is Noetherian and regular over such a Dedekind ring). For every integer $m \geq 0$, the map*

$$\mathcal{H}\mathcal{V}_{2n}(\text{Spec } R) \rightarrow \mathcal{H}\mathcal{V}_{2n}(\text{Spec } R[t_1, \dots, t_m])$$

is a bijection.

Proof For any integer $n \geq 1$, the group Sp_{2n} is a split reductive R -group scheme (and, by definition, linear). Therefore, applying Theorem 3.2.5, it suffices to demonstrate the result with R replaced by $R_{\mathfrak{m}}$. Since $R_{\mathfrak{m}}$ is local, every finitely generated projective module over $R_{\mathfrak{m}}$ is free. By the assumption on R and [9, Theorem 5.2.1], we know

that, for any integer m , every finitely generated projective $R_m[t_1, \dots, t_m]$ -module is free. Applying [37, Corollary I.4.1.2], we conclude that every symplectic space over $R_m[t_1, \dots, t_m]$ is isometric to the hyperbolic space of a free module. In particular, every symplectic space over $R_m[t_1, \dots, t_m]$ is extended from R_m . \square

A formalism for homotopy invariance We recall a formalism introduced by Colliot-Thélène and Ojanguren; the following result is a slight extension of [21, Théorème 1.1].

Proposition 3.3.4 *Fix an infinite base field k . Suppose F is a functor from the category of k -algebras to the category of pointed sets with the following properties:*

(P1) *The functor F commutes with filtered inductive limits of rings with flat transition morphisms.*

(P2) *For every extension field L/k and every integer $n \geq 0$, the restriction map*

$$F(L[t_1, \dots, t_n]) \rightarrow F(L(t_1, \dots, t_n))$$

has trivial kernel.

(P3) *The functor F has weak affine Nisnevich excision, ie for any smooth k -algebra A , any étale A -algebra B and any element $f \in A$ such that $A/fA \cong B/fB$, the map*

$$\ker(F(A) \rightarrow F(A_f)) \rightarrow \ker(F(B) \rightarrow F(B_f))$$

is a surjection.

If B is the localization of a smooth k -algebra at a maximal ideal, then, setting $K_B = \text{Frac}(B)$, for any integer $n \geq 0$ the restriction map

$$F(B[t_1, \dots, t_n]) \rightarrow F(K_B(t_1, \dots, t_n))$$

has trivial kernel.

Proof Set $d := \dim B$ and write \mathfrak{m} for the maximal ideal of B . Suppose that

$$\xi_0 \in \ker(F(B[t_1, \dots, t_n]) \rightarrow F(K_B(t_1, \dots, t_n))).$$

Let ξ be the image of ξ_0 in $F(K_B[t_1, \dots, t_n])$. Then, by assumption, ξ lies in the kernel of $F(K_B[t_1, \dots, t_n]) \rightarrow F(K_B(t_1, \dots, t_n))$. By (P2), we conclude that ξ is trivial.

Using (P1), we conclude that there is an element $g \in \mathfrak{m} \setminus 0$ such that ξ_0 restricts trivially to $F(B_g[t_1, \dots, t_n])$. Then, by Knus [37, Corollary VIII.3.2.5], there exist a polynomial ring $L[x_1, \dots, x_d]$, a maximal ideal $\mathfrak{n} \subset L[x_1, \dots, x_d]$, a local

essentially étale morphism $\varphi: A \rightarrow B$ (where $A = L[x_1, \dots, x_d]_n$) and an element $f \in \mathfrak{m}$ such that $\varphi(f) = ug$ for u a unit in $B_{\mathfrak{m}}$ and φ induces an isomorphism $A/fA \xrightarrow{\sim} B/gB$. By (P3), we conclude that there exists an element $\xi'_0 \in \ker(\mathbf{F}(A[t_1, \dots, t_n]) \rightarrow \mathbf{F}(A_f[t_1, \dots, t_n]))$ mapping to ξ_0 . However, ξ'_0 is also evidently in $\ker(\mathbf{F}(A[t_1, \dots, t_n]) \rightarrow \mathbf{F}(K_A(t_1, \dots, t_n)))$. Thus, it suffices to establish the result in the case where B is the localization of a polynomial ring at a maximal ideal, which is precisely [21, Proposition 1.5]. \square

Isotropic reductive groups If k is a field, a reductive k -group scheme will be called *anisotropic* if it contains no k -subgroup isomorphic to \mathbf{G}_m . We take the following definition for isotropic reductive k -group, but we caution the reader that our definition differs from that of Borel [19, Definition V.20.1]; we choose this definition because it better suits our eventual applications.

Definition 3.3.5 If k is a field, a reductive k -group scheme G will be called *isotropic* if each of the almost k -simple components of the derived group of G contains a k -subgroup scheme isomorphic to \mathbf{G}_m .

Remark 3.3.6 See Borel [19, Section V.20] or Gille [31, Section 9.1] for further discussion of isotropic reductive groups. In general, the existence of a *noncentral* split multiplicative k -subgroup is equivalent to the existence of a parabolic k -subgroup by the dynamic construction described just before Lemma 3.1.5. In particular, isotropic reductive k -groups admit proper parabolic subgroups.

Theorem 3.3.7 If k is an infinite field, and G is an isotropic reductive k -group (see Definition 3.3.5), then, for any smooth k -algebra A and any integer $n \geq 0$, the map

$$H_{\text{Nis}}^1(\text{Spec } A, G) \rightarrow H_{\text{Nis}}^1(\text{Spec } A[t_1, \dots, t_n], G)$$

is a bijection.

Proof We have to show that every Nisnevich locally trivial G -torsor \mathcal{P} over the ring $A[t_1, \dots, t_n]$ is extended from A . After Corollary 3.2.6, it suffices to show that, for every maximal ideal \mathfrak{m} of A , the G -torsor $\mathcal{P}_{\mathfrak{m}}$ over $A_{\mathfrak{m}}[t_1, \dots, t_n]$ is extended from $A_{\mathfrak{m}}$; we will show that in fact $\mathcal{P}_{\mathfrak{m}}$ is trivial.

We claim that the functor $A \mapsto H_{\text{Nis}}^1(\text{Spec } A, G)$ from k -algebras to pointed sets satisfies the axioms (P1)–(P3) of Proposition 3.3.4. Axiom (P1) is a consequence of our finite presentation hypotheses by way of Lemma 2.3.3. Axiom (P2) uses the

hypothesis that G is isotropic and follows from [21, Proposition 2.4 and Theorem 2.5] (note that our definition of isotropic reductive k -group coincides with that used in [21, Section 2, page 103]). Axiom (P3) is a formal consequence of the fact that $H_{\text{Nis}}^1(-, G) \cong \pi_0(B\mathbf{Tors}_{\text{Nis}}(G))$ where $B\mathbf{Tors}_{\text{Nis}}(G)$ satisfies affine Nisnevich excision (see Section 2.3). By the conclusion of Proposition 3.3.4, it suffices to show \mathcal{P}_m becomes trivial over $\text{Frac}(A_m)(t_1, \dots, t_n)$, but this follows immediately from the fact that a field has no nontrivial Nisnevich covering sieves. \square

Remark 3.3.8 At least if k is an infinite perfect field, Theorem 3.3.7 admits a converse: if G is a reductive k -group such that $H_{\text{Nis}}^1(-, G)$ is \mathbb{A}^1 -invariant on Sm_k^{aff} , then G is isotropic; see Balwe and Sawant [14, Theorem 1]. In fact, for G reductive, the following three conditions are equivalent:

- (i) G is isotropic (in the sense of Definition 3.3.5).
- (ii) $H_{\text{Nis}}^1(-, G)$ is \mathbb{A}^1 -invariant on smooth affine k -schemes.
- (iii) $R_{\text{Nis}} \text{Sing}^{\mathbb{A}^1} G$ is \mathbb{A}^1 -invariant.

The implication (i) \implies (ii) is Theorem 3.3.7, (ii) \implies (iii) is a special case of Theorem 2.4.2, and (iii) \implies (i) is [14, Theorem 4.7].

4 Applications

In this section, we collect a number of applications of the results established so far. Section 4.1 discusses representability results for Nisnevich locally trivial torsors. As mentioned in Remark 3, that representability results should hold for torsors under SL_n and Sp_{2n} was observed by Schlichting [57, Remark 6.23]; we simply observe in these cases that the expected classical geometric objects yield models for the representing spaces. In Section 4.2 we establish that for various classes of homogeneous spaces for reductive groups applying the singular construction produces an \mathbb{A}^1 -local space. Section 4.3 establishes strong \mathbb{A}^1 -invariance of homotopy sheaves of the singular construction of a reductive group under suitable additional hypotheses. Finally, Section 4.4 studies a purely algebraic problem using our techniques, namely nilpotence of nonstable K_1 -functors.

4.1 Affine representability results for torsors

Let $\text{Gr}_{n,n+N}$ be the usual Grassmannian parametrizing n -dimensional subspaces of an $(n+N)$ -dimensional vector space. Let $\widetilde{\text{Gr}}_{n,n+N}$ be the complement of the

zero section in the total space of the determinant of the tautological vector bundle on $\text{Gr}_{n,n+N}$. The space $\widetilde{\text{Gr}}_{n,n+N}$ parametrizes rank n subspaces of the $(n+N)$ -dimensional vector space equipped with a specified trivialization of their determinant. We set $\widetilde{\text{Gr}}_n := \text{colim}_N \widetilde{\text{Gr}}_{n,n+N}$, where the transition maps are the same as those in the definition Gr_n . With these definitions, we can establish a geometric representability result for oriented vector bundles.

Theorem 4.1.1 *Suppose k is ind-smooth over a Dedekind ring with perfect residue fields. Then, for any $X \in \text{Sm}_k^{\text{aff}}$, and any integer $n \geq 1$, there is a bijection*

$$\mathcal{V}_n^o(X) \cong [X, \widetilde{\text{Gr}}_n]_{\mathbb{A}^1}$$

that is functorial in X .

Proof Recall from Example 2.3.4 and the discussion preceding Theorem 3.3.1 that, for any integer $n \geq 1$, there is a functorial bijection of the form $\mathcal{V}_n^o(X) \cong H_{\text{Nis}}^1(X, \text{SL}_n)$. Combining Theorems 2.3.5 and 3.3.1, we conclude that, under the stated hypotheses on k , $H_{\text{Nis}}^1(X, \text{SL}_n) \cong [X, \text{BSL}_n]_{\mathbb{A}^1}$ for any smooth affine k -scheme X .

Using the notation of Morel and Voevodsky [45, Section 4.2], the space $B_{\text{gm}}(\text{SL}_n, i)$ (attached to the defining inclusion $i: \text{SL}_n \hookrightarrow \text{GL}_n$) is precisely $\widetilde{\text{Gr}}_n$. Therefore, combining the results of [45, Section 4.2], and using the fact that all SL_n -torsors are Zariski (and thus Nisnevich) locally trivial, we conclude that the map $\widetilde{\text{Gr}}_n \rightarrow \text{BSL}_n$ classifying the universal SL_n -torsor over $\widetilde{\text{Gr}}_n$ is an \mathbb{A}^1 -weak equivalence. \square

If we let H be the standard 2-dimensional hyperbolic space, then we can consider the symplectic vector space $H^{\oplus N}$. Panin and Walter construct a scheme $\text{HGr}_{n,n+N}$ that parametrizes rank $2n$ symplectic subspaces of $H^{\oplus(n+N)}$ and we set $\text{HGr}_n := \text{colim}_N \text{HGr}_{n,n+N}$ [51]. Alternatively, HGr can be described as the colimit

$$\text{colim}_N \text{Sp}_{2(n+N)} / (\text{Sp}_{2n} \times \text{Sp}_{2N}).$$

Using these definitions, we are now able to establish a geometric representability theorem for symplectic vector bundles.

Theorem 4.1.2 *Suppose k is ind-smooth over a Dedekind ring with perfect residue fields. Then, for any $X \in \text{Sm}_k^{\text{aff}}$, there is a bijection*

$$\mathcal{H}\mathcal{V}_{2n}(X) \cong [X, \text{HGr}_n]_{\mathbb{A}^1}$$

that is functorial in X .

Proof Proceeding as in the proof of [Theorem 4.1.1](#), we combine [Example 2.3.4](#) and the discussion preceding [Theorem 3.3.3](#) to conclude that there is a functorial bijection of the form $\mathcal{H}\mathcal{V}_{2n}(X) \cong H_{\text{Nis}}^1(X, \text{Sp}_{2n})$. Combining [Theorems 2.3.5](#) and [3.3.3](#), we conclude that, under the stated hypotheses on k , for any smooth affine k -scheme X , $H_{\text{Nis}}^1(X, \text{Sp}_{2n}) \cong [X, B\text{Sp}_{2n}]_{\mathbb{A}^1}$. Finally, by the proof of [\[50, Theorem 8.2\]](#), we can conclude that HGr_n is \mathbb{A}^1 -weakly equivalent to $B\text{Sp}_{2n}$, and thus, for any smooth k -scheme X , $[X, \text{HGr}_n]_{\mathbb{A}^1} \cong [X, B\text{Sp}_{2n}]_{\mathbb{A}^1}$. □

We now establish [Theorem 1](#).

Theorem 4.1.3 *Suppose k is an infinite field, and G is an isotropic reductive k -group (see [Definition 3.3.5](#)). For any smooth affine k -scheme X , there is a functorial bijection*

$$H_{\text{Nis}}^1(X, G) \cong [X, BG]_{\mathbb{A}^1}.$$

Proof Combine [Theorems 2.3.5](#) and [3.3.7](#). □

Remark 4.1.4 In [Theorem 4.1.3](#), the isotropy condition on G cannot be weakened; see [Remark 3.3.8](#).

4.2 Affine representability results for some homogeneous spaces

Let Q_{2n-1} be the smooth affine quadric over \mathbb{Z} defined by $\sum_i x_i y_i = 1$. There is a standard identification $\text{SL}_n/\text{SL}_{n-1} \xrightarrow{\sim} Q_{2n-1}$. Let Q_{2n} be the smooth affine quadric over \mathbb{Z} defined by $\sum_i x_i y_i = z(z + 1)$ (in [Asok, Doran and Fasel \[4\]](#), it is shown that Q_{2n} is \mathbb{A}^1 -weakly equivalent to $\mathbb{P}^{1 \wedge n}$ over $\text{Spec } \mathbb{Z}$). In particular, there are isomorphisms $Q_2 \cong \text{SL}_2/\mathbf{G}_m$ and $Q_4 \cong \text{Sp}_4/(\text{Sp}_2 \times \text{Sp}_2)$ over $\text{Spec } \mathbb{Z}$. If R is a ring in which 2 is invertible, then Q_{2n} is isomorphic over R to the quadric defined by the standard hyperbolic form $\sum_i x_i y_i + z^2 = 1$. It then follows from [Lemma 3.1.7](#) that Q_{2n} is isomorphic over R to the homogeneous space $\text{SO}_{2n+1}/\text{SO}_{2n}$.

Theorem 4.2.1 *If R is a ring that is ind-smooth over a Dedekind ring with perfect residue fields, then Q_{2n-1} is \mathbb{A}^1 -naive. In particular, for any smooth affine R -scheme X , there is a functorial bijection*

$$\pi_0(\text{Sing}^{\mathbb{A}^1} Q_{2n-1})(X) \xrightarrow{\sim} [X, Q_{2n-1}]_{\mathbb{A}^1}.$$

Proof The scheme Q_{2n-1} is isomorphic over $\text{Spec } \mathbb{Z}$ to the homogeneous space $\text{GL}_n/\text{GL}_{n-1}$. Since all torsors for GL_{n-1} are Zariski locally trivial, it follows that $\text{GL}_n \rightarrow Q_{2n-1}$ is Zariski locally trivial (in fact, one can just write down an explicit trivialization). Using [9, Theorem 5.2.1] we may apply [Theorem 2.4.2](#) to conclude. \square

Theorem 4.2.2 *If either (a) $n \leq 2$, and R is a ring that is ind-smooth over a Dedekind ring with perfect residue fields, or (b) $n \geq 3$ and R is an infinite field having characteristic unequal to 2, then Q_{2n} is \mathbb{A}^1 -naive. In particular, under either set of hypotheses, for any smooth affine R -scheme X , there is a functorial bijection*

$$\pi_0(\text{Sing}^{\mathbb{A}^1} Q_{2n})(X) \xrightarrow{\sim} [X, Q_{2n}]_{\mathbb{A}^1}.$$

Proof For $n = 1$ consider the identification $Q_2 \cong \text{SL}_2/\mathbf{G}_m$. Affine homotopy invariance holds for \mathbf{G}_m -torsors over an arbitrary regular base, and for torsors under $\text{SL}_2 \cong \text{Sp}_2$ by assumption. The result follows immediately from [Theorem 2.4.2](#). Similarly, for $n = 2$ consider the identification $Q_4 \cong \text{Sp}_4/(\text{Sp}_2 \times \text{Sp}_2)$. Again, by assumption we may combine [Theorems 3.3.3](#) and [2.4.2](#) to conclude.

For $n \geq 3$ we proceed slightly differently. The SO_{2n} -torsor $\text{SO}_{2n+1} \rightarrow Q_{2n}$ is still Zariski locally trivial by [Lemma 3.1.7](#). Since SO_m is split for $m \geq 3$, we may apply [Theorem 3.3.7](#) to conclude that $H_{\text{Nis}}^1(-, \text{SO}_m)$ is \mathbb{A}^1 -invariant on Sm_R^{aff} for any integer $m \geq 3$. Then we apply [Theorem 2.4.2](#) to conclude. \square

Remark 4.2.3 If $X = \text{Spec } A$, then a map $f: X \rightarrow Q_{2n}$ yields an ideal $I \subset A$ and a surjection $\omega: (A/I)^{\oplus n} \rightarrow I/I^2$; the ideal I is the ideal generated by the images of x_1, \dots, x_n, z in the coordinate presentation of the quadric. The class of f in $\pi_0(\text{Sing}^{\mathbb{A}^1} Q_{2n})(X)$ depends only on the pair (I, ω) and is called the ‘‘Segre class’’ of (I, ω) ; see Fasel [28, Theorem 2.0.2]. When X is smooth over an infinite field, the Segre class provides an obstruction to lifting ω to a surjection $A^{\oplus n} \rightarrow I$ [28, Theorem 3.2.8].

Zariski fiber bundles with affine space fibers If F is a fixed S -scheme, we will say that an S -morphism $\pi: E \rightarrow B$ is a *Zariski fiber bundle of S -schemes with fibers isomorphic to F* if there exist an S -scheme U , a Zariski covering morphism $U \rightarrow B$ and an isomorphism $\varphi: U \times_B E \xrightarrow{\sim} U \times_S F$ over U . The following result, which generalizes a result of Morel [44, Theorem 8.9(2)], applies to affine vector bundle torsors (also called Jouanolou–Thomason devices; see Weibel [63, Definition 4.2 and Proposition 4.4]).

Lemma 4.2.4 *Suppose $B \in \text{Sm}_S$, and $\pi: E \rightarrow B$ is a Zariski fiber bundle of S -schemes with fibers isomorphic to \mathbb{A}_S^n . For any $X = \text{Spec } R \in \text{Sm}_S^{\text{aff}}$, the induced map*

$$\text{Sing}^{\mathbb{A}^1} E(X) \rightarrow \text{Sing}^{\mathbb{A}^1} B(X)$$

is an acyclic Kan fibration. Moreover, E is \mathbb{A}^1 -naive if and only if B is \mathbb{A}^1 -naive.

Proof By Goerss and Jardine [32, Theorem I.11.2], it suffices to show that, for any integer $n \geq 0$, given a diagram of the form

$$\begin{array}{ccc} \partial\Delta_R^n & \longrightarrow & E \\ \downarrow & & \downarrow \pi \\ \Delta_R^n & \longrightarrow & B \end{array}$$

there is a morphism $\Delta_R^n \rightarrow E$ making both resulting triangles commute.

Given a diagram as above, there is an induced map $\partial\Delta_R^n \rightarrow \Delta_R^n \times_B E$. By the assumption on π , the pullback $\pi': \Delta_R^n \times_B E \rightarrow \Delta_R^n$ makes the ring of functions on $\Delta_R^n \times_B E$ into a locally polynomial algebra over $R[t_1, \dots, t_n]$ in the sense of Bass, Connell and Wright [16, Theorem 4.4]. Therefore, by [16, Theorem 4.4], we conclude that π' is a geometric vector bundle over Δ_R^n , ie the spectrum of a symmetric algebra over Δ_R^n . Now, if $\mathcal{E} \rightarrow \Delta_R^n$ is a geometric vector bundle, then the inclusion map $\partial\Delta_R^n \rightarrow \Delta_R^n$ induces a surjective map $\text{Hom}(\Delta_R^n, \mathcal{E}) \rightarrow \text{Hom}(\partial\Delta_R^n, \mathcal{E})$. Therefore, the lift we hoped to construct is guaranteed to exist.

For the second statement, let \tilde{E} and \tilde{B} be Nisnevich-local \mathbb{A}^1 -invariant replacements of E and B , respectively, and consider the commutative square of simplicial presheaves

$$\begin{array}{ccc} \text{Sing}^{\mathbb{A}^1} E & \longrightarrow & \tilde{E} \\ \downarrow & & \downarrow \\ \text{Sing}^{\mathbb{A}^1} B & \longrightarrow & \tilde{B}. \end{array}$$

Since the left vertical map is a weak equivalence on affines, the right vertical map is a weak equivalence. It follows that the upper horizontal map is a weak equivalence on affines if and only if the lower horizontal map is. □

Example 4.2.5 If $X \in \text{Sm}_S^{\text{aff}}$ is an affine scheme, then any finitely presented Zariski fiber bundle of S -schemes $\pi: E \rightarrow X$ with fibers isomorphic to affine spaces is actually a vector bundle by the result of Bass, Connell and Wright mentioned above [16]; this

result was obtained independently by Suslin [60]. On the other hand, if X is not affine, then even if π admits a section, it may not be isomorphic to a vector bundle; see Iarrobino [35, Theorem 1] for an example with $X = \mathbb{P}^1$.

Homogeneous spaces with nonreductive stabilizers The following result extends and simplifies the proof of a theorem of Morel [44, Theorem 8.9]; in particular, we allow the case $n = 2$.

Corollary 4.2.6 *If R is a ring that is ind-smooth over a Dedekind ring with perfect residue fields, then $\mathbb{A}^n \setminus 0$ is \mathbb{A}^1 -naive. In particular, for any smooth affine R -scheme X , there is a canonical bijection*

$$\pi_0 \operatorname{Sing}^{\mathbb{A}^1}(\mathbb{A}^n \setminus 0)(X) \xrightarrow{\sim} [X, \mathbb{A}^n \setminus 0]_{\mathbb{A}^1}.$$

Proof The map $\operatorname{SL}_n \rightarrow \mathbb{A}^n \setminus 0$ given by “projection onto the first column” factors through a map $\operatorname{SL}_n/\operatorname{SL}_{n-1} \rightarrow \mathbb{A}^n \setminus 0$; this map is a Zariski fiber bundle with fibers isomorphic to affine spaces. By Lemma 4.2.4, it suffices to show that $\operatorname{SL}_n/\operatorname{SL}_{n-1}$ is \mathbb{A}^1 -naive. This follows from Theorem 4.2.1 via the standard isomorphism $\operatorname{SL}_n/\operatorname{SL}_{n-1} \cong Q_{2n-1}$ (send a matrix in SL_n to the first row and first column of its inverse). \square

Lemma 4.2.7 *Let X be a simplicial set and $k \geq 0$. If X has the right lifting property with respect to the inclusion $\partial\Delta^m \subset \Delta^m$ for every $m \leq k + 1$, then X is k -connected.*

Proof A simplicial set X is k -connected if and only if the Kan complex

$$\operatorname{cosk}_{k+1} \operatorname{Ex}^\infty X$$

is contractible, or equivalently has the right lifting property with respect to $\partial\Delta^m \subset \Delta^m$ for all m . By adjunction, this is the case if and only if $\operatorname{Ex}^\infty X$ has the right lifting property with respect to $\partial\Delta^m \subset \Delta^m$ for $m \leq k + 1$. By definition of Ex^∞ , it suffices to show that X itself has the right lifting property with respect to $\operatorname{sd}^r(\partial\Delta^m) \subset \operatorname{sd}^r(\Delta^m)$ for all r and all $m \leq k + 1$. In fact, X has the right lifting property with respect to any monomorphism between $(k + 1)$ -skeletal simplicial sets, since such a monomorphism is a transfinite composition of pushouts of $\partial\Delta^m \subset \Delta^m$ for $m \leq k + 1$. \square

Proposition 4.2.8 *Let $n, k \geq 0$ and let R be a commutative ring such that the Bass stable range of $R[t_0, \dots, t_k]$ is at most n . Then the simplicial set $\operatorname{Sing}^{\mathbb{A}^1}(\mathbb{A}^n \setminus 0)(R)$ is k -connected. In particular, if R is Noetherian of Krull dimension d , $\operatorname{Sing}^{\mathbb{A}^1}(\mathbb{A}^n \setminus 0)(R)$ is $(n - d - 2)$ -connected.*

Proof By Lemma 4.2.7, it suffices to show that the map

$$\mathrm{Um}_n(\Delta_R^m) \rightarrow \mathrm{Um}_n(\partial\Delta_R^m)$$

is surjective for all $m \leq k + 1$, where $\mathrm{Um}_n(X) = \mathrm{Hom}(X, \mathbb{A}^n \setminus 0)$ is the set of unimodular rows of length n in $\mathcal{O}(X)$. By assumption, the Bass stable range of Δ_R^{k+1} is at most n . It follows that the Bass stable range of Δ_R^m is at most n , for all $m \leq k + 1$. Now the result is a special case of the following more general statement, which follows easily from the definition of Bass stable range: if X is an affine scheme of Bass stable range $\leq n$ and $Y \subset X$ is a finitely presented closed subscheme, then the map $\mathrm{Um}_n(X) \rightarrow \mathrm{Um}_n(Y)$ is surjective. \square

Remark 4.2.9 Under the assumption of Corollary 4.2.6, the set $\pi_0 \mathrm{Sing}^{\mathbb{A}^1}(\mathbb{A}^n \setminus 0)(X)$ for $n \geq 3$ has a concrete description due to Fasel [27, Theorem 2.1]. Indeed, it is the quotient of the set $\mathrm{Um}_n(X)$ of unimodular rows of length n by the action of the subgroup $E_n(X) \subset \mathrm{SL}_n(X)$ generated by elementary shearing matrices. In loc. cit., it is assumed that R is a field, but the proof works more generally using a result of Lindel and Popescu [53, Proposition 2.1]. Taking $X = \mathcal{Q}_{2n-1}$, we obtain a bijection

$$[\mathbb{A}^n \setminus 0, \mathbb{A}^n \setminus 0]_{\mathbb{A}^1} \cong \mathrm{Um}_n(\mathcal{Q}_{2n-1})/E_n(\mathcal{Q}_{2n-1}).$$

By Corollary 4.2.6, we have $[S^1, \mathbb{A}^n \setminus 0]_{\mathbb{A}^1, * } \cong \pi_1 \mathrm{Sing}^{\mathbb{A}^1}(\mathbb{A}^n \setminus 0)(R)$, and Proposition 4.2.8 shows that this group is trivial if n is at least the Bass stable range of $R[t_0, t_1]$. In that case, we may therefore identify $[\mathbb{A}^n \setminus 0, \mathbb{A}^n \setminus 0]_{\mathbb{A}^1}$ with the set of maps in the pointed \mathbb{A}^1 -homotopy category. Note that $\mathrm{colim}_n [\mathbb{A}^n \setminus 0, \mathbb{A}^n \setminus 0]_{\mathbb{A}^1, * }$ is the set of endomorphisms of the motivic sphere spectrum over the ring R .

The following result is Theorem 4.

Theorem 4.2.10 *If k is an infinite field, G is an isotropic reductive k -group (see Definition 3.3.5) and $P \subset G$ is a parabolic k -subgroup possessing an isotropic Levi factor (eg if G is split), then G/P is \mathbb{A}^1 -naive. In particular, for any smooth affine k -scheme X , there is a functorial bijection*

$$\pi_0(\mathrm{Sing}^{\mathbb{A}^1} G/P)(X) \xrightarrow{\sim} [X, G/P]_{\mathbb{A}^1}.$$

Remark 4.2.11 Given a reductive k -group and a nontrivial parabolic subgroup $P \subset G$, it is not obvious that P has a Levi factor. Nevertheless, as mentioned in Remarks 3.1.6, our hypotheses guarantee that P has a Levi factor. If L is a Levi factor for P , then L may itself be anisotropic.

Proof Lemma 3.1.5(ii) implies that $G \rightarrow G/L$ is generically trivial. Since k is assumed infinite and L is reductive, we claim $G \rightarrow G/L$ is actually Zariski locally trivial. An elementary argument for Zariski local triviality of $G \rightarrow G/L$ sketched in Remark 3.1.6(2), but alternatively we can use [21, Théorème 2.1], to which, in a moment, implicit appeal will be made.

By Theorem 2.4.2, whose hypotheses hold by Theorem 3.3.7, we conclude that G/L is \mathbb{A}^1 -naive. By Lemma 3.1.5(iii), $G/L \rightarrow G/P$ is a composition of Zariski fiber bundles with affine space fibers. Hence, G/P is also \mathbb{A}^1 -naive, by Lemma 4.2.4. \square

The above result can be significantly strengthened at the expense of further restrictions on the groups under consideration.

Theorem 4.2.12 *Suppose R is ind-smooth over a Dedekind ring with perfect residue fields (for example, R is Noetherian and regular over such a Dedekind ring). If $G \cong \mathrm{GL}_n$ or Sp_{2n} , and if $P \subset G$ is a standard parabolic subgroup, then G/P is \mathbb{A}^1 -naive. In particular, for any smooth affine R -scheme X , there is a functorial bijection*

$$\pi_0(\mathrm{Sing}^{\mathbb{A}^1} G/P)(X) \xrightarrow{\sim} [X, G/P]_{\mathbb{A}^1}.$$

Proof Assume first that $R = \mathbb{Z}$. If $P \subset G$ is a standard parabolic with Levi factor L , then L is itself a special group in the sense of Grothendieck–Serre, ie all étale locally trivial torsors are Zariski locally trivial. Thus, the map $G \rightarrow G/L$ in Lemma 3.1.5(ii) is automatically Zariski locally trivial. One sees that the map $G/L \rightarrow G/P$ is a Zariski fiber bundle with affine space fibers by combining Lemma 3.1.5(iii) with the fact that all finitely generated projective \mathbb{Z} -modules are free. By extending scalars to R , it follows that corresponding statements hold for the resulting group scheme over R .

With these modifications, the proof is essentially identical to that of Theorem 4.2.10; however, instead of appealing to Theorem 3.3.7, we use Theorem 3.3.3 or [9, Theorem 5.2.1] to establish the necessary homotopy invariance statement. \square

Example 4.2.13 Theorem 4.2.12 applies if $P \subset \mathrm{GL}_n$ is a maximal parabolic subgroup, in which case $G/P \cong \mathrm{Gr}_{m,n}$ for some integer $m \leq n$.

4.3 Affine representability for nonstable K –theory and strong \mathbb{A}^1 –invariance results

Suppose G is a smooth linear R –group scheme. For any integer $i \geq 1$, one can define Karoubi–Villamayor–style nonstable K –theory functors attached to G by means of the formula

$$\mathrm{KV}_{i+1}^G(U) := \pi_i(\mathrm{Sing}^{\mathbb{A}^1} G)(U)$$

In this form, the definition goes back to Jardine [36, Theorem 3.8], but had precursors in the work of Krusemeyer [38, Section 3]; see Wendt [64] for a more detailed analysis of such functors in the context of \mathbb{A}^1 –homotopy theory. As a straightforward application of our results, we obtain \mathbb{A}^1 –representability results for nonstable KV–functors.

Theorem 4.3.1 *If k is an infinite field, and G is an isotropic reductive k –group (in the sense of Definition 3.3.5), then G is \mathbb{A}^1 –naive. In particular, for any smooth affine k –scheme U , there are canonical isomorphisms*

$$\mathrm{KV}_{i+1}^G(U) \cong [S^i \wedge U_+, G]_{\mathbb{A}^1, *}$$

Proof Apply Theorem 2.4.2 with $H = e$ (Theorem 3.3.7 ensures the hypotheses are satisfied). □

Remark 4.3.2 Results such as the above were studied initially by Morel [44, Theorem 8.1] and Moser [46] (see also [65, Theorem 5.3]) for G a general split group, and by the third author and K Völkel in the isotropic reductive case [62]. These results depend crucially on first establishing homotopy invariance for nonstable K_1 –functors via “elementary matrix” techniques. As a consequence these proofs do not easily extend to the important case where G has semisimple rank 1, which was treated separately by Moser. Our proof above makes no such assumption on the homotopy invariance of nonstable K_1 –functors. As a consequence, Theorem 4.3.1 can also be used to slightly uniformize the proof of [13, Theorem 3.4].

We can also establish the strong \mathbb{A}^1 –invariance of the sheafifications of the nonstable K_1 –presheaves attached to arbitrary isotropic reductive k –groups with k infinite.

Theorem 4.3.3 *Suppose k is an infinite field, and G is an isotropic reductive k –group (in the sense of Definition 3.3.5). For any integer $n \geq 0$, the following statements hold:*

- (i) *The Zariski sheaf $a_{\mathrm{Zar}}\pi_n(\mathrm{Sing}^{\mathbb{A}^1} G)$ is a Nisnevich sheaf.*
- (ii) *The sheaf $a_{\mathrm{Zar}}\pi_n(\mathrm{Sing}^{\mathbb{A}^1} G)$ is strongly \mathbb{A}^1 –invariant.*

Proof We begin by recalling some key results of Morel [44, Chapter 6]. If \mathcal{X} is a Nisnevich-local and \mathbb{A}^1 -invariant pointed simplicial presheaf on Sm_k , the sheaf $a_{\text{Nis}}\pi_1(\mathcal{X})$ is strongly \mathbb{A}^1 -invariant by [44, Theorem 6.1]. Moreover, the map $a_{\text{Zar}}\pi_1(\mathcal{X}) \rightarrow a_{\text{Nis}}\pi_1(\mathcal{X})$ is an isomorphism by [44, Corollary 6.9(2)] (the standing assumption that $a_{\text{Nis}}\pi_0(\mathcal{X})$ is trivial is not used in the proof).

By Theorems 2.3.5(i) and 3.3.7, under the stated hypotheses on k , the simplicial presheaf $R_{\text{Zar}}\text{Sing}^{\mathbb{A}^1}B_{\text{Nis}}G$ is Nisnevich-local and \mathbb{A}^1 -invariant. Applying the results of the previous paragraph to the simplicial presheaf

$$\mathcal{X} = \mathbf{R}\Omega^n R_{\text{Zar}}\text{Sing}^{\mathbb{A}^1}B_{\text{Nis}}G,$$

we conclude that

$$a_{\text{Zar}}\pi_{n+1}(\text{Sing}^{\mathbb{A}^1}B_{\text{Nis}}G)$$

is a strongly \mathbb{A}^1 -invariant Nisnevich sheaf of groups for any $n \geq 0$. By Corollary 2.2.2, the map

$$\pi_n(\text{Sing}^{\mathbb{A}^1}\mathbf{R}\Omega B_{\text{Nis}}G) \rightarrow \pi_{n+1}(\text{Sing}^{\mathbb{A}^1}B_{\text{Nis}}G)$$

is an isomorphism on affines, and hence it becomes an isomorphism after Zariski sheafification. Finally, we conclude the proof by observing that $G \simeq \mathbf{R}\Omega B_{\text{Nis}}G$ by Lemma 2.3.2(iii). \square

Remark 4.3.4 The results from [44, Chapter 6] used in the proof of Theorem 4.3.3 do not require k to be perfect. If the base field k is in addition perfect, then, provided $a_{\text{Zar}}\pi_n(\text{Sing}^{\mathbb{A}^1}G)$ is abelian, we can use [44, Theorem 5.46] to conclude that it is strictly \mathbb{A}^1 -invariant. The assumption that k is infinite in the above statement only appears because of our appeal to Theorem 3.3.7. To remove this restriction, we would need homotopy invariance for torsors under isotropic reductive groups over finite fields.

If G is a reductive k -group, we can define $G(k)^+$ to be the normal subgroup of $G(k)$ generated by the k -points of subgroups of G isomorphic to \mathbb{G}_a . The Whitehead group of G is defined by the formula

$$W(k, G) := G(k)/G(k)^+;$$

we refer the reader to Gille's survey [30] for more details about Whitehead groups. In particular, Tits showed that $W(k, G)$ detects whether $G(k)$ is projectively simple. Results of Margaux allow us to connect nonstable K_1 -functors (as above) with Whitehead groups. More precisely, one has the following result:

Proposition 4.3.5 *Suppose k is an infinite field, and G is an isotropic reductive k -group (in the sense of Definition 3.3.5). For any finitely generated separable extension L/k , there are canonical isomorphisms*

$$\pi_0(\text{Sing}^{\mathbb{A}^1} G)(L) \cong W(L, G).$$

functorial with respect to field extensions. Moreover, the assignment $L \mapsto W(L, G)$ extends to a strongly \mathbb{A}^1 -invariant sheaf on Sm_k .

Proof The first statement follows from Margaux [40, Theorem 3.10] (see also Gille [30, Section 4.3]) and only requires G to be isotropic in the sense of Borel [19, Definition V.20.1]. The second statement follows from the strong \mathbb{A}^1 -invariance of $a_{\text{Zar}}\pi_0(\text{Sing}^{\mathbb{A}^1} G)$ established in Theorem 4.3.3(2). □

Whitehead groups are also related to arithmetic questions, eg regarding R -equivalence in $G(k)$ (see Gille [30, Section 7] for a discussion of R -equivalence in the context under consideration).

Corollary 4.3.6 *Let k be an infinite field and G a semisimple, simply connected, absolutely almost simple, isotropic k -group, and set $\mathbf{G} := a_{\text{Zar}}\pi_0(\text{Sing}^{\mathbb{A}^1} G)$. Then*

- (i) *for any finitely generated separable extension L/k , there is an isomorphism of the form $\mathbf{G}(L) \cong G(L)/R$,*
- (ii) *the contracted sheaf \mathbf{G}_{-1} is trivial, and*
- (iii) *if k is furthermore perfect and G has classical type, then \mathbf{G} is strictly \mathbb{A}^1 -invariant.*

Proof The first statement follows from Proposition 4.3.5 and [30, Théorème 7.2].

For the second statement, recall that $\mathbf{G}_{-1}(U) = \ker((id, 1)^*: \mathbf{G}(U \times \mathbf{G}_m) \rightarrow \mathbf{G}(U))$. As \mathbf{G} is strongly \mathbb{A}^1 -invariant by Theorem 4.3.3, \mathbf{G}_{-1} is also strongly \mathbb{A}^1 -invariant by Morel [44, Lemma 2.32]. In particular, it is an unramified sheaf, which implies that the map $\mathbf{G}(X) \rightarrow \mathbf{G}(k(X))$ is injective for any irreducible smooth scheme X . By [30, Theorem 5.8], we conclude that $\mathbf{G}(k(U)) \rightarrow \mathbf{G}(k(U \times \mathbf{G}_m))$ is a bijection and thus that $\mathbf{G}_{-1}(U)$ is trivial for any $U \in \text{Sm}_k$.

For the final statement, if k is furthermore perfect, it suffices by [44, Theorem 5.46] to show that \mathbf{G} is an abelian group valued functor. Because \mathbf{G} is unramified, it suffices to check abelianness on sections over extensions of the base field. By (i), if G has classical type, this follows from a result of Chernousov and Merkurjev [30, Théorème 7.7]. □

Remark 4.3.7 The statement $\mathbf{G}_{-1} = 0$ of [Corollary 4.3.6\(ii\)](#) is equivalent to the assertion that \mathbf{G} is a birational sheaf. If G is not simply connected, then the sheaf \mathbf{G} is not, in general, birational. For example suppose G is a split semisimple group having nontrivial algebraic fundamental group Π (in the sense of Chevalley groups). If we let $\mathcal{H}_{\text{ét}}^1(\Pi)$ be the Nisnevich sheaf associated with the presheaf $U \mapsto H_{\text{ét}}^1(U, \Pi)$, then $\mathbf{G} \cong \mathcal{H}_{\text{ét}}^1(\Pi)$, which is not birational.

4.4 Nilpotence for nonstable K_1 functors

In this section, we include one more sample application of our results: we give a uniform proof of some nilpotence results for nonstable K_1 -functors discussed in the previous section; such nilpotence results have been studied for instance by Bak [\[11\]](#) and Bak, Hazrat and Vavilov [\[12\]](#). The main result of this section is [Theorem 4.4.3](#), which solves [\[12, Problem 6\]](#) in a number of cases of interest (see [Remark 4.4.4](#) for more details). The approach we pursue has the benefit that it is conceptually simple (modeled on classical topological results) and applies to rather general isotropic reductive k -groups. The tradeoff to this generality is that, unlike [\[12\]](#), we are forced to restrict attention to smooth k -algebras with k an infinite field.

We use the following notation/terminology. If (\mathcal{X}, x) is a pointed simplicial presheaf on Sm_k , we will say that \mathcal{X} is Nisnevich-connected if $a_{\text{Nis}}\pi_0(\mathcal{X})$ is trivial and, given an integer $n \geq 1$, we will say that \mathcal{X} is Nisnevich n -connected if $a_{\text{Nis}}\pi_i(\mathcal{X}, x)$ is trivial for $i \leq n$.

Now, suppose \mathcal{G} is a simplicial presheaf of group-like h -spaces on Sm_k (h -group for short) pointed by the identity. In that case, there is an induced morphism $\mathcal{G} \rightarrow a_{\text{Nis}}\pi_0\mathcal{G}$; this morphism is a morphism of h -groups. Write \mathcal{G}^0 for the homotopy fiber of \mathcal{G} , so that there is a homotopy fiber sequence of the form

$$\mathcal{G}^0 \rightarrow \mathcal{G} \rightarrow a_{\text{Nis}}\pi_0\mathcal{G}.$$

By construction, \mathcal{G}^0 is a Nisnevich-connected h -group. Using this notation, we can adapt arguments of Whitehead [\[66, Corollary 2.12\]](#) to establish an abstract nilpotence result.

Proposition 4.4.1 *Assume k is a Noetherian ring of finite Krull dimension, and suppose \mathcal{G} is a Nisnevich-local simplicial presheaf of h -groups on Sm_k (pointed by the identity).*

(i) For any $X \in \text{Sm}_k$, there is an exact sequence of groups of the form

$$1 \rightarrow [X, \mathcal{G}^0] \rightarrow [X, \mathcal{G}] \rightarrow a_{\text{Nis}}\pi_0(\mathcal{G})(X).$$

(ii) If $X \in \text{Sm}_k$ has Krull dimension $\leq d$, then $[X, \mathcal{G}^0]$ is nilpotent of class $\leq d$.

Proof Point (i) is immediate from the long exact sequence of maps into a homotopy fiber sequence and the fact that $a_{\text{Nis}}\pi_0(\mathcal{G})$ is 0-truncated.

For (ii), it suffices to assume $\mathcal{G} = \mathcal{G}^0$ is Nisnevich-connected. In that case, $\mathcal{G}^{\wedge n}$ is Nisnevich n -connected. Indeed, this follows from the corresponding connectivity estimate for smash products of simplicial sets by checking on stalks. Now, a straightforward obstruction theory argument (see Morel [44, Lemma B.5]) using the connectivity estimate we just mentioned shows that $[X, \mathcal{G}^{\wedge n}] = *$ if $\dim X \leq n$. To conclude, we simply observe that every n -fold commutator in $[X, \mathcal{G}]$ factors as $X \rightarrow \mathcal{G}^{\wedge n} \rightarrow \mathcal{G}$ (here, we use the assumption that \mathcal{G} is an h -group and thus has a strict identity). \square

Remark 4.4.2 The result above is rather general. Indeed, as is evident from the proof, it holds for simplicial h -group objects in the local homotopy theory of simplicial presheaves on a site for which Postnikov towers converge.

Now suppose G is an isotropic reductive k -group in the sense of Definition 3.3.5. Following Petrov and Stavrova [52], for any commutative k -algebra R and any parabolic k -subgroup $P \subset G$, we define the elementary subgroup $E_P(R)$ as the subgroup of $G(R)$ generated by the R -points of the unipotent radical of P and the R -points of the unipotent radical of its opposite. A priori, $E_P(R)$ depends on P and $E_P(R)$ need not be a normal subgroup of G . However, Theorem 1 of [52] guarantees that if each semisimple normal subgroup of G has rank ≥ 2 , then $E_P(R)$ is both independent of P and normal in $G(R)$; under these hypotheses we define $E(R) := E_P(R)$ for any choice of proper parabolic and define $K_1^G(R) := G(R)/E(R)$.

We can also consider $G^0(R) \subset G(R)$, the subset of $G(R)$ consisting of matrices g for which there exists $g(t) \in G(R[t])$ with $g(0) = 1$ and $g(1) = g$; this subgroup is evidently normal. By construction, $E_P(R) \subset G^0(R)$ and $\text{KV}_1^G(R) = G(R)/G^0(R)$. Therefore there is a short exact sequence of groups

$$1 \rightarrow G^0(R)/E(R) \rightarrow K_1^G(R) \rightarrow \text{KV}_1^G(R) \rightarrow 1.$$

Theorem 4.4.3 Suppose k is an infinite field, G is an isotropic reductive k -group (in the sense of Definition 3.3.5) and R is a smooth k -algebra of dimension d .

- (i) If for every finitely generated separable extension L/k the Whitehead group $W(L, G)$ is trivial (abelian), then $\mathrm{KV}_1^G(R)$ is (an extension of an abelian group by) a nilpotent group of class $\leq d$.
- (ii) If furthermore k is perfect, for every finitely generated separable extension L/k the Whitehead group $W(L, G)$ is trivial (abelian) and every semisimple normal subgroup of G has rank ≥ 2 , then $K_1^G(R)$ is (an extension of an abelian group by) a nilpotent group of class $\leq d$.

Proof Let $\mathcal{G} = R_{\mathrm{Zar}} \mathrm{Sing}^{\mathbb{A}^1} G$. According to [Theorem 4.3.3](#), the Nisnevich sheaf $a_{\mathrm{Nis}}\pi_0(\mathcal{G})$ is strongly \mathbb{A}^1 -invariant. By [Proposition 4.3.5](#), the group of sections $a_{\mathrm{Nis}}\pi_0(\mathcal{G})(L)$ over finitely generated extensions L/k coincides with $W(L, G)$. In particular, the assumption that $W(L, G)$ is trivial (abelian) for every finitely generated separable extension L/k implies that the sheaf $a_{\mathrm{Nis}}\pi_0(\mathcal{G})$ is trivial (abelian).

By [Theorem 4.3.1](#) and [Proposition 2.1.3](#), \mathcal{G} is Nisnevich-local and $\mathrm{KV}_1^G(R) = [\mathrm{Spec} R, \mathcal{G}]$. Then (i) follows immediately from [Proposition 4.4.1](#).

Consider the exact sequence appearing before the statement gives a surjective map $K_1^G(R) \rightarrow \mathrm{KV}_1^G(R)$. Under the additional hypotheses in (ii), it follows immediately from a result of Stavrova [[59](#), Theorem 1.3] that this surjection is an isomorphism and (ii) follows from (i). \square

Remark 4.4.4 Combined with known structural results about $W(-, G)$ (viewed as a functor on the category of finitely generated extensions of the base field), the above result solves a problem posed by Bak, Hazrat and Vavilov [[12](#), Problem 6] in a number of new cases. For example, in [[30](#), Théorème 6.1], Gille summarizes results of Chernousov and Platonov that detail situations where $W(-, G)$ is trivial for all finitely generated separable extensions L/k . See [Corollary 4.3.6\(iii\)](#) for hypotheses that guarantee $W(-, G)$ is an abelian group valued functor on the category of (eg if G has classical type). Furthermore, it has been conjectured that $W(-, G)$ always takes values in abelian groups.

References

- [1] **S Anantharaman**, *Schémas en groupes, espaces homogènes et espaces algébriques sur une base de dimension 1*, from “Sur les groupes algébriques”, Mém. Soc. Math. France 33, Soc. Math. France, Paris (1973) 5–79 [MR](#)
- [2] **M Artin**, *Versal deformations and algebraic stacks*, Invent. Math. 27 (1974) 165–189 [MR](#)

- [3] **M Artin, A Grothendieck, J L Verdier**, *Théorie des topos et cohomologie étale des schémas, Tome 1: Théorie des topos, Exposés I–IV (SGA 4_I)*, Lecture Notes in Math. 269, Springer (1972) [MR](#)
- [4] **A Asok, B Doran, J Fasel**, *Smooth models of motivic spheres and the clutching construction*, Int. Math. Res. Not. 2017 (2017) 1890–1925 [MR](#)
- [5] **A Asok, J Fasel**, *Algebraic vector bundles on spheres*, J. Topol. 7 (2014) 894–926 [MR](#)
- [6] **A Asok, J Fasel**, *A cohomological classification of vector bundles on smooth affine threefolds*, Duke Math. J. 163 (2014) 2561–2601 [MR](#)
- [7] **A Asok, J Fasel**, *Splitting vector bundles outside the stable range and \mathbb{A}^1 -homotopy sheaves of punctured affine spaces*, J. Amer. Math. Soc. 28 (2015) 1031–1062 [MR](#)
- [8] **A Asok, J Fasel**, *Euler class groups and motivic stable cohomotopy*, preprint (2016) [arXiv](#)
- [9] **A Asok, M Hoyois, M Wendt**, *Affine representability results in \mathbb{A}^1 -homotopy theory, I: vector bundles*, Duke Math. J. 166 (2017) 1923–1953 [MR](#)
- [10] **A Asok, M Hoyois, M Wendt**, *Generically split octonion algebras and \mathbb{A}^1 -homotopy theory*, preprint (2017) [arXiv](#)
- [11] **A Bak**, *Nonabelian K -theory: the nilpotent class of K_1 and general stability*, K -Theory 4 (1991) 363–397 [MR](#)
- [12] **A Bak, R Hazrat, N Vavilov**, *Localization-completion strikes again: relative K_1 is nilpotent by abelian*, J. Pure Appl. Algebra 213 (2009) 1075–1085 [MR](#)
- [13] **C Balwe, A Sawant**, *R -equivalence and \mathbb{A}^1 -connectedness in anisotropic groups*, Int. Math. Res. Not. 2015 (2015) 11816–11827 [MR](#)
- [14] **C Balwe, A Sawant**, *\mathbb{A}^1 -connectedness in reductive algebraic groups*, Trans. Amer. Math. Soc. 369 (2017) 5999–6015 [MR](#)
- [15] **J Barge, J Lannes**, *Suites de Sturm, indice de Maslov et périodicité de Bott*, Progress in Mathematics 267, Birkhäuser, Basel (2008) [MR](#)
- [16] **H Bass, E H Connell, D L Wright**, *Locally polynomial algebras are symmetric algebras*, Invent. Math. 38 (1976) 279–299 [MR](#)
- [17] **P Belmans, A J de Jong**, et al., *The Stacks project*, electronic reference, Columbia University (2005–) Available at <http://stacks.math.columbia.edu>
- [18] **A Białyński-Birula**, *Rationally trivial homogeneous principal fibrations of schemes*, Invent. Math. 11 (1970) 259–262 [MR](#)
- [19] **A Borel**, *Linear algebraic groups*, 2nd edition, Graduate Texts in Mathematics 126, Springer (1991) [MR](#)
- [20] **C Cazanave**, *Algebraic homotopy classes of rational functions*, Ann. Sci. Éc. Norm. Supér. 45 (2012) 511–534 [MR](#)

- [21] **J-L Colliot-Thélène, M Ojanguren**, *Espaces principaux homogènes localement triviaux*, Inst. Hautes Études Sci. Publ. Math. 75 (1992) 97–122 [MR](#)
- [22] **B Conrad**, *Reductive group schemes*, from “Autour des schémas en groupes, I”, Panor. Synthèses 42/43, Soc. Math. France, Paris (2014) 93–444 [MR](#)
- [23] **B Conrad, O Gabber, G Prasad**, *Pseudo-reductive groups*, New Mathematical Monographs 17, Cambridge Univ. Press (2010) [MR](#)
- [24] **M Demazure, A Grothendieck**, *Schémas en groupes, Tome II: Groupes de type multiplicatif, et structure des schémas en groupes généraux, Exposés VIII–XVIII (SGA 3_{II})*, Lecture Notes in Math. 152, Springer (1970) [MR](#)
- [25] **M Demazure, A Grothendieck**, *Schémas en groupes, Tome III: Structure des schémas en groupes réductifs, Exposés XIX–XXVI (SGA 3_{III})*, Lecture Notes in Math. 153, Springer (1970) [MR](#)
- [26] **D Dugger, S Hollander, D C Isaksen**, *Hypercovers and simplicial presheaves*, Math. Proc. Cambridge Philos. Soc. 136 (2004) 9–51 [MR](#)
- [27] **J Fasel**, *Some remarks on orbit sets of unimodular rows*, Comment. Math. Helv. 86 (2011) 13–39 [MR](#)
- [28] **J Fasel**, *On the number of generators of ideals in polynomial rings*, Ann. of Math. 184 (2016) 315–331 [MR](#)
- [29] **R Fedorov, I Panin**, *A proof of the Grothendieck–Serre conjecture on principal bundles over regular local rings containing infinite fields*, Publ. Math. Inst. Hautes Études Sci. 122 (2015) 169–193 [MR](#)
- [30] **P Gille**, *Le problème de Kneser–Tits*, from “Séminaire Bourbaki, 2007/2008”, Astérisque 326, Soc. Mat. de France, Paris (2009) Exposé 983, 39–81 [MR](#)
- [31] **P Gille**, *Sur la classification des schémas en groupes semi-simples*, from “Autour des schémas en groupes, III”, Panor. Synthèses 47, Soc. Math. France, Paris (2015) 39–110 [MR](#)
- [32] **P G Goerss, J F Jardine**, *Simplicial homotopy theory*, Progress in Mathematics 174, Birkhäuser, Basel (1999) [MR](#)
- [33] **A Grothendieck**, *Éléments de géométrie algébrique, IV: Étude locale des schémas et des morphismes de schémas, IV*, Inst. Hautes Études Sci. Publ. Math. 32 (1967) 5–361 [MR](#)
- [34] **S Hollander**, *A homotopy theory for stacks*, Israel J. Math. 163 (2008) 93–124 [MR](#)
- [35] **A Iarrobino**, *An algebraic fibre bundle over \mathbf{P}_1 that is not a vector bundle*, Topology 12 (1973) 229–232 [MR](#)
- [36] **J F Jardine**, *On the homotopy groups of algebraic groups*, J. Algebra 81 (1983) 180–201 [MR](#)

- [37] **M-A Knus**, *Quadratic and Hermitian forms over rings*, Grundle. Math. Wissen. 294, Springer (1991) [MR](#)
- [38] **MI Krusemeyer**, *Fundamental groups, algebraic K-theory, and a problem of Abhyankar*, Invent. Math. 19 (1973) 15–47 [MR](#)
- [39] **T Y Lam**, *Serre’s problem on projective modules*, Springer (2006) [MR](#)
- [40] **B Margaux**, *The structure of the group $G(k[t])$: variations on a theme of Soulé*, Algebra Number Theory 3 (2009) 393–409 [MR](#)
- [41] **JS Milne**, *Étale cohomology*, Princeton Mathematical Series 33, Princeton Univ. Press (1980) [MR](#)
- [42] **J Milnor, D Husemoller**, *Symmetric bilinear forms*, Ergeb. Math. Grenzgeb. 73, Springer (1973) [MR](#)
- [43] **F Morel**, *On the Friedlander–Milnor conjecture for groups of small rank*, from “Current developments in mathematics, 2010” (D Jerison, B Mazur, T Mrowka, W Schmid, R Stanley, S-T Yau, editors), International, Somerville, MA (2011) 45–93 [MR](#)
- [44] **F Morel**, \mathbb{A}^1 -*algebraic topology over a field*, Lecture Notes in Mathematics 2052, Springer (2012) [MR](#)
- [45] **F Morel, V Voevodsky**, \mathbb{A}^1 -*homotopy theory of schemes*, Inst. Hautes Études Sci. Publ. Math. 90 (1999) 45–143 [MR](#)
- [46] **L-F Moser**, \mathbb{A}^1 -*locality results for linear algebraic groups*, in preparation
- [47] **L-F Moser**, *Rational triviale Torseure und die Serre–Grothendiecksche Vermutung*, Diplomarbeit, Ludwig-Maximilians-Universität München (2008) Available at <http://www.mathematik.uni-muenchen.de/~lfmoser/da.pdf>
- [48] **Y A Nisnevich**, *Espaces homogènes principaux rationnellement triviaux et arithmétique des schémas en groupes réductifs sur les anneaux de Dedekind*, C. R. Acad. Sci. Paris Sér. I Math. 299 (1984) 5–8 [MR](#)
- [49] **I Panin**, *Proof of Grothendieck–Serre conjecture on principal G -bundles over regular local rings containing a finite field*, preprint (2014) [arXiv](#)
- [50] **I Panin, C Walter**, *On the motivic commutative ring spectrum \mathbf{BO}* , preprint (2010) [arXiv](#)
- [51] **I Panin, C Walter**, *Quaternionic Grassmannians and Pontryagin classes in algebraic geometry*, preprint (2010) [arXiv](#)
- [52] **V A Petrov, A K Stavrova**, *Elementary subgroups in isotropic reductive groups*, Algebra i Analiz 20 (2008) 160–188 [MR](#) In Russian; translated in *St. Petersburg Math. J.* 20 (2009) 625–644
- [53] **D Popescu**, *Polynomial rings and their projective modules*, Nagoya Math. J. 113 (1989) 121–128 [MR](#)

- [54] **D Quillen**, *Projective modules over polynomial rings*, Invent. Math. 36 (1976) 167–171 [MR](#)
- [55] **MS Raghunathan**, *Principal bundles on affine space*, from “C P Ramanujam: a tribute” (K G Ramanathan, editor), Tata Inst. Fund. Res. Studies in Math. 8, Springer (1978) 187–206 [MR](#)
- [56] **A Ramanathan**, *Deformations of principal bundles on the projective line*, Invent. Math. 71 (1983) 165–191 [MR](#)
- [57] **M Schlichting**, *Euler class groups and the homology of elementary and special linear groups*, Adv. Math. 320 (2017) 1–81 [MR](#)
- [58] **T A Springer, F D Veldkamp**, *Octonions, Jordan algebras and exceptional groups*, Springer (2000) [MR](#)
- [59] **A Stavrova**, *Homotopy invariance of non-stable K_1 -functors*, J. K-Theory 13 (2014) 199–248 [MR](#)
- [60] **A A Suslin**, *Locally polynomial rings and symmetric algebras*, Izv. Akad. Nauk SSSR Ser. Mat. 41 (1977) 503–515 [MR](#) In Russian; translated in *Math. USSR-Izv.* 11 (1977) 472–484
- [61] **R W Thomason**, *Equivariant resolution, linearization, and Hilbert’s fourteenth problem over arbitrary base schemes*, Adv. in Math. 65 (1987) 16–34 [MR](#)
- [62] **K Voelkel, M Wendt**, *On \mathbb{A}^1 -fundamental groups of isotropic reductive groups*, C. R. Math. Acad. Sci. Paris 354 (2016) 453–458 [MR](#)
- [63] **C A Weibel**, *Homotopy algebraic K -theory*, from “Algebraic K -theory and algebraic number theory” (MR Stein, R K Dennis, editors), Contemp. Math. 83, Amer. Math. Soc., Providence, RI (1989) 461–488 [MR](#)
- [64] **M Wendt**, *\mathbb{A}^1 -homotopy of Chevalley groups*, J. K-Theory 5 (2010) 245–287 [MR](#)
- [65] **M Wendt**, *Rationally trivial torsors in \mathbb{A}^1 -homotopy theory*, J. K-Theory 7 (2011) 541–572 [MR](#)
- [66] **G W Whitehead**, *On mappings into group-like spaces*, Comment. Math. Helv. 28 (1954) 320–328 [MR](#)

Department of Mathematics, University of Southern California
Los Angeles, CA, United States

Department of Mathematics, Massachusetts Institute of Technology
Cambridge, MA, United States

Fakultät Für Mathematik, Universität Duisburg-Essen
Essen, Germany

asok@usc.edu, hoyois@mit.edu, matthias.wendt@uni-due.de

Proposed: Haynes R Miller

Received: 13 July 2016

Seconded: Mark Behrens, Stefan Schwede

Revised: 25 April 2017

