

4
5 **ON A NONLINEAR FRACTIONAL BOUNDARY VALUE PROBLEM INCLUDING**
6 **HADAMARD FRACTIONAL DERIVATIVE WITH P-LAPLACIAN**

7
8 TUGBA SENLIK CERDIK

9
10 **ABSTRACT.** This paper presents the existence results of the p-Laplacian Hadamard fractional boundary
11 value problem. We analyze the existence of multiple positive solutions for the fractional boundary value
12 problem with the nonlinear term which involves the derivative term with the help of a fixed point theorem.
13 An example is considered to describe the validity of the main result.

14
15
16 **1. Introduction**

17
18 Fractional calculus has been a widely studied area of research in recent decades, especially because
19 of its ability to describe complex issues in various areas of science and engineering, by the help of
20 arbitrary order approach. Also, with respect to integer order derivatives, fractional order derivatives are
21 more suitable to describe the memory and inherent properties of many kinds of materials and processes.
22 Fractional order derivative is a global operator, that can serve as a tool for various applied phenomena
23 such as mathematical biology [27], control theory [30], dynamical process [31], *etc.* For the newest
24 studies on fractional calculus, we refer to some books [1, 17, 18, 19, 23, 25, 28]. Moreover, many
25 scientists have introduced the existence results of fractional boundary value problems whereby many
26 various techniques, such as the mixed monotone operator theory, Banach's contraction mapping principle,
27 see [5, 7, 10, 12, 14, 22, 26, 34, 36, 38, 39].

28 Most common fractional derivative types that have been researched in fractional calculus field are
29 Riemann-Liouville and Caputo fractional derivatives. Moreover, Hadamard fractional derivative which
30 was demonstrated by Hadamard [13] has recently received attention from researchers. Hadamard fractional
31 derivative and integral are different because their definitions include logarithmic function of arbitrary order.
32 Within this field, discovering new generalizations for the existing fractional derivatives has always been a
33 major focus of research. Using these generalized operators, we will provide new opportunities to develop
34 existing results from theoretical and applied point of views. In [8], the authors introduced some operators
35 that include Hadamard derivatives and presented the advantage of their approach through some examples.
36 [8] showed a new approach based on linear integro-differential operators with logarithmic kernel via
37 Hadamard fractional calculus for a generalization of the Lomnitz logarithmic creep law. Hadamard
38 fractional calculus can show the mathematical essential of this creep law more exactly in this subject [8].
39 In addition, the works in [1, 4, 18] contributed significantly to the development of Hadamard derivatives
40 within fractional calculus. At the same time, the subject of the fractional differential equations with
41 Hadamard-type derivative has gained a lot of popularity due to their capability to modeling real world
42 phenomena. To examine some of the recent developments on the Hadamard fractional boundary value
43 problem (HFBVP for short), see [2, 3, 24, 29, 32, 35, 37]. Furthermore, the analysis of solutions with
44 p-Laplacian operator for the HFBVP is still an area of further research and many researchers have shown
45 more interest to the HFBVP including p-Laplacian operator, see [11, 15, 16, 21, 33]. Also, the function ρ
46 which involves Hadamard fractional derivative has been examined by only a few researchers, see [6, 11].
47 For a detail monograph on Hadamard fractional calculus and integral, see [1].

In [21], Li and Lin studied the following HFBVP involving the p-Laplacian operator

$$\begin{cases} \mathfrak{D}^\beta(\phi_p(\mathfrak{D}^\alpha u(z))) = \rho(z, u(z)), & 2 < \alpha \leq 3, 1 < \beta \leq 2, 1 < z < e, \\ u(1) = u'(1) = u'(e) = 0, \mathfrak{D}^\alpha u(1) = \mathfrak{D}^\alpha u(e) = 0, \end{cases}$$

where $\rho \in \mathcal{C}([1, e] \times [0, +\infty), [0, +\infty))$. They obtained the existence and uniqueness of the positive solution of the problem.

In [15], Xu *et.al.* examined the following problem

$$\begin{cases} -\mathfrak{D}^\alpha(\phi_p(\mathfrak{D}^\beta u(z))) = \rho(z, u(z)), & z \in (1, T), \\ u(1) = \delta u(1) = \delta u(e) = u(e) = \mathfrak{D}^\beta u(1) = 0, \mathfrak{D}^\beta u(e) = b\mathfrak{D}^\beta u(\eta), \end{cases}$$

in which $1 < \alpha \leq 2, 3 < \beta \leq 4, \rho \in \mathcal{C}([1, e] \times [0, \infty), [0, \infty))$ and δ means delta derivative. By using fixed point theorems, the existence results for the HFBVP is ensured.

In light of the papers mentioned above, we generate some new results about the following problem involving Hadamard fractional derivative

$$(1.1) \quad \begin{cases} {}^H\mathfrak{D}_{1^+}^{\sigma_1}(\phi_p({}^H\mathfrak{D}_{1^+}^{\sigma_2}\kappa(z))) = \rho(z, \kappa(z), {}^H\mathfrak{D}_{1^+}^{\sigma_2}\kappa(z)), & 2 < \sigma_1, \sigma_2 \leq 3, z \in (1, e), \\ \kappa(1) = \kappa'(1) = {}^H\mathfrak{D}_{1^+}^{\sigma_2}\kappa(1) = (\phi_p({}^H\mathfrak{D}_{1^+}^{\sigma_2}\kappa(1)))' = 0, \\ \phi_p({}^H\mathfrak{D}_{1^+}^{\sigma_2}\kappa(e)) = \sum_{i=1}^k a_i \phi_p({}^H\mathfrak{D}_{1^+}^{\sigma_2}\kappa(\eta_i)) + \int_1^e \phi_p({}^H\mathfrak{D}_{1^+}^{\sigma_2}\kappa(z))g(z)\frac{dz}{z}, \\ \kappa(e) = \sum_{j=1}^q b_j \kappa(\xi_j) + \int_1^e \kappa(z)h(z)\frac{dz}{z}, \end{cases}$$

in which $5 < \sigma_1 + \sigma_2 \leq 6$, ${}^H\mathfrak{D}_{1^+}^{\sigma_1}$ and ${}^H\mathfrak{D}_{1^+}^{\sigma_2}$ are the Hadamard fractional derivative, ϕ_p is the p-Laplacian, i.e., $\phi_p(s) = |s|^{p-2}s$, with $s \in \mathbb{R}, p > 1, \phi_p^{-1} = \phi_q$, and $\frac{1}{p} + \frac{1}{q} = 1. a_i \geq 0, b_j \geq 0 (i = 1, 2, \dots, k), (j = 1, 2, \dots, q), 1 < \eta_1 < \eta_2 < \dots < \eta_k < e$ and $1 < \xi_1 < \xi_2 < \dots < \xi_q < e$.

Let us give the necessary assumptions below:

(C1) $\rho \in \mathcal{C}([1, e] \times [0, \infty) \times (-\infty, \infty), (0, \infty)), g, h \in \mathcal{C}([1, e], [0, \infty))$,

(C2) $\sum_{i=1}^k a_i (\log \eta_i)^{\sigma_1-1} + \int_1^e g(z) (\log z)^{\sigma_1-1} \frac{dz}{z} < 1, \sum_{j=1}^q b_j (\log \xi_j)^{\sigma_2-1} + \int_1^e h(z) (\log z)^{\sigma_2-1} \frac{dz}{z} < 1$.

In this study, by using the theory of fixed point under appropriate conditions, we aim to ensure the multiple positive solutions for a HFBVP involving p-Laplacian operator. The function ρ which involves of derivative operator ${}^H\mathfrak{D}_{1^+}^{\sigma_2}$ is studied by only a few scientists. In comparison to [15],[21], our problem includes the Hadamard fractional derivative in the nonlinear term and the boundary conditions are more complicated because of that the boundary conditions consist of the linear combination of the unknown function and integral boundary condition. Also, Hadamard derivative in nonlinear term is complicated, so it requires a more careful study and this type of derivative is a growing area of research.

2. Preliminaries

To meet the requirements in the next part, the fundamental principles of Hadamard fractional calculus are shown here.

Definition 2.1. [25] The Hadamard fractional derivative of fractional order ϖ for a function $c : [1, \infty) \rightarrow \mathbb{R}$ is expressed as

$${}^H\mathfrak{D}_{1^+}^\varpi c(z) = \frac{1}{\Gamma(m-\varpi)} \left(z \frac{d}{dz}\right)^m \int_1^t \left(\log \frac{z}{s}\right)^{m-\varpi-1} c(s) \frac{ds}{s}, \quad m-1 < \varpi < m, m = [\varpi] + 1,$$

where $[\varpi]$ shows the integer part of the real number ϖ and $\log(\cdot) = \log_e(\cdot)$.

Definition 2.2. [25] The Hadamard fractional integral of order ϖ for a function $c : [1, \infty) \rightarrow \mathbb{R}$ is expressed as

$${}^H I_{1+}^{\varpi} c(z) = \frac{1}{\Gamma(\varpi)} \int_1^z (\log \frac{z}{s})^{\varpi-1} c(s) \frac{ds}{s}, \quad \varpi > 0,$$

in the event that the associated integral exists.

Lemma 2.1. [25] Let $\varpi > 0$ and $a \in \mathcal{C}[1, \infty) \cap L^1[1, \infty)$, then the solution of Hadamard fractional differential equation ${}^H \mathcal{D}_{1+}^{\varpi} a(z) = 0$ is expressed as

$$a(z) = \sum_{j=1}^m t_j (\log z)^{\varpi-j},$$

then,

$${}^H I_{1+}^{\varpi} {}^H \mathcal{D}_{1+}^{\varpi} a(z) = a(z) + \sum_{j=1}^m t_j (\log z)^{\varpi-j},$$

in which $t_j \in \mathbb{R}, j = 1, 2, \dots, m, m-1 < \varpi < m, m = [\varpi] + 1$.

Here, we introduce $\mathfrak{X} = \{ \kappa \in \mathcal{C}[1, e] : {}^H \mathcal{D}_{1+}^{\sigma_2} \kappa \in \mathcal{C}[1, e] \}$ by the norm

$$\| \kappa \| = \max \left\{ \max_{z \in [1, e]} |\kappa(z)|, \max_{z \in [1, e]} |{}^H \mathcal{D}_{1+}^{\sigma_2} \kappa(z)| \right\}.$$

Hence, \mathfrak{X} is a Banach space. Demonstrate a cone M by

$$M = \{ \kappa \in \mathfrak{X} : \kappa(z) \geq 0, \forall z \in [1, e] \}.$$

Lemma 2.2. Let $r \in \mathcal{C}[1, e]$ be given, then $c \in \mathfrak{X}$ is a solution of

$$(2.1) \quad \begin{cases} {}^H \mathcal{D}_{1+}^{\sigma_1} c(z) = r(z), & 2 < \sigma_1 \leq 3, \quad z \in (1, e), \\ c(1) = c'(1) = 0, & c(e) = \sum_{i=1}^k a_i c(\eta_i) + \int_1^e g(z) c(z) \frac{dz}{z}, \end{cases}$$

provided that c satisfies the integral equation

$$c(z) = - \int_1^e H(z, s) r(s) \frac{ds}{s}, \quad z \in [1, e],$$

where

$$(2.2) \quad \begin{aligned} H(z, s) &= H_1(z, s) + H_2(z, s), \\ H_1(z, s) &= h(z, s) + \sum_{i=1}^k \frac{a_i (\log z)^{\sigma_1-1}}{\Delta} h(\eta_i, s), \\ H_2(z, s) &= \frac{(\log z)^{\sigma_1-1}}{\Delta_1} \int_1^e H_1(z, s) g(z) \frac{dz}{z}, \end{aligned}$$

with

$$(2.3) \quad h(z, s) = \frac{1}{\Gamma(\sigma_1)} \begin{cases} (\log z)^{\sigma_1-1} (1 - \log s)^{\sigma_1-1} - (\log z - \log s)^{\sigma_1-1}, & 1 \leq s \leq z \leq e, \\ (\log z)^{\sigma_1-1} (1 - \log s)^{\sigma_1-1}, & 1 \leq z \leq s \leq e, \end{cases}$$

$$\text{and } \Delta = 1 - \sum_{i=1}^k a_i (\log \eta_i)^{\sigma_1-1}, \Delta_1 = \Delta - \int_1^e g(z) (\log z)^{\sigma_1-1} \frac{dz}{z}.$$

Proof. From Lemma 2.1, the problem (2.1) can be shown by

$$c(z) = \frac{1}{\Gamma(\sigma_1)} \int_1^z (\log \frac{z}{s})^{\sigma_1-1} r(s) \frac{ds}{s} + t_1 (\log z)^{\sigma_1-1} + t_2 (\log z)^{\sigma_1-2} + t_3 (\log z)^{\sigma_1-3},$$

for $t_1, t_2, t_3 \in \mathbb{R}$. Using $c(1) = c'(1) = 0$, we get $t_2 = 0, t_3 = 0$ and

$$(2.4) \quad c(z) = \frac{1}{\Gamma(\sigma_1)} \int_1^z (\log \frac{z}{s})^{\sigma_1-1} r(s) \frac{ds}{s} + t_1 (\log z)^{\sigma_1-1}.$$

If we apply the boundary condition $c(e) = \sum_{i=1}^k a_i c(\eta_i) + \int_1^e g(z)c(z) \frac{dz}{z}$, from (2.4), we have

$$(2.5) \quad t_1 = \frac{1}{\Delta} \left(-\frac{1}{\Gamma(\sigma_1)} \int_1^e (\log \frac{e}{s})^{\sigma_1-1} r(s) \frac{ds}{s} + \sum_{i=1}^k \frac{a_i}{\Gamma(\sigma_1)} \int_1^{\eta_i} (\log \frac{\eta_i}{s})^{\sigma_1-1} r(s) \frac{ds}{s} + \int_1^e g(z)c(z) \frac{dz}{z} \right).$$

Inserting (2.5) into (2.4),

$$\begin{aligned} c(z) &= \frac{1}{\Gamma(\sigma_1)} \int_1^z (\log \frac{z}{s})^{\sigma_1-1} r(s) \frac{ds}{s} - \frac{(\log z)^{\sigma_1-1}}{\Delta \Gamma(\sigma_1)} \int_1^e (\log \frac{e}{s})^{\sigma_1-1} r(s) \frac{ds}{s} \\ &\quad + \sum_{i=1}^k \frac{a_i (\log z)^{\sigma_1-1}}{\Delta \Gamma(\sigma_1)} \int_1^{\eta_i} (\log \frac{\eta_i}{s})^{\sigma_1-1} r(s) \frac{ds}{s} + \frac{(\log z)^{\sigma_1-1}}{\Delta} \int_1^e g(z)c(z) \frac{dz}{z} \\ &= \frac{1}{\Gamma(\sigma_1)} \int_1^z (\log \frac{z}{s})^{\sigma_1-1} r(s) \frac{ds}{s} - \frac{(\log z)^{\sigma_1-1}}{\Gamma(\sigma_1)} \int_1^e (\log \frac{e}{s})^{\sigma_1-1} r(s) \frac{ds}{s} \\ &\quad - \sum_{i=1}^k \frac{a_i (\log z)^{\sigma_1-1} (\log \eta_i)^{\sigma_1-1}}{\Delta \Gamma(\sigma_1)} \int_1^e (\log \frac{e}{s})^{\sigma_1-1} r(s) \frac{ds}{s} \\ &\quad + \sum_{i=1}^k \frac{a_i (\log z)^{\sigma_1-1}}{\Delta \Gamma(\sigma_1)} \int_1^{\eta_i} (\log \frac{\eta_i}{s})^{\sigma_1-1} r(s) \frac{ds}{s} + \frac{(\log z)^{\sigma_1-1}}{\Delta} \int_1^e g(z)c(z) \frac{dz}{z} \\ &= - \int_1^e h(z,s)r(s) \frac{ds}{s} - \sum_{i=1}^k \frac{a_i (\log z)^{\sigma_1-1}}{\Delta} \int_1^e h(\eta_i,s)r(s) \frac{ds}{s} + \frac{(\log z)^{\sigma_1-1}}{\Delta} \int_1^e g(z)c(z) \frac{dz}{z} \\ &= - \int_1^e H_1(z,s)r(s) \frac{ds}{s} + \frac{(\log z)^{\sigma_1-1}}{\Delta} \int_1^e g(z)c(z) \frac{dz}{z}. \end{aligned}$$

Furthermore,

$$\begin{aligned} \int_1^e g(z)c(z) \frac{dz}{z} &= - \int_1^e g(z) \int_1^e H_1(z,s)r(s) \frac{ds}{s} \frac{dz}{z} \\ &\quad + \frac{1}{\Delta} \int_1^e g(z) (\log z)^{\sigma_1-1} \frac{dz}{z} \int_1^e g(z)c(z) \frac{dz}{z}, \end{aligned}$$

which ensure

$$\int_1^e g(z)c(z) \frac{dz}{z} = -\frac{\Delta}{\Delta_1} \int_1^e g(z) \int_1^e H_1(z,s)r(s) \frac{ds}{s} \frac{dz}{z},$$

Hence,

$$\begin{aligned} c(z) &= - \int_1^e H_1(z,s)r(s) \frac{ds}{s} - \int_1^e H_2(z,s)r(s) \frac{ds}{s} \\ &= - \int_1^e H(z,s)r(s) \frac{ds}{s}. \end{aligned}$$

□

Lemma 2.3. For $c \in \mathcal{C}[1, e]$ and $\kappa \in \mathfrak{X}$, the following problem

$$(2.6) \quad \begin{cases} {}^H\mathcal{D}_{1^+}^{\sigma_2} \kappa(z) = \phi_q(c(z)), & 2 < \sigma_2 \leq 3, \quad z \in (1, e), \\ \kappa(1) = \kappa'(1) = 0, \quad \kappa(e) = \sum_{j=1}^q b_j \kappa(\xi_j) + \int_1^e h(z) \kappa(z) \frac{dz}{z} \end{cases}$$

is equivalent to,

$$\kappa(z) = - \int_1^e K(z, s) \phi_q(c(s)) \frac{ds}{s}, \quad z \in [1, e],$$

where

$$(2.7) \quad K(z, s) = K_1(z, s) + K_2(z, s),$$

$$K_1(z, s) = k(z, s) + \sum_{j=1}^q \frac{b_j (\log z)^{\sigma_2 - 1}}{\Delta^*} k(\xi_j, s),$$

$$K_2(z, s) = \frac{(\log z)^{\sigma_2 - 1}}{\Delta_1^*} \int_1^e K_1(z, s) h(z) \frac{dz}{z},$$

with

$$(2.8) \quad k(z, s) = \frac{1}{\Gamma(\sigma_2)} \begin{cases} (\log z)^{\sigma_2 - 1} (1 - \log s)^{\sigma_2 - 1} - (\log z - \log s)^{\sigma_2 - 1}, & 1 \leq s \leq z \leq e, \\ (\log z)^{\sigma_2 - 1} (1 - \log s)^{\sigma_2 - 1}, & 1 \leq z \leq s \leq e, \end{cases}$$

$$\text{and } \Delta^* = 1 - \sum_{j=1}^q b_j (\log \xi_j)^{\sigma_2 - 1}, \quad \Delta_1^* = \Delta^* - \int_1^e h(z) (\log z)^{\sigma_2 - 1} \frac{dz}{z}.$$

Set $n_{\sigma_i}(z) = (\log z)^{\sigma_i - 1} (1 - \log z)$, $m_{\sigma_i}(z) = (1 - \log z)^{\sigma_i - 1} \log z$, for $\sigma_i > 2$, $z \in [1, e]$, $i = 1, 2$.

Lemma 2.4. [35] $h(z, s), k(z, s)$ introduced by (2.3) and (2.8) have the below features:

(i) $h(z, s)$ and $k(z, s)$ are continuous functions and $h(z, s) \geq 0$, $k(z, s) \geq 0$ for any $z, s \in [1, e]$;

(ii) $n_{\sigma_1}(z) m_{\sigma_1}(s) \leq \Gamma(\sigma_1) h(z, s) \leq (\sigma_1 - 1) m_{\sigma_1}(s)$, for any $z, s \in [1, e]$,

(iii) $n_{\sigma_2}(z) m_{\sigma_2}(s) \leq \Gamma(\sigma_2) k(z, s) \leq (\sigma_2 - 1) m_{\sigma_2}(s)$, for any $z, s \in [1, e]$.

Lemma 2.5. Let $\mu_1 = \frac{\sigma_1 - 1}{\Gamma(\sigma_1)} + \sum_{i=1}^k \frac{a_i (\sigma_1 - 1)}{\Delta \Gamma(\sigma_1)}$, $\mu_2 = \frac{\sigma_2 - 1}{\Gamma(\sigma_2)} + \sum_{j=1}^q \frac{b_j (\sigma_2 - 1)}{\Delta^* \Gamma(\sigma_2)}$, $\mu_1^* = 1 + \frac{1}{\Delta_1} \int_1^e g(z) \frac{dz}{z}$, $\mu_2^* = 1 + \frac{1}{\Delta_1^*} \int_1^e h(z) \frac{dz}{z}$ and $V_1 = \sum_{i=1}^k \frac{a_i n_{\sigma_1}(\eta_i)}{\Delta \Gamma(\sigma_1)}$, $V_2 = \sum_{j=1}^q \frac{b_j n_{\sigma_2}(\xi_j)}{\Delta^* \Gamma(\sigma_2)}$. Then, the functions $H(z, s)$ and $K(z, s)$ introduced by (2.2) and (2.7) ensure the following features:

(i) $H(z, s)$, $K(z, s)$ are continuous and $H(z, s) \geq 0$, $K(z, s) \geq 0$, for $(z, s) \in [1, e] \times [1, e]$;

(ii) $V_1 (\log z)^{\sigma_1 - 1} m_{\sigma_1}(s) \leq H(z, s) \leq \mu_1 \mu_1^* m_{\sigma_1}(s)$, for $(z, s) \in [1, e] \times [1, e]$;

(iii) $V_2 (\log z)^{\sigma_2 - 1} m_{\sigma_2}(s) \leq K(z, s) \leq \mu_2 \mu_2^* m_{\sigma_2}(s)$, for $(z, s) \in [1, e] \times [1, e]$.

Proof. It can be shown that (i) ensures. To show (ii), for $(z, s) \in [1, e] \times [1, e]$, we get,

$$\begin{aligned}
 H(z, s) &= H_1(z, s) + H_2(z, s) \\
 &= h(z, s) + \sum_{i=1}^k \frac{a_i(\log z)^{\sigma_1-1}}{\Delta} h(\eta_i, s) \\
 &\quad + \frac{(\log z)^{\sigma_1-1}}{\Delta_1} \int_1^e H_1(z, s) g(z) \frac{dz}{z} \\
 &\leq \frac{(\sigma_1 - 1)m_{\sigma_1}(s)}{\Gamma(\sigma_1)} + \sum_{i=1}^k \frac{a_i(\sigma_1 - 1)m_{\sigma_1}(s)}{\Delta\Gamma(\sigma_1)} \\
 &\quad + \frac{1}{\Delta_1} \int_1^e (h(z, s) + \sum_{i=1}^k \frac{a_i}{\Delta} h(\eta_i, s)) g(z) \frac{dz}{z} \\
 &\leq \frac{(\sigma_1 - 1)m_{\sigma_1}(s)}{\Gamma(\sigma_1)} + \sum_{i=1}^k \frac{a_i(\sigma_1 - 1)m_{\sigma_1}(s)}{\Delta\Gamma(\sigma_1)} \\
 &\quad + \frac{1}{\Delta_1} \int_1^e \left(\frac{(\sigma_1 - 1)m_{\sigma_1}(s)}{\Gamma(\sigma_1)} + \sum_{i=1}^k \frac{a_i(\sigma_1 - 1)m_{\sigma_1}(s)}{\Delta\Gamma(\sigma_1)} \right) g(z) \frac{dz}{z} \\
 &= \mu_1 \mu_1^* m_{\sigma_1}(s),
 \end{aligned}$$

and

$$\begin{aligned}
 H(z, s) &\geq \sum_{i=1}^k \frac{a_i(\log z)^{\sigma_1-1}}{\Delta} h(\eta_i, s) \\
 &\geq (\log z)^{\sigma_1-1} \sum_{i=1}^k \frac{a_i n_{\sigma_1}(\eta_i) m_{\sigma_1}(s)}{\Delta\Gamma(\sigma_1)} \\
 &= V_1 (\log z)^{\sigma_1-1} m_{\sigma_1}(s).
 \end{aligned}$$

In a similar manner, we observe that (iii) holds. \square

Using Lemma 2.2 and Lemma 2.3, let $A : M \rightarrow \mathfrak{X}$ be an operator by

$$A\kappa(z) = \int_1^e K(z, s) \phi_q \left(\int_1^e H(s, \tau) \rho(\tau, \kappa(\tau), {}^H \mathfrak{D}_{1+}^{\sigma_2} \kappa(\tau)) \frac{d\tau ds}{\tau s} \right).$$

Clearly, if κ is a fixed point of the operator A , then κ represents a solution for the problem (1.1).

Lemma 2.6. *Let (C1) and (C2) hold. Then, $A : M \rightarrow M$ is a completely continuous operator.*

Proof. Let us demonstrate that $A(M) \subset M$. $H(z, s)$, $K(z, s)$, ρ are continuous functions. Thus, A is continuous. Lemma 2.5 and the selection of ρ satisfy that $A(z) \geq 0$ for $z \in [1, e]$. Hence, $A : M \rightarrow M$. Furthermore, by the Arzela–Ascoli theorem, it is obtained that A is a completely continuous operator. \square

Now, the fixed point theorem is demonstrated, which is fundamental in our main result’s proof.

Theorem 2.1. [20] *Let \mathfrak{X} be a Banach space, $M \subseteq \mathfrak{X}$ a cone of \mathfrak{X} . Set*

$$\begin{aligned}
 M_y &= \{ \kappa \in M : \|\kappa\| < y \}, \\
 M(\zeta, j_1, j_2) &= \{ \kappa \in M : j_1 \leq \zeta(\kappa), \|\kappa\| \leq j_2 \}.
 \end{aligned}$$

Assume $A : \overline{M}_y \rightarrow \overline{M}_y$ be a completely continuous operator and ζ be a nonnegative, continuous, concave functional on M with $\zeta(\kappa) \leq \|\kappa\|$ for all $\kappa \in \overline{M}_y$. If there exists $0 < p < n < d \leq y$ such that the following conditions hold:

$$(B_1) \{ \kappa \in M(\zeta, n, d) : \zeta(\kappa) > n \} \neq \emptyset \text{ and } \zeta(A\kappa) > n \text{ for all } \kappa \in M(\zeta, n, d);$$

- 1 (B₂) $\|A\kappa\| < p$ for all $\|\kappa\| \leq p$;
- 2 (B₃) $\zeta(A\kappa) > n$ for $\kappa \in M(\zeta, n, y)$ with $\|A\kappa\| > d$.

3 Then A has at least three positive solutions κ_j for $j \in \{1, 2, 3\}$ in \overline{M}_y satisfying

4
5
6
7
$$\|\kappa_1\| < p, \zeta(\kappa_2) > n, p < \|\kappa_3\| \text{ with } \zeta(\kappa_3) < n.$$

8
9 For simplicity, let

10
11
12
$$B = \mu_2 \mu_2^* \phi_q(\mu_1 \mu_1^*) \int_1^e m_{\sigma_2}(s) \phi_q \left(\int_1^e m_{\sigma_1}(\tau) \frac{d\tau}{\tau} \right) \frac{ds}{s},$$

13
14
$$C = \phi_q(\mu_1 \mu_1^*) \phi_q \left(\int_1^e m_{\sigma_1}(\tau) \frac{d\tau}{\tau} \right),$$

15
16
17
$$F = V_2(\log \eta)^{\sigma_2-1} \phi_q(V_1(\log \eta)^{\sigma_1-1}) \int_{\eta}^e m_{\sigma_2}(s) \frac{ds}{s} \phi_q \left(\int_{\eta}^e m_{\sigma_1}(\tau) \frac{d\tau}{\tau} \right),$$

18
19
20
$$N = V_2(\log \eta)^{\sigma_2-1} \phi_q(V_1(\log \eta)^{\sigma_1-1}) \int_{\eta}^e m_{\sigma_2}(s) \frac{ds}{s}.$$

21
22 **3. Existence Theorem**

23
24 Here, we ensure the required conditions to get at least three positive solutions for the HFBVP (1.1).
25 Firstly, we introduce the nonnegative, continuous, concave functional $\Lambda : M \rightarrow [0, +\infty)$ given by

26
27
$$\Lambda(\kappa) = \min_{z \in [\eta, e]} |\kappa(z)|, \quad \forall \kappa \in M,$$

28
29 where $\eta \in (1, e)$. It is obvious that $\Lambda(\kappa) \leq \|\kappa\|$ for all $\kappa \in M$.

30
31
32 **Theorem 3.1.** Consider that (C1)-(C2) are satisfied and there exist $0 < h < L < K < r$ and $\frac{K(\log \eta)^{\sigma_2}}{\Gamma(\sigma_2+1)} > L$,
33 $K \geq N^{-1} L \max\{1, \mu_2 \mu_2^* \int_1^e m_{\sigma_2}(s) \frac{ds}{s}\} \phi_q(\mu_1 \mu_1^*)$, $\frac{L}{F} < \min\{\frac{r}{B}, \frac{r}{C}\}$ such that the following hypotheses on ρ
34 are satisfied.

- 35
36 (i) $\rho(z, \kappa, \iota) > \phi_p(\frac{L}{F})$ for $(z, \kappa, \iota) \in [\eta, e] \times [L, K] \times [-K, 0]$,
- 37
38 (ii) $\rho(z, \kappa, \iota) \leq \min\{\phi_p(\frac{r}{B}), \phi_p(\frac{r}{C})\}$ for $(z, \kappa, \iota) \in [1, e] \times [0, r] \times [-r, 0]$,
- 39
40 (iii) $\rho(z, \kappa, \iota) < \min\{\phi_p(\frac{h}{B}), \phi_p(\frac{h}{C})\}$ for $(z, \kappa, \iota) \in [1, e] \times [0, h] \times [-h, 0]$.

41
42
43 Then, the HFBVP (1.1) has at least three positive solutions κ_j for $j \in \{1, 2, 3\}$ satisfying

44
45
46
47
$$\|\kappa_1\| < h, \Lambda(\kappa_2) > L, h < \|\kappa_3\| \text{ with } \Lambda(\kappa_3) < L.$$

48
49 **Proof.** We will ensure that the conditions of Theorem 2.1 will be achieved. First, we will demonstrate
50 $A : \overline{M}_r \rightarrow \overline{M}_r$. If $\kappa \in \overline{M}_r$, then $\|\kappa\| \leq r$, using Lemma 2.5 and assumption (ii), for $z \in [1, e]$, we get

$$\begin{aligned}
 |A\kappa(z)| &= \left| \int_1^e K(z,s) \phi_q \left(\int_1^e H(s,\tau) \rho(\tau, \kappa(\tau), {}^H \mathfrak{D}_{1+}^{\sigma_2} \kappa(\tau)) \frac{d\tau}{\tau} \right) \frac{ds}{s} \right| \\
 &\leq \mu_2 \mu_2^* \int_1^e m_{\sigma_2}(s) \phi_q \left(\int_1^e \mu_1 \mu_1^* m_{\sigma_1}(\tau) \rho(\tau, \kappa(\tau), {}^H \mathfrak{D}_{1+}^{\sigma_2} \kappa(\tau)) \frac{d\tau}{\tau} \right) \frac{ds}{s} \\
 &\leq \frac{r}{B} \mu_2 \mu_2^* \phi_q(\mu_1 \mu_1^*) \int_1^e m_{\sigma_2}(s) \phi_q \left(\int_1^e m_{\sigma_1}(\tau) \frac{d\tau}{\tau} \right) \frac{ds}{s} \\
 &= r,
 \end{aligned}$$

and

$$\begin{aligned}
 |{}^H \mathfrak{D}_{1+}^{\sigma_2} \kappa(z)| &= \left| -\phi_q \left(\int_1^e H(s,\tau) \rho(\tau, \kappa(\tau), {}^H \mathfrak{D}_{1+}^{\sigma_2} \kappa(\tau)) \frac{d\tau}{\tau} \right) \right| \\
 &\leq \phi_q(\mu_1 \mu_1^*) \phi_q \left(\int_1^e m_{\sigma_1}(\tau) \rho(\tau, \kappa(\tau), {}^H \mathfrak{D}_{1+}^{\sigma_2} \kappa(\tau)) \frac{d\tau}{\tau} \right) \\
 &\leq \frac{r}{C} \phi_q(\mu_1 \mu_1^*) \phi_q \left(\int_1^e m_{\sigma_1}(\tau) \frac{d\tau}{\tau} \right) \\
 &= r.
 \end{aligned}$$

Thus, $A : \overline{M_r} \rightarrow \overline{M_r}$. By (iii) and similarly to the proof above, it can be obtained that $A : \overline{M_h} \rightarrow M_h$.

Now, we ensure that condition (B1) of Theorem 2.1 is obtained. By choosing $\kappa(z) = \frac{K(\log z)^{\sigma_2}}{\Gamma(\sigma_2+1)}$ for $z \in [1, e]$, we obtain, $\max_{z \in [1, e]} |\kappa(z)| = \frac{K}{\Gamma(\sigma_2+1)}$, $\max_{z \in [1, e]} |\mathfrak{D}^{\sigma_2} \kappa(z)| = K$ and $\Lambda(\kappa) = \min_{z \in [\eta, e]} |\kappa(z)| = \frac{K(\log \eta)^{\sigma_2}}{\Gamma(\sigma_2+1)} > L$ and $\|\kappa\| = K$. Hence, $\kappa(z) = \frac{K(\log z)^{\sigma_2}}{\Gamma(\sigma_2+1)} \in M(\Lambda, L, K)$, that is,

$$\{\kappa \in M(\Lambda, L, K) : \Lambda(\kappa) > L\} \neq \emptyset.$$

If $\kappa \in M(\Lambda, L, K)$, then $L \leq \kappa(z) \leq K$ and $-K \leq \mathfrak{D}^{\sigma_2} \kappa(z) \leq 0$ for any $z \in [\eta, e]$. Using (i), we ensure

$$\begin{aligned}
 \Lambda(A\kappa) &= \min_{z \in [\eta, e]} A\kappa(z) \\
 &\geq V_2(\log \eta)^{\sigma_2-1} \int_{\eta}^e m_{\sigma_2}(s) \phi_q \left(\int_{\eta}^e V_1(\log s)^{\sigma_1-1} m_{\sigma_1}(\tau) \rho(\tau, \kappa(\tau), {}^H \mathfrak{D}_{1+}^{\sigma_2} \kappa(\tau)) \frac{d\tau}{\tau} \right) \frac{ds}{s} \\
 &\geq V_2(\log \eta)^{\sigma_2-1} \phi_q(V_1(\log \eta)^{\sigma_1-1}) \int_{\eta}^e m_{\sigma_2}(s) \frac{ds}{s} \phi_q \left(\int_{\eta}^e m_{\sigma_1}(\tau) \rho(\tau, \kappa(\tau), {}^H \mathfrak{D}_{1+}^{\sigma_2} \kappa(\tau)) \frac{d\tau}{\tau} \right) \\
 &> \frac{L}{F} V_2(\log \eta)^{\sigma_2-1} \phi_q(V_1(\log \eta)^{\sigma_1-1}) \int_{\eta}^e m_{\sigma_2}(s) \frac{ds}{s} \phi_q \left(\int_{\eta}^e m_{\sigma_1}(\tau) \frac{d\tau}{\tau} \right) \\
 &= L.
 \end{aligned}$$

This proves condition (B1) of Theorem 2.1.

Lastly, we will prove the condition (B3) of Theorem 2.1. If $\kappa \in M(\Lambda, L, r)$ and $\|A\kappa\| > K$, for $z \in [1, e]$,

$$\begin{aligned}
 |A\kappa(z)| &= \left| \int_1^e K(z,s) \phi_q \left(\int_1^e H(s,\tau) \rho(\tau, \kappa(\tau), {}^H \mathfrak{D}_{1+}^{\sigma_2} \kappa(\tau)) \frac{d\tau}{\tau} \right) \frac{ds}{s} \right| \\
 &\leq \mu_2 \mu_2^* \phi_q(\mu_1 \mu_1^*) \int_1^e m_{\sigma_2}(s) \phi_q \left(\int_1^e m_{\sigma_1}(\tau) \rho(\tau, \kappa(\tau), {}^H \mathfrak{D}_{1+}^{\sigma_2} \kappa(\tau)) \frac{d\tau}{\tau} \right) \frac{ds}{s} \\
 &= \mu_2 \mu_2^* \phi_q(\mu_1 \mu_1^*) \int_1^e m_{\sigma_2}(s) \frac{ds}{s} \phi_q \left(\int_1^e m_{\sigma_1}(\tau) \rho(\tau, \kappa(\tau), {}^H \mathfrak{D}_{1+}^{\sigma_2} \kappa(\tau)) \frac{d\tau}{\tau} \right),
 \end{aligned}$$

1 and
2

$$\begin{aligned}
 |{}^H\mathcal{D}_{1^+}^{\sigma_2} \kappa(z)| &= \left| -\phi_q \left(\int_1^e H(s, \tau) \rho(\tau, \kappa(\tau), {}^H\mathcal{D}_{1^+}^{\sigma_2} \kappa(\tau)) \frac{d\tau}{\tau} \right) \right| \\
 &\leq \phi_q(\mu_1 \mu_1^*) \phi_q \left(\int_1^e m_{\sigma_1}(\tau) \rho(\tau, \kappa(\tau), {}^H\mathcal{D}_{1^+}^{\sigma_2} \kappa(\tau)) \frac{d\tau}{\tau} \right).
 \end{aligned}$$

7 Thus, we ensure

$$\|A\kappa\| \leq \max\{1, \mu_2 \mu_2^*\} \int_1^e m_{\sigma_2}(s) \frac{ds}{s} \} \phi_q(\mu_1 \mu_1^*) \phi_q \left(\int_1^e m_{\sigma_1}(\tau) \rho(\tau, \kappa(\tau), {}^H\mathcal{D}_{1^+}^{\sigma_2} \kappa(\tau)) \frac{d\tau}{\tau} \right).$$

11 Therefore,

$$\begin{aligned}
 \Lambda(A\kappa) &= \min_{z \in [\eta, e]} A\kappa(z) \\
 &\geq V_2(\log \eta)^{\sigma_2-1} \int_1^e m_{\sigma_2}(s) \phi_q \left(\int_1^e V_1(\log s)^{\sigma_1-1} m_{\sigma_1}(\tau) \rho(\tau, \kappa(\tau), {}^H\mathcal{D}_{1^+}^{\sigma_2} \kappa(\tau)) \frac{d\tau}{\tau} \right) \frac{ds}{s} \\
 &\geq V_2(\log \eta)^{\sigma_2-1} \phi_q(V_1(\log \eta)^{\sigma_1-1}) \int_{\eta}^e m_{\sigma_2}(s) \frac{ds}{s} \phi_q \left(\int_1^e m_{\sigma_1}(\tau) \rho(\tau, \kappa(\tau), {}^H\mathcal{D}_{1^+}^{\sigma_2} \kappa(\tau)) \frac{d\tau}{\tau} \right) \\
 &\geq \frac{N}{\max\{1, \mu_2 \mu_2^*\} \int_1^e m_{\sigma_2}(s) \frac{ds}{s} \} \phi_q(\mu_1 \mu_1^*)} \|A\kappa\| \\
 &> \frac{NK}{\max\{1, \mu_2 \mu_2^*\} \int_1^e m_{\sigma_2}(s) \frac{ds}{s} \} \phi_q(\mu_1 \mu_1^*)} \\
 &\geq L.
 \end{aligned}$$

27 Then, (B3) of Theorem 2.1 is also verified. Then, all conditions of Theorem 2.1 are confirmed, the
28 problem (1.1) has at least three positive solutions such that

$$\|\kappa_1\| < h, \quad L < \Lambda(\kappa_2) \text{ and } h < \|\kappa_3\| \text{ with } \Lambda(\kappa_3) < L.$$

31 □

32 **Example 3.1.** Let us consider the following HFBVP

$$(3.1) \quad \begin{cases}
 {}^H\mathcal{D}_{1^+}^{14/5}(\phi_2({}^H\mathcal{D}_{1^+}^{14/5} \kappa(z))) = \rho(z, \kappa(z), {}^H\mathcal{D}_{1^+}^{14/5} \kappa(z)), & z \in (1, e), \\
 \kappa(1) = \kappa'(1) = {}^H\mathcal{D}_{1^+}^{14/5} \kappa(1) = (\phi_2({}^H\mathcal{D}_{1^+}^{14/5} \kappa(1)))' = 0, \\
 \phi_2({}^H\mathcal{D}_{1^+}^{14/5} \kappa(e)) = \phi_2({}^H\mathcal{D}_{1^+}^{14/5} \kappa(e^{1/2})) + 3\phi_2({}^H\mathcal{D}_{1^+}^{14/5} \kappa(e^{1/4})) + \int_1^e \phi_2({}^H\mathcal{D}_{1^+}^{14/5} \kappa(z)) \frac{dz}{z}, \\
 \kappa(e) = \frac{1}{3} \kappa(e^{1/2}) + \frac{2}{3} \kappa(e^{1/4}) + \int_1^e \kappa(z) \frac{dz}{z},
 \end{cases}$$

41 where $p = 2$, $\sigma_1, \sigma_2 = \frac{14}{5}$, $k = q = 2$, $\eta_1 = e^{1/2}$, $\eta_2 = e^{1/4}$, $\xi_1 = e^{1/2}$, $\xi_2 = e^{1/4}$, $a_1 = 1$, $a_2 = 3$, $b_1 = \frac{1}{3}$,
42 $b_2 = \frac{2}{3}$, $\eta = e^{7/10}$, $g(z) = h(z) = 1$ for $z \in [1, e]$ and

$$\rho(z, \kappa, \iota) = \begin{cases}
 e^{-10z} + \frac{2\kappa}{300} + \frac{|\iota|}{10^5}, & \kappa \in [0, 0.3], \\
 e^{-10z} + (10\kappa - 3)(10^5 - 0.002) + 0.002 + \frac{|\iota|}{10^5}, & \kappa \in [0.3, 0.4], \\
 e^{-10z} + \frac{1000\kappa - 400}{899999.6} + 10^5 + \frac{|\iota|}{10^5}, & \kappa \in [0.4, \infty),
 \end{cases}$$

48 for $z \in [1, e]$, $\iota \in [0, \infty)$. After calculating directly, we obtain $B \approx 5.9268$, $C \approx 8.8956$, $N \approx 0.0002$,
49 $F \approx 0.000004$. If $h = 0.3$, $L = 0.4$, $K = 200000$ and $r = 900000$,

50

$$\rho(z, \kappa, \iota) > \phi_p\left(\frac{L}{F}\right) \approx 100000 \text{ for } (z, \kappa, \iota) \in [e^{7/10}, e] \times [0.4, 200000] \times [-200000, 0],$$

$$\rho(z, \kappa, \iota) \leq \min\left\{\phi_p\left(\frac{r}{B}\right), \phi_p\left(\frac{r}{C}\right)\right\} \approx 101173 \text{ for } (z, \kappa, \iota) \in [1, e] \times [0, 900000] \times [-900000, 0],$$

$$\rho(z, \kappa, \iota) < \min\left\{\phi_p\left(\frac{h}{B}\right), \phi_p\left(\frac{h}{C}\right)\right\} \approx 0.0034 \text{ for } (z, \kappa, \iota) \in [1, e] \times [0, 0.3] \times [-0.3, 0].$$

i.e., ρ satisfies the conditions of Theorem 3.1. By Theorem 2.1, the problem (3.1) has at least three positive solutions κ_j for $j \in \{1, 2, 3\}$ with

$$\|\kappa_1\| < 0.3, \Lambda(\kappa_2) > 0.4, \quad 0.3 < \|\kappa_3\| \text{ with } \Lambda(\kappa_3) < 0.4.$$

Conclusion

This study examined the existence results of fractional boundary value problem by the Hadamard fractional derivative. With the help of the Green's function's properties and the fixed point theory, we obtained the appropriate conditions to ensure the existence results for the HFBVP including p-Laplacian operator. To our knowledge, few researchers worked the HFBVP with p-Laplacian operator. Moreover, the function ρ which consists of the Hadamard derivative operator ${}^H\mathcal{D}_{1+}^{\sigma_2}$ has been studied by only a few researchers.

References

- [1] B. Ahmad, A. Alsaedi, S.K. Ntouyas, J. Tariboon, Hadamard Type Fractional Differential Equations, Inclusions and Inequalities, Cham, Switzerland: Springer International Publishing, 2017.
- [2] B. Ahmad, S.K. Ntouyas, A. Alsaedi, New results for boundary value problems of hadamard- type fractional differential inclusion and integral boundary conditions, Bound. Value Probl., 2013:275. DOI:10.1186/1687-2770-2013-275.
- [3] B. Ahmad, S.K. Ntouyas, A fully Hadamard type integral boundary value problem of a coupled system of fractional differential equations, Fractional Calculus and Applied Analysis, 17.2 (2014): 348-360.
- [4] P.L. Butzer, A. A. Kilbas, J.J. Trujillo, Mellin transform analysis and integration by parts for Hadamard-type fractional integrals, J. Math. Anal. Appl. (2002) 1–15. Vol. 270.
- [5] A. Cabada, G. Wang, Positive solutions of nonlinear fractional differential equations with integral boundary value conditions, J.Math.Anal.Appl., 389 (2012), 403-411.
- [6] C. Ciftci, F. Yuruk Deren, Analysis of p-Laplacian Hadamard fractional boundary value problems with the derivative term involved in the nonlinear term, Mathematical Methods in the Applied Sciences, 2023, DOI:10.1002/mma.9028.
- [7] F.T. Fen, I.Y. Karaca, O.B. Ozen, Positive solutions of boundary value problems for p-Laplacian fractional differential equations, Filomat, 31(5) (2017), 1265–1277.
- [8] R. Garra, F. Polito, On Some Operators Involving Hadamard Derivatives. Integral Transform Spec. Funct. 2013, 24, 773–782.
- [9] R. Garra, F. Mainardi, G. Spada, A generalization of the Lomnitz logarithmic creep law via Hadamard fractional calculus Chaos Solitons Fractals, 102 (2017), pp. 333-338.
- [10] J.R. Graef, L. Kong, Q. Kong, M. Wang, Uniqueness of positive solutions of fractional boundary value problems with non-homogeneous integral boundary condition, Fract. Calc. Appl. Anal., 15:509-528.
- [11] L. Guo, L. Liu, Y. Wang, Maximal and minimal iterative positive solutions for p-Laplacian Hadamard fractional differential equations with the derivative term contained in the nonlinear term, AIMS Mathematics 6.11 (2021): 12583-12598.
- [12] M. Gunendi, I. Yaslan, Positive solutions of higher-order nonlinear multi-point fractional equations with integral boundary conditions, Fractional Calculus and Applied Analysis (2016) Volume 19: Issue 4.
- [13] J. Hadamard, Essai sur l'etude des fonctions donnees par leur developpment de Taylor, J. Pure Appl. Math. 4 (1892), no. 8, 101-186.
- [14] X. Hao, H.Sun, L. Liu, D. Wang, Positive solutions for semipositone fractional integral boundary value problem on the half-line. RACSAM 113, 3055–3067 (2019). <https://doi.org/10.1007/s13398-019-00673-w>.
- [15] X. Jiafa, J. Jiang, D. O'Regan, Positive solutions for a class of p-Laplacian Hadamard fractional-order three-point boundary value problems, Mathematics (2020), 8(3), 308.
- [16] J. Jiang, D. O'Regan, J. Xu, Y. Cui, Positive Solutions for a Hadamard Fractional p-Laplacian Three-Point Boundary Value Problem. Mathematics 2019, 7, 439. <https://doi.org/10.3390/math7050439>.
- [17] A. A. Kilbas, H.M. Srivastava, J.J. Trujillo, Theory and applications of fractional differential equations, in: North-Holland Mathematics Studies, vol. 204, Elsevier Science B.V, Amsterdam, 2006.
- [18] A.A. Kilbas, Hadamard-type fractional calculus, Journal of the Korean Mathematical Society 38.6 (2001): 1191-1204.

- 1
2 [19] V. Lakshmikantham, S. Leela, J.V. Devi, Theory of Fractional Dynamic Systems, Cambridge Academic Publishers,
3 Cambridge, 2009.
- 4 [20] R.W. Leggett, L.R. Williams, Multiple positive fixed points of nonlinear operators on ordered Banach spaces, Indiana Univ.
5 Math. J. 28, 673-688 (1979).
- 6 [21] Y. Li, S. Lin, Positive Solution for the Nonlinear Hadamard Type Fractional Differential Equation with-Laplacian, Journal
7 of Function Spaces and Applications 2013 (2013), <https://doi.org/10.1155/2013/951643>.
- 8 [22] X. Liu, M. Jia, The positive solutions for integral boundary value problem of fractional p-Laplacian equation with mixed
9 derivatives, Mediterranean Journal of Mathematics, 14.2 (2017): 1-17.
- 10 [23] K.S. Miller, B. Ross, An Introduction to the Fractional Calculus and Differential Equations, John Wiley, New York, 1993.
- 11 [24] K. Pei, G. Wang, Y. Sun, Successive iterations and positive extremal solutions for a Hadamard type fractional integro-
12 differential equations on infinite domain, Applied Mathematics and Computation, 312 (2017) 158-168.
- 13 [25] I. Podlubny, Fractional Differential Equations, Academic Press, San Diego, 1999.
- 14 [26] K.R. Prasad, M. Khuddush, P. Veeraiah, Countably many positive solutions for singular R-L fractional order bvp with R-S
15 integral boundary conditions, Nonlinear Studies (2020), Vol. 27, No. 4, pp. 1075-1089, 202.
- 16 [27] F.A. Rihan, Numerical modeling of fractional order biological systems. Abstr Appl Anal 2013.
- 17 [28] S.G. Samko, A.A. Kilbas, O.I. Marichev, Fractional Integrals and Derivatives, Theory and Applications, Gordon and
18 Breach, Yverdon, 1993.
- 19 [29] T. Senlik Cerdik, F. Yoruk Deren, New results for higher-order Hadamard-type fractional differential equations on the half
20 line, Mathematical Methods in the Applied Sciences 45.4 (2022): 2315-2330.
- 21 [30] V.E. Tarasov, Fractional dynamics: application of fractional calculus to dynamics of particles, fields and media. Heidel-
22 berg/Beijing: Springer/Higher Education Press; 2010.
- 23 [31] B.M. Vintagre, I. Podlybni, A. Hernandez, V. Feliu, Some approximations of fractional order operators used in control
24 theory and applications. Fract Calc Appl Anal 2000;3(3):231-48.
- 25 [32] G. Wang, K. Pei, R.P. Agarwal, L. Zhang, B. Ahmad, Nonlocal Hadamard fractional boundary value problem with
26 Hadamard integral and discrete boundary conditions on a half-line, Journal of computational and applied mathematics, 343
27 (2018): 230-239.
- 28 [33] G. Wang, T. Wang, On a nonlinear Hadamard type fractional differential equation with p-Laplacian operator and strip
29 condition, J. Nonlinear Sci. Appl. 9 (2016): 5073-5081.
- 30 [34] Y. Wang, S. Liang, Q. Wang, Existence results for fractional differential equations with integral and multi-point boundary
31 conditions, Boundary Value Problems (2018) 2018:4.
- 32 [35] W. Yang, Positive solutions for singular Hadamard fractional differential system with four-point coupled boundary
33 conditions, Journal of Applied Mathematics and Computing, 49.1 (2015): 357-381.
- 34 [36] A. Yang, H. Wang, Positive solutions for higher-order nonlinear fractional differential equation with integral boundary
35 condition, Electronic Journal of Qualitative Theory of Differential Equations, 2011, No. 1, 1-15.
- 36 [37] F. Yoruk Deren, T. Senlik Cerdik, Extremal Positive Solutions for Hadamard Fractional Differential Systems on an Infinite
37 Interval. Mediterr. J. Math. 20, 158 (2023), <https://doi.org/10.1007/s00009-023-02369-3>.
- 38 [38] F. Yoruk Deren, T. Senlik Cerdik, R.P. Agarwal, Existence Criteria of Positive Solutions for Fractional p-Laplacian
39 Boundary Value Problems, Filomat, (2020) 34, no 11.
- 40 [39] X. Zhao, Y. Liu, H. Pang, Iterative positive solutions to a coupled fractional differential system with the multistrip and
41 multipoint mixed boundary conditions, Advances in Difference Equations (2019) 2019:389.

42 DEPARTMENT OF MATHEMATICS, İSTANBUL BEYKENT UNIVERSITY, TÜRKİYE

43 *Email address:* tugbacerdik@beykent.edu.tr

44

45

46

47

48

49

50