

SOME RESULTS ON NONNEGATIVELY CURVED MANIFOLDS

WALTER A. POOR, JR.

Cheeger and Gromoll have proven [1] that every complete Riemannian manifold M of nonnegative sectional curvature has a compact totally convex totally geodesic submanifold S , which they called a *soul* of M . Furthermore, they have proven that M is homeomorphic to the normal bundle $\nu(S)$ of S in the tangent bundle TM of M ; for M simply connected and of dimension greater than five, using the open h -cobordism theorem they proved that M and $\nu(S)$ are diffeomorphic. This result greatly simplifies the study of noncompact complete manifolds of nonnegative curvature since it partially reduces the problem to the study of vector bundles over compact manifolds.

Our first result will be to prove the differentiable soul theorem, that is, we will prove that M and $\nu(S)$ are diffeomorphic (without dimension or connectivity assumptions). The diffeomorphism to be constructed will be rather closely adapted to the geometric situation.

In preparation for the statement of the second result, we recall that if χ is the Euler form on an oriented complete even-dimensional Riemannian manifold M , then $\int_M \chi$ (if it exists) is called the total curvature of M . It has long been known [4] that the total curvature of a complete surface of nonnegative curvature is less than or equal to one; since in this case $K = 2\pi\chi$, where K is the sectional curvature, the result is usually written

$$\int_M K \leq 2\pi .$$

Our second result will be to prove that for an oriented complete nonnegatively curved Riemannian manifold M of dimension 4, the total curvature exists and is bounded between zero and the Euler characteristic of M . This obviously generalizes both the above result for surfaces and the generalized Gauss-Bonnet theorem for four-dimensional oriented compact manifolds of nonnegative curvature. In addition, for oriented complete nonnegatively curved Riemannian manifolds of even dimension greater than 4, the argument reduces the question to the so-called algebraic Hopf conjecture. This implies a partial result in dimension 6.

The results are related in the sense that both rest heavily on the global existence of certain convex functions; the analysis of these functions, which are in general only continuous, is the central technical aspect of the work.

Finally we give some applications.

This paper presents the essence of the author's doctoral dissertation at the State University of New York at Stony Brook under the direction of Detlef Gromoll. Here the author would like to express his thanks to Professor Gromoll and also to the State University of New York at Stony Brook for its support.

1. Preliminaries

Given a C^∞ manifold $M = M^n$, $TM \xrightarrow{\pi} M$ will denote the tangent bundle. For p in M , $M_p := \pi^{-1}(p)$ is the tangent space to M at p . $\Gamma\xi$ will be the set of C^∞ sections in the bundle $\xi \rightarrow M$; particular examples are $FM := \Gamma(M \times \mathbb{R}) = \{C^\infty \text{ functions on } M\}$, and $\mathfrak{b}M := \Gamma TM = \{C^\infty \text{ vector fields on } M\}$. ∇ will always denote the covariant derivative operator $\mathfrak{b}M \times \Gamma\xi \rightarrow \Gamma\xi$ in whatever bundle ξ we are discussing; \exp will be the exponential map of ∇ .

If M is a Riemannian manifold, then $\langle \cdot, \cdot \rangle$ will be the first fundamental form, and $\nabla: \mathfrak{b}M \times \mathfrak{b}M \rightarrow \mathfrak{b}M$ will denote the Levi-Civita connection. $T_1M := \{u \in TM \mid \|u\| = 1\}$ is the (unit) tangent sphere bundle, and for p in M , $S_p^{n-1} := M_p \cap T_1M$ is the (unit) tangent sphere to M at p . Given nonzero u and v in M_p , $\sphericalangle(u, v)$ will denote the angle in $[0, \pi]$ between u and v .

For p and q in M , $\rho(p, q)$ is the metric distance from p to q . For $r > 0$ and p in M , $B_r(p)$ is the open metric ball $\{q \in M \mid \rho(p, q) < r\}$ with boundary equal to the metric sphere $S_r(p) := \{q \in M \mid \rho(p, q) = r\}$. A geodesic is said to be *normal* if it is parametrized by arc length. If there exists a unique normal geodesic $c: [0, \rho(p, q)] \rightarrow M$ from p to q , it will be called the *minimal connection* from p to q .

Let χ_M be the Euler characteristic of M .

In addition to the construction of a soul of M , we will also need a few of the technical facts which Cheeger and Gromoll used in the proof of the continuous soul theorem; this section consists mainly of the necessary material from [1], which we reformulate for our purposes.

A nonempty subset C of M will be said to be *totally convex* if for any p and q in C and any geodesic $c: [0, 1] \rightarrow M$ from p to q , c lies in C . C is *strongly convex* if for any p and q in C , there is a minimal connection c from p to q and c lies in C .

Lemma 1. *Let C be a closed totally convex subset of an arbitrary Riemannian manifold M . Then C has the structure of an imbedded k -dimensional submanifold of M with smooth totally geodesic interior N and (possibly nonsmooth) boundary $\partial C = \bar{N} - N$.*

Cheeger and Gromoll actually proved a stronger version of the lemma, but this is all we need.

For a closed totally convex set C and for $a \geq 0$, let ${}^a C := \{p \in M \mid \rho(p, C) \leq a\} = \overline{\bigcup_{p \in C} B_a(p)}$; in addition, if $\partial C \neq \emptyset$, let $C^a := \{p \in C \mid \rho(p, \partial C) \geq a\}$. Notice that C^a may be empty.

Lemma 2. *Let M have nonnegative curvature, and let C be a closed totally convex subset of M with nonempty boundary. If C^a is nonempty, then C^a is a closed totally convex set. If $a_{\max} := \max \{a \in \mathbb{R} \mid C^a \neq \emptyset\} = \max_{p \in C} \rho(p, \partial C)$, then $\dim C^{a_{\max}} < \dim C$.*

Lemma 3 (standard). *For every compact subset K of M there is a number $\varepsilon > 0$ such that for all p in K and r in $(0, \varepsilon)$:*

- (1) *the metric ball $B_r(p)$ is the diffeomorphic strongly convex image under \exp_p of the open Euclidean ball of radius r about 0 in M_p ;*
- (2) *if $c: [0, u] \rightarrow B_r(p)$ is a nonconstant geodesic, $c_0: [0, 1] \rightarrow B_r(p)$ is the minimal geodesic from p to $c(0)$, and $\langle \dot{c}(0), \dot{c}_0(1) \rangle \geq 0$, then the function $s \mapsto \rho(c(s), p)$ is strictly increasing on $[0, u]$.*

From now on, we will always assume that M is a complete noncompact Riemannian manifold of nonnegative curvature.

We will only use parts (1) and (2) of the next lemma; part (3) is presented because it gives the flavor of the proof of the continuous soul theorem and is a model for some of the work we must do.

Lemma 4. *Let K be a compact totally convex set in M with nonempty boundary, and let $0 < \varepsilon \leq a_{\max}$, where ε is a constant as in Lemma 3, and $a_{\max} = \max \{\rho(p, \partial K) \mid p \in K\}$. Then*

- (1) *there exists $\delta \in (0, \varepsilon)$ such that $\rho(p, K^{a'}) < \varepsilon$ for all p in K^a if $0 \leq a \leq a' \leq a_{\max}$ and $a' - a < \delta$;*
- (2) *there exists a “continuous geodesic contraction” of K^a onto $K^{a'}$ if $0 \leq a \leq a' \leq a_{\max}$ and $a' - a < \delta$;*
- (3) *given $0 \leq a < a' < a_{\max}$ such that $a' - a < \delta$, there is a homeomorphism of $\partial K^a \times [0, 1]$ onto $K^a - \text{int } K^{a'}$.*

We will now describe the constructions in parts (2) and (3) explicitly.

Given p in $K^a - K^{a'}$, there is a closest point q in $K^{a'}$; q is unique by the first variation formula and Lemma 3. Define $h(p) := q$. h is continuous by uniqueness of q and the continuity of $p \mapsto \rho(p, K^{a'})$. Thus we can contract K^a onto $K^{a'}$ along the minimal connections from points on ∂K^a to the closest points on $K^{a'}$. We will occasionally refer to h as “the contraction” onto $K^{a'}$.

Let $a' < a'' < a_{\max}$ so that $a'' - a < \delta$. Let \tilde{h} be the contraction onto $K^{a''}$, and for p in ∂K^a let $g_p: [0, 1] \rightarrow K^a$ be the unique minimal geodesic such that $g_p(0) = p$ and $g_p(1) = \tilde{h}(p)$. Define $\Phi: \partial K^a \rightarrow (0, 1)$ such that $g_p(\Phi(p)) \in \partial K^{a'}$; Φ is continuous. Define $F: \partial K^a \times [0, 1] \rightarrow K^a - \text{int } K^{a'}$ such that $(p, t) \mapsto g_p(\Phi(p)t)$. F is the homeomorphism.

Let $\{C_t\}_{t \geq 0}$ be a family of nonempty compact totally convex sets such that

- (1) $\dim C_t \begin{cases} < \dim M & \text{if } t = 0, \\ = \dim M & \text{if } t > 0; \end{cases}$

- (2) $t_2 \geq t_1 \Rightarrow C_{t_1} = C_{t_2}^{t_2-t_1}$,
- (3) $\bigcup_{t \geq 0} C_t = M$.

Note that for all $t > 0$, $C_t^t = C_0$.

Cheeger and Gromoll constructed this family as follows: a normal geodesic $c: [0, \infty) \rightarrow M$ is called a *ray* if each segment of c is minimal between its end-points. It is straightforward to see that every point in any noncompact complete Riemannian manifold has at least one ray through it. If c is a ray, let $B_c := \bigcup_{t > 0} B_t(c(t))$. B_c is called the *open half-space* with respect to c , and its complement $M - B_c$ is totally convex in our case.

Now pick p in M . For $t \geq 0$, let $L_t := \bigcap_c (M - B_{c_t})$, where the intersection is taken over all rays c emanating from p and c_t is the ray determined by $c_t(s) := c(t + s)$, $0 \leq s < \infty$. Each L_t is compact and totally convex, and p is in ∂L_0 .

If $\dim L_0 < \dim M$, then let $C_t = L_t$ for all $t \geq 0$. If $\dim L_0 = \dim M$, let $a_0 := \max_{q \in L_0} \rho(q, \partial L_0)$ and

$$C_t = \begin{cases} L_0^{a_0-t} & \text{for } 0 \leq t \leq a_0, \\ L_{t-a_0} & \text{for } a_0 < t. \end{cases}$$

If $\partial C_0 = \emptyset$, then let $S = C_0$. If $\partial C_0 \neq \emptyset$, let $K_0 := C_0$ and $a_0 := \max_{p \in K_0} \rho(p, \partial K_0)$; thus $K_1 := K_0^{a_0}$ is a totally convex set of lower dimension than K_0 . Iteration yields a sequence $K_0 \supset K_1 \supset \dots \supset K_J$ of compact totally convex sets such that $K_{i+1} := K_i^{a_i}$, where $a_i := \max \{ \rho(p, \partial K_i) \mid p \in K_i \}$, and such that $\partial K_J = \emptyset$; $J < \dim M$. Let $S := K_J$.

Cheeger and Gromoll called S a *soul* of M ; it is unique up to the choice of p . They proved that M is homeomorphic to the normal bundle of S , and stated that the construction could be smoothed out so as to give a diffeomorphism. We have carried out this process in detail, and consider this approach to be unnecessarily complicated. Here we present a modified construction which is simpler and also more satisfying geometrically.

2. The differentiable soul theorem

Let M be a complete noncompact Riemannian manifold of nonnegative sectional curvature.

Theorem A. *M is diffeomorphic to $\nu(S)$.*

The exponential map restricted to $\nu(S)$ is a smooth surjective map onto M , which fails to be injective. The map we obtain is modelled (roughly speaking) on the exponential map, and in fact we may think of it as being a smooth bijective “bent exponential map”.

We will actually prove the following theorem; by standard arguments from differential topology it is equivalent to Theorem A. For these arguments, as

well as for several arguments used in the proof of the following theorem, we refer the reader to references [8] and [10].

Theorem B. *Let $M_0 = S$. Then there exists an increasing sequence M_1, M_2, \dots of nonempty compact submanifolds with boundary such that*

- (1) M_1 is diffeomorphic to a disc bundle in $\nu(S)$,
- (2) $M_{i+1} - \text{int } M_i$ is diffeomorphic to $\partial M_i \times [0, 1]$ for $i \geq 1$, and
- (3) M is the (disjoint) union $S \cup (\cup_{i \geq 0} (M_{i+1} - M_i))$.

This result will be proven in several steps. First we make a few remarks.

Recall that we have an expanding sequence $\{C_i\}_{i \geq 0}$ of nonempty totally convex compact sets and a decreasing sequence $C_0 = K_0 \supset K_1 \supset \dots \supset K_J = S$ as constructed in § 1.

Cheeger and Gromoll construct a continuous “broken exponential map” by locally adjusting the exponential map restricted to $\nu(S)$ to get a locally injective map; by using the above sequences of sets to keep track of the resulting “continuous flow”, they end up with a homeomorphism from $\nu(S)$ to M .

The local arguments are of three types. First, given a sufficiently small $r > 0$ they construct a homeomorphism between ${}^r K_0$ and the disc bundle $\nu_r(S)$ of radius r . This argument takes into account the difference in dimension between K_i and K_{i+1} at each step. Then they use another argument to show that ${}^r C_0$ is homeomorphic to $C_r = C_1^{1-r}$. The third argument uses the exhaustive sequence $\{C_i\}$ to extend this homeomorphism globally.

Simply smoothing out these constructions directly is not very efficient, and so we have modified the whole approach.

First we present an interesting fact.

Lemma 5. *Let K be a compact totally convex subset of M . Let $d: M \rightarrow R$ such that $d(p) := \rho(p, K)$. Then there exists an open neighborhood U of K such that $d|_{U-K}$ is C^1 .*

Proof. Consider the set ${}^1 K$. Choose ε in $(0, 1]$ for ${}^1 K$ as in Lemma 3. Then ε also satisfies the conclusion of Lemma 3 for the smaller set ${}^\varepsilon K$. Let $U := \text{int } {}^\varepsilon K$. Let $h: U \rightarrow K$ be the “continuous geodesic contraction” onto K given in Lemma 4(3); $h(p)$ is the unique point in K such that $\rho(p, K) = \rho(p, h(p)) = d(p)$.

Let $p \in A := U - K$. Define $f_p: A \rightarrow R$ such that $f_p(q) := \rho(q, h(p))$. Clearly we have $f_p \geq d$; in addition, $f_p(p) = d(p)$, and f_p is C^∞ on a neighborhood of p in A . Let $\Delta \subset M_{h(p)}$ be the supporting hyperplane for K orthogonal to the minimal connection from p to $h(p)$ (cf. [1]). Let $\Delta_\varepsilon := \{v \in \Delta \mid \|v\| < \varepsilon\}$. Then $\exp_{h(p)} \Delta_\varepsilon$ is a local supporting hypersurface for K . For $q \in A$, let $g_p(q) := \rho(q, \exp_{h(p)} \Delta_\varepsilon)$. Then $g_p \leq d$, $g_p(p) = d(p)$, and g_p is C^∞ on a neighborhood of p in A .

By the squeezing principle, d is (once) differentiable at p , and in fact

$$d_{*|p} = f_{p*|p} = g_{p*|p} .$$

To see this, let c be a C^∞ curve in A such that $c(0) = p$. For small positive t ,

$$\frac{1}{t}[g_p \circ c(t) - g_p(p)] \leq \frac{1}{t}[d \circ c(t) - d(p)] \leq \frac{1}{t}[f_p \circ c(t) - f_p(p)],$$

and therefore,

$$(g_p \circ c)'(0) \leq \lim_{t \rightarrow 0^+} \frac{1}{t}[d \circ c(t) - d(p)] \leq (f_p \circ c)'(0).$$

Similarly,

$$(f_p \circ c)'(0) \leq \lim_{t \rightarrow 0^-} \frac{1}{t}[d \circ c(t) - d(p)] \leq (g_p \circ c)'(0).$$

Thus $(f_p \circ c)'(0) = (g_p \circ c)'(0)$, and therefore $(d \circ c)'(0)$ exists and equals $(f_p \circ c)'(0)$.

Since the assignment $p \mapsto h(p)$ is continuous, so is $p \mapsto \nabla f_p(p) = \nabla d(p)$ (see the remark below); thus d is C^1 on $U - K$.

Remark. Given $p \in U - K$, let γ_p be the minimal connection from p to $h(p) \in K$. Then

$$\nabla d \circ \gamma_p(t) = \nabla f_p \circ \dot{\gamma}_p(t) = -\dot{\gamma}_p(t),$$

for $0 \leq t < d(p)$. In particular, ∇d is a continuous autoparallel unit vector field on $U - K$. In addition, $h(p) = \exp_p(-d(p)\nabla d(p))$.

Lemma 6. Let K be a compact totally convex subset of M with nonempty boundary, and let $a_{\max} := \max\{\rho(p, \partial K) \mid p \in K\}$. Choose $\varepsilon \in (0, 1]$ as in the proof of Lemma 5, and choose $r \in (0, \varepsilon)$. Define $d: [0, a_{\max}] \times {}^r K \rightarrow R$ such that $d(a, p) = d_a(p) := \rho(p, K^a)$, and set $(\nabla d)(a, p) = \nabla d_a(p)$. Then ∇d is uniformly continuous on the set

$$A := \{(a, p) \in [0, a_{\max}] \times {}^r K \mid p \in {}^r[K^a] - \text{int } {}^{r/4}[K^a]\}.$$

Proof. First we prove that d is continuous on $[0, a_{\max}] \times {}^r K$; we already know that for each a , d_a is continuous on ${}^r K$ and C^1 on ${}^r[K^a] - K^a$.

Let $\{(a_i, p_i)\}$ be a sequence in $[0, a_{\max}] \times {}^r K$ converging to (a, p) . Choose a sequence $\{q_i \in K^{a_i}\}$ such that $d(a_i, p_i) = \rho(p_i, q_i)$. Let q be a limit point of this sequence; such a q exists since K is compact. Then $\lim_i d(a_i, p_i) = \lim_i \rho(p_i, q_i) = \rho(p, q) \geq d(a, p)$. Now choose $\bar{q} \in K^a$ such that $\rho(p, \bar{q}) = d(a, p)$, and let $\{\bar{q}_i\}$ be a sequence such that $\bar{q}_i \in K^{a_i}$ and $\lim_i \bar{q}_i = \bar{q}$. For each i , $d(a_i, p_i) \leq \rho(p_i, \bar{q}_i)$, and therefore $\lim_i d(a_i, p_i) \leq \lim_i \rho(p_i, \bar{q}_i) = \rho(p, \bar{q}) = d(a, p)$. Thus d is continuous.

But then since $A = d^{-1}[\frac{1}{4}r, r]$, A must be compact. Define $h: A \rightarrow K$ such

that $h(a, p)$ is the unique point q in K^a for which $\rho(p, q) = d(a, p)$. By continuity of d , h is also continuous. But

$$h(a, p) = \exp_p (-d(a, p)\nabla d(a, p)) ,$$

and thus ∇d is continuous on the set A .

We are now ready to prove Theorem B.

Choose $\varepsilon_1 \in (0, 1)$ for the set C_1 as in Lemma 3. Choose $\tilde{\delta}_1 \in (0, \varepsilon)$ as in Lemma 4. Let $0 < r < \tilde{\delta}_1$, and choose δ_1 for $\varepsilon = \frac{1}{4}r$ and the set C_1 as in Lemma 4. Recall that $K_0 = C_0$ and $K_{i+1} = K_i^{a_i}$, where $a_i = \max \{\rho(p, \partial K_i) \mid p \in K_i\}$.

Since ∇d is uniformly continuous on the set

$$A := \{(a, p) \in [0, a_0] \times {}^r K_0 \mid p \in {}^r [K_0^a] - \text{int } {}^{r/4} [K_0^a]\} ,$$

there exists $\xi > 0$ such that if (a, p) and $(b, p) \in A$ and $|a - b| < \xi$, then $\angle(\nabla d_a(p), \nabla d_b(p)) < \frac{1}{8}\pi$; we may assume that $\xi < \delta_1$.

Choose numbers $0 = b_0 < b_1 < \dots < b_k = a_0$ such that $b_{i+1} - b_i < \xi$, $0 \leq i \leq k - 1$. Let $L_0 = K_0$, $L_1 = K_0^{b_1}$, \dots , $L_k = K_0^{b_k} = K_1$. Continue similarly with K_1, \dots, K_{j-1} (choosing a different value of ξ at each step, if necessary) to get a decreasing sequence

$$K_0 = L_0 \supset L_1 \supset \dots \supset L_k = K_1 \supset \dots \supset K_2 \supset \dots \supset L_\alpha = S$$

such that for $1 \leq i \leq \alpha$, $L_i = L_{i-1}^{a_i}$ for some sufficiently small $a_i > 0$, and

$${}^{r/4} L_{i-1} \subset {}^{r/2} L_i \subset {}^{r/2} L_{i-1} .$$

Let $d_i: {}^r L_i \rightarrow R$ be the distance function from ${}^r L_i$ to L_i . Then each d_i is C^1 off of L_i , and if $p \in {}^{r/2} L_{i-1} - {}^{r/2} L_i$, then $\angle(\nabla d_i(p), \nabla d_{i-1}(p)) < \frac{1}{8}\pi$.

We can choose a C^∞ function \tilde{d}_i on ${}^r L_i$ such that

- (1) $|d_i - \tilde{d}_i| < \frac{1}{8}r/\alpha$,
- (2) $\nabla \tilde{d}_i \neq 0$ and $\angle(\nabla d_i, \nabla \tilde{d}_i) < \frac{1}{8}\pi$ off of ${}^{r/8} L_i$.

For $1 \leq i \leq \alpha$, let N_i be the compact submanifold $\left\{ p \in {}^r L_i \mid \tilde{d}_i(p) \leq \frac{4\alpha - 2i + 1}{4\alpha} r \right\}$ with C^∞ boundary $H_i = \tilde{d}_i^{-1} \left(\frac{4\alpha - 2i + 1}{4\alpha} r \right)$. Then

$${}^o L_i \subset \text{int } N_i \subset N_i \subset \text{int } {}^r L_i ,$$

where $\sigma = (\frac{1}{2}(2\alpha - i)/\alpha)r$ and $\tau = (\frac{1}{2}(2\alpha - i + 1)/\alpha)r$. Also, $\nabla \tilde{d}_i \perp H_i$. Thus H_i is also transverse to $\nabla \tilde{d}_{i-1}$, $1 \leq i \leq \alpha$, by the way we chose everything. Therefore the gradient flow of \tilde{d}_{i-1} determines (after reparametrization) a diffeomorphism $H_i \times [0, 1] \cong N_{i-1} - \text{int } N_i$, $2 \leq i \leq \alpha$.

In addition, $\nabla \tilde{d}_\alpha$ determines a diffeomorphism between $\partial({}^{r/2} S) \times [0, 1]$ and

$N_\alpha - \text{int } \tau^{1/2}S$. By the choice of r , the exponential map is a diffeomorphism between the normal disc bundle $\nu_{\tau^{1/2}}(S)$ of radius $\frac{1}{2}r$ and $\tau^{1/2}S$.

Set

$$M_1 := \tau^{1/2}S, \quad M_2 := N_\alpha, \dots, \quad M_{\alpha+1} := N_1.$$

$M_{\alpha+1}$ is a compact submanifold with boundary such that

$$\tau^1 L_1 \subset M_{\alpha+1} \subset \tau L_1,$$

where $r_1 = (\frac{1}{2}(2\alpha - 1)/\alpha)r$. We will now show how to find the larger submanifolds required by the theorem.

Choose $\varepsilon_2 \in (0, 1)$ for the set C_2 as in Lemma 3, $\tilde{\delta}_2 \in (0, \varepsilon_2)$ for C_2 as in Lemma 4, and let $0 < r_2 < \min\{\tilde{\delta}_2, r_1\}$; finally, choose δ_2 for $\varepsilon = r_2$ as in Lemma 4 for the set C_2 .

Let κ be an integer greater than $1/\delta_2$. Then $(r_1 - r_2)/\kappa < 1/\kappa < \delta_2$. For $j = 0, 1, \dots, \kappa$, let $\eta(j) := r_1 - j(r_1 - r_2)/\kappa$. As j runs from 0 to κ , $\eta(j)$ decreases from r_1 to r_2 ; let $\ell(j) = j/\kappa$. We have the following inclusions:

$$C_{\ell(j-1)} \subset C_{\ell(j)} \subset \eta^{(j)}C_{\ell(j-1)} \subset \eta^{(j-1)}C_{\ell(j-1)} \subset \eta^{(j)}C_{\ell(j)}.$$

By the way we have chosen everything, we can now use the appropriate distance functions as before to obtain compact submanifolds M_i with boundary for $i = \alpha + 2, \dots, \alpha + \kappa - 1$, such that

$$\eta^{(j)}C_{\ell(j)} \subset M_{\alpha+j+2} \subset \eta^{(j+1)}C_{\ell(j+1)}$$

for $j = 0, \dots, \kappa - 1$, and having the properties required in the theorem.

$M_{\alpha+\kappa-1}$ is now close to $\tau^2 C_1 \subset C_2$. Pick all the appropriate constants for C_3 and iterate. This completes the argument.

3. Total curvature of M

In this section we will prove that for a complete noncompact oriented Riemannian manifold M of dimension 4, the total curvature of M exists and is bounded between 0 and the Euler characteristic of M .¹ In addition, if the dimension of M is $2k$, $k \geq 3$, then our argument reduces the problem to the algebraic Hopf conjecture; in particular, this implies a partial result in dimension 6.

The calculus we employ is not new, but does not appear in the literature in this exact form. We start by presenting classical material to help fix the notation.

Let π^*TM be the pullback of TM over itself under the map π :

¹ This result has also been proved by R. Walter using a different approach.

$$\begin{array}{ccc}
 \pi^*TM & \longrightarrow & TM \\
 \downarrow & & \downarrow \pi \\
 TM & \xrightarrow{\pi} & M .
 \end{array}$$

We may realize π^*TM as the vertical bundle VM in TTM : for $v \in TM$, the fiber in VM over v is $\pi_*|_v^{-1}(0) \subset TTM$. For u in M_p , let $u_\#$ denote the canonical identification of M_p with $(M_p)_u \subset VM$ by parallel transport, that is, for $v \in M_p$, $u_\#v := \left. \frac{d}{dt} \right|_{t=0} (u + tv)$. Then we may identify π^*TM with $V M$ using $\#$

as follows: for $(u, v) \in \pi^*TM$, u and v are in the same tangent space to M , so $u_\#v$ makes sense, and $(u, v) \leftrightarrow u_\#v$ is the required bundle identification.

Suppose $\xi \rightarrow M$ is a vector bundle over M ; let $A^j\xi$ be the associated bundle over M for which the fiber at p in M is the vector space of alternating j -linear forms on the fiber ξ_p in ξ over p .

Given $\omega \in A^j\xi$ and $\mu \in A^\ell\xi$, both over the same point p in M , we have as usual the $(j + \ell)$ -form

$$\omega \wedge \mu := \frac{1}{j! \ell!} \sum_{\sigma} \text{sgn}(\sigma) (\omega \otimes \mu) \circ \sigma ,$$

where the sum is taken over the symmetric group on $j + \ell$ letters.

Recall that $\mathfrak{b}M = \Gamma TM$, and let $A^r(\xi)$ be the FM -module of alternating r -linear forms on $\mathfrak{b}M$ with values in $\Gamma\xi$. If $\nabla: \mathfrak{b}M \times \Gamma\xi \rightarrow \Gamma\xi$ is a covariant derivative operator in ξ , then we have the standard exterior derivative operator $d^\nabla: A^r(\xi) \rightarrow A^{r+1}(\xi)$ with respect to ∇ defined by

$$[d^\nabla \omega](X) := \nabla_X \omega$$

for $\omega \in A^0(\xi)$, and by

$$\begin{aligned}
 [d^\nabla \omega](X_0, \dots, X_r) &:= \sum_i (-1)^i \nabla_{X_i} (\omega(X_0, \dots, \hat{X}_i, \dots, X_r)) \\
 &+ \sum_{i < j} (-1)^{i+j} \omega([X_i, X_j], X_0, \dots, \hat{X}_i, \dots, \hat{X}_j, \dots, X_r)
 \end{aligned}$$

for $\omega \in A^r(\xi)$, $r \geq 1$.

Notice that for an arbitrary covariant derivative ∇ in an arbitrary bundle ξ , $d^\nabla d^\nabla$ does not have to vanish; a bundle ξ is said to be *flat* with respect to ∇ if and only if $d^\nabla d^\nabla = 0$. Example: $M \times R$ is flat with respect to its canonical connection.

We will be using modules of the form $A^r(A^j\xi)$, and we need a sort of double wedge product. Given $\phi \in A^r(A^j\xi)$, $\psi \in A^s(A^\ell\xi)$, and $X_1, \dots, X_{r+s} \in \mathfrak{b}M$, we define the form $\phi \wedge \psi(X_1, \dots, X_{r+s}) \in \Gamma A^{j+\ell}\xi$ to be $(r!s!)^{-1} \sum_{\sigma} \text{sgn}(\sigma) \phi(X_{\sigma_1},$

$\dots, X_{\sigma_r}) \wedge \psi(X_{\sigma_{r+1}}, \dots, X_{\sigma_{r+s}})$, where as usual the sum is taken over the symmetric group on $r + s$ letters.

d^r and \wedge have the following usual properties:

- (1) d^r is natural: $f^*(d^r\omega) = d^{r*}f^*\omega$.
- (2) \wedge is associative.
- (3) For $\phi \in A^r(A^j\xi)$ and $\psi \in A^s(A^k\xi)$, $\phi \wedge \psi = (-1)^{rs+jk}\psi \wedge \phi$.
- (4) For ϕ and ψ as in (3), $d^r(\phi \wedge \psi) = (d^r\phi) \wedge \psi + (-1)^r\phi \wedge d^r\psi$.

We will use the canonical identification of functions on M^n with n -forms to convert elements of $A^r(A^nTM)$ into ordinary differential r -forms on M ; this will allow us to integrate such (r, n) -forms without further comment. More generally, any (r, r) -form restricted to an r -plane can be thought of as a number; an example of this occurs in the next paragraph.

A curvature form on a Euclidean vector space \mathcal{V} is a 2-form Ω on \mathcal{V} with values in 2-forms on \mathcal{V} such that Ω satisfies the Jacobi identity in the first three arguments. Assume that \mathcal{V} is even-dimensional, $\dim \mathcal{V} = 2j$. The algebraic Hopf conjecture is that if $-\Omega$ is nonnegative (resp. positive) on all 2-planes in \mathcal{V} , then $(-\Omega)^j$ is nonnegative (resp. positive) on \mathcal{V} .

Notice that $-\Omega$ restricted to 2-planes in \mathcal{V} is just the usual sectional curvature function. The conclusion of the Hopf conjecture is true under certain extra algebraic conditions which have no geometric significance in our situation.

Define the curvature form $\Omega \in A^2(A^2TM)$ on M with respect to ∇ by

$$\Omega(X, Y)(U, V) := \langle d^r d^r U(X, Y), V \rangle = \langle R(X, Y)U, V \rangle$$

for vector fields X, Y, U and V on M , where R is the usual curvature tensor on M . The Bianchi identity now takes the form

$$d^r \Omega = 0 .$$

Pull \langle , \rangle and ∇ back to $\pi^*TM = VM$, and let $A = A(VM) = A(\pi^*TM)$. Let I be the "position vector field" on TM along π :

$$\begin{array}{ccc} VM & \longrightarrow & TM \\ \downarrow & \nearrow I & \downarrow \pi \\ TM & \longrightarrow & M \end{array} \quad I(v) := v .$$

Let $\eta \in A^0(A^1)$ be the canonical 1-form defined by

$$\eta(s) := \langle \pi^*I, s \rangle$$

for s in VM . Suppose that $s = u_{\#}v$; then

$$\eta(s) = \langle \pi^*I, s \rangle|_u = \langle (\pi^*D)u, u_{\#}v \rangle = \langle u_{\#}u, u_{\#}v \rangle = \langle u, v \rangle$$

by the definition of \langle , \rangle on TM as the pullback of \langle , \rangle on M .

Next we define the connection 1-form on VM by

$$\theta := d^v \eta \in A^1(A^1) .$$

For $v \in TTM$ and $s \in VM$, $\theta(v)(s) = (\nabla_v \eta)(s) = \langle \nabla_v \pi^* I, s \rangle$. We now mention some facts which will be needed below.

(1) If v is vertical on TM , then $\nabla_v I = v$, and therefore $\theta(s)(t) = \langle s, t \rangle$ for s and t in VM .

(2) If v is tangent to T_1M , then $\theta(v)(\pi^* I) = 0$.

(3) Consider $d^v \theta \in A^2(A^1)$; for u and $v \in TTM$ and $s \in VM$,

$$d^v \theta(u, v)(s) = \pi^* \Omega(u, v)(\pi^* I, s) .$$

(4) $\pi^* \Omega(u, v) = 0$ for vertical vectors $u, v \in TTM$.

For $1 \leq i \leq k$, let

$$\begin{aligned} \Pi_i &:= \eta \wedge \theta^{2i-1} \wedge \pi^* \Omega^{k-i} \in A^{n-1}(A^n) , \\ c_i &:= \frac{(-1)^{k+i+1}(i-1)!}{\pi^k 2^{k-i+1}(k-i)!(2i-1)!} . \end{aligned}$$

Define

$$\Pi := \sum_{i=1}^k c_i \Pi_i .$$

The key points in the classical Chern argument [2] are the following: first of all, on T_1M ,

$$d^v \Pi = \frac{(-1)^k}{k! (2\pi)^k} \pi^* \Omega^k ;$$

second, on the unit sphere S_p^{n-1} in M_p , Π is a constant times the volume form, and in particular

$$\int_{S_p^{n-1}} \Pi = -1 .$$

These equalities are proven by choosing special bases and then applying the definitions and facts (1)–(4).

Define the Euler form on M by

$$\chi := \frac{(-1)^k}{k! (2\pi)^k} \Omega^k \in A^n(A^n) .$$

$\pi^* \chi$ is exact on T_1M . χ is often called the Chern integrand.

Let H be an $(n - 1)$ -dimensional submanifold of M equal to a level set $f^{-1}(c)$ for some f in FM and some regular value c of f . The Hessian form of f at p is the symmetric bilinear form $h_p: M_p \times M_p \rightarrow R$ defined by $h_p(u, v) := \langle \nabla_u \nabla f, v \rangle$ for u and v in M_p . Notice that for p in H , h_p restricted to $H_p \times H_p$ is just $\|\nabla f(p)\|$ times the second fundamental form of H at p with respect to the unit normal field $\nabla f / \|\nabla f\|$ on H ; call this restricted unit normal field X .

We want to estimate $\int_H X^* \Pi$. To do this we first evaluate $g(p)$, where g is the function on H determined by the identification

$$[X^* \Pi]_p \simeq g(p) \cdot d_p(\text{vol}_H) ,$$

where $d(\text{vol}_H)$ is the volume form on H . Let $u_0 := X_p$, and let u_1, \dots, u_{n-1} be the principal curvature directions of H at p with principal curvatures $\lambda_1, \dots, \lambda_{n-1}$ respectively. Then

$$\begin{aligned} g(p) &= g(p) \cdot d_p(\text{vol}_H)(u_1, \dots, u_{n-1}) \\ &= X^* \Pi(u_1, \dots, u_{n-1})(u_0, \dots, u_{n-1}) \\ &= \sum_{i=1}^k c_i X^*(\gamma \wedge \theta^{2i-1} \wedge \Omega^{k-i})(u_1, \dots, u_{n-1})(u_0, \dots, u_{n-1}) \\ &= \sum c_i \sum \text{sgn}(\sigma) \text{sgn}(\tau) \frac{1}{[(n - 2i)!]^2} \langle X, u_{\tau_0} \rangle \langle \nabla_{X_{\sigma_1} u_{\sigma_1}} X_{\#} X, X_{\#} u_{\tau_1} \rangle \\ &\quad \dots X^* \pi^* \Omega^{k-i}(u_{\sigma_{2i}}, \dots, u_{\sigma_{n-1}})(u_{\tau_{2i}}, \dots, u_{\tau_{n-1}}) , \end{aligned}$$

where the inner sum is taken over all $\sigma \in S_{n-1}$ acting on $\{1, \dots, n - 1\}$, and $\tau \in S_n$ acting on $\{0, \dots, n - 1\}$, and everything is evaluated at p . Since $\langle X, u_{\tau_0} \rangle = \delta_{0, \tau_0}$ by the definition of the u_i , and

$$\langle \nabla_{X_{\sigma_1} u_{\sigma_1}} X_{\#} X, X_{\#} u_{\tau_1} \rangle = \langle \nabla_{u_{\sigma_1}} X, u_{\tau_1} \rangle = \delta_{\sigma_1, \tau_1} \lambda_{\sigma_1} ,$$

we have

$$g(p) = \sum_1^k (2i - 1)! c_i \sum \lambda_{j_1} \dots \lambda_{j_{2i-1}} \Omega^{k-i} ,$$

where Ω^{k-i} is evaluated on $u_{\ell_1}, \dots, u_{\ell_{n-2i}}$, and the inner sum is taken over all $j_1 < \dots < j_{2i-1}$ and $\ell_1 < \dots < \ell_{n-2i}$ such that $\{j_1, \dots, j_{2i-1}, \ell_1, \dots, \ell_{n-2i}\} = \{1, \dots, n - 1\}$.

If the boundaries of the sets C_t were C^2 for all sufficiently large t , the remaining work would be simple; unfortunately, the boundary of a totally convex set is in general only continuous, and the set of points of non-differentiability can in fact be dense in the boundary. The nicest way around the difficulty would be to approximate each set by a totally convex set with C^2 boundary; so far we have been unable to prove that this is possible in general. For

our purposes it is sufficient to have an approximating set with a differentiable boundary which is only close to being totally convex.

Recall that on a manifold a convex function is a continuous function $f: M \rightarrow R$ such that for every geodesic c in M , $f \circ c$ is a convex function on R in the standard sense, that is, given real numbers a and b ,

$$f \circ c(sa + (1 - s)b) \leq sf \circ c(a) + (1 - s)f \circ c(b)$$

for $0 \leq s \leq 1$. All the sublevel sets of a convex function are totally convex.

In our situation, the function $f: M \rightarrow R$ such that

$$f(p) = \begin{cases} 0 & \text{if } p \in C_0, \\ t & \text{if } p \in \partial C_t \text{ for } t > 0 \end{cases}$$

is convex. This follows since on C_t , $[p \mapsto -\rho(p, \partial C_t)]$ is a convex function (see [1]), and $f(p) = t - \rho(p, \partial C_t)$ for p in C_t .

In attempting to find a C^∞ convex approximation for f one is naturally led to the idea of convolving f with a C^∞ bump function. The result fails to be convex, but is close in the sense of the following lemma.

Lemma 7. *Given K compact in M , choose $\varepsilon > 0$ as in Lemma 3. Suppose $f: M \rightarrow R$ is convex. Then there exist an open neighborhood U of K and a family $\{f_r \in FU \mid r \in (0, \varepsilon)\}$ such that*

- (1) *the functions $f_r|_K$ converge uniformly to $f|_K$ as $r \rightarrow 0$,*
- (2) *f_r approaches f monotonically from above,*
- (3) *$\varinjlim_{r \rightarrow 0} (\inf_{u \in T_1 K} h_{f_r}(u, u)) \geq 0$ where h_{f_r} is the Hessian form of f_r .*

A complete proof of this result has been given by Greene and Wu in [5], so we will give no more than the explicit construction of the functions f_r for later use.

Let $\mu: R \rightarrow R$ be a nonnegative C^∞ bump function with support in $[-1, 1]$ which is constant near zero and such that

$$\int_{R^n} \mu(\|v\|) dv = 1.$$

Define

$$f_r(p) := \frac{1}{r^n} \int_{M_p} f \circ \exp(v) \mu\left(\frac{\|v\|}{r}\right) dv.$$

Theorem. *Let M be a complete oriented Riemannian manifold of non-negative sectional curvature and even dimension $2k$. Assume that the algebraic Hopf conjecture holds on M . Then the total curvature of M exists and is bounded between zero and the Euler characteristic of M .*

Proof. For each positive integer i , let $\varepsilon_i > 0$ be a constant as chosen in

Lemma 3 for the set C_i in the filtration of M by compact totally convex sets C_t . Fix a positive integer $i \geq 2$; let $\{f_r \mid r \in (0, \varepsilon_i)\}$ be a sequence of C^∞ functions on a neighborhood U of C_i as given by Lemma 7 for the function f indexing the sets C_t .

First we prove that for r sufficiently small, f_r has no critical points off of a neighborhood of C_0 in C_1 , and $f_r^{-1}(-\infty, i]$ is homeomorphic to C_i .

Choose a positive $\delta < \min\{\varepsilon_i, 1\}$ for the set C_i as in Lemma 4. Let $i - \delta < a' < a < i$. Then we have a continuous geodesic contraction of C_i onto $C_{a'}$ and the distance function d from C_i to $C_{a'}$ is C^1 on $C_i - C_{a'}$. For each p in $\partial C_{a'}$, there is a unique geodesic γ_p through p such that $\gamma_p(0) = p$ and $\dot{\gamma}_p(0) = \nabla d(p)$; γ_p is a reparametrization of one of the geodesics given by the contraction of C_i onto $C_{a'}$. $\partial C_{a'}$ is transverse to γ_p , and therefore f is strictly increasing along γ_p . By continuity, for each p there is a cone W_p of normal geodesics through p about γ_p such that f is strictly increasing along each c in W_p . We may assume that there is some positive number α such that for all p , W_p is the set of all normal geodesics on the interval $[-\beta, 1]$ which have $c(0) = p$ and $\angle(\dot{c}(0), \dot{\gamma}_p(0)) \leq \alpha$, where β is some positive number less than $a - a'$. Let $W := \bigcup_{p \in \partial C_{a'}} W_p$. Then $\{f \circ c|_{[-\beta, 1]} \mid c \in W\}$ is a continuous compact family of strictly increasing functions, all of which are convex and have the value a at 0. All of the difference quotients

$$\frac{1}{t}[f \circ c(t) - a], \quad t \neq 0, c \in W,$$

are positive, and it is a straightforward proof that the difference quotients for $t > 0$ have a uniform positive lower bound κ . Then by monotonicity of the difference quotients of a convex function,

$$\frac{f \circ c(t) - f \circ c(s)}{t - s} \geq \kappa$$

if $0 \leq s < t$ and $c \in W$.

Furthermore, there exists a positive number r_0 such that for all c in W , the difference quotients of f along every curve of the form $[t \mapsto \exp_{c(t)} V(t)]$ are greater than or equal to $\frac{1}{2}\kappa$, where each V is a parallel vector field along c for which $\|V(t)\| < r_0$. We may assume that $r_0 < \varepsilon_i$.

Now let $c = \gamma_p$ for some p in $\partial C_{a'}$. We want to calculate $(f_r \circ c)'(t)$ for $t > 0$. Now $r^n(f_r \circ c(s) - f_r \circ c(t))$ is equal to

$$\begin{aligned} & \int_{M_{c(s)}} f \circ \exp_{c(s)}(v) \mu\left(\frac{\|v\|}{r}\right) dv - \int_{M_{c(t)}} f \circ \exp_{c(t)}(v) \mu\left(\frac{\|v\|}{r}\right) dv \\ & = \int_{M_{c(t)}} (f \circ \exp_{c(s)} V(s) - f \circ \exp_{c(t)} V(t)) \mu\left(\frac{\|v\|}{r}\right) dv, \end{aligned}$$

where V is the parallel vector field along c with $V(t) = v$. Thus, if $r < r_0$, then

$$f_r \circ c(s) - f_r \circ c(t) \geq \frac{1}{2}(s - t)\kappa ,$$

and therefore $(f_r \circ c)'(t) \geq \frac{1}{2}\kappa$. But this means that $\langle \nabla f_r, \nabla d \rangle$ is positive for $r < r_0$; notice that by the same argument we can prove that off of a small neighborhood of C_0 , the derivative of f_r in a direction transverse to the boundary of each set C_i is positive for sufficiently small r . In particular, for small r we may use the gradient flow of some distance function d to obtain a homeomorphism between C_i and $\mathcal{E}_r := f_r^{-1}(-\infty, i]$. This is a crucial step in the proof of the theorem since it allows us to equate the Euler characteristic of \mathcal{E} with the Euler characteristic of M :

$$\chi_{\mathcal{E}_r} = \chi_{C_i} = \chi_S = \chi_M .$$

We will also use the fact that f_r has no critical points near ∂C_i since that means that $\mathcal{E}_r = f_r^{-1}(i)$ is a submanifold.

From here on the proof is almost identical to what it would have been if the boundary of C_i had been C^2 .

We may approximate ∇f_r by a C^∞ vector field V_r which equals ∇f_r near $\partial \mathcal{E}_r$ and has only isolated zeros. By the Poincaré-Hopf Theorem [9], the sum of the indices of the zeros of V_r is just the Euler characteristic of \mathcal{E}_r , that is, χ_M . Let $Y_r := V_r / \|V_r\|$ on $\mathcal{E}_r - \{\text{zeros of } V_r\}$.

Then

$$\int_{\mathcal{E}_r} \chi = \int_{\partial \mathcal{E}_r} Y_r^* \Pi + \chi_M .$$

But

$$C_i = \bigcup_{r>0} \overline{\mathcal{E}_r}$$

since $f_r \geq f$, and then

$$\int_{C_i} \chi = \lim_{r \rightarrow 0} \int_{\mathcal{E}_r} \chi .$$

In particular, the limit of $\int_{\partial \mathcal{E}_r} Y_r^* \Pi$ exists as $r \rightarrow 0$. As before, $\int_{\partial \mathcal{E}_r} Y_r^* \Pi = \int_{\partial \mathcal{E}_r} g_r(p) \cdot d_p(\text{vol}_{\partial \mathcal{E}_r})$, and

$$g_r(p) = - \sum_{m=1}^k \frac{(m-1)!}{2^{k-m+1} \pi^k (k-i)!} \sum \lambda_{j_1} \cdots \lambda_{j_{2m-1}} (-\Omega)^{k-i} ,$$

where $(-\Omega)^{k-i}$ is evaluated on the plane spanned by v_{ℓ_1} through $v_{\ell_{n-2m}}$, and

the inner sum is taken over all $j_1 < \dots < j_{2m-1}$ and $\ell_1 < \dots < \ell_{n-2m}$ such that $\{j_1, \dots, j_{2m-1}, \ell_1, \dots, \ell_{n-2m}\} = \{1, \dots, n-1\}$. Here v_1, \dots, v_{n-1} are the principal curvature directions of ∂E_r at p with corresponding principal curvatures $\lambda_1, \dots, \lambda_{n-1}$. By the choice of the functions f_r and the fact that the Hessian form of f_r at p is just $\|\nabla f_r(p)\|$ times the second fundamental form of ∂E_r at p , we see that $\lim_{r \rightarrow 0} \lambda_{j_1} \dots \lambda_{j_{2m-1}} \geq 0$. Since we have assumed that the Hopf conjecture is true on M , $(-\Omega)^{k-i}$ is nonnegative. The result is that $\lim_{r \rightarrow 0} g_r(p) \leq 0$, and consequently,

$$\lim_{r \rightarrow 0} \int_{\partial E_r} Y_r^* H \leq 0 .$$

Here we use the fact that $\{\text{vol}(\partial E_r)\}_{r>0}$ is bounded, which is easily proven using the basic Hausdorff measure theory. But then,

$$\int_{C_i} \chi = \lim_{r \rightarrow 0} \int_{E_r} \chi = \lim_{r \rightarrow 0} \int_{\partial E_r} Y_r^* H + \chi_M \leq \chi_M .$$

Again by the Hopf conjecture, $0 \leq \int_{C_i} \chi \leq \int_{C_{i+1}} \chi$ for all i . Therefore $\int_M \chi$ exists and $0 \leq \int_M \chi \leq \chi_M$.

4. Some remarks

Cohn-Vossen classified the complete noncompact surfaces of nonnegative curvature [4]; for such a surface M , the soul S is either a point or is diffeomorphic to S^1 . If S is a point, then M is diffeomorphic to R^2 , and Cohn-Vossen proved that the total curvature is bounded between 0 and 1; these estimates are sharp. If S is diffeomorphic to S^1 , then M is isometric to one of the two flat line bundles over S and has zero total curvature.

Our result gives a new proof of the Cohn-Vossen total curvature result.

Cheeger and Gromoll classified the complete noncompact 3-manifolds of nonnegative curvature using the soul theorem [1]; the soul is diffeomorphic to one of the following: a point, S^1, S^2, RP^2 , the flat torus, or the flat Klein bottle. In addition, they proved that if the soul S of an arbitrary complete Riemannian manifold of nonnegative curvature has codimension 1, then M is isometrically a flat line bundle over S .

Gromoll and Meyer proved that if the curvature of M is everywhere positive, then the soul is a point (see [7]). From this they proved, for $n \neq 3$ or 4, that M and R^n are diffeomorphic; our result implies directly that for any manifold for which the soul is a point, the manifold is diffeomorphic to R^n and the total curvature, if it exists, is bounded above by 1.

We will assume that all our manifolds are oriented; by considering the orientation covering, similar information can be gotten for nonoriented mani-

folds. No attempt will be made to classify anything, but we will make some comments about the cases of dimensions 4 and 6.

Dimension 4. The only possible souls with positive Euler characteristic are a point and S^2 . Thus we see immediately that $0 \leq \int_M \chi \leq 2$.

If the soul is diffeomorphic to S^1 , then the total curvature of M is zero.

Suppose $\dim S = 2$. If S is the flat torus, then M is locally isometrically trivial over S (see [1]), and in such a case the Chern integrand is identically zero. If S is diffeomorphic to S^2 and the total curvature of M is positive, then M is not isometrically the product $S^2 \times R^2$; M may still be the product of S^2 and some nonnegatively curved surface diffeomorphic to R^2 , and there are examples where M is not even locally isometrically a product, e.g., TS^2 .

If $\dim S = 3$, then χ_M is of course zero, as is the case whenever the soul of M is odd-dimensional. In addition, in this case M is isometrically a flat line bundle over S with zero Chern integrand. Despite these restrictions there are still many possibilities for M , since there are already 6 flat compact orientable 3-manifolds [11].

Dimension 6. Here we cannot in general say that the total curvature exists, but since the boundary terms involve only the first and second powers of the curvature form, we do know that

$$\overline{\lim}_{i \rightarrow \infty} \int_{C_i} \chi \leq \chi_M .$$

Even if the total curvature does exist, without more information we have no guarantee that it will be nonnegative. The number of manifolds here will be quite large since in the flat case alone there are very many manifolds of dimension less than six. Notice however that if S is flat, then the Chern integrand is identically zero on M since locally M has a flat factor [1].

References

- [1] J. Cheeger & D. Gromoll, *On the structure of complete manifolds of nonnegative curvature*, Ann. of Math. **96** (1972) 413–443.
- [2] S. S. Chern, *A simple intrinsic proof of the Gauss-Bonnet formula for closed Riemannian manifolds*, Ann. of Math. **45** (1944) 747–752.
- [3] —, *On curvature and characteristic classes of a Riemannian manifold*, Abh. Math. Sem. Univ. Hamburg **20** (1956) 117–126.
- [4] S. Cohn-Vossen, *Kürzeste Wege und Totalkrümmung auf Flächen*, Composito Math. **2** (1935) 69–133.
- [5] R. Greene & H. Wu, *On the subharmonicity and plurisubharmonicity of geodesically convex functions*, Indiana Univ. Math. J. **22** (1973) 641–654.
- [6] D. Gromoll, W. Klingenberg & W. Meyer, *Riemannsche Geometrie im Grossen*, Springer, Berlin, 1968.
- [7] D. Gromoll & W. Meyer, *On complete open manifolds of positive curvature*, Ann. Math. **90** (1969) 75–90.
- [8] J. Milnor, *Morse theory*, Annals of Math. Studies, No. 51, Princeton University Press, Princeton, 1963.

- [9] ———, *Topology from the differentiable viewpoint*, University Press of Virginia, Charlottesville, 1965.
- [10] J. Munkres, *Elementary differential topology*, Annals of Math. Studies, No. 54, Princeton University Press, Princeton, 1963.
- [11] J. Wolf, *Spaces of constant curvature*, McGraw-Hill, New York, 1966.

STATE UNIVERSITY OF NEW YORK, STONY BROOK