

On a certain property of a Riemannian space admitting a special concircular scalar field

Dedicated to Professor Yoshie Katsurada on her 60th birthday

By Tsunehira KOYANAGI

§ 0. Introduction.

The purpose of the present paper is to investigate the property of a Riemannian space which admits a scalar field Φ characterised by the property

$$(0.1) \quad \Phi_{k;l} = \rho \Phi g_{kl}, \quad \rho = \text{non-zero constant},$$

(such a scalar field Φ is called the special concircular scalar field in this paper) where $\Phi_k \stackrel{\text{def}}{=} \Phi_{;k}$ and g_{kl} means the metric tensor of the space. In § 1, we consider a Riemannian space with certain special curvature tensor, and prove the property that the space is of constant curvature. Next, in § 2, we give some corollaries of it.

The author wishes to express to Prof. Y. Katsurada and Dr. T. Nagai his very sincere thanks for their kind guidance.

§ 1. Riemannian space with certain special curvature tensor.

We suppose an n -dimensional Riemannian space M ($n \geq 3$) of class C^r ($r \geq 3$) which has local coordinates x^i and admits the special concircular scalar field Φ defined by the equation (0.1). First, substituting the relation obtained from (0.1) into the Ricci identity

$$2\Phi_{i;[j;k]} = -R^a{}_{ijk} \Phi_a,$$

we have

$$(1.1) \quad \rho(\Phi_k g_{ij} - \Phi_j g_{ik}) = -R^a{}_{ijk} \Phi_a,$$

from which, by covariant differentiation with respect to x^l ,

$$\rho(\Phi_{k;l} g_{ij} - \Phi_{j;l} g_{ik}) = -R^a{}_{ijk;l} \Phi_a - R^a{}_{ijk} \Phi_{a;l},$$

and consequently, inserting the relation (0.1), we obtain

$$(1.2) \quad \rho \Phi \left\{ \rho(g_{kl} g_{ij} - g_{jl} g_{ik}) + R_{lijk} \right\} = -R^a{}_{ijk;l} \Phi_a.$$

Moreover, on differentiating (1.2) covariantly with respect to x^h , we have

$$\begin{aligned} \rho \Phi_h \{ \rho (g_{kl} g_{ij} - g_{jl} g_{ik}) + R_{lijk} \} + \rho \Phi R_{lijk;h} \\ = -R^a_{ijk;l;h} \Phi_a - R^a_{ijk;l} \Phi_{a;h}, \end{aligned}$$

and inserting the values of $\Phi_{i;j}$ given by (0.1) we obtain

$$(1.3) \quad \begin{aligned} \rho \Phi_h \{ \rho (g_{kl} g_{ij} - g_{jl} g_{ik}) + R_{lijk} \} \\ = -R^a_{ijk;l;h} \Phi_a - \rho \Phi (R_{lijk;h} + R_{hlijk;l}), \end{aligned}$$

so that multiplying the expression (1.3) by Φ^h and summing with respect to h we get

$$(1.4) \quad \begin{aligned} \rho (\Phi_h \Phi^h) \{ \rho (g_{kl} g_{ij} - g_{jl} g_{ik}) + R_{lijk} \} \\ = -R^a_{ijk;l;h} \Phi_a \Phi^h - \rho \Phi (R_{lijk;h} + R_{hlijk;l}) \Phi^h. \end{aligned}$$

Also, multiplying both sides of (1.3) by Φ^l and summing with respect to l we have

$$\begin{aligned} \rho \Phi_h \{ \rho (\Phi_k g_{ij} - \Phi_j g_{ik}) + R_{lijk} \Phi^l \} \\ = -R^a_{ijk;l;h} \Phi_a \Phi^l - \rho \Phi (R_{lijk;h} + R_{hlijk;l}) \Phi^l. \end{aligned}$$

From (1.1), it is evident that the left hand side of these equations are equal to zero, and consequently we get

$$-R^a_{ijk;l;h} \Phi_a \Phi^l - \rho \Phi (R_{lijk;h} + R_{hlijk;l}) \Phi^l = 0,$$

from which, by interchanging the indices h and l , we obtain

$$(1.5) \quad -R^a_{ijk;h;l} \Phi_a \Phi^h - \rho \Phi (R_{hlijk;l} + R_{lijk;h}) \Phi^h = 0.$$

And subtracting from (1.4) the equation (1.5), we find

$$(1.6) \quad \begin{aligned} \rho (\Phi_h \Phi^h) \{ \rho (g_{kl} g_{ij} - g_{jl} g_{ik}) + R_{lijk} \} \\ = (R^a_{ijk;h;l} - R^a_{ijk;l;h}) \Phi_a \Phi^h. \end{aligned}$$

Suppose that our space has the curvature tensor satisfying $R^a_{ijk;[h;l]} = 0$ ([2], p. 222). Then, ρ defined by (0.1) being different from zero, the equation (1.6) can be written as follows:

$$(1.7) \quad (\Phi_h \Phi^h) \{ \rho (g_{kl} g_{ij} - g_{jl} g_{ik}) + R_{lijk} \} = 0.$$

We assume, moreover, that there exists no open set U such that $\Phi = \text{constant}$ at any point of it. And then it follows that there exists no open set V such that $\Phi_i = 0$ at any point of it. Under these assumptions, from (1.7), we obtain the following relation:

$$R_{lijk} = \rho (g_{ij} g_{ik} - g_{ik} g_{ij}),$$

that is, our space is of constant curvature. Hence we have the following

THEOREM. *Let M be a Riemannian space of dimension n which has the curvature tensor satisfying*

$$(1.8) \quad R_{hijk;[z;m]} = 0$$

and admits the special concircular scalar field Φ defined by (0.1). Then M is of constant curvature.

§ 2. Some corollaries.

Suppose that a Riemannian space M is symmetric. Then it is evident that the condition (1.8) is satisfied. Therefore we have

COROLLARY 1. *Let M be a symmetric Riemannian space which admits the special concircular scalar field Φ . Then M is of constant curvature. ([1])*

Next we consider an n -dimensional Einstein space $M(n > 2)$ which has the scalar curvature $R \neq 0$ and admits a proper conformal Killing vector field ξ^t , that is, ξ^t satisfies an equations :

$$(2.1) \quad \mathcal{L}_{\xi} g_{ij} \equiv \xi_{i;j} + \xi_{j;i} = 2\phi g_{ij} \quad ([2], \text{ p. } 32),$$

where $\mathcal{L}_{\xi} g_{ij}$ means the Lie derivative of the metric tensor g_{ij} with respect to ξ^t . Then the Lie derivative of the curvature tensor $R^h{}_{ijk}$ with respect to the conformal Killing vector field ξ^t is given by

$$(2.2) \quad \mathcal{L}_{\xi} R^h{}_{ijk} = \delta_j^h \phi_{i;k} - \delta_k^h \phi_{i;j} + g_{ik} \phi_{;j}^h - g_{ij} \phi_{;k}^h, \quad ([2], \text{ p. } 160)$$

where $\phi_{;i} \stackrel{\text{def}}{=} \phi_{;i}$, $\phi^i \stackrel{\text{def}}{=} g^{ij} \phi_{;j}$ and δ_j^h is the Kronecker delta. Since M is an Einstein space, we have

$$(2.3) \quad R_{ij} = \frac{R}{n} g_{ij} \quad (R = \text{constant}),$$

where R_{ij} is the Ricci tensor and R the scalar curvature. On making use of (2.2) and (2.3), after some calculations we obtain the following result :

$$(2.4) \quad \phi_{i;j} = -\frac{R}{n(n-1)} \phi g_{ij}$$

and, remembering the round brackets of (2.3) and the assumption $R \neq 0$, we have $-\frac{R}{n(n-1)} = \text{non-zero constant}$.

From (2.4) we can see that the Einstein space M admitting the proper conformal Killing vector field ξ^t must always admit the scalar field ϕ , which

is the special concircular scalar field. Hence we have the following corollary :

COROLLARY 2. *Let M be an n -dimensional Einstein space ($n > 2$) which has the scalar curvature $R \neq 0$, the curvature tensor such that $R_{h\ell jk;[l;m]} = 0$ and admits a proper conformal Killing vector field ξ^i . Then M is of constant curvature.*

On the other hand, multiplying both sides of (1.6) by g^{ij} and summing with respect to i and j , we get

$$\rho(\Phi_h \Phi^h) \{ \rho(n-1)g_{kl} + R_{kl} \} = (R_{ak;h;l} - R_{ak;l;h}) \Phi^a \Phi^h .$$

When we think of the vector field Φ_h as being provided with the assumption with respect to Φ in the manner described in §1, we have the following

PROPOSITION. *Let M be a Riemannian space of dimension n which has the Ricci tensor such that*

$$(2.5) \quad R_{ak;[h;l]} = 0$$

and admits the special concircular scalar field Φ . Then M is an Einstein space.

Now, suppose that a Riemannian space M is Ricci symmetric (defined by $R_{ij;k} = 0$). Then it is evident that the condition (2.5) is satisfied, and so that we obtain

COROLLARY. *Let M be a Ricci symmetric space of dimension n admitting a special concircular scalar field Φ . Then M is an Einstein space.*

Department of Mathematics
Hokkaido University

References

- [1] T. SUMITOMO: Projective and conformal transformations in compact Riemannian manifolds, *Tensor (N.S.)*, 9 (1959), 113-135.
- [2] K. YANO: *The theory of Lie derivatives and its applications*, (North-Holland, 1957).

(Received August 30, 1971)