

Structure Theory of Rack-Bialgebras

Alexandre C¹, Bordemann M², Rivière S³ and Wagemann F^{3*}

¹Department of Mathematics, Université de Strasbourg, 4 Rue Blaise Pascal, 67081 Strasbourg, France

²Laboratoire de Mathématiques, Informatique et Applications, Université de Mulhouse, France

³Department of Mathematics, Université de Nantes, 1 Quai de Tourville, 44035 Nantes Cedex 1, France

Abstract

In this paper we focus on a certain self-distributive multiplication on coalgebras, which leads to so-called rack bialgebra. Inspired by semi-group theory (adapting the Suschkewitsch theorem), we do some structure theory for rack bialgebras and cocommutative Hopf dialgebras. We also construct canonical rack bialgebras (some kind of enveloping algebras) for any Leibniz algebra and compare to the existing constructions.

We are motivated by a differential geometric procedure which we call the Serre functor: To a pointed differentiable manifold with multiplication is associated its distribution space supported in the chosen point. For Lie groups, it is well-known that this leads to the universal enveloping algebra of the Lie algebra. For Lie racks, we get rack-bialgebras, for Lie digroups, we obtain cocommutative Hopf dialgebras.

Keywords: Coalgebras; Cocommutative Hopf dialgebras; Canonical rack bialgebras; Manifolds; Drinfeld center

Introduction

All manifolds considered in this manuscript are assumed to be Hausdorff and second countable. Basic Lie theory relies heavily on the fundamental links between associative algebras, Lie algebras and groups. Some of these links are the passage from an associative algebra A to its underlying Lie algebra A^{Lie} which is the vector space A with the bracket $[a, b] := ab - ba$. On the other hand, to any Lie algebra \mathfrak{g} one may associate its universal enveloping algebra $U(\mathfrak{g})$ which is associative. Groups arise as groups of units in associative algebras. To any group G , one may associate its group algebra KG which is associative. The theme of the present article is to investigate links of this kind for more general objects than groups, namely for racks and digroups.

Recall that a pointed rack is a pointed set (X, e) together with a binary operation $\triangleright : X \times X \rightarrow X$ such that for all $x \in X$, the map $y \mapsto x \triangleright y$ is bijective and such that for all $x, y, z \in X$, the self-distributivity and unit relations

$$x \triangleright (y \triangleright z) = (x \triangleright y) \triangleright (x \triangleright z), e \triangleright x = x, \text{ and } x \triangleright e = e$$

are satisfied. Imitating the notion of a Lie group, the smooth version of a pointed rack is called Lie rack.

An important class of examples of racks are the so-called augmented racks [1]. An augmented rack is the data of a group G , a G -set X and a map $p: X \rightarrow G$ such that for all $x \in X$ and all $g \in G$,

$$p(g \cdot x) = gp(x)g^{-1}.$$

The set X becomes then a rack by setting $x \triangleright y := p(x) \cdot y$. In fact, augmented racks are the Drinfeld center (or the Yetter-Drinfeld modules) in the monoidal category of G -sets over the (set-theoretical) Hopf algebra G , see for example [2]. Any rack may be augmented in many ways, for example by using the canonical morphism to its associated group or to its group of bijections or to its group of automorphisms.

In order to formalize the notion of a rack, one needs the diagonal map $diag : X \rightarrow X \times X$ given by $x \mapsto (x, x)$. Then the self-distributivity relation reads in terms of maps $m \circ (id_M \times m)$,

$$= m \circ (m \times m) \circ (id_M \times \tau_{M,M} \times id_M) \circ (diag_M \times id_M \times id_M)$$

Axiomatizing this kind of structure, one may start with a coalgebra C and look for rack operations on this fixed coalgebra [3,4].

A natural framework where this kind of structure arises (as we show in Section 3) is by taking point-distributions (i.e. applying the Serre functor) over (resp. to) the pointed manifold given by a Lie rack. We dub the arising structure as rack bialgebra.

Lie racks are intimately related to Leibniz algebra \mathfrak{h} , i.e. a vector space \mathfrak{h} with a bilinear bracket $[\cdot, \cdot]: \mathfrak{h} \otimes \mathfrak{h} \rightarrow \mathfrak{h}$ such that for all $X, Y, Z \in \mathfrak{h}$, $[X, -]$ acts as a derivation:

$$[X, [Y, Z]] = [[X, Y], Z] + [Y, [X, Z]]. \tag{1}$$

Indeed, Kinyon showed that the tangent space at $e \in H$ of a Lie rack H carries a natural structure of a Leibniz algebra, generalizing the relation between a Lie group and its tangent Lie algebra [5]. Conversely, every (finite dimensional real or complex) Leibniz algebra \mathfrak{h} may be integrated into a Lie rack $R_{\mathfrak{h}}$ (with underlying manifold \mathfrak{h}) using the rack product.

$$X \blacktriangleright Y := e^{ad_X}(Y), \tag{2}$$

noting that the exponential of the inner derivation ad_X for each $X \in \mathfrak{h}$ is an automorphism.

Another closely related algebraic structure is that of dialgebras. A dialgebra is a vector space D with two (bilinear) associative operations $\vdash : D \times D \rightarrow D$ and $\dashv : D \times D \rightarrow D$ which satisfy three compatibility relations, namely for $a, b, c \in D$:

$$(a \vdash b) \dashv c = (a \dashv b) \vdash c, a \dashv (b \vdash c) = a \dashv (b \dashv c), (a \dashv b) \dashv c = a \dashv (b \vdash c)$$

*Corresponding author: Wagemann F, Department of Mathematics, Université de Nantes, 1 Quai de Tourville, 44035 Nantes Cedex 1, France, Tel: 0276645089, E-mail: Friedrich.Wagemann@univ-nantes.fr

Received July 17, 2016; Accepted November 19, 2016; Published November 30, 2016

Citation: Alexandre C, Bordemann M, Rivière S, Wagemann F (2016) Structure Theory of Rack-Bialgebras. J Generalized Lie Theory Appl 10: 244. doi:10.4172/1736-4337.1000244

Copyright: © 2016 Alexandre C, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

A dialgebra D becomes a Leibniz algebras via the formula $[a, b] = a \dashv b - b \dashv a$.

In this sense \dashv and \dashv are two halves of a Leibniz bracket. Loday and Guichot have defined an enveloping dialgebra of a Leibniz algebra [6,7].

One main point of this paper is the link between rack bialgebras and cocommutative Hopf dialgebras. In Theorem 2.5, we adapt Suschkewitsch's Theorem in semi-group theory to the present context. The classical result (see Appendix B) treats semi-groups with a left unit e and right inverses (analogous results in the left-context), called right groups. Suschkewitsch shows that such a right group Γ decomposes as a product $\Gamma = \Gamma e \times E$ where E is the set of all idempotent elements.

Its incarnation here shows that a cocommutative right Hopf algebra \mathcal{H} decomposes as a tensor product $\mathcal{H}1 \otimes E_{\mathcal{H}}$ where $E_{\mathcal{H}}$ is the subspace of generalized idempotents.

Furthermore, we will show in Theorem 2.6 how to associate to any augmented rack bialgebra an augmented cocommutative Hopf dialgebra. In Theorem 2.7, we investigate what Suschkewitsch's decomposition gives for a cocommutative Hopf dialgebra A . It turns out that A decomposes as a tensor product $E_A \otimes H_A$ of E_A with H_A which may be identified to the associative quotient A_{ass} of A . This result permits to show that the Leibniz algebra of primitives in A is a hemi-semi-direct product, and thus always split. In this way we arrive once again at the result that Lie digroups may serve only to integrate split Leibniz algebras which has already been observed by Covez in his master thesis [8].

Let us comment on the content of the paper:

All our bialgebra notion are based on the standard theory of coalgebras, some features of which as well as our notions are recalled in Appendix A. Rack bialgebras and augmented rack bialgebras are studied in Section 2. Connected, cocommutative Hopf algebras give rise to a special case of rack bialgebras. In Section 2.2, we associate to any Leibniz algebra \mathfrak{h} an augmented rack bialgebra $UAR^{\infty}(\mathfrak{h})$ and study the functorial properties of this association. This rack bialgebra plays the role of an enveloping algebra in our context. It turns out that a truncated, non-augmented version $UR(\mathfrak{h})$ is a left adjoint of the functor of primitives $Prim$.

We also study the "group-algebra" functor associating to a rack X its rack bialgebra $K[X]$. Like in the classical framework, $K[-]$ is left adjoint to the functor $Slike$ associating to a track bialgebra its rack of set-like elements. The relation between rack bialgebras and the other algebraic notion discussed in this paper is summarized in the diagram (see the end of Section 2.2) of categories and functors:

$$\begin{array}{c} Lie@ <lex> [r]^U @^{\dashv} -> [d]^{\dagger} Hopf @^{\dashv} -> [d]^{\dagger} [I]^{Prim} @ <lex> [r]^{Slike} Grp [I]^{K[-]} \\ @^{\dashv} -> [d]^{\dagger} Leib @ <lex> [r]^{UAR^{\infty}} RackBialg [I]^{Prim} @ <lex> [r]^{Slike} Racks [I]^{K[-]} \end{array}$$

In Section 2.3, we develop the structure theory for rack bialgebras and cocommutative Hopf dialgebras, based on Suschkewitsch's Theorem. Section 2.3 contains Theorem 2.5, Theorem 2.6 and Theorem 2.7 whose content we have described above.

Recollecting basic knowledge about the Serre functor F is the subject of Section 3. In particular, we show in Section 3.2 that F is a strong monoidal functor from the category of pointed manifolds \mathcal{Mf}^* to the category of coalgebras, based on some standard material on coalgebras (Appendix A). In Section 3.3, we apply F to Lie groups, Lie semi-groups, Lie digroups, and to Lie racks and augmented Lie racks, and study the additional structure which we obtain on the coalgebra.

The case of Lie racks motivates the notion of rack bialgebra.

Recall that for a Leibniz algebra \mathfrak{h} , the vector space \mathfrak{h} becomes a Lie rack $R_{\mathfrak{h}}$ with the rack product.

$$X \blacktriangleright Y = e^{ad_X}(Y).$$

In Theorem 3.8, we show that the rack bialgebra $UAR^{\infty}(\mathfrak{h})$ associated to \mathfrak{h} coincides with the rack bialgebra $F(R_{\mathfrak{h}})$.

Several Bialgebras

In the following, let K be an associative commutative unital ring containing all the rational numbers. The symbol \otimes will always denote the tensor product of K -modules over K . For any coalgebra (C, Δ) over K , we shall use Sweedler's notation $\Delta(a) = \sum_{(a)} a^{(1)} \otimes a^{(2)}$ for any $a \in A$. See also Appendix 4 for a survey on definitions and notations in coalgebra theory.

The following sections will all deal with the following type of nonassociative bialgebra: Let $(B, \Delta, \varepsilon, 1, \mu)$ be a K -module such that $(B, \Delta, \varepsilon, 1)$ is a coassociative counital coaugmented coalgebra (a C^3 -coalgebra), and such that the linear map $\mu: B \otimes B \rightarrow B$ (the multiplication) is a morphism of C^3 -coalgebras (it satisfies in particular $\mu(1 \otimes 1) = 1$). We shall call this situation a nonassociative C^3I -bialgebra (where I stands for 1 being an idempotent for the multiplication μ). For another nonassociative C^3I -bialgebra $(B', \Delta', \varepsilon', 1', \mu')$ a K -linear map $\phi: B \rightarrow B'$ will be called a morphism of nonassociative C^3I -bialgebras iff it is a morphism of C^3 -coalgebras and is multiplicative in the usual sense $\phi(\mu(a \otimes b)) = \mu'(\phi(a) \otimes \phi(b))$ for all $a, b \in B$. The nonassociative C^3I -bialgebra $(B, \Delta, \varepsilon, 1)$ is called left-unital (resp. right-unital) iff for all $a \in B$ $\mu(1 \otimes a) = a$ (resp. $\mu(a \otimes 1) = a$).

Moreover, consider the associative algebra $A := \text{Hom}_K(B, B)$ equipped with the composition of K -linear maps, and the identity map id_B as the unit element. There is an associative convolution multiplication $*$ in the K -module $\text{Hom}_K(B, A)$ of all K -linear maps $B \rightarrow \text{Hom}_K(B, B)$, see Appendix 4, eqn (103) for a definition with $\text{id}_{B, \varepsilon}$ as the unit element. For a given nonassociative C^3I -bialgebra $(B, \Delta, 1, \mu)$ we can consider the map μ as a map $B \rightarrow \text{Hom}_K(B, B)$ in two ways: as left multiplication map $L^{\mu}: b \mapsto (b \mapsto \mu(b \otimes b'))$ or as right multiplication map $R^{\mu}: b \mapsto (b' \mapsto \mu(b' \otimes b))$. We call $(B, \varepsilon, 1, \mu)$ a left-regular (resp. right-regular) nonassociative C^3I -bialgebra iff the map L^{μ} (resp. the map R^{μ}) has a convolution inverse, i.e. iff there is a K -linear map $\mu': B \otimes B \rightarrow B$ (resp. $\mu'': B \otimes B \rightarrow B$) such that $L^{\mu} = L^{\mu'} \circ \text{id}_{B, \varepsilon} = L^{\mu'} * L^{\mu}$ (resp. $R^{\mu} = R^{\mu''} \circ \text{id}_{B, \varepsilon} = R^{\mu''} * R^{\mu}$), or on elements $a, b \in B$ for the left regular case:

$$\sum_{(a)} \mu(a^{(1)} \otimes \mu'(a^{(2)} \otimes b)) = \varepsilon(a)b = \mu'(a^{(1)} \otimes \mu(a^{(2)} \otimes b)). \quad (3)$$

Note that every associative unital Hopf algebra $(H, \Delta, \varepsilon, 1, \mu, S)$ (where S denotes the antipode, i.e. the convolution inverse of the identity map in $\text{Hom}_K(H, H)$) is right- and left-regular by setting $\mu' = \mu^{\circ}(S \otimes \text{id}_H)$ and $\mu'' = \mu^{\circ}(\text{id}_H \otimes S)$.

Lemma 2.1: Let $(B, \Delta, \varepsilon, 1, \mu)$ be a nonassociative C^3I -bialgebra.

1. If B is left-regular (resp. right-regular), then the corresponding K -linear map $\mu': B \otimes B \rightarrow B$ is unique, and in case Δ is cocommutative, μ' is map of C^3 -coalgebras.

2. If $(B, \Delta, \varepsilon, 1, \mu)$ is left-unital (rep. right-unital) and its underlying C^3 -coalgebra is connected, then $(B, \Delta, \varepsilon, 1, \mu)$ is always left-regular (resp. right-regular).

Proof: 1. In any monoid (in particular in the convolution monoid) two-sided inverses are always unique. Moreover, as can easily be

checked, a K -linear map $\phi: B \otimes B \rightarrow B$ is a morphism of coalgebras iff for each $b \in B$.

$$\sum_{(b)} (L^\phi(b^{(1)}) \otimes L^\phi(b^{(2)})) \circ \Delta = \Delta \circ L^\phi(b). \quad (4)$$

(and analogously for right multiplications). Both sides of the preceding equation, seen as maps of b , are in $\text{Hom}_K(B, \text{Hom}_K(B, B \otimes B))$. Since $\text{Hom}_K(B, B \otimes B)$ is an obvious right $\text{Hom}_K(B, B)$ -module, the K -module $\text{Hom}_K(B, \text{Hom}_K(B, B \otimes B))$ is a right $\text{Hom}_K(B, \text{Hom}_K(B, B))$ -module with respect to the convolution. Define the K -linear map $F^\mu: B \rightarrow \text{Hom}_K(B, B \otimes B)$ by:

$$F^\mu(b) := \sum_{(b)} (L^\mu(b^{(1)}) \otimes L^\mu(b^{(2)})) \circ \Delta - \Delta \circ L^\mu(b).$$

Using eqn (4), the fact that L^μ is a convolution inverse of L , and the cocommutativity of Δ , we get:

$$F^\mu * L^\mu = 0, \text{ hence } 0 = F^\mu * L^\mu * L^\mu = F^\mu * (\text{id}_B \varepsilon) = F^\mu$$

and μ' preserves comultiplications. A similar reasoning where $B \otimes B$ is replaced by K shows that μ' preserves counits. Finally, it is obvious that $L^{\mu'}(1)$ is the inverse of the K -linear map $L^\mu(1)$, and since the latter fixes 1 so does the former. The reasoning for right-regular bialgebras is completely analogous.

2. For left-unital bialgebras we get $L^\mu(1) = \text{id}_B$, and the generalized Takeuchi-Sweedler argument, see Appendix 4, shows that L^μ has a convolution inverse. Right-unital bialgebras are treated in an analogous manner.

Note that any C^3 -coalgebra $(C, \Delta, \varepsilon, 1)$ becomes a left-unital (resp. right-unital) associative C^3I -bialgebra by equipping with the left-trivial (resp. right-trivial) multiplication.

$$\mu_0(a \otimes b) := \varepsilon(a)b \text{ (resp. } \mu_0(a \otimes b) := \varepsilon(b)a). \quad (5)$$

We shall call an element $c \in B$ a generalized idempotent iff $\sum_{(c)} c^{(1)} c^{(2)} = c$. Moreover $c \in B$ will be called a generalized left (resp. right) unit element iff for all $b \in B$ we have $cb = \varepsilon(c)b$ (resp. $bc = \varepsilon(c)b$).

Rack bialgebras, augmented rack bialgebras and Leibniz algebras

Definition 2.1: A rack bialgebra $(B, \Delta, \varepsilon, 1, \mu)$ is a nonassociative C^3I -bialgebra (where we write for all $a, b \in B$ $\mu(a \otimes b) =: a \triangleright b$) such that the following identities hold for all $a, b, c \in B$

$$1 \triangleright a = a, \quad (6)$$

$$a \triangleright 1 = \varepsilon(a)1, \quad (7)$$

$$a \triangleright (b \triangleright c) = \sum_{(a)} (a^{(1)} \triangleright b) \triangleright (a^{(2)} \triangleright c). \quad (8)$$

The last condition (8) is called the self-distributivity condition.

Note that we do not demand that the C^3 -coalgebra B should be cocommutative nor connected.

Example 2.1: Any C^3 coalgebra $(C, \Delta, \varepsilon, 1)$ carries a trivial rack bialgebra structure defined by the left-trivial multiplication

$$a \triangleright b := \varepsilon(a)b \quad (9)$$

which in addition is easily seen to be associative and left-unital, but in general not unital.

Another method of constructing rack bialgebras is gauging: Let $(B, \Delta, \varepsilon, 1, \mu)$ a rack bialgebra –where we write $\mu(a \otimes b) =: a \triangleright b$ for all $a, b \in B$ –, and let $f: B \rightarrow B$ a morphism of C^3 -coalgebras such that for all $a, b \in B$

$$f(a \triangleright b) = \overline{a \triangleright (f(b))}, \quad (10)$$

i.e. f is μ -equivariant. It is a routine check that $(B, \Delta, \varepsilon, 1, \mu_f)$ is a rack bialgebra where for all $a, b \in B$ the multiplication is defined by

$$\mu_f(a \otimes b) := a \triangleright_f b := (f(a)) \triangleright b. \quad (11)$$

We shall call $(B, \Delta, \varepsilon, 1, \mu_f)$ the f -gauge of $(B, \Delta, \varepsilon, 1, \mu)$.

Example 2.2: Let $(H, \Delta, \varepsilon, 1, \mu, S)$ be a cocommutative Hopf algebra over K . Then it is easy to see (cf. also the particular case $B=H$ and $\Phi = \text{id}_H$ of Proposition 2.1) that the new multiplication $\mu(H \otimes H) \rightarrow H$, written $\mu(h \otimes h') = h \triangleright h'$, defined by the usual adjoint representation

$$h \triangleright h' := \text{ad}_h(h') := \sum_{(h)} h^{(1)} h'(S(h^{(2)})), \quad (12)$$

equips the C^4 -coalgebra $(H, \Delta, \varepsilon, 1, \mu)$ with a rack bialgebra structure.

In general, the adjoint representation does not seem to preserve the coalgebra structure if no cocommutativity is assumed.

Example 2.3: Recall that a pointed set (X, e) is a pointed rack in case there is a binary operation $\triangleright: X \times X \rightarrow X$ such that for all $x \in X$, the map $y \mapsto x \triangleright y$ is bijective and such that for all $x, y, z \in X$, the self-distributivity and unit relations:

$$x \triangleright (y \triangleright z) = (x \triangleright y) \triangleright (x \triangleright z), \quad e \triangleright x = x, \text{ and } x \triangleright e = e$$

are satisfied. Then there is a natural rack bialgebra structure on the vector space $K[X]$ which has the elements of X as a basis. $K[X]$ carries the usual coalgebra structure such that all $x \in X$ are set-like: $\Delta(x) = x \otimes x$ for all $x \in X$. The product μ is then induced by the rack product. By functoriality, μ is compatible with ε and e .

Observe that this construction differs slightly from the construction, Section 3.1.

More generally there is the following structure:

Definition 2.2: An augmented rack bialgebra over K is a quadruple $(B, \Phi, H, 1)$ consisting of a C^3 -coalgebra $(B, \Delta, \varepsilon, 1)$, of a cocommutative (!) Hopf algebra $(H, \Delta_H, \varepsilon_H, 1_H, \mu_H, S)$, of a morphism of C^3 -coalgebras $\Phi: B \rightarrow H$, and of a left action $! : H \otimes B \rightarrow B$ of H on B which is a morphism of C^3 -coalgebras (i.e. B is a H -module-coalgebra) such that for all $h \in H$ and $a \in B$

$$h.1 = \varepsilon_H(h)1 \quad (13)$$

$$(h.a) = \text{ad}_k(\Phi(a)) \quad (14)$$

Where ad denotes the usual adjoint representation for Hopf algebras, see e.g. eqn (12).

We shall define a morphism $(B, \Phi, H, 1) \rightarrow (B', \Phi', H', 1')$ of augmented rack bialgebras to be a pair (φ, ψ) of K -linear maps where $\varphi: (B, \Delta, \varepsilon, 1) \rightarrow (B', \Delta', \varepsilon', 1')$ is a morphism of C^3 -coalgebras, and $\psi: H \rightarrow H'$ is a morphism of Hopf algebras such that the obvious diagrams commute:

$$\Phi' \circ \varphi = \psi \circ \Phi, \text{ and } 1' \circ (\varphi \otimes \varphi) = \varphi \circ 1 \quad (15)$$

An immediate consequence of this definition is the following:

Proposition 2.1: Let $(B, \Phi, H, 1)$ be an augmented rack bialgebra. Then the C^3 -coalgebra $(B, \varepsilon, 1)$ will become a left-regular rack bialgebra by means of the multiplication:

$$a \triangleright b := \Phi(a).b \quad (16)$$

for all $a, b \in B$. In particular, each Hopf algebra H becomes an augmented rack bialgebra via $(H, \text{id}_H, H, \text{ad})$. In general, for each augmented rack bialgebra the map $\Phi: B \rightarrow H$ is a morphism of rack bialgebras.

Proof: For a proof of this proposition [9].

Example 2.4: Exactly in the same way as a pointed rack gives rise to a rack bialgebra $K[X]$, an augmented pointed rack $p: X \rightarrow G$ gives rise to an augmented rack bialgebra $p: K[X] \rightarrow K[G]$.

Remark 2.1: Motivated by the fact that the augmented racks $p: X \rightarrow G$ are exactly the Yetter-Drinfeld modules over the (set-theoretical) Hopf algebra G , we may ask about the relation of augmented rack bialgebras to Yetter-Drinfeld modules, and more generally of rack bialgebras to the Yang-Baxter equation. For these subjects.

The link to Leibniz algebras is contained in the following:

Proposition 2.2: Let $(B, \Delta, \varepsilon, 1, \mu)$ be a rack bialgebra over K .

1. Then its K -submodule of all primitive elements, $\text{Prim}(B) =: \mathfrak{h}$, (see eqn (101) of Appendix 4) is a subalgebra with respect to μ (written $a \triangleright b$) satisfying the (left) Leibniz identity:

$$x \triangleright (y \triangleright z) = (x \triangleright y) \triangleright z + y \triangleright (x \triangleright z) \quad (17)$$

for all $x, y, z \in \mathfrak{h} = \text{Prim}(B)$. Hence the pair $(\mathfrak{h}, [\cdot, \cdot])$ with $[x, y] := x \triangleright y$ for all $x, y \in \mathfrak{h}$ is a Leibniz algebra over K . Moreover, every morphism of rack bialgebras maps primitive elements to primitive elements and thus induces a morphism of Leibniz algebras.

2. More generally, \mathfrak{h} and each subcoalgebra of order $k \in \mathbb{N}$, $B_{(k)}$ (see eqn (102)) is stable by left-multiplications with every $a \in B$. In particular, each $B_{(k)}$ is a rack subbialgebra of $(B, \Delta, \varepsilon, 1, \mu)$.

Proof: 2. Let $x \in \mathfrak{h}$ and $a \in B$. Since μ is a morphism of C^3 -coalgebras and x is primitive, we get:

$$\begin{aligned} \Delta(a \triangleright x) &= \sum_{(a)} (a^{(1)} \triangleright x) \otimes (a^{(2)} \triangleright \mathbf{1}) + \sum_{(a)} (a^{(1)} \triangleright \mathbf{1}) \otimes (a^{(2)} \triangleright x) \\ &\stackrel{(7)}{=} \sum_{(a)} ((a^{(1)} \varepsilon(a^{(2)})) \triangleright x) \otimes \mathbf{1} + \sum_{(a)} \mathbf{1} \otimes ((\varepsilon(a^{(1)})) a^{(2)} \triangleright x) \\ &= (a \triangleright x) \otimes \mathbf{1} + \mathbf{1} \otimes (a \triangleright x), \end{aligned}$$

whence $a \triangleright x$ is primitive. For the statement on the $B_{(k)}$, we proceed by induction: For $k=0$, this is clear. Suppose the statement is true until $k \in \mathbb{N}$, and let $x \in B_{(k+1)}$. Then:

$$\begin{aligned} &(a \triangleright x) - (a \triangleright x) \otimes \mathbf{1} - \mathbf{1} \otimes (a \triangleright x) \\ &= \sum_{(a)(x)} ((a^{(1)} \triangleright x^{(1)}) \otimes (a^{(2)} \triangleright x^{(2)}) - (a^{(1)} \triangleright x) \otimes (a^{(2)} \triangleright \mathbf{1}) \\ &- (a^{(1)} \triangleright \mathbf{1}) \otimes (a^{(2)} \triangleright x)) \\ &= (\Delta(a)) \triangleright (\Delta(x) - x \otimes \mathbf{1} - \mathbf{1} \otimes x) \\ &= \sum_{(a)(x)} (a^{(1)} \triangleright x^{(1)}) \otimes (a^{(2)} \triangleright x^{(2)}) \end{aligned}$$

where we have used the extended multiplication (still denoted \triangleright) $\triangleright: (B \otimes B) \otimes (B \otimes B) \rightarrow (B \otimes B)$ and set:

$$\Delta(x) - x \otimes \mathbf{1} - \mathbf{1} \otimes x =: \sum_{(x)} x^{(1')} \otimes x^{(2')} \in B_{(k)} \otimes B_{(k)}$$

by the definition of $B_{(k+1)}$, see Appendix 4. By the induction hypothesis, all the terms $a^{(1)} \triangleright x^{(1)}$ and $a^{(2)} \triangleright x^{(2)}$ are in $B_{(k)}$, when $\Delta(a \triangleright x) - (a \triangleright x) \otimes \mathbf{1} - \mathbf{1} \otimes (a \triangleright x)$ is in $B_{(k)} \otimes B_{(k)}$, implying that $a \triangleright x$ is in $B_{(k+1)}$.

1. It follows from 2. that \mathfrak{h} is a subalgebra with respect to μ . Let $x, y, z \in \mathfrak{h}$. Then since x is primitive, it follows from $(x) = x \otimes \mathbf{1} + \mathbf{1} \otimes x$ and the self-distributivity identity (8) that:

$$x \triangleright (y \triangleright z) = (x \triangleright y) \triangleright (\mathbf{1} \triangleright z) + (\mathbf{1} \triangleright y) \triangleright (x \triangleright z) \stackrel{(6)}{=} (x \triangleright y) \triangleright z + y \triangleright (x \triangleright z).$$

proving the left Leibniz identity. The morphism statement is clear, since each morphism of rack bialgebras is a morphism of C^3 -coalgebras and preserves primitives.

Leibniz algebras have been invented by A. M. Blokh in 1965, and then rediscovered by J.-L. Loday in 1992 in the search of an explanation for the absence of periodicity in algebraic K-Theory [10,11].

As an immediate consequence, we get that the functor Prim induces a functor from the category of all rack bialgebras over K to the category of all Leibniz algebras over K .

Remark 2.2: Define set-like elements to be elements a in a rack bialgebra B such that $\Delta(a) = a \otimes a$. Thanks to the fact that \triangleright is a morphism of coalgebras, the set of set-like elements $\text{Slike}(B)$ is closed under \triangleright . In fact, $\text{Slike}(B)$ is a rack, and one obtains in this way a functor $\text{Slike}: \text{RackBialg} \rightarrow \text{Racks}$.

Proposition 2.3: The functor of set-like $\text{Slike}: \text{RackBialg} \rightarrow \text{Racks}$ has the functor $K[-]: \text{Racks} \rightarrow \text{RackBialg}$ (see Example 2.3) as its left-adjoint.

Proof: This follows from the adjointness of the same functors, seen as functors between the categories of pointed sets and of C^1 -coalgebras, observing that the C^1 -coalgebra morphism induced by a morphism of racks respects the rack product.

Observe that the restriction of $\text{Slike}: \text{RackBialg} \rightarrow \text{Racks}$ to the subcategory of connected, cocommutative Hopf algebras Hopf (where the Hopf algebra is given the rack product defined in eqn (12)) gives the usual functor of group-like elements.

(Augmented) rack bialgebras for any Leibniz algebra

Let $(\mathfrak{h}, [\cdot, \cdot])$ be a Leibniz algebra over K , i.e. \mathfrak{h} is a K -module equipped with a K -linear map $[\cdot, \cdot]: \mathfrak{h} \otimes \mathfrak{h} \rightarrow \mathfrak{h}$ satisfying the (left) Leibniz identity (1).

Recall first that each Lie algebra over K is a Leibniz algebra giving rise to a functor from the category of all Lie algebras to the category of all Leibniz algebras.

Furthermore, recall that each Leibniz algebra has two canonical K -submodules:

$$Q(\mathfrak{h}) := \{x \in \mathfrak{h} \mid \exists N \in \mathbb{N} \setminus \{0\}, \exists \lambda_1, \dots, \lambda_N \in K, \exists x_1, \dots, x_N \text{ such that } x = \sum_{r=1}^N \lambda_r [x_r, x_r]\}, \quad (18)$$

$$z(\mathfrak{h}) := \{x \in \mathfrak{h} \mid \forall y \in \mathfrak{h}: [x, y] = 0\}. \quad (19)$$

It is well-known and not hard to deduce from the Leibniz identity that both $Q(\mathfrak{h})$ and $z(\mathfrak{h})$ are two-sided abelian ideals of $(\mathfrak{h}, [\cdot, \cdot])$, that $Q(\mathfrak{h}) \subset z(\mathfrak{h})$, and that the quotient Leibniz algebras:

$$\bar{\mathfrak{h}} := \mathfrak{h} / Q(\mathfrak{h}) \quad \text{and} \quad \mathfrak{g}(\mathfrak{h}) := \mathfrak{h} / z(\mathfrak{h}) \quad (20)$$

are Lie algebras. Since the ideal $Q(\mathfrak{h})$ is clearly mapped into the ideal $Q(\mathfrak{h}')$ by any morphism of Leibniz algebras $\mathfrak{h} \rightarrow \mathfrak{h}'$ (which is a priori not the case for $z(\mathfrak{h})$!), there is an obvious functor $\mathfrak{h} \rightarrow \bar{\mathfrak{h}}$ from the category of all Leibniz algebras to the category of all Lie algebras.

In order to perform the following constructions of rack bialgebras for any given Leibniz algebra $(\mathfrak{h}, [\cdot, \cdot])$, choose first a two-sided ideal $z \subset \mathfrak{h}$ such that:

$$Q(\mathfrak{h}) \subset z \subset z(\mathfrak{h}), \quad (21)$$

Let \mathfrak{g} denote the quotient Lie algebra \mathfrak{h}/z , and let $p: \mathfrak{h} \rightarrow \mathfrak{g}$ be the natural projection. The data of $z \subset \mathfrak{h}$, i.e. of a Leibniz algebra \mathfrak{h} together with an ideal z such that $Q(\mathfrak{h}) \subset z \subset z(\mathfrak{h})$, could be called an augmented Leibniz algebra. Thus we are actually associating an augmented rack bialgebra to every augmented Leibniz algebra. In fact, we will see that

this augmented rack bialgebra does not depend on the choice of the ideal \mathfrak{z} and therefore refrain from introducing augmented Leibniz algebras in a more formal way.

The Lie algebra \mathfrak{g} naturally acts as derivations on \mathfrak{h} by means of (for all $x, y \in \mathfrak{h}$)

$$P(x).y := [x, y] := \text{ad}_x(y) \tag{22}$$

because $\mathfrak{z} \subset \mathfrak{z}(\mathfrak{h})$. Note that:

$$\mathfrak{h} / \mathfrak{z}(\mathfrak{h}) \cong \{ \text{ad}_x \in \text{Hom}_K(\mathfrak{h}, \mathfrak{h}) \mid x \in \mathfrak{h} \}. \tag{23}$$

as Lie algebras.

Consider now the C^5 -coalgebra $(B=S(\mathfrak{h}), \Delta, \epsilon, 1)$ which is actually a commutative cocommutative Hopf algebra over K with respect to the symmetric multiplication \bullet . The linear map $p: \mathfrak{h} \rightarrow \mathfrak{g}$ induces a unique morphism of Hopf algebras:

$$\tilde{\Phi} = S(p) : S(\mathfrak{h}) \rightarrow S(\mathfrak{g}) \tag{24}$$

satisfying

$$\tilde{\Phi}(x_1 \bullet \dots \bullet x_k) = p(x_1) \bullet \dots \bullet p(x_k) \tag{25}$$

for any nonnegative integer k and $x_1, \dots, x_k \in \mathfrak{h}$. In other words, the association $S: V \rightarrow S(V)$ is a functor from the category of all K -modules to the category of all commutative unital C^5 -coalgebras. Consider now the universal enveloping algebra $U(\mathfrak{g})$ of the Lie algebra \mathfrak{g} . Since $\mathbb{Q} \subset K$ by assumption, the Poincaré-Birkhoff-Witt Theorem (in short: PBW) holds [12]. More precisely, the symmetrisation map $\omega: S(\mathfrak{g}) \rightarrow U(\mathfrak{g})$, defined by:

$$\omega(\mathbf{1}_{S(\mathfrak{g})}) = \mathbf{1}_{U(\mathfrak{g})}, \text{ and } \omega(\xi_1 \bullet \dots \bullet \xi_k) = \frac{1}{k!} \sum_{\sigma \in S_k} \xi_{\sigma(1)} \dots \xi_{\sigma(k)}, \tag{26}$$

is an isomorphism of C^5 -coalgebras (in general not of associative algebras) [13]. We now need an action of the Hopf algebra $H=U(\mathfrak{g})$ on B , and an intertwining map $\Phi: B \rightarrow U(\mathfrak{g})$. In order to get this, we first look at \mathfrak{g} -modules: The K -module \mathfrak{h} is a \mathfrak{g} -module by means of eqn (22), the Lie algebra \mathfrak{g} is a \mathfrak{g} -module via its adjoint representation, and the linear map $p: \mathfrak{h} \rightarrow \mathfrak{g}$ is a morphism of \mathfrak{g} -modules since p is a morphism of Leibniz algebras. Now $S(\mathfrak{h})$ and $S(\mathfrak{g})$ are \mathfrak{g} -modules in the usual way, i.e. for all $k \in \mathbb{N} \setminus \{0\}$, $\xi, \xi_1, \dots, \xi_k \in \mathfrak{g}$, and $x_1, \dots, x_k \in \mathfrak{h}$

$$\xi.(x_1 \bullet \dots \bullet x_k) := \sum_{r=1}^k x_1 \bullet \dots \bullet (\xi.x_r) \bullet \dots \bullet x_k, \tag{27}$$

$$\xi.(\xi_1 \bullet \dots \bullet \xi_k) := \sum_{r=1}^k \xi_1 \bullet \dots \bullet [\xi.\xi_r] \bullet \dots \bullet \xi_k, \tag{28}$$

and of course $\xi.1_{S(\mathfrak{h})} = 0$ and $\xi.1_{S(\mathfrak{g})} = 0$. Recall that $U(\mathfrak{g})$ is a \mathfrak{g} -module via the adjoint representation $\text{ad}_\xi(u) = \xi.u = \xi u - u \xi$ (for all $\xi \in \mathfrak{g}$ and all $u \in U(\mathfrak{g})$).

It is easy to see that the map $\tilde{\Phi}$ (25) is a morphism of \mathfrak{g} -modules, and it is well-known that the symmetrization map ω (26) is also a morphism of \mathfrak{g} -modules. Define the K -linear map $\Phi: S(\mathfrak{g}) \rightarrow U(\mathfrak{g})$ by the composition:

$$\Phi := \omega \circ \tilde{\Phi}. \tag{29}$$

Then Φ is a map of C^5 -coalgebras and a map of \mathfrak{g} -modules. Thanks to the universal property of the universal enveloping algebra, it follows that $S(\mathfrak{h})$ and $U(\mathfrak{g})$ are left $U(\mathfrak{g})$ -modules, via (for all $\xi_1, \dots, \xi_k \in \mathfrak{g}$, and for all $a \in S(\mathfrak{h})$)

$$(\xi_1 \dots \xi_k).a = \xi_1.(\xi_2.(\dots \xi_k.a) \dots) \tag{30}$$

and the usual adjoint representation (12) (for all $u \in U(\mathfrak{g})$)

$$\text{ad}_{x_1 \dots x_k}(u) = (\text{ad}_{x_1} \circ \dots \circ \text{ad}_{x_k})(u), \tag{31}$$

and that Φ intertwines the $U(\mathfrak{g})$ -action on $C=S(\mathfrak{h})$ with the adjoint action of $U(\mathfrak{g})$ on itself.

Finally it is a routine check using the above identities (27) and (12) that $S(\mathfrak{h})$ becomes a module coalgebra.

We can resume the preceding considerations in the following:

Theorem 2.1: Let $(\mathfrak{h}, [\cdot, \cdot])$ be a Leibniz algebra over K , let \mathfrak{z} be a two-sided ideal of \mathfrak{h} such that $Q(\mathfrak{h}) \subset \mathfrak{z} \subset Z(\mathfrak{h})$, let \mathfrak{g} denote the quotient Lie algebra $\mathfrak{h}/\mathfrak{z}$ by \mathfrak{g} , and let $p: \mathfrak{h} \rightarrow \mathfrak{g}$ be the canonical projection.

1. Then there is a canonical $U(\mathfrak{g})$ -action ℓ on the C^5 -coalgebra $B:=S(\mathfrak{h})$ (making it into a module coalgebra leaving invariant 1) and a canonical lift of p to a map of C^5 -coalgebras, $\Phi: S(\mathfrak{h}) \rightarrow U(\mathfrak{g})$ such that eqn (14) holds.

Hence the quadruple $(S(\mathfrak{h}), \Phi, U(\mathfrak{g}), \ell)$ is an augmented rack bialgebra whose associated Leibniz algebra is equal to $(\mathfrak{h}, [\cdot, \cdot])$ (independently of the choice of \mathfrak{z}).

The resulting rack multiplication μ of $S(\mathfrak{h})$ (written $\mu(a \otimes b) = a \triangleright b$) is also independent on the choice of \mathfrak{z} and is explicitly given as follows for all positive integers k, l and $x_1, \dots, x_k, y_1, \dots, y_l \in \mathfrak{h}$:

$$(x_1 \bullet \dots \bullet x_k) \triangleright (y_1 \bullet \dots \bullet y_l) = \frac{1}{k! l!} \sum_{\sigma \in S_k} (\text{ad}_{x_{\sigma(1)}}^s \circ \dots \circ \text{ad}_{x_{\sigma(k)}}^s)(y_1 \bullet \dots \bullet y_l) \tag{32}$$

where ad_x^s denotes the action of the Lie algebra $\mathfrak{h}/\mathfrak{z}(\mathfrak{h})$ (see eqn (23)) on $S(\mathfrak{h})$ according to eqn (27).

2. In case $\mathfrak{z}=Q(\mathfrak{h})$, the construction mentioned in 1. is a functor $\mathfrak{h} \rightarrow \text{UAR}^\infty(\mathfrak{h})$ from the category of all Leibniz algebras to the category of all augmented rack bialgebras associating to \mathfrak{h} the rack bialgebra:

$$\text{UAR}^\infty(\mathfrak{h}) := (S(\mathfrak{h}), \Phi, U(\mathfrak{g}), \ell)$$

and to each morphism f of Leibniz algebras the pair $(S(f), U(\bar{f}))$ where \bar{f} is the induced Lie algebra morphism.

3. For each nonnegative integer k , the above construction restricts to each subcoalgebra of order k , $S(\mathfrak{h})_{(k)} = \bigoplus_{r=0}^k S^r(\mathfrak{h})$, to define an augmented rack bialgebra $(S(\mathfrak{h})_{(k)}, \Phi_{(k)}, U(\mathfrak{g}), \ell|_{U(\mathfrak{g}) \otimes S(\mathfrak{h})_{(k)}})$ which in case $\mathfrak{z}=Q(\mathfrak{h})$ defines a functor $\mathfrak{h} \rightarrow \text{UAR}_{(k)}^\infty(\mathfrak{h}) := (\text{UAR}^\infty(\mathfrak{h}))_{(k)}$ from the category of all Leibniz algebras to the category of all augmented rack bialgebras.

Remark 2.3: This theorem should be compared to Proposition 3.5. [3]. In [3], the authors work with the vector space $N := K \oplus \mathfrak{h}$, while we work with the whole symmetric algebra on the Leibniz algebra. In some sense, we extend their Proposition 3.5 “to all orders”. However, as we shall see below, N is already enough to obtain a left-adjoint to the functor of primitives.

The above rack bialgebra associated to a Leibniz algebra \mathfrak{h} can be seen as one version of an enveloping algebra of \mathfrak{h} .

Definition 2.3: Let \mathfrak{h} be a Leibniz algebra. We will call the augmented rack bialgebra $(S(\mathfrak{h}), \Phi, U(\mathfrak{g}), \ell)$ the enveloping algebra of infinite order of \mathfrak{h} . As such, it will be denoted by $\text{UAR}^\infty(\mathfrak{h})$.

This terminology is justified, for example, by the fact that \mathfrak{h} is identified to the primitives in $S(\mathfrak{h})$ (cf Proposition 2.2). This is also justified by the following theorem the goal of which is to show that the enveloping algebra $\text{UAR}^\infty(\mathfrak{h})$ fits into the following diagram of functors:

$$\text{Lie}[r]^U[d]^i \text{Hopf}[d]^i \text{Leib}[r] \xrightarrow{\text{UAR}^\infty} \text{RackBial}$$

Here, i is the embedding functor of Lie algebras into Leibniz algebras, and j is the embedding functor of the category of connected, cocommutative Hopf algebras into the category of rack bialgebras, using the adjoint action (see eqn (12)) as a rack product.

Theorem 2.2: Let \mathfrak{g} be a Lie algebra. The PBW isomorphism $U(\mathfrak{g}) \cong S(\mathfrak{g})$ induces an isomorphism of functors:

$$j \circ U \cong \text{UAR} \circ i.$$

Proof: The enveloping algebra $\text{UAR}^\circ(\mathfrak{h})$ is by definition the functorial version of the rack bialgebra $S(\mathfrak{h})$, i.e. associated to the ideal $Q(\mathfrak{h})$. But in case \mathfrak{h} is a Lie algebra, $Q(\mathfrak{h}) = \{0\}$. Then the map p is simply the identity, and $\text{UAR}^\circ(\mathfrak{h}) = j(U(\mathfrak{h}))$.

As a relatively easy corollary we obtain from the preceding construction the computation of universal rack bialgebras. More precisely, we look for a left adjoint functor for the functor Prim , seen as a functor from the category of all rack bialgebras to the category of all Leibniz algebras. For a given Leibniz algebra $(\mathfrak{h}, [,])$ define the subcoalgebra of order 1 of the first component of $\text{UAR}_{(1)}(\mathfrak{h})$ (see the third statement of Theorem 2.1), i.e.

$$\text{UR}(\mathfrak{h}) := \text{UAR}^\circ(\mathfrak{h})_{(1)} : K \oplus \mathfrak{h} \quad (33)$$

With $1 := 1_K$ which is rack subbialgebra according to Proposition 2.2. Its structure reads for all $\lambda, \lambda' \in K$ and for all $x, y \in \mathfrak{h}$

$$\Delta(\lambda 1 + x) = \lambda 1 \otimes 1 + x \otimes 1 + 1 \otimes x, \quad (34)$$

$$\varepsilon(\lambda 1 + x) = \lambda, \quad (35)$$

$$\mu((\lambda 1 + x) \otimes (\lambda' 1 + x')) = \lambda \lambda' 1 + \lambda x' + [x, x']. \quad (36)$$

For the particular case of a Lie algebra $(\mathfrak{h}, [,])$, the above construction can be found in [3]. Moreover, for any other Leibniz algebra $(\mathfrak{h}', [,])'$ and any morphism of Leibniz algebras $f: \mathfrak{h} \rightarrow \mathfrak{h}'$ define the K -linear map $\text{UR}(f): \text{UR}(\mathfrak{h}) \rightarrow \text{UR}(\mathfrak{h}')$ as the first component of $\text{UAR}_{(1)}(f)$ (cf. the third statement of Theorem 2.1) by

$$\text{UR}(f)(\lambda \hat{1} + x) = \lambda \hat{1}' + f(x), \quad (37)$$

which is clearly a morphism of rack bialgebras. Hence UR is a functor from the category of all Leibniz algebras to the category of all rack bialgebras. Now let $(C, \Delta_C, \varepsilon_C, 1_C, \mu_C)$ be a rack bialgebra, and let $f: \mathfrak{h} \rightarrow \text{Prim}(C)$ be a morphism of Leibniz algebras. Define the K -linear map $\hat{f}: \text{UR}(\mathfrak{h}) \rightarrow C$ by

$$\hat{f}(\lambda \hat{1} + x) = \lambda 1_C + f(x), \quad (38)$$

and it is again a routine check that it defines a morphism of rack bialgebras. Moreover, due to the almost trivial coalgebra structure of $\text{UR}(\mathfrak{h})$, it is clear that any morphism of rack bialgebras $\text{UR}(\mathfrak{h}) \rightarrow C$ is of the above form and is uniquely determined by $\text{Prim}(\hat{f}) = f$. Hence we have shown the following:

Theorem 2.3: There is a left adjoint functor, UR , for the functor Prim (associating to each Rack bialgebra its Leibniz algebra of all primitive elements). For a given Leibniz algebra $(\mathfrak{h}, [,])$, the object $\text{UR}(\mathfrak{h})$ – which we shall call the Universal Rack Bialgebra of the Leibniz algebra $(\mathfrak{h}, [,])$ – has the usual universal properties.

Next we can refine the above universal construction by taking into account the augmented rack bialgebra structure of $\text{UAR}^\circ(\mathfrak{h})$ to define another universal object. Consider the more detailed category of all augmented rack bialgebras. Again, the functor Prim applied to the coalgebra B (and not to the Hopf algebra H) gives a functor from the

first category to the category of all Leibniz algebras, and we seek again a left adjoint of this functor, called UAR . Hence, a natural candidate for a universal augmented rack bialgebra associated to a given Leibniz algebra \mathfrak{h} is:

$$\text{UAR}(\mathfrak{h}) := \text{UAR}_{(1)}(\mathfrak{h}) = (K \oplus \mathfrak{h}, \Phi_{(1)}, U(\bar{\mathfrak{h}}), \text{ad}^s |_{U(\bar{\mathfrak{h}}) \otimes (K \oplus \mathfrak{h})}). \quad (39)$$

The third statement of Theorem 2.1 tells us that this is a well-defined augmented rack bialgebra, and that UAR is a functor from the category of all Leibniz algebras to the category of all augmented rack bialgebras. Now let (B', Φ', H', I') be an augmented rack bialgebra, and let $f: \mathfrak{h} \rightarrow \text{Prim}(B')$ be a morphism of Leibniz algebras. Clearly, as has been shown in Theorem 2.3, the map $\hat{f}: \text{UR}(\mathfrak{h}) \rightarrow B'$ given by eqn (37) is a morphism of rack bialgebras. Observe that the morphism of C^3 -coalgebras Φ' sends the Leibniz subalgebra $\text{Prim}(B')$ of B' into K -submodule of all primitive elements of the Hopf algebra H' , $\text{Prim}(H')$, which is known to be a Lie subalgebra of H' equipped with the commutator Lie bracket $[,]_{H'}$. Moreover this restriction is a morphism of Leibniz algebras. Indeed, for any $x', y' \in \text{Prim}(B')$ we have

$$\begin{aligned} \Phi'([x', y']) &= \Phi'(x' \triangleright y') = \Phi'((\Phi'(x'), y')) = \text{ad}_{\Phi'(x')}(\Phi'(y')) \\ &= \Phi'(x')\Phi'(y') - \Phi'(y')\Phi'(x') = [\Phi'(x'), \Phi'(y')]_{H'}. \end{aligned}$$

It follows that the two-sided ideal $Q(\text{Prim}(B'))$ of the Leibniz algebra $\text{Prim}(B')$ is in the kernel of the restriction of Φ' to $\text{Prim}(B')$, whence the map $\bar{\Phi}'$ induces a well-defined K -linear morphism of Lie algebras $\bar{\Phi}': \text{Prim}(B') \rightarrow \text{Prim}(H')$. It follows that the composition $\bar{\Phi}' \circ \bar{f}: \mathfrak{h} \rightarrow \text{Prim}(H') \subset H'$ is a morphism of Lie algebras, and by the universal property of universal enveloping algebras there is a unique morphism of associative unital algebras $\psi := U(\bar{\Phi}' \circ \bar{f}): U(\bar{\mathfrak{h}}) \rightarrow H'$. But we have for all $\xi_1, \dots, \xi_k \in \bar{\mathfrak{h}}$

$$\begin{aligned} \Delta_{H'}(\psi(\xi_1 \dots \xi_k)) &= \Delta_{H'}(\psi(\xi_1) \dots \psi(\xi_k)) = \Delta_{H'}(\xi_1) \dots \Delta_{H'}(\xi_k) \\ &= (\psi(\xi_1) \otimes 1_{H'} + 1_{H'} \otimes \psi(\xi_1)) \dots (\psi(\xi_k) \otimes 1_{H'} + 1_{H'} \otimes \psi(\xi_k)) \\ &= (\psi \otimes \psi)(\xi_1 \otimes 1_{U(\bar{\mathfrak{h}})} + 1_{U(\bar{\mathfrak{h}})} \otimes \xi_1) \dots (\psi \otimes \psi)(\xi_k \otimes 1_{U(\bar{\mathfrak{h}})} + 1_{U(\bar{\mathfrak{h}})} \otimes \xi_k) \\ &= (\psi \otimes \psi)(\Delta_{U(\bar{\mathfrak{h}})}(\xi_1)) \dots (\psi \otimes \psi)(\Delta_{U(\bar{\mathfrak{h}})}(\xi_k)) \\ &= (\psi \otimes \psi)(\Delta_{U(\bar{\mathfrak{h}})}(\xi_1 \dots \xi_k)) \end{aligned}$$

since ψ maps primitives to primitives whence ψ is a morphism of coalgebras. It is easy to check that ψ preserves counits, whence ψ is a morphism of C^5 -Hopf-algebras. For all $\lambda \in K$ and $x \in \mathfrak{h}$ we get:

$$\begin{aligned} (\psi \circ \Phi_{(1)})(\lambda \hat{1} + x) &= \psi(\lambda 1_{U(\bar{\mathfrak{h}})} + p(x)) = \lambda 1_{H'} + (\bar{\Phi}' \circ \bar{f})(p(x)) \\ &= \lambda 1_{H'} + \Phi'(f(x)) = \Phi'(\hat{f}(\lambda \hat{1} + x)), \end{aligned}$$

showing the first equation $\psi \circ \Phi_{(1)} = \Phi' \circ \hat{f}$ of the morphism equation (15). Moreover for all $\lambda \in K, x \in \mathfrak{h}$, and $u \in U(\bar{\mathfrak{h}})$ we get

$$\hat{f}(u(\lambda \hat{1} + x)) = \hat{f}(\lambda \varepsilon_{U(\bar{\mathfrak{h}})}(u) \hat{1} + u x) = \lambda \varepsilon_{U(\bar{\mathfrak{h}})}(u) 1_C + f(u x)$$

Let $x_1, \dots, x_k \in \mathfrak{h}$ such that $u = p(x_1) \dots p(x_k)$. Then

$$\begin{aligned} f(u x) &= f([x_1, [x_2, \dots [x_k, x] \dots]]) = f(x_1) \triangleright f(x_2) \triangleright \dots \triangleright f(x_k) \triangleright f(x) \dots \\ &= (\Phi'(f(x_1)) \dots \Phi'(f(x_k))) \cdot (f(x)) \\ &= (\psi(p(x_1)) \dots \psi(p(x_k))) \cdot (f(x)) = (\psi(u)) \cdot (f(x)), \end{aligned}$$

and therefore

$$\hat{f}(u(\lambda \hat{1} + x)) = (\psi(u)) \cdot (\hat{f}(\lambda \hat{1} + x))$$

showing the second equation $\psi \circ \Phi_{(1)} = \Phi' \circ \hat{f}$ of the morphism equation (15). It follows that the pair (\hat{f}, ψ) is a morphism of augmented rack bialgebras. We therefore have the following

Theorem 2.4: There is a left adjoint functor, UAR, for the functor Prim (associating to each augmented rack bialgebra its Leibniz algebra of all primitive elements). For a given Leibniz algebra $(\mathfrak{h}, [,])$, the object $\text{UAR}(\mathfrak{h})$ –which we shall call the Universal Augmented Rack Bialgebra of the Leibniz algebra $(\mathfrak{h}, [,])$ – has the usual universal properties.

The relationship between the different notions (taking into account also Remark (2.2)) is resumed in the following diagram:

$$\begin{array}{c} \text{Lie} @ < \text{lex} > [r]^{\text{U}} @ \langle - \rangle [d]^{\text{P}} \text{Hopf} @ \langle - \rangle [d]^{\text{P}} [l]^{\text{Prim}} @ < \text{lex} > \\ [r]^{\text{Slike}} \text{Grp} [l]^{\text{K}[-]} @ \langle - \rangle [d]^{\text{L}} \text{Leib} @ < \text{lex} > [r]^{\text{UAR}^{\infty}} \\ \text{RackBialg} [l]^{\text{Prim}} @ < \text{lex} > [r]^{\text{Slike}} \text{Racks} [l]^{\text{K}[-]} \end{array}$$

where UAR is not left-adjoint to Prim, while UR is, but does not render the square commutative. There is a similar diagram for augmented notions.

Relation with bar-unital di(co)algebras

In the beginning of the nineties the ‘enveloping structure’ associated to Leibniz algebras has been the structure of dialgebras. We shall show in this section that rack bialgebras and certain cocommutative Hopf dialgebras are strongly related.

Left-unital bialgebras and right Hopf algebras: Let $(B, \Delta, \varepsilon, 1, \mu)$ be a nonassociative left-unital C^3 -bialgebra. It will be called left-unital C^3 -bialgebra iff μ is associative. In general, $(B, \Delta, \varepsilon, 1)$ need not be unital, i.e. we do not have in general $a1=a$. However, it is easy to see that the sub-module $B1$ of B is a C^3 -subcoalgebra of $(B, \Delta, \varepsilon, 1)$, and a subalgebra of (B, μ) such that $(B1, \Delta', \varepsilon', 1, \mu')$ is a unital (i.e. left-unital and right-unital) bialgebra. Here Δ', ε' , and μ' denote the obvious restrictions and corestrictions.

In a completely analogous way right-unital C^3 -bialgebras are defined.

A left-unital (resp. right unital) cocommutative C^3 -bialgebra $(B, \Delta, \varepsilon, 1, \mu)$ will be called a cocommutative right Hopf algebra (resp. a cocommutative left Hopf algebra), $(B, \Delta, \varepsilon, 1, \mu, S)$, iff there is a right antipode S (resp. left antipode S), i.e. there is a K -linear map $S: B \rightarrow B$ which is a morphism of C^3 -coalgebras $(B, \Delta, \varepsilon, 1)$ to itself such that

$$\text{id}^* S = 1\varepsilon \text{ (resp. } S^* \text{id} = 1) \tag{40}$$

where $*$ denotes the convolution product (see Appendix 4 for definitions). It will become clear a posteriori that right or left antipodes are always unique, see Lemma 2.2.

A first class of examples is of course the well-known class of all cocommutative Hopf algebras $(H, \Delta, \varepsilon, 1, \mu, S)$ for which 1 is a unit element, and S is a right and left antipode.

Secondly it is easy to check that every C^4 -coalgebra $(C, \Delta, \varepsilon, 1)$ equipped with the left-trivial multiplication (resp. right trivial multiplication) μ_0 (see eqn (5)) and trivial right antipode (resp. trivial left antipode) S_0 defined by $S_0(x) = \varepsilon(x)1$ for all $x \in C$ (in both cases) is a cocommutative right Hopf algebra (resp. cocommutative left Hopf algebra) called the cocommutative left-trivial right Hopf algebra (resp. right-trivial left Hopf algebra) defined by the C^4 -coalgebra $(C, \Delta, \varepsilon, 1)$.

We have the following elementary properties showing in particular that each right (resp. left) antipode is unique:

Lemma 2.2: Let $(\mathcal{H}, \Delta, \varepsilon, 1, \mu, S)$ be a cocommutative right Hopf algebra.

- 1. $S^*(S^{\circ}S) = 1\varepsilon$, $S^{\circ}(S^{\circ}S) = \text{id}^* 1\varepsilon$, $S^* 1\varepsilon = S$ and $S^{\circ}S^{\circ}S = S$, which for each

$a \in \mathcal{H}$ implies $\sum_{(a)} a^{(1)} S(a^{(2)}) = 1\varepsilon(a) = \sum_{(a)} S(a^{(1)}) a^{(2)} 1$. It follows that right antipodes are unique.

- 2. For all $a, b \in \mathcal{H}: S(ab) = S(b)S(a)$.

3. For any element $c \in \mathcal{H}$, c is a generalized idempotent if and only if $c = \sum_{(c)} S(c^{(1)}) c^{(2)}$ iff there is $x \in \mathcal{H}$ with $c = \sum_{(x)} S(x^{(1)}) x^{(2)}$, and all these three statements imply that c is a generalized left unit element.

Proof: 1. Since S is a coalgebra morphism, it preserves convolutions when composing from the right. This gives the first equation from statement 2.2. Hence the elements id, S , and $S^{\circ}S$ satisfy the hypotheses of the elements a, b, c of Lemma 5.1 in the left-unital convolution semigroup $(\text{Hom}_K(\mathcal{H}, \mathcal{H}), *, 1\varepsilon)$, whence the second and third equations of statement 2.2. are immediate, and the fourth follows from composing the second from the right with S and using the third. Clearly S is unique according to Lemma 5.1.

2. Again the elements $\mu, S^{\circ}\mu$ and $(\text{id}^* 1\varepsilon)^{\circ}\mu$ satisfy the hypotheses on the elements a, b, c of Lemma 5.1 in the left-unital convolution semigroup $(\text{Hom}_K(\mathcal{H} \otimes \mathcal{H}, \mathcal{H}), *, 1\varepsilon \otimes \varepsilon)$ (using the fact that μ is a morphism of coalgebras) whence $S^{\circ}\mu$ is the unique right inverse of μ . A computation shows that also $\mu^{\circ} \tau^{\circ} (S^{\circ}S)$ is a right inverse of μ , whence we get statement 2.2. by uniqueness of right inverses (Lemma 5.1).

3. The second statement obviously implies the third, and it is easy to see by straight-forward computations that the third statement implies the first and the second. Conversely, if $c \in \mathcal{H}$ is a generalized idempotent, i.e. $c = \sum_{(c)} S(c^{(1)}) c^{(2)}$, we get –since Δ is a morphism of C^3 -coalgebras– that

$$\begin{aligned} \sum_{(c)} S(c^{(1)}) c^{(2)} &= \sum_{(c)} S(c^{(1)}) ((\mu \circ \Delta)(c))^{(2)} = \sum_{(c)} S(c^{(1)}) (\mu \circ \Delta)(c^{(2)}) \\ &= \sum_{(c)} S(c^{(1)}) c^{(2)} c^{(3)} = \sum_{(c)} S(c^{(1)}) c^{(2)} 1 c^{(3)} \stackrel{\text{Lemma 2.2.1.}}{=} c, \end{aligned}$$

and all the three statements are equivalent. In order to see that every such element c is a generalized left unit element pick $y \in \mathcal{H}$ and

$$cy = \sum_{(x)} S(x^{(1)}) x^{(2)} y = \sum_{(x)} S(x^{(1)}) x^{(2)} 1 y \stackrel{\text{Lemma 2.2.1.}}{=} \varepsilon(x) y = \varepsilon(c) y$$

since obviously $\varepsilon(c) = \varepsilon(x)$, so c is a generalized left unit element.

There is the following right Hopf algebra analogue of the Suschkewitsch decomposition theorem for right groups (see Appendix 5):

Theorem 2.5: Let $(\mathcal{H}, \Delta, \varepsilon, 1, \mu, S)$ a cocommutative right Hopf algebra. Then the following holds:

- 1. The K -submodule $(\mathcal{H}1, \Delta|_{\mathcal{H}1}, \varepsilon|_{\mathcal{H}1}, \mu|_{\mathcal{H}1}, \mu|_{\mathcal{H}1} \otimes_{\mathcal{H}1}, S|_{\mathcal{H}1})$ is a unital Hopf subalgebra of $(\mathcal{H}, \Delta, \varepsilon, 1, \mu, S)$.
- 2. The K -submodule $E_{\mathcal{H}} := \{x \in \mathcal{H} | x \text{ is a generalized idempotent}\}$ is a right Hopf subalgebra of \mathcal{H} equal to the left-trivial right Hopf algebra defined by the C^4 -coalgebra $(E_{\mathcal{H}}, \Delta|_{E_{\mathcal{H}}}, \varepsilon|_{E_{\mathcal{H}}}, 1\varepsilon|_{E_{\mathcal{H}}})$.
- 3. The map

$$\Psi : \mathcal{H} \rightarrow \mathcal{H}1 \otimes E_{\mathcal{H}} : x \mapsto \sum_{(x)} x^{(1)} 1 \otimes S(x^{(2)}) x^{(3)}$$

is an isomorphism of right Hopf algebras whose inverse Ψ^{-1} is the restriction of the multiplication map.

Proof: 1. It is easy to see that $\mathcal{H}1$ equipped with all the restrictions is a unital bialgebra. Note that for all $a \in \mathcal{H}$

$$(S|_{\mathcal{H}1} * \text{id}_{\mathcal{H}1})(a1) = ((S^* \text{id}_{\mathcal{H}})(a))1 = (S^* \text{id}_{\mathcal{H}} * (1\varepsilon))(a)$$

$$\stackrel{\text{Lemma 2.2}}{=} (\mathbf{1}\varepsilon)(a) = (\mathbf{1}\varepsilon)(a\mathbf{1}) = (\mathbf{1}\varepsilon|_{\mathcal{H}\mathbf{1}})(a\mathbf{1})$$

Whence $S|_{\mathcal{H}\mathbf{1}}$ is also a left antipode. It follows that $\mathcal{H}\mathbf{1}$ is a Hopf algebra.

2. Since the property of being a generalized idempotent is a K -linear condition, it follows that $E_{\mathcal{H}}$ is a K -submodule. Moreover since each $c \in E_{\mathcal{H}}$ is of the general form $c = \sum_{(x)} S(x^{(1)})x^{(2)}$, $x \in \mathfrak{t}$, and since the map $\iota: \mathcal{H} \rightarrow \mathcal{H}$ defined by $\iota(x) = \sum_{(x)} S(x^{(1)})x^{(2)}$ is an idempotent morphism of C^3 -coalgebras, we get

$$\Delta(c) = \Delta(\iota(c)) = (\iota \otimes \iota)(\Delta(x))$$

showing that $E_{\mathcal{H}}$ is a C^3 -subcoalgebra of \mathcal{H} . Furthermore, since every element of $E_{\mathcal{H}}$ is a generalized left unit element (Lemma 2.2, 3.), the restriction of the multiplication of μ of \mathcal{H} to $E_{\mathcal{H}} \otimes E_{\mathcal{H}}$ is left trivial. Finally,

$$S(c) = S(\iota(c)) = \sum_{(c)} S(S(c^{(1)})c^{(2)}) \stackrel{\text{Lemma 2.2,2.}}{=} \sum_{(c)} S(c^{(1)})S(S(c^{(2)}))$$

$$\stackrel{\text{Lemma 2.22,1.}}{=} \sum_{(c)} S(c^{(1)})c^{(2)}\mathbf{1} \stackrel{\text{Lemma 2.22,2,1.}}{=} \varepsilon(c)\mathbf{1} = S_0(c),$$

showing that the restriction of S to $E_{\mathcal{H}}$ is the trivial right antipode.

3. It is clear from the two preceding statements that Ψ is a well-defined linear map into the tensor product of two cocommutative right Hopf algebras. We have for all $x \in \mathcal{H}$

$$(\mu \circ \Psi)(x) = \sum_{(x)} x^{(1)}\mathbf{1}S(x^{(2)})x^{(3)} \stackrel{\text{Lemma 2.2,1.}}{=} \sum_{(x)} \varepsilon(x^{(1)})x^{(2)} = x$$

and for all $a \in \mathcal{H}$, $c \in E_{\mathcal{H}}$ the term $(\Psi^0\mu)(a\mathbf{1} \otimes c)$ is equal to

$$\sum_{(a)(c)} (a^{(1)}c^{(1)}\mathbf{1}) \otimes (S(c^{(2)})S(a^{(2)})a^{(3)}c^{(3)}) \\ = \sum_{(c)} (ac^{(1)}\mathbf{1}) \otimes (S(c^{(2)})c^{(3)}) = \sum_{(c)} (a\mathbf{1}) \otimes (S(c^{(1)})c^{(2)}) = (a\mathbf{1}) \otimes c$$

because all the terms $S(a^{(2)})a^{(3)}$ and the components $c^{(1)}, \dots$ of iterated comultiplications of generalized idempotents can be chosen in $E_{\mathcal{H}}$ (since the latter has been shown to be a subcoalgebra), and are thus generalized left unit elements (Lemma 2.2, 3.). Hence Ψ is a K -linear isomorphism. Moreover, it is easy to see from its definition that Ψ is a morphism of C^3 -coalgebras.

Next we compute for all $a, a' \in \mathcal{H}$ and $c, c' \in E_{\mathcal{H}}$:

$$\Psi^{-1}(((a\mathbf{1}) \otimes c)((a'\mathbf{1}) \otimes c')) = \Psi^{-1}((a\mathbf{1}a') \otimes \varepsilon(c)c') = (c)aa'c',$$

and -since c is a generalized left unit element-

$$\Psi^{-1}(((a\mathbf{1}) \otimes c) \Psi^{-1}((a'\mathbf{1}) \otimes c')) = acac' = \varepsilon(c)aa'c',$$

showing that Ψ^{-1} and hence Ψ is a morphism of left-unital algebras. Finally we obtain

$$(S|_{\mathcal{H}\mathbf{1}} \otimes S_0)(\Psi(x)) = \sum_{(x)} (S(x^{(1)})\mathbf{1}) \otimes (\varepsilon(S(x^{(2)})x^{(3)})\mathbf{1}) = S(x)\mathbf{1} \otimes \mathbf{1},$$

$$\Psi(S(x)) = \sum_{(x)} (S(x^{(1)})\mathbf{1}) \otimes (S(S(x^{(2)}))S(x^{(3)})) = S(x)\mathbf{1} \otimes \mathbf{1},$$

thanks to Lemma 2.2, and intertwines right antipodes.

Note that the K -submodule of all generalized left unit elements of a right Hopf algebra \mathcal{H} is given by $K\mathbf{1} \oplus (\Phi^{-1}(\mathcal{H}\mathbf{1} \otimes E_{\mathcal{H}}^+))$ and thus in general much bigger than the submodule $E_{\mathcal{H}}$ of all generalized idempotents.

As it is easy to see that every tensor product $H \otimes C$ of a unital cocommutative Hopf algebra H and a C^3 -coalgebra C (equipped with

the left-trivial multiplication and the trivial right antipode) is a right Hopf algebra, it is a fairly routine check -using the preceding Theorem 2.5- that the category of all cocommutative right Hopf algebras is equivalent to the product category of all cocommutative Hopf algebras and of all C^3 -coalgebras.

In the sequel, we shall need the dual left Hopf algebra version where all the formulas have to be put in reverse order: Here every left Hopf algebra is isomorphic to $C \otimes H$.

Dialgebras and Rack Bialgebras: Recall that a dialgebra over K is a K -module D equipped with two associative multiplications $\vdash, \dashv: A \otimes A \rightarrow A$ (written $ab \mapsto a \vdash b$ and $ab \mapsto a \dashv b$) satisfying for all $a, b, c \in A$:

$$(a \vdash b) \vdash c = (a \vdash b) \vdash c, \tag{41}$$

$$a \dashv (b \vdash c) = a \dashv (b \vdash c), \tag{42}$$

$$(a \vdash b) \dashv c = a \vdash (b \dashv c). \tag{43}$$

An element 1 of A is called a bar-unit element of the dialgebra (A, \vdash, \dashv) and $(A, 1, \vdash, \dashv)$ is called a bar-unital dialgebra iff in addition the following holds

$$1 \vdash a = a, \tag{44}$$

$$a \dashv 1 = a, \tag{45}$$

for all $a \in A$. Moreover, we shall call a bar-unital dialgebra $(A, 1, \vdash, \dashv)$ balanced iff in addition for all $a \in A$

$$a \vdash 1 = 1 \dashv a. \tag{46}$$

Clearly each associative algebra is a dialgebra upon setting $\vdash = \dashv$ equal to the given multiplication. The class of all (bar-unital and balanced) dialgebras forms a category where morphisms preserve both multiplications and map the initial bar-unit to the target bar-unit.

These algebras had been introduced to have a sort of 'associative analogue' for Leibniz algebras. More precisely, there is the following important fact, which can easily be checked:

Proposition 2.4: Let (A, \vdash, \dashv) be a dialgebra. Then the K -module A equipped with the bracket $[,]: A \otimes A \rightarrow A$, written $[a, b]$,

$$[a, b] := a \vdash b - b \dashv a \tag{47}$$

is a Leibniz algebra, denoted by A^- .

In fact, this construction is well-known to give rise to a functor $A \rightarrow A^-$ from the category of all dialgebras to the category of all Leibniz algebras in complete analogy to the obvious functor from the category of all associative algebras to the category of all Lie algebras.

An important construction of (bar-unital) dialgebras is the following:

Example 2.5: Let $(B, 1_B)$ be a unital associative algebra over K , and let A be a K -module which is a B -bimodule, i.e. there are K -linear maps $B \otimes A \rightarrow A$ and $A \otimes B \rightarrow A$ (written $(b \otimes x) \mapsto bx$ and $(x \otimes b) \mapsto xb$) equipping A with the structure of a left B -module and a right B -module such that $(bx)b' = b(xb')$ for all $b, b' \in B$ and for all $x \in B$. Suppose in addition that there is a bimodule map $\Phi: A \rightarrow B$, i.e. $\Phi(bxb') = b\Phi(x)b'$ for all $b, b' \in B$ and for all $x \in A$. Then it is not hard to check that the two multiplications $\vdash, \dashv: A \otimes A \rightarrow A$ defined by

$$x \vdash y := \Phi(x)y \text{ and } x \dashv y := x\Phi(y) \tag{48}$$

equip A with the structure of a dialgebra. If in addition there is an element $1 \in A$ such that $\Phi(1) = 1_B$, then $(A, 1, \vdash, \dashv)$ will be a bar-unital dialgebra. We shall call this structure (A, Φ, B) an augmented dialgebra.

In fact, every dialgebra (A, \vdash, \dashv) arises in that fashion: Consider the K -submodule $I \subset A$ whose elements are linear combinations of arbitrary product expressions

$$p(a_1, \dots, a_{r-1}, (a_r \vdash b_r - a_r \dashv b_r), a_{r+1}, \dots, a_n)$$

(where all reasonable parentheses and symbols \vdash and \dashv are allowed) for any two strictly positive integer $r \leq n$, and $a_1, \dots, a_n, b_r \in A$. It follows that the quotient module A/I is equipped with an associative multiplication induced by both \vdash and \dashv . Let A_{ass}^1 be equal to A/I if A is bar-unital: In that case, the bar-unit 1 of A projects on the unit element of A/I ; and let A_{ass}^1 be equal to $A/I \otimes K$ (adjoining a unit element) in case A does not have a bar-unit. Thanks to the defining equations (41), (42), (43), it can be shown by induction that for any strictly positive integer n , any $a_1, \dots, a_n \in A$, and any product expression made of the preceding elements upon using \vdash or \dashv

$$p(a_1, \dots, a_n) \vdash a = a_1 \vdash \dots \vdash a_n \vdash a = (a_1 \dashv \dots \dashv a_n) \vdash a, \\ a \dashv p(a_1, \dots, a_n) = a \dashv a_1 \dashv \dots \dashv a_n \dashv a = a \dashv (a_1 \vdash \dots \vdash a_n),$$

proving in particular that I acts trivially from the left (via \vdash) and from the right (via \dashv) on A such that there is a well-defined A_{ass}^1 -bimodule structure on A such that the natural map $\Phi_A: A \rightarrow A_{\text{ass}}^1$ is a bimodule morphism. Hence $(A, \Phi_A, A_{\text{ass}}^1)$ is always an augmented dialgebra, and the assignment $A \rightarrow (A, \Phi_A, A_{\text{ass}}^1)$ is known to be a faithful functor.

Note also that this construction allows to adjoin a bar-unit to a dialgebra (A, \vdash, \dashv) : Consider the K -module $\tilde{A} := A \oplus A_{\text{ass}}^1$ with the obvious A_{ass}^1 -bimodule structure $\alpha \cdot (b + \beta) = \alpha \cdot b + \alpha \beta$ and $(b + \beta) \cdot \alpha = b \cdot \alpha + \beta \alpha$ for all $\alpha, \beta \in A_{\text{ass}}^1$ and $b \in A$. Observe that the obvious map $\Phi_{\tilde{A}}: \tilde{A} \rightarrow A_{\text{ass}}^1$ defined by $\Phi_{\tilde{A}}(b + \beta) = \Phi_A(b) + \beta$ is an A_{ass}^1 -bimodule map, and that $\mathbf{1} = \mathbf{1}_{A_{\text{ass}}^1}$ is a bar-unit. The bar-unital augmented dialgebra $(\tilde{A}, \Phi_{\tilde{A}}, A_{\text{ass}}^1)$ is easily seen to be balanced. There are nonbalanced bar-unital dialgebras as can be seen from the augmented bar-unital dialgebra example $(B \otimes B, \mathbf{1}_B \otimes \mathbf{1}_B, \mu_B, B)$ where $(B, \mathbf{1}_B, \mu_B)$ is any unital associative algebra and the bimodule action is defined by $b \cdot (b_1 \otimes b_2) \cdot b' := (bb_1) \otimes (b_2 b')$ for all $b, b', b_1, b_2 \in B$.

Again, in case the dialgebra $(A, \vdash, \dashv, \mathbf{1})$ is bar-unital and balanced, note that $A \dashv \mathbf{1} = \mathbf{1} \dashv A$ is an associative unital subalgebra A' of A whose multiplication is induced by both \vdash and \dashv , i.e. $a' \vdash b' = a' \dashv b'$ for all $a', b' \in A'$. Since the K -linear map $\pi_A: A \rightarrow A': a \mapsto a \dashv \mathbf{1} \dashv a$ descends to a surjective morphism of associative algebras $A_{\text{ass}}^1 \rightarrow A'$ by the above, it is clear that the ideal I contains the kernel of π_A . On the other hand, if $a \in \text{Ker}(\pi_A)$ then $0 = \pi(a) = \mathbf{1} \dashv a$, and obviously $a = \mathbf{1} \dashv a - \mathbf{1} \dashv a \in I$, thus inducing a useful isomorphism $A_{\text{ass}}^1 \cong A'$, and thus a subalgebra injection $i_A: A_{\text{ass}}^1 \rightarrow A: \Phi_A(a) \mapsto a \dashv \mathbf{1}$ which is a right inverse to the projection Φ_A , i.e. $\Phi_A \circ i_A = \text{id}_{A_{\text{ass}}^1}$.

In this work, we also have to take into account coalgebra structures and thus define the following:

Definition 2.4: Let $(A, \Delta, \varepsilon, \mathbf{1})$ be cocommutative C^3 -coalgebra (a C^4 -coalgebra) and two K -linear maps $\vdash, \dashv: A \otimes A \rightarrow A$. Then $(A, \Delta, \varepsilon, \mathbf{1}, \vdash, \dashv)$ will be called a cocommutative bar-unital di-coalgebra if and only if

1. $(A, \mathbf{1}, \vdash, \dashv)$ is a bar-unital balanced dialgebra.
2. Both \vdash and \dashv are morphisms of C^3 -coalgebras.

If in addition there is a morphism of C^3 -coalgebras $S: A \rightarrow A$ such that $(A, \Delta, \varepsilon, \mathbf{1}, \vdash, S)$ is a cocommutative right Hopf algebra and $(A, \Delta, \varepsilon, \mathbf{1}, \dashv, S)$ is a cocommutative left Hopf algebra, then $(A, \Delta, \varepsilon, \mathbf{1}, \vdash, \dashv, S)$ is called a cocommutative Hopf dialgebra.

We have used a relatively simple notion of one single compatible coalgebra structure motivated from differential geometry, see Section 3. In contrast to that, F. Goichot uses two a priori different coalgebra structures. Moreover, a slightly more general context would have been to demand the existence of two different antipodes, a right antipode S for \vdash , and a left antipode S' for \dashv . The theory –including the classification in terms of ordinary Hopf algebras– could have been done as well, but we have refrained from doing so since it is not hard to see that such a more general Hopf dialgebra is balanced iff $S=S'$. This fact is crucial in the following refinement of Proposition 2.4:

Proposition 2.5: Let $(A, \Delta, \varepsilon, \mathbf{1}, \vdash, \dashv, S)$ be cocommutative Hopf dialgebra. Then the submodule of all primitive elements of A , $\text{Prime}(A)$, is a Leibniz subalgebra of A equipped with the bracket (47).

Proof: Let $x, y \in A$ be primitive. Then, using that \vdash and \dashv are morphisms of coalgebras, we get

$$\Delta(x \vdash y - y \dashv x) = \mathbf{1} \otimes (x \vdash y - y \dashv x) + (x \vdash y - y \dashv x) \otimes \mathbf{1} \\ + (x \vdash \mathbf{1}) \otimes y + y \otimes (x \vdash \mathbf{1}) - y \otimes (\mathbf{1} \dashv x) - (\mathbf{1} \dashv x) \otimes y \\ = \mathbf{1} \otimes (x \vdash y - y \dashv x) + (x \vdash y - y \dashv x) \otimes \mathbf{1}$$

because A is balanced, and therefore $x \vdash y - y \dashv x$ is primitive.

The first relationship with rack bialgebras is the following simple generalization of a cocommutative Hopf algebra equipped with the adjoint representation:

Proposition 2.6: Let $(A, \Delta, \varepsilon, \mathbf{1}, \vdash, \dashv, S)$ be cocommutative Hopf dialgebra. Define the following multiplication

$$\mu: A \otimes A \rightarrow A \text{ by} \\ \mu(a \otimes b) := a \triangleright b := \sum_{(a)} (a^{(1)} \vdash b) \dashv (S(a^{(2)})). \quad (49)$$

Then we have the following:

1. The map \triangleright defines on the K -module A two left module structures, one with respect to the algebra $(A, \mathbf{1}, \vdash)$, and one with respect to the algebra $(A, \mathbf{1}, \dashv)$, making the Hopf-dialgebra $(A, \Delta, \varepsilon, \mathbf{1}, \vdash, \dashv, S)$ a module-Hopf dialgebra, i.e.

$$a \triangleright (b \triangleright c) = (a \vdash b) \triangleright c = (a \dashv b) \triangleright c \quad (50)$$

$$\Delta(a \triangleright b) = \sum_{(a), (b)} (a^{(1)} \triangleright b^{(1)}) \otimes (a^{(2)} \triangleright b^{(2)}) \quad (51)$$

$$a \triangleright (b \vdash c) = \sum_{(a)} (a^{(1)} \triangleright b) \vdash (a^{(2)} \triangleright c) \quad (52)$$

$$a \triangleright (b \dashv c) = \sum_{(a)} (a^{(1)} \triangleright b) \dashv (a^{(2)} \triangleright c) \quad (53)$$

2. $(A, \Delta, \varepsilon, \mathbf{1}, \mu)$ is a cocommutative rack bialgebra.

Proof: 1. First of all we have

$$\mu = \mu^{-\circ} (\mu \vdash \otimes S) \circ (\text{id}_A \otimes \tau_{A,A}) \circ (\Delta \otimes \text{id}_A)$$

where $\mu \vdash$ and $\mu \dashv$ stand for the multiplication maps \vdash and \dashv , and this is clearly a composition of morphisms of C^3 -coalgebras whence μ is a morphism of C^3 -coalgebras proving eqn (51). Next, there is clearly $\mathbf{1} \triangleright b = b$ for all $b \in A$, and, since the dialgebra is balanced, we get for all $a \in A$

$$a \triangleright \mathbf{1} = \sum_{(a)} (a^{(1)} \vdash \mathbf{1}) \dashv (S(a^{(2)})) = \sum_{(a)} \mathbf{1} \dashv a^{(1)} \dashv (S(a^{(2)})) \\ = \sum_{(a)} \mathbf{1} \dashv (a^{(1)} \vdash (S(a^{(2)}))) = \varepsilon(a) \mathbf{1} \dashv \mathbf{1} = \varepsilon(a) \mathbf{1}.$$

Next, let $a, b, c \in A$. Then

$$\begin{aligned}
 a \triangleright (b \triangleright c) &= \sum_{(a),(b)} (a^{(1)} \vdash ((b^{(1)} \vdash c) \dashv S(b^{(2)}))) \dashv S(a^{(2)}) \\
 &= \sum_{(a),(b)} (((a^{(1)} \vdash b^{(1)}) \vdash c) \dashv (S(b^{(2)}) \dashv S(a^{(2)}))) \\
 &= \sum_{(a),(b)} (((a^{(1)} \vdash b^{(1)}) \vdash c) \dashv (S(b^{(2)}) \dashv S(a^{(2)}))) \\
 &= \sum_{(a),(b)} (((a^{(1)} \dashv b^{(1)}) \vdash c) \dashv (S(a^{(2)} \dashv b^{(2)}))) \\
 &= (a \dashv b) \triangleright c, \tag{54}
 \end{aligned}$$

Proving eqs (50). Next, for all $a, a', a'' \in A$, we get

$$\begin{aligned}
 \sum_{(a)} (a^{(1)} \triangleright a') \dashv (a^{(2)} \triangleright a'') &= \sum_{(a)} ((a^{(1)} \vdash a') \dashv S(a^{(2)})) \dashv ((a^{(3)} \vdash a'') \dashv S(a^{(4)})) \\
 &= \sum_{(a)} (a^{(1)} \vdash a') \dashv ((S(a^{(2)}) \dashv a^{(3)}) \dashv (a'' \dashv S(a^{(4)}))) \\
 &= \sum_{(a)} (a^{(1)} \vdash a') \dashv ((\varepsilon(a^{(2)}) \mathbf{1}) \dashv (a'' \dashv S(a^{(3)}))) \\
 &= \sum_{(a)} (a^{(1)} \vdash (a' \dashv a'')) \dashv S(a^{(2)}) = a \triangleright (a' \dashv a''),
 \end{aligned}$$

where –in the second to last equation– we have used the left antipode identity for the case \dashv and the fact that $(a)S(a^{(1)}) a^{(2)}$ is a generalized left unit element for the case \vdash . It follows that (A, \dashv) is an A -module-algebra proving eqs (52) and (53).

2. It remains to prove self-distributivity: For all $a, b, c \in A$, we get

$$\sum_{(a)} (a^{(1)} \triangleright b) \triangleright (a^{(2)} \triangleright c) = \sum_{(a)} ((a^{(1)} \triangleright b) \dashv a^{(2)}) \triangleright c,$$

and in the end

$$\begin{aligned}
 \sum_{(a)} (a^{(1)} \triangleright b) \dashv a^{(2)} &= \sum_{(a)} ((a^{(1)} \vdash b) \dashv S(a^{(2)})) \dashv a^{(3)} \\
 &= \sum_{(a)} (a^{(1)} \vdash b) \dashv (S(a^{(2)}) \dashv a^{(3)}) \\
 &= \sum_{(a)} (a^{(1)} \vdash b) \dashv (\varepsilon(a^{(2)}) \mathbf{1}) = a \dashv b
 \end{aligned}$$

proving the self-distributivity identity.

The next theorem relates augmented cocommutative rack bialgebras with cocommutative Hopf dialgebras:

Theorem 2.6: Let (B, Φ_B, H, ℓ) be a cocommutative augmented rack bialgebra. Then the K -module $(B \otimes H, \Delta_{B \otimes H}, \varepsilon_B \otimes \varepsilon_H, \mathbf{1}_B \otimes \mathbf{1}_H, \Phi, H)$ will be an augmented cocommutative Hopf dialgebra by means of the following definitions. Here we use Example 2.5 and take $h, h' \in H$ and $b \in B$:

1. $\Phi: B \otimes H \rightarrow H: (b \otimes h) \mapsto \Phi(b \otimes h) := \Phi B(b)h$.
2. $h' \cdot (b \otimes h) := \Sigma(h')((h')^{(1)} \cdot b) \otimes ((h')^{(2)} h)$ and $(b \otimes h) \cdot h' := b \otimes (hh')$.
3. $S(b \otimes h) := \mathbf{1}_B \otimes S_H(\Phi B(b)h)$.

Moreover, the Leibniz bracket on the K -module of all primitive elements of $B \otimes H$, $a := \text{Prim}(B) \otimes \text{Prim}(H)$, is computed as follows for all $x, y \in \text{Prim}(B)$ and all ξ, η in the Lie algebra $\text{Prim}(H)$ (writing x and ξ for the more precise $x \otimes \mathbf{1}_H$ and $\mathbf{1}_B \otimes \xi$)

$$[x + \xi, y + \eta] = ([x, y] + \cdot y) + ([\Phi B(x), \eta] + [\xi, \eta]) \tag{55}$$

where each bracket is of the form (47)¹. Note that this Leibniz algebra is split over the Lie subalgebra $\text{Prim}(H)$, the complementary two-sided ideal $\{x - \Phi B(x) \mid x \in \text{Prim}(B)\}$ being in the left center of \mathfrak{a} .

¹ For an explicit formula, see the end of the proof of the theorem.

Proof: It is clear from the definitions that condition 2.6 defines a H -bimodule structure on $C \otimes H$ making it into a module C^3 -coalgebra. Moreover, we compute for all $h, h', h'' \in H$ and $b \in B$

$$\begin{aligned}
 \Phi(h' \cdot (b \otimes h) \cdot h'') &= \sum_{(h')} \Phi(((h')^{(1)} \cdot b) \otimes ((h')^{(2)} h h'')) = \sum_{(h')} \Phi_B((h')^{(1)} \cdot b)(h')^{(2)} h h'' \\
 &= \sum_{(h')} \text{ad}_{(h')^{(1)}}(\Phi_B(b))(h')^{(2)} h h'' = \sum_{(h')} (h')^{(1)} \Phi_B(b) S_H((h')^{(2)})(h')^{(3)} h h'' \\
 &= \sum_{(h')} (h')^{(1)} \Phi_B(b) \varepsilon_H((h')^{(2)}) h h'' = h' \Phi_B(b) h h'' = h' \Phi(b \otimes h) h'',
 \end{aligned}$$

whence Φ is a morphism of H -bimodules. Next, we get for all $b \in B$ and $h \in H$:

$$\begin{aligned}
 (\text{id}_{B \otimes H} * \dashv S)(b \otimes h) &= \sum_{(b)(h)} (b^{(1)} \otimes h^{(1)}) \dashv S(b^{(2)} \otimes h^{(2)}) \\
 &= \sum_{(b)(h)} \Phi(b^{(1)} \otimes h^{(1)}) \cdot (\mathbf{1}_B \otimes S_H(\Phi_B(b^{(2)} h^{(2)}))) \\
 &= \sum_{(b)(h)} (\Phi_B(b^{(1)}) h^{(1)}) \cdot (\mathbf{1}_B \otimes S_H(\Phi_B(b^{(2)} h^{(2)}))) \\
 &= \sum_{(b)(h)} \varepsilon_H(\Phi_B(b^{(1)}) h^{(1)}) \mathbf{1}_B \otimes (\Phi_B(b^{(2)} h^{(2)}) S_H(\Phi_B(b^{(3)} h^{(3)}))) \\
 &= \mathbf{1}_B \otimes (\varepsilon_B(b) \varepsilon_H(h)) \mathbf{1}_H = (\varepsilon_B \otimes \varepsilon_H)(b \otimes h) (\mathbf{1}_B \otimes \mathbf{1}_H),
 \end{aligned}$$

proving the right antipode identity, and

$$\begin{aligned}
 (S * \dashv \text{id}_{B \otimes H})(b \otimes h) &= \sum_{(b)(h)} S(b^{(1)} \otimes h^{(1)}) \dashv (b^{(2)} \otimes h^{(2)}) \\
 &= \sum_{(b)(h)} (\mathbf{1}_B \otimes S_H(\Phi_B(b^{(1)}) h^{(1)})) \cdot (\Phi_B(b^{(2)} h^{(2)})) \\
 &= \sum_{(b)(h)} (\mathbf{1}_B \otimes S_H(\Phi_B(b^{(1)}) h^{(1)})) \cdot (\Phi_B(b^{(2)} h^{(2)})) \\
 &= \sum_{(b)(h)} \mathbf{1}_B \otimes (S_H(\Phi_B(b^{(1)}) h^{(1)}) \Phi_B(b^{(2)} h^{(2)})) \\
 &= \mathbf{1}_B \otimes (\varepsilon_B(b) \varepsilon_H(h)) \mathbf{1}_H = (\varepsilon_B \otimes \varepsilon_H)(b \otimes h) (\mathbf{1}_B \otimes \mathbf{1}_H),
 \end{aligned}$$

proving the left antipode identity. Finally for all $h \in H$ we get

$$h \cdot (\mathbf{1}_B \otimes \mathbf{1}_H) = \sum_{(h)} (\varepsilon_H(h^{(1)}) \mathbf{1}_B) \otimes h^{(2)} = \mathbf{1}_B \otimes h = (\mathbf{1}_B \otimes \mathbf{1}_H) \cdot h$$

implying that the bar-unital dialgebra is balanced.

Formula (55) is straight-forward:

$$\begin{aligned}
 &[x \otimes \mathbf{1}_H + \mathbf{1}_B \otimes \xi, y \otimes \mathbf{1}_H + \mathbf{1}_B \otimes \eta] \\
 &= (x \otimes \mathbf{1}_H) \vdash (y \otimes \mathbf{1}_H) - (y \otimes \mathbf{1}_H) \vdash (x \otimes \mathbf{1}_H) \\
 &+ (x \otimes \mathbf{1}_H) \vdash (\mathbf{1}_B \otimes \eta) - (\mathbf{1}_B \otimes \eta) \vdash (x \otimes \mathbf{1}_H) \\
 &+ (\mathbf{1}_B \otimes \xi) \vdash (y \otimes \mathbf{1}_H) - (y \otimes \mathbf{1}_H) \vdash (\mathbf{1}_B \otimes \xi) \\
 &+ (\mathbf{1}_B \otimes \xi) \vdash (\mathbf{1}_B \otimes \eta) - (\mathbf{1}_B \otimes \eta) \vdash (\mathbf{1}_B \otimes \xi) \\
 &= (\Phi_B(x) \cdot y) \otimes \mathbf{1}_H + y \otimes \Phi_B(x) - y \otimes \Phi_B(x) \\
 &+ \varepsilon_B(x) \mathbf{1}_B \otimes \eta + \mathbf{1}_B \otimes (\Phi_B(x) \eta) - \mathbf{1}_B \otimes (\eta \Phi_B(x)) \\
 &+ (\xi \cdot y) \otimes \mathbf{1}_H + y \otimes \xi - y \otimes \xi \\
 &= \varepsilon_H(\xi) \mathbf{1}_B \otimes \eta + \mathbf{1}_B \otimes (\xi \eta) - \mathbf{1}_B \otimes (\eta \xi) \\
 &= [x, y] \otimes \mathbf{1}_H + \mathbf{1}_B \otimes [\Phi_B(x), \eta] + (\xi \cdot y) \otimes \mathbf{1}_H + \mathbf{1}_B \otimes [\xi, \eta],
 \end{aligned}$$

because primitives are killed by counits, and the formula is proved.

A lengthy, but straight-forward reasoning shows that the above construction assigning $(B, \Phi_B, H, \ell) \rightarrow (B \otimes H, \Phi, H)$ defines a covariant functor from the category of all cocommutative rack bialgebras to the category of all cocommutative Hopf dialgebras.

A particular case of the preceding theorem is obtained by picking any C^4 -coalgebra $(B, \Delta_B, \varepsilon_B, 1_B)$ such that there is any H -module coalgebra structure l on B (such that $h.1_B = \varepsilon_H(h)1_B$) and by choosing the trivial map $\Phi_B(b) = \varepsilon_B(b)1_H$. It follows that (B, Φ_B, H, l) is an augmented rack bialgebra with left-trivial multiplication. It turns out that the Hopf dialgebra $B \otimes H$ formed out of this is already isomorphic to the general cocommutative Hopf dialgebra:

Theorem 2.7: Let $(A, \Delta, \varepsilon, 1, \dashv, \dashv, S)$ be a cocommutative Hopf dialgebra. Let E_A be the C^3 -subcoalgebra of all generalized idempotent elements² with respect to \dashv , and let $H_A = 1 \dashv A$ be the Hopf subalgebra according to the Suschkewitsch decomposition of the left Hopf algebra $(A, \Delta, \varepsilon, 1, \dashv, \dashv, S)$, see Theorem 2.5. Then we have:

1. By means of the Suschkewitsch isomorphism $A \rightarrow E_A \otimes H_A$ for the left Hopf algebra $(A, \Delta, \varepsilon, 1, \dashv, \dashv, S)$, we can transfer the cocommutative dialgebra structure of A to $E_A \otimes H_A$: There is a well-defined left module-coalgebra action l of H_A on E_A defined by (for all $c \in E_{H_A}, h \in H_A, a \in A$ such that $h = 1 \dashv a$)

$$(c \otimes h) \dashv' (c' \otimes h') = \varepsilon(c) \sum_{(h)} (h^{(1)} \cdot c') \otimes (h^{(2)} h'), \tag{56}$$

and the transferred multiplications \dashv' and \dashv'' and the antipode S' on $E_A \otimes H_A$ read (for all $c, c' \in E_{H_A}, h, h' \in H_A$)

$$(c \otimes h) \dashv' (c' \otimes h') = \varepsilon(c) \sum_{(h)} (h^{(1)} \cdot c') \otimes (h^{(2)} h'), \tag{57}$$

$$(c \otimes h) \dashv'' (c' \otimes h') = \varepsilon(c') c(hh'), \tag{58}$$

$$S'(c \otimes h) = \varepsilon(c) 1_{E_A} \otimes (S_{H_A}(h)). \tag{59}$$

2. The K -linear map $A \rightarrow 1 \dashv A : a \mapsto 1 \dashv a$ descends to an isomorphism of associative algebras $A_{\text{ass}} \cong H_A$.

3. S.Covez, 2006: The Leibniz subalgebra $\text{Prim}(A)$ of A (equipped with the Leibniz bracket (47)) is a split semidirect sum out of the two-sided ideal $\text{Prim}(E_A) \subset z(\text{Prim}(A))$ and the Lie subalgebra $\text{Prim}(H_A)$, i.e. for all $z, z' \in \text{Prim}(E_A)$ and $\xi, \xi' \in \text{Prim}(H_A)$, we have

$$[z + \xi, z' + \xi'] = \xi.z + [\xi, \xi']. \tag{60}$$

4. Let (B, Φ_B, H, l) be a cocommutative augmented rack bialgebra. Then for the Hopf dialgebra $B \otimes H$ of Theorem 2.6, we get that the Hopf subalgebra $H_B \otimes_{H_A} H$ equals $1_B \otimes H \cong H$, and

$$E_{B \otimes H} = \left\{ \sum_{(b)} b^{(1)} \otimes (S_H(\Phi_B(b^{(2)}))) \in B \otimes H \mid b \in B \right\}$$

which is isomorphic to B as a C^4 -coalgebra of BH .

Proof: 1. Note first that the right hand side of eqn (56) is just $a \triangleright c$ of Proposition 2.6 which had been shown to be a left module-Hopf dialgebra action of (A, \dashv) and of (A, \dashv) on A . Observe that for all $a, a' \in A$

$$(1 \dashv a) \triangleright a' = 1 \triangleright (a \triangleright a') = a \triangleright a'$$

whence the H_A -action l is well-defined on A . Moreover we compute for all $h \in H_A$ and all $a, a', a'' \in A$ such that

$$\begin{aligned} h = 1 \dashv a: \\ \sum_{(h)} (h^{(1)} \cdot a') \dashv (h^{(2)} \cdot a'') &= \sum_{(a)} (a^{(1)} \triangleright a') \dashv (a^{(2)} \triangleright a'') \\ &\stackrel{(53)}{=} a \triangleright (a' \dashv a'') = h.(a' \dashv a'') \end{aligned}$$

whence (A, \dashv) is also a H_A -module-algebra. Now let $c \in E_{H_A}$. By definition, c is a generalized idempotent (w.r.t. \dashv), hence $c = \sum_{(c)} c^{(1)} \dashv c^{(2)}$, and thus for all $h \in H$

$$h.c = h. \sum_{(c)} c^{(1)} \dashv c^{(2)} = \sum_{(h),(c)} (h^{(1)} \cdot c^{(1)}) \dashv (h^{(2)} \cdot c^{(2)}) = \sum_{(h,c)} (h.c)^{(1)} \dashv (h.c)^{(2)}$$

Whence $h.c$ is also in E_A , and E_A is a H_A -submodule of A .

Recall the Suschkewitsch decomposition of the left Hopf algebra $(A, \Delta, \varepsilon, 1, \dashv, \dashv, S)$ where one can use Theorem 2.5 and dualize all the formulas:

$$\Psi : A \rightarrow E_A \otimes H_A : a \mapsto \sum_{(a)} (a^{(1)} \dashv S(a^{(2)})) \otimes (1 \dashv a^{(3)}).$$

$$\Psi^{-1} : E_A \otimes H_A \rightarrow A : (c \otimes (1 \dashv a)) \mapsto c \dashv a.$$

Formulas (58) and (59) consequences of Theorem 2.5. The only formula which remains to be shown is eqn (57). Note first that every generalized idempotent $c \in E_{H_A}$ (w.r.t. \dashv) is also a generalized idempotent with respect to \dashv . Indeed, since all the components $c^{(1)}$ and $c^{(2)}$ in $\Delta(c) = \sum_{(c)} c^{(1)} \otimes c^{(2)}$ can be chosen in E_{H_A} , we get

$$\begin{aligned} \sum_{(c)} c^{(1)} \dashv c^{(2)} &= \sum_{(c)} (c^{(1)} \dashv S(c^{(2)})) \dashv c^{(3)} = \sum_{(c)} (c^{(1)} \dashv S(c^{(2)})) \dashv c^{(3)} \\ &= \sum_{(c)} \varepsilon(c^{(1)}) 1 \dashv c^{(2)} = c. \end{aligned}$$

Next for all $c, c' \in E_{H_A}, h, h' \in H_A$ and $a, a' \in A$ such that $1 \dashv a = h$ and $1 \dashv a' = h'$, we get –since c is a generalized left unit element (w.r.t. \dashv) thanks to 3. in Lemma 2.2–

$$\begin{aligned} (c \otimes h) \dashv' (c' \otimes h') &= \Psi(\Psi^{-1}(c \otimes h)) \dashv \Psi^{-1}(c' \otimes h') = \Psi((c \dashv a) \dashv (c' \dashv a')) \\ &= \Psi((c \dashv a) \dashv (c' \dashv a')) = \varepsilon(c) \Psi(a \dashv (c' \dashv a')) \end{aligned}$$

and this is equal to

$$\begin{aligned} \varepsilon(c) \sum_{(a),(c'),(a')} ((a^{(1)} \dashv ((c')^{(1)} \dashv (a')^{(1)})) \dashv (S(a^{(2)} \dashv ((c')^{(2)} \dashv (a')^{(2)}))) \\ \otimes (1 \dashv (a^{(3)} \dashv ((c')^{(3)} \dashv (a')^{(3)}))) \\ = \varepsilon(c) \sum_{(a),(c'),(a')} (a^{(1)} \dashv ((c')^{(1)} \dashv ((a')^{(1)} \dashv S((a')^{(2)})) \dashv S((c')^{(2)} \dashv S(a^{(2)}))) \\ \otimes (1 \dashv ((a^{(3)} \dashv (c')^{(3)} \dashv (a')^{(3)}))) \\ = \varepsilon(c) \sum_{(a),(c')} (a^{(1)} \dashv (((c')^{(1)} \dashv S((c')^{(2)})) \dashv S(a^{(2)}))) \\ \otimes (1 \dashv (a^{(3)} \dashv a')) \\ = \varepsilon(c) \sum_{(a)} (a^{(1)} \dashv (c' \dashv S(a^{(2)}))) \otimes (1 \dashv (a^{(3)} \dashv a')) \\ = \varepsilon(c) \sum_{(h)} (h^{(1)} \cdot c') \otimes (h^{(2)} h') \end{aligned}$$

proving eqn (57).

2. Clear for any bar-unital balanced dialgebra.

3. Straight-forward computation using $\text{Prim}(\mathcal{H}) = \text{Prim}(E_{\mathcal{H}}) \otimes \text{Prim}(H_{\mathcal{H}})$ where the latter is well-known to be a Lie algebra and the former is abelian.

4. For each $b \in b$ and $h \in H$, we get

$$(1_B \otimes 1_H) \dashv (b \otimes h) = (1_B \otimes (\Phi_B(b)h)),$$

Proving the first statement. Moreover

$$\begin{aligned} \sum_{(b),(h)} (b^{(1)} \otimes h^{(1)}) \dashv (S(b^{(2)} \otimes h^{(2)})) &= \sum_{(b),(h)} b^{(1)} \otimes h^{(1)} S_H(h^{(2)}) S_H(\Phi_B(b^{(2)})) \\ &= \sum_{(b)} b^{(1)} \otimes S_H(\Phi_B(b^{(2)})), \end{aligned}$$

Proving the form of the generalized idempotents, and since the

²Recall that this means $c = (-^{\circ} \Delta)(c)$.

K -linear map $B \rightarrow B \otimes H$ given by $(\text{id} \otimes B \otimes (S_H \circ \Phi_B)) \circ \Delta$ is an injective morphism of C^3 -coalgebra, the statement is proved.

The third statement had been proved by Simon Covez in his Master thesis in the differential geometric context of digroups, compare with Section 3.

Example 2.6: As an example, let us compute the Suschkewitsch decomposition for the augmented rack bialgebra $K[X]$ where $p: X \rightarrow G$ is an augmented pointed rack, see Example 2.4. By the above theorem, part 4., its associated cocommutative augmented Hopf dialgebra decomposes as $B \otimes H$, where the Hopf algebra $H = K[G]$ is the standard group algebra and $B = K[X]$. The generalized idempotents are in this case

$$E_B \otimes_{H} = \{b \otimes p(b)^{-1} | b \in B\}.$$

We finish this section with a formula relating universal algebras: The functor associating to any dialgebra A its Leibniz algebra A^- via eqn (47) is well-known to have a left adjoint associating to any Leibniz algebra $(\mathfrak{h}, [, \cdot])$ its (in general non bar-unital) universal enveloping dialgebra $\text{Ud}(\mathfrak{h})$ associated to \mathfrak{h} defined by

$$\text{Ud}(\mathfrak{h}) = \mathfrak{h} \otimes \text{U}(\bar{\mathfrak{h}}). \quad (61)$$

But also in the category of bar-unital balanced dialgebras, there is such a left adjoint: To any Leibniz algebra $(\mathfrak{h}, [, \cdot])$, we associate its universal balanced bar-unital enveloping dialgebra $\widetilde{\text{Ud}}(\mathfrak{h})$

$$\widetilde{\text{Ud}}(\mathfrak{h}) = \text{UAR}(\mathfrak{h}) \otimes \text{U}(\bar{\mathfrak{h}}). \quad (62)$$

Before proving the theorem, we note that $\widetilde{\text{Ud}}(\mathfrak{h}) = \text{Ud}(\mathfrak{h}) \oplus \text{U}(\bar{\mathfrak{h}})$ is obtained by adjoining a balanced bar-unit to $\text{Ud}(\mathfrak{h})$.

Theorem 2.8: For any Leibniz algebra $(\mathfrak{h}, [, \cdot])$, the assignment $\mathfrak{h} \rightarrow \widetilde{\text{Ud}}(\mathfrak{h})$ defines a left adjoint functor to the functor associating to any bar-unital balanced dialgebra its commutator Leibniz algebra.

Proof: Clearly, $\widetilde{\text{Ud}}(\mathfrak{h})$ is the cocommutative Hopf dialgebra associated to the universal augmented rack bialgebra $(\text{UAR}(\mathfrak{h}), \Phi_{\mathfrak{h}}, \text{U}(\bar{\mathfrak{h}}), \ell)$ (cf. Theorem 2.6) which in turn is associated to the Leibniz algebra $(\mathfrak{h}, [, \cdot])$ (cf. Theorem 2.4). Since both assignments are functorial, it follows that the assignment $\mathfrak{h} \rightarrow \widetilde{\text{Ud}}(\mathfrak{h})$ is a functor. It remains to prove the universal property: Let $(\mathfrak{h}, [, \cdot])$ be a Leibniz algebra, let $(A, 1, \vdash, \dashv)$ a bar-unital balanced dialgebra, and let $\phi: \mathfrak{h} \rightarrow A^-$ be a morphism of Leibniz algebras. It follows that the K -linear map $\Phi_A \circ \phi: \mathfrak{h} \rightarrow A_{\text{ass}}^1$ vanishes on the two-sided ideal $Q(\mathfrak{h})$ and descends to a morphism $\bar{\phi}$ of the quotient Lie algebra $\bar{\mathfrak{h}}$ to A_{ass}^1 with its commutator Lie bracket such that $\bar{\phi} \circ p = \Phi_A \circ \phi$. Hence there is a unique morphism $\text{U}(\bar{\phi}): \text{U}(\bar{\mathfrak{h}}) \rightarrow A_{\text{ass}}^1$ of unital associative algebras extending $\bar{\phi}$. Define the K -linear map $\hat{\phi}: \widetilde{\text{Ud}}(\mathfrak{h}) = \text{U}(\bar{\mathfrak{h}}) \oplus \text{Ud}(\mathfrak{h}) \rightarrow A$ by (for all $u, v \in \text{U}(\bar{\mathfrak{h}})$ and $x \in \mathfrak{h}$):

$$\hat{\phi}(u) = i_A(\text{U}(\bar{\phi})(u)) \text{ and } \hat{\phi}(x \otimes v) = \phi(x) + i_A(\text{U}(\bar{\phi})(v))$$

where we recall the natural injection of unital algebras $i_A: A_{\text{ass}}^1 \rightarrow A$ given by $i_A(\Phi_A(a)) = 1 \dashv a$ for all $a \in A$. We shall show that $\hat{\phi}$ is a morphism of augmented dialgebras: We compute for all $u, u', u'' \in \text{U}(\bar{\mathfrak{h}})$, using that i_A and $\text{U}(\bar{\phi})$ are morphisms of unital associative algebras and that in the image of i_A , we can use the multiplication symbols \vdash and \dashv arbitrarily:

$$\hat{\phi}(u'uu'') = i_A(\text{U}(\bar{\phi})(u')) \vdash i_A(\text{U}(\bar{\phi})(u)) + i_A(\text{U}(\bar{\phi})(u'')) = \hat{\phi}(u') \vdash \hat{\phi}(u) \dashv \hat{\phi}(u'')$$

Showing the fact that $\hat{\phi}$ preserves the bimodule structures on the first component of $\widetilde{\text{Ud}}(\mathfrak{h})$. Next we have for all $x_1, x \in \mathfrak{h}$

$$\hat{\phi}(p(x_1).x) = \hat{\phi}(x_1, x) = \phi(x_1) \vdash \phi(x) - \phi(x) - \phi(x) \phi(x_1)$$

$$= i_A(\bar{\phi}(p(x_1))) \vdash \phi(x) - \phi(x) + i_A(\bar{\phi}(p(x_1)))$$

and by induction on k in $u' = p(x_1) \cdots p(x_k) \in \text{U}(\bar{\mathfrak{h}})$ and $x_1, \dots, x_k \in \mathfrak{h}$, we prove

$$\hat{\phi}(u'.x) = \sum_{(u')} i_A(\bar{\phi}(u'^{(1)})) \vdash \phi(x) + (\bar{\phi}(S(u'^{(2)}))).$$

Now, for all $u', u'', v \in \text{U}(\bar{\mathfrak{h}})$ and $x \in \mathfrak{h}$, we get:

$$\begin{aligned} &= \sum_{(u')} \hat{\phi}((u')^{(1)}.x) + i_A(\text{U}(\bar{\phi})((u')^{(2)}vu'')) \\ &= \sum_{(u')} \hat{\phi}((u')^{(1)}.x) + i_A(\text{U}(\bar{\phi})((u')^{(2)}vu'')) \\ &= \sum_{(u')} i_A(\text{U}(\bar{\phi})((u')^{(1)})) \vdash \phi(x) + i_A(\text{U}(\bar{\phi})(S((u')^{(2)})(u')^{(3)}vu'')) \\ &= i_A(\text{U}(\bar{\phi})(u')) \vdash \phi(x) + i_A(\text{U}(\bar{\phi})(v)) + i_A(\text{U}(\bar{\phi})(u'')) \\ &= i_A(\text{U}(\bar{\phi})(u')) \vdash \hat{\phi}(x \otimes v) + i_A(\text{U}(\bar{\phi})(u'')) \end{aligned}$$

Showing the fact that $\hat{\phi}$ preserves the bimodule structures on the second component of $\widetilde{\text{Ud}}(\mathfrak{h})$. Hence $\hat{\phi}$ is a morphism of bar-unital (augmented) dialgebras. The uniqueness of $\hat{\phi}$ follows from the universal property of $\text{U}(\bar{\mathfrak{h}})$

Coalgebra Structures for Pointed Manifolds with Multiplication

In this section, the symbol \mathbb{K} denotes either the field of all real numbers, \mathbb{R} , or the field of all complex numbers, \mathbb{C} . We define here the monoidal category of pointed manifolds, and exhibit the Serre functor sending a pointed manifold to the coalgebra of point-distributions supported in the distinguished point. We recall further that this is a strong monoidal functor. Further down, we will study Lie (semi) groups, Lie racks, and Lie digroups as examples of this construction, motivating geometrically the notions of a rack bialgebra and of a Hopf dialgebra.

Pointed manifolds with multiplication(s)

Recall first the category of all pointed manifolds $\mathcal{M}f^*$ whose objects consist of pairs (M, e) where M is a non-empty differentiable manifold and e is an element of M and whose morphisms $(M, e) \rightarrow (M', e')$ are given by all smooth maps $\phi: M \rightarrow M'$ of the underlying manifolds such that $\phi(e) = e'$. Recall that the cartesian product \times makes $\mathcal{M}f^*$ into a monoidal category by setting $(M, e_1) \times (N, e_2) := (M \times N, (e_1, e_2))$ with the one-point set $(\{pt\}, pt)$ as unit object and the usual associators, left-unit and right-unit identifications borrowed from the category of sets [14]. This monoidal category is symmetric by means of the usual (tensor) flip map $\tau_T = \tau_{T(M,N)}: M \times N \rightarrow N \times M: (x, y) \mapsto (y, x)$ where the pair of distinguished points is also interchanged.

By simply forgetting about the differentiable structure we get the category of pointed sets.

Recall that a pointed manifold with multiplication is a triple (M, e, m) where (M, e) is a pointed manifold, and $m: (M, e) \times (M, e) \rightarrow (M, e)$ is a smooth map of pointed manifolds, i.e. is a smooth map $M \times M \rightarrow M$ such that $m(e, e) = e$. Moreover, a pointed manifold with multiplication will be called left-regular (resp. right-regular) if all the left (resp. right) multiplication maps $y \mapsto m(x, y)$ (resp. $y \mapsto m(y, x)$) are diffeomorphisms. Morphisms of pointed manifolds with multiplication $(M, e, m) \rightarrow (M', e', m')$ are smooth maps of pointed manifolds $\phi: (M, e) \rightarrow (M', e')$ such that

$$\phi \circ m = m' \circ (\phi \times \phi). \quad (63)$$

The obvious generalization are a finite number of maps $M^n \rightarrow M$ with $n \geq 1$ arguments.

Again by forgetting about differentiable structures, we get the category of pointed sets with multiplication.

Coalgebra Structure for distributions supported in one point

For any pointed manifold (M, e) recall the \mathbb{K} -vector space

$$\begin{aligned} \mathcal{E}'_e(M) &:= \\ \{T: C^\infty(M, \mathbb{K}) \rightarrow \mathbb{K} \mid T \text{ is a continuous linear map and } \text{supp}(T) = \{e\}\} \end{aligned} \quad (64)$$

of all distributions supported in the singleton $\{e\}$ [15]. Now let $\phi: M \rightarrow M'$ be a smooth map such that $\phi: (e) \rightarrow e'$. For any distribution $S \in \mathcal{E}'_e(M)$ and any smooth function $f \in C^\infty(M', \mathbb{K})$ the well-known prescription

$$(\phi_* S)(f) := S(f \circ \phi) \quad (65)$$

gives a well-defined distribution $\phi_* S$ on the target manifold M' supported in $e' = \phi(e)$, and the map $\phi^*: \mathcal{E}'_{e'}(M') \rightarrow \mathcal{E}'_e(M)$ is a \mathbb{K} -linear map (which is continuous). Clearly for three pointed manifolds (M, e) , (M', e') , and (M'', e'') with smooth maps $\phi: (M, e) \rightarrow (M', e')$ and $\psi: (M', e') \rightarrow (M'', e'')$ we get

$$(\psi \circ \phi)_* = \psi_* \circ \phi_* \quad \text{and} \quad (\text{id}_M)_* = \text{id}_{\mathcal{E}'_e(M)}. \quad (66)$$

This defines a covariant functor $F: \mathcal{M}^{\text{pt}} \rightarrow \mathbb{K}\text{Vect}$ to the category of all \mathbb{K} -vector spaces by associating to any pointed manifold (M, e) the \mathbb{K} -vector space $F(M, e) := \mathcal{E}'_e(M)$, and to any smooth map $(M, e) \rightarrow (M', e')$ the linear map $F(\phi) := \phi^*: \mathcal{E}'_{e'}(M') \rightarrow \mathcal{E}'_e(M)$. We call this functor Serre functor in tribute to the predominant role it plays in [16]. It is one of the main objects of this article.

There is, however, much more structure in this functor: First any distribution space $\mathcal{E}'_e(M)$ contains a canonical linear form $\varepsilon = \varepsilon_e: \mathcal{E}'_e(M) \rightarrow \mathbb{K}$ defined by

$$\varepsilon_e(T) := T(1), \quad (67)$$

Where 1 denotes the constant function $M \rightarrow \mathbb{K}$ whose only value is equal to $1 \in \mathbb{K}$. Moreover each space $\mathcal{E}'_e(M)$ contains a canonical element $1_e = 1_e$ defined by the well-known delta distribution

$$1_e = \delta_e: f \mapsto f(e), \quad (68)$$

and we clearly have

$$\varepsilon_e(1_e) = 1. \quad (69)$$

Moreover, both ε_e and 1_e are natural in the following sense: let $\phi: (M, e) \rightarrow (M', e')$ be a smooth map of pointed differentiable manifolds. Then it is straight-forward to check that

$$\varepsilon_e \circ \phi_* = \varepsilon_{e'} \quad \text{and} \quad \phi^*(1_{e'}) = 1_e. \quad (70)$$

Recall the well-known tensor product or direct product of two distributions: More generally, let M and N be two differentiable manifolds, and let $S \in \mathcal{D}'(M)$ and $T \in \mathcal{D}'(N)$ be two distributions (where the symbol $\mathcal{D}'(M)$ denotes the continuous dual space of the test function space $\mathcal{D}(M)$ of all smooth \mathbb{K} -valued functions with compact support) [17]. Let $f: M \times N \rightarrow \mathbb{K}$ be a smooth function with compact support $K \subset M \times N$, and let $K_1 := \text{pr}_M(K) \subset M$, $K_2 := \text{pr}_N(K) \subset N$, whence K is a subset of the compact set $K_1 \times K_2$. Let $T^{(2)}: \mathcal{D}(M \times N) \rightarrow \text{Fun}(M, \mathbb{K})$ be the following map: For each $x \in M$, let $f_x \in \mathcal{D}(N)$ be the partial function $y \mapsto f(x, y)$. Then we set

$$(T^{(2)}(f))(x) = T(f_x).$$

The superscript (2) means here that we see f as a function of its second variable only, when applying the distribution T .

Upon using the approximation theorem of any distribution by a sequence of regular distributions, one can show that $(T^{(2)}(f)): M \rightarrow \mathbb{K}$ is a smooth function having compact support in K_1 . It is clear that $T^{(2)}: \mathcal{D}(M \times N) \rightarrow \mathcal{D}(M)$ is linear, and it can be shown by the same approximation theorem that $T^{(2)}$ is continuous. It follows that the map

$$(S, T) \mapsto (f \mapsto S(T^{(2)}(f)))$$

is a well-defined \mathbb{K} -bilinear map $\mathcal{D}'(M)\mathcal{D}'(N) \rightarrow \mathcal{D}'(M \times N)$, and there is thus a unique linear map (where \otimes denotes the usual algebraic tensor product over \mathbb{K})

$$F_{2M, 2N}: \mathcal{D}(M) \otimes \mathcal{D}(N) \rightarrow \mathcal{D}(M \times N) \quad (71)$$

such that for all $f \in \mathcal{D}(M \times N)$ we have

$$(F_{2M, 2N}(S \otimes T))(f) = S(T^{(2)}(f)).$$

Note also that it can be shown that the right hand side is equal to $T(S^{(1)}(f))$ where the notation is self-explanatory. Furthermore, it is not hard to see that for two distributions supported in one point, i.e. $S \in \mathcal{E}'_{e_1}(M)$ and $T \in \mathcal{E}'_{e_2}(N)$ the distribution $F_{2M, 2N}(S \otimes T)$ is supported in (e_1, e_2) , i.e. is an element of $\mathcal{E}'_{(e_1, e_2)}(M \times N)$. We shall denote the restriction of the map $F_{2M, 2N}$ to $\mathcal{E}'_{e_1}(M) \otimes \mathcal{E}'_{e_2}(N)$ by the same symbol $F_{2M, 2N}$. For three pointed manifolds (M, e_1) , (N, e_2) , and (P, e_3) , let $\alpha_{M, N, P}: M \times (N \times P) \rightarrow (M \times N) \times P$ be the usual associator for the monoidal category of all sets, and for three vector spaces V, W, X over \mathbb{K} , let $v_{V, W, X}: V \otimes (W \otimes X) \rightarrow (V \otimes W) \otimes X$ be the well-known associator for the monoidal category of all vector spaces. By using the definitions, it is not hard to see that the following identity holds

$$\begin{aligned} & F_{2(M \times N), P} \circ (F_{2M, 2N} \otimes \text{id}_{\mathcal{E}'_{e_3}(P)}) \circ \beta_{\mathcal{E}'_{e_1}(M), \mathcal{E}'_{e_2}(N), \mathcal{E}'_{e_3}(P)} \\ &= (\alpha_{M, N, P})_* \circ F_{2M, (N \times P)} \circ (\text{id}_{\mathcal{E}'_{e_1}(M)} \otimes F_{2N, P}) \end{aligned} \quad (72)$$

hence eqn (3) of [15]. In the same vein, the two diagrams in eqn (4) of [15] are satisfied upon setting $F_0 = \varepsilon_{\text{pt}}$ and $\lambda: \mathbb{K} \otimes V \rightarrow V$ and $\rho: V \otimes \mathbb{K} \rightarrow V$ the usual left-unit and right-unit identifications in the monoidal category of vector spaces.

Let (M', e'_1) and (N', e'_2) two other pointed differentiable manifolds, and let $\phi: (M, e) \rightarrow (M', e'_1)$ and $\psi: (N, e_2) \rightarrow (N', e'_2)$ two smooth maps of pointed differentiable manifolds. It is a straight-forward check that the map $F_{2M, 2N}$ to is natural in the following sense

$$F_{2M', 2N'} \circ (\phi_* \otimes \psi_*) = (\phi \times \psi)_* \circ F_{2M, 2N}. \quad (73)$$

Moreover, note that the map $F_0 = \varepsilon_{\text{pt}}$ (see eqn (67)) defines an isomorphism of $\mathcal{E}'_{\text{pt}}(\{\text{pt}\})$ to \mathbb{K} which had already been seen to be natural.

As a result, the functor F is a monoidal functor in the sense of [15]. Moreover, since the category \mathcal{M}^{pt} is even a symmetric monoidal category by means of the canonical flip map $\tau_{M, N}: M \times N \rightarrow N \times M: (x, y) \rightarrow (y, x)$, see e.g. [15], and the monoidal category $\mathbb{K}\text{-vect}$ is also symmetric, it is not hard to see that the monoidal functor is also symmetric, see e.g. [15] for definitions.

We shall now show that the monoidal functor F is strong, i.e. that $F_0 = \varepsilon_{\text{pt}}$ and $F_{2M, 2N}$ are isomorphisms. This is clear for ε_{pt} . Recall that for each distribution T in $\mathcal{E}'_e(V)$ (where V is a nonempty open set in \mathbb{R}^m containing the point e), there is nonnegative integer l (called the order of the distribution) such that

$$T = \sum_{r=0}^l \sum_{\mathbf{k} \in \mathbb{N}^m, |\mathbf{k}|=r} c_{\mathbf{k}} \frac{\partial^{\mathbf{k}}}{\partial x_{\mathbf{k}}} (\delta_e)$$

Where $c_{\mathbf{k}} \in \mathbb{K}$ for each multi-index \mathbf{k} . In a slightly more algebraic

manner we can express this as follows: let E be a finite-dimensional real vector space, let $V \subset E$ be an open set containing $e \in E$. Then we have the following linear isomorphism $S(E) \rightarrow \mathcal{E}'(V)$ given by $\Phi_S(1) = \delta_e$ and for any positive integer k and vectors $w^{(1)}, \dots, w^{(k)} \in E$ and $f \in \mathcal{C}(V, \mathbb{K})$

$$\Phi_S(w^{(1)} \bullet \dots \bullet w^{(k)})(f) = \frac{\partial^k (f(e + s_1 w^{(1)} + \dots + s_k w^{(k)}))}{\partial s_1 \dots \partial s_k} \Big|_{s_1=0, \dots, s_k=0} \quad (74)$$

where \bullet denotes the commutative multiplication in the symmetric algebra, see Appendix A. Using the fact that the inclusion map $t_{U_\alpha}: U_\alpha \rightarrow M$ of any chart domain of M such that $e \in U_\alpha$ defines an isomorphism $t_{U_\alpha}: \mathcal{E}'(U_\alpha) \rightarrow \mathcal{E}'_e(M)$, and that any chart $\phi_\alpha: U_\alpha \rightarrow V_\alpha \subset \mathbb{R}^m$ defines an isomorphism $\varphi_{\alpha*}: \mathcal{E}'(U_\alpha) \rightarrow \mathcal{E}'_{\varphi_\alpha(e)}(V_\alpha)$, we can conclude that there is a linear isomorphism

$$\Phi_\alpha =: t_{U_\alpha} \circ \varphi_{\alpha*}^{-1} \circ t_{V_\alpha}^{-1} \circ \Phi_S: S(\mathbb{R}^m) \rightarrow \mathcal{E}'_e(M) \quad (75)$$

with the symmetric coalgebra $S(\mathbb{R}^m)$ on \mathbb{R}^m (see Appendix A) computed as follows

$$w^{(1)} \bullet \dots \bullet w^{(k)} \mapsto (f \mapsto \sum_{i_1, \dots, i_k=1}^m \frac{\partial^k (f|_{U_\alpha} \circ \varphi_\alpha^{-1})}{\partial x_{i_1} \dots \partial x_{i_k}} (\varphi_\alpha(e))_{w_{(i_1)} \dots w_{(i_k)}}). \quad (76)$$

where we write $\mathbb{R}^m \ni w_{(j)} = \sum_{i=1}^m w_{(j)i} e_i$ where all the $w_{(j)i}$ are real numbers and e_1, \dots, e_m is the canonical base of \mathbb{R}^m . Note that for the particular case of M being an open set V of \mathbb{R}^m and the chart ϕ_α being the identity map the map Φ_α (see eqs (75) and (76)) coincides with the canonical map Φ_S , see eqn (??).

For two pointed manifolds (M, e_1) and (N, e_2) and given charts (U_α, ϕ_α) of M such that $e_1 \in U_\alpha$ and $(\tilde{U}_\beta, \tilde{\phi}_\beta)$ of N such that $e_2 \in \tilde{U}_\beta$, we thus have linear isomorphisms $\Phi_\alpha: S(\mathbb{R}^m) \rightarrow \mathcal{E}'_{e_1}(M)$, $\tilde{\Phi}_\beta: S(\mathbb{R}^n) \rightarrow \mathcal{E}'_{e_2}(N)$, and $\Phi_{\alpha\beta}: S(\mathbb{R}^{m+n}) \rightarrow \mathcal{E}'_{(e_1, e_2)}(M \times N)$ (upon using the product chart $(U_\alpha \times \tilde{U}_\beta, \varphi_\alpha \times \tilde{\varphi}_\beta)$). Using the above definitions, one can compute that

$$F_{2M, N} \circ (\Phi_\alpha \otimes \tilde{\Phi}_\beta) = \Phi_{\alpha, \beta} \circ \Theta_{m, n}$$

Where $\Theta_{m, n}: S(\mathbb{R}^m) \otimes S(\mathbb{R}^n) \rightarrow S(\mathbb{R}^{m+n})$ denotes the natural isomorphism of commutative associative unital algebras induced by the obvious inclusions $\mathbb{R}^m \rightarrow \mathbb{R}^{m+n}$ (first m coordinates) and $\mathbb{R}^n \rightarrow \mathbb{R}^{m+n}$ (last n coordinates). It follows that the natural map $F_{2M, N}$ is equal to $\Phi_{\alpha, \beta} \circ \Theta_{m, n} \circ (\Phi_\alpha^{-1} \otimes \tilde{\Phi}_\beta^{-1})$ and is thus a linear isomorphism, whence the functor F is a strong monoidal functor.

In order to define more structure, let us consider the well-known diagonal map $\text{diag}_M: M \rightarrow M \times M$ defined by

$$\text{diag}_M(x) = (x, x)$$

for all $x \in M$. Clearly, diag_M is a smooth map of pointed manifolds $(M, e) \rightarrow (M \times M, (e, e))$. Moreover, the diagonal map is clearly natural in the sense that

$$\text{diag}_M \circ \varphi = (\varphi \times \varphi) \circ \text{diag}_M$$

for any smooth map $\varphi: (M, e) \rightarrow (M', e')$ of pointed differentiable manifolds. In other words, the class of diagonal maps diag_M constitutes a natural transformation from the identity functor to the diagonal functor $(M, e) \rightarrow (M \times M, (e, e))$ and $\varphi \rightarrow (\varphi \times \varphi)$. Define the following linear map $\Delta = \Delta_e = \Delta_{(M, e)}: \mathcal{E}'_e(M) \rightarrow \mathcal{E}'_e(M) \otimes \mathcal{E}'_e(M)$ by

$$\Delta_e := F_{2M, M}^{-1} \circ \text{diag}_{M*} \quad (77)$$

This definition has avatars with more than two tensor factors. Indeed, observe that the naturality relation (73) implies for $\phi = \text{id}_M$ and $\psi = \text{diag}_M$ that

$$\begin{aligned} & (\text{id}_M \times \text{diag}_M) \\ &= F_{2M, (M \times M)} \circ (\text{id}_{\mathcal{E}'_e(M)} \times F_{2M, M}) \circ (\text{id}_{\mathcal{E}'_e(M)} \times \Delta_e) \circ F_{2M, M}^{-1} \end{aligned}$$

Similarly, we have relations of this type for any number of tensor factors.

In the following, we invite the reader to look again at Appendix A for definitions and notations about coalgebras.

We have the following

Theorem 3.1 With the above notations: 1. The \mathbb{K} -vector space $\mathcal{E}'_e(M)$ equipped with the linear maps Δ_e (cf. eqn (77)), ε_e (cf. eqn (67)), and 1_e (cf. eqn (68)) is a C^5 -coalgebra which is (non canonically) isomorphic to the standard symmetric coalgebra

$$(S(\mathbb{R}^m), \varepsilon, \Delta, 1).$$

2. The above strong symmetric monoidal functor F extends to a functor –also denoted by F – from $\mathcal{M}f^*$ to the symmetric monoidal category of C^5 -coalgebras over \mathbb{K} .

3. The subspace of all primitive elements of the coaugmented coalgebra $\mathcal{E}'_e(M)$ is natural isomorphic to the tangent space $T_e(M)$. Moreover for each smooth map $\varphi: (M, e) \rightarrow (M', e')$ of pointed manifolds the coalgebra morphism $\varphi_*: \mathcal{E}'_e(M) \rightarrow \mathcal{E}'_{e'}(M')$ induces the tangent map $T_e: T_e(M) \rightarrow T_{e'}(M')$.

Proof: 1. Coassociativity of Δ_e : This follows from the coassociativity-diagram of diag_{M*} by first taking the induced diagram between distribution spaces which reads then

$$(\alpha_{M, M, M}) \circ (\text{id}_M \times \text{diag}_M) \circ \text{diag}_M = (\text{diag}_M \times \text{id}_M) \circ \text{diag}_M.$$

Starting for example on the left hand side, one replaces the map diag_M by $F_{2M, N} \circ \Delta_e$ and also the map $(\text{id}_M \times \text{diag}_M)$ by

$$F_{2M, (M \times M)} \circ (\text{id}_{\mathcal{E}'_e(M)} \times F_{2M, M}) \circ (\text{id}_{\mathcal{E}'_e(M)} \times \Delta_e) \circ F_{2M, M}^{-1}.$$

Now one observes that one may apply the relation (72) on the left hand side. One obtains

$$\begin{aligned} & F_{2(M \times M), M} \circ (F_{2M, M} \times \text{id}_{\mathcal{E}'_e(M)}) \circ \beta \circ (\text{id}_{\mathcal{E}'_e(M)} \times \Delta_e) \circ \Delta_e \\ &= F_{2(M \times M), M} \circ (F_{2M, M} \times \text{id}_{\mathcal{E}'_e(M)}) \circ (\Delta_e \times \text{id}_{\mathcal{E}'_e(M)}) \circ \Delta_e. \end{aligned}$$

One deduces coassociativity.

2. Cocommutativity of Δ_e : This follows from the symmetry of F (already noted before) and the cocommutativity of diag_{M*} .

3. Counitality of Δ_e : This follows from the counitality of diag_{M*} , i.e.

$$(\text{proj}_e \times \text{id}) \circ \text{diag}_M = \text{incl}_e^1, \text{ and } (\text{id} \times \text{proj}_e) \circ \text{diag}_M = \text{incl}_e^2,$$

Where $\text{proj}_e: M \rightarrow \{e\}$, $\text{incl}_e^1: M \rightarrow \{e\} \times M$ and $\text{incl}_e^2: M \rightarrow M \times \{e\}$ are the canonical maps. Indeed, these equations induce the corresponding equations between distribution spaces, and translating direct products into tensor products (and thus $(\text{proj}_e)_*$ into ε and diag_{M*} into Δ_e), one obtains counitality.

4. Connectedness of Δ_e : The coalgebra $\mathcal{E}'_e(M)$ is isomorphic to the symmetric algebra $S(\mathbb{R}^n)$, and the latter is connected.

This shows part (1) of the statement, as the isomorphism to the standard symmetric coalgebra has been shown above.

The only thing which has to be shown for the second statement is the preservation of the coalgebra structure on the level of morphisms, which is clear.

For the third part, consider the linear map $T_e M \rightarrow \text{Prim}(e'_e(M))$ (see Appendix A for the definition of the primitives $\text{Prim}(C)$ of a coalgebra C) defined by

$$v \mapsto (f \mapsto df_e(v)):$$

Indeed the right hand side is clearly in $e'_e(M)$, and the Leibniz rule for the derivative shows that this is in $\text{Prim}(e'_e(M))$. Moreover the above map is clearly injective, and since $\text{Prim}(S(\mathbb{R}^n)) = \mathbb{R}^n$ and $\dim(T_e M) = n$ it follows that the above map is an isomorphism of real vector spaces. The naturality is a simple computation.

The last statement means that the composed functor $\text{Prim} \circ F$ of the Serre functor F and the functor associating to any coalgebra C its space of primitive elements, $\text{Prim}(C)$, is naturally isomorphic to the tangent functor T , associating to any pointed differentiable manifold (M, e) its tangent space $T_e M$.

Remark 3.1: There is neither a canonically defined (i.e. not depending on the choice of a chart) projection from the coalgebra to its primitives, so the coalgebras $e'_e(M)$ are isomorphic to the cofree $S(\mathbb{R}^m)$, but in general not naturally, nor a canonically defined commutative multiplication (the classical convolution of distributions of compact support which needs the additive vector space structure).

Remark 3.2: Note also the disjoint union $\cup_x \in_{M_x} (M)_{(k)}$ carries the structure of a smooth vector bundle over M : Its smooth sections coincide with the space of all differential operators of order k .

Remark 3.3: In case $U \subset \mathbb{R}^m$ and $V \subset \mathbb{R}^n$ are pointed open sets, the coalgebra morphism ϕ , of a smooth map $\phi: U \rightarrow V$ of pointed manifolds is isomorphic to the coalgebra morphism $S(\mathbb{R}^m) \rightarrow S(\mathbb{R}^n)$ induced by the jet of infinite order of ϕ at the distinguished point e of U , $j^\infty(\phi)e$, for further information [18]. The functorial equation $(\phi \circ \psi) = \phi \circ \psi$, can be computed out of the chain rule for higher derivatives.

Pointed manifolds with multiplication and their associated bialgebras

We can now apply the Serre functor defined in the preceding Section 3.2 to pointed manifolds with multiplication:

Theorem 3.2: Let (M, e, m) be a pointed manifold with multiplication. Then the C^3 -coalgebra $e'_e(M)$ carries a multiplication, i.e. a linear map $\mu = m \circ F_{2, M, M}: e'_e(M) \otimes e'_e(M) \rightarrow e'_e(M)$ which is a morphism of C^3 -coalgebras.

In case m is left-unital (resp. right unital), the nonassociative C^3 -algebra $e'_e(M)$ is left regular (resp. right regular)

Proof: The map μ exists and is linear by functoriality. We have trivially

$$\text{diag}_M \circ m = (m \times m) \circ (\text{id}_M \times \tau_{M, M} \times \text{id}_M) \circ (\text{diag}_M \times \text{diag}_M),$$

and this shows that μ is a morphism of coalgebras by translating diag_M into Δ_e using as before the maps of type F_2 . The regularity statements are a consequence of the connectedness of the C^3 -coalgebra $e'_e(M)$, see Lemma 2.1.

In the following, we shall enumerate some important (sub) categories of pointed differentiable manifolds with multiplications.

Lie groups and universal enveloping algebras: Let $(G, m, e, (-)^{-1})$ a Lie group. The following theorem is well-known:

Theorem 3.3: The associated coalgebra with multiplication μ of the Lie group $(G, m, e, (-)^{-1})$ is an associative unital bialgebra (in fact, a Hopf algebra) isomorphic to the universal enveloping algebra of the Lie algebra $\mathfrak{g} = T_e G$ of G .

We just indicate the isomorphism: For any $\xi \in \mathfrak{g}$, let ξ^+ denote the left invariant vector field $\xi^+(g) := T_e L_g(\xi)$ generated by its value $\xi \in \mathfrak{g} = T_e G$. Then the map $\Phi_U: U(\mathfrak{g}) \rightarrow F(G)$ is given by (for all $k \in \mathbb{N}$, $\xi_1, \dots, \xi_k \in \mathfrak{g}$ and $f \in C^\infty(G, \mathbb{K})$)

$$\Phi_U(\xi_1 \cdots \xi_k)(f) = ((\mathbb{L}_{\xi_1^+} \circ \cdots \circ \mathbb{L}_{\xi_k^+})(f'))(e) \quad (78)$$

Where \mathbb{L}_X denotes the Lie derivative in the direction of the vector field X .

Note that the identities for the inverse map $g \rightarrow g^{-1}$ can be written as

$$m \circ ((-)^{-1} \times \text{id}_G) \circ \text{diag}_G = (g \rightarrow e) = m \circ (\text{id}_G \times (-)^{-1}) \circ \text{diag}_G, \quad (79)$$

and an application of the functor F gives the convolution identities for the antipode, defined by $S = ((-)^{-1})$.

Lie semigroups and Lie monoids: It is easy to see but presumably less known that the result of the preceding subsection remains true for a Lie monoid (G, m, e) :

Theorem 3.4: 1. The associated coalgebra with multiplication μ of the Lie monoid (G, m, e) is an associative unital bialgebra (in fact a Hopf algebra) isomorphic to the universal enveloping algebra of a Lie algebra $\mathfrak{g} \cong T_e G$.

2. The associated coalgebra with multiplication μ of the right Lie group $(G, m, e, (-)^{-1})$ is a right Hopf algebra.

In order to see the first statement note that it is clear that the associated coalgebra $C := F(G)$ carries an associative unital multiplication $\mu = m \circ F_{2G, G}$. The fact that the coalgebra is always connected implies by the Takeuchi-Sweedler argument (see Appendix 4) that the identity map id_C has a convolution inverse, and is thus a Hopf algebra. Since the coalgebra C is connected and cocommutative, it follows from the Cartier-Milnor-Moore Theorem that the Hopf algebra $F(G)$ is isomorphic to the universal enveloping algebra over the Lie subalgebra \mathfrak{g} of its primitive elements which is equal to $T_e G$.

The second statement is an immediate consequence of the functorial properties of F .

(Lie) dimonoids and digroups: Recall that a Lie dimonoid (see e.g. Lod 2001) is a pointed differentiable manifold (D, e) equipped with two smooth associative multiplications $D \times D \rightarrow D$, written $(x, y)x \dashv y$ and $(x, y)x \dashv y$ (and preserving points, i.e. $e \dashv e = e \dashv e$), such that the dialgebra conditions eqs (41), (42), (43), (44), and (45) hold for all $x, y, z \in D$ and e (replacing 1): Hence (D, \dashv, e) is a left unital Lie semigroup and (D, \dashv, e) is a right unital Lie semigroup, and as for dialgebras, we shall say bar-unital dimonoid to stress the fact that the bar-unit e is among the data for the dimonoid.

Let us call a Lie dimonoid $(D, e, \dashv, \bar{\dashv})$ balanced iff in addition for all $x \in D$ the analogue of eqn (46) holds, i.e. $x \bar{\dashv} e = e \bar{\dashv} x$. Any Lie monoid (G, e, m) is a Lie dimonoid by setting $\bar{\dashv} = \dashv = m$.

Another class of examples is obtained by the following important augmented dimonoid construction (cf Example 2.5): Let G be a Lie group, let (D, e_D) be a pointed differentiable manifold, let G smoothly act on the left and on the right of M (written $(g, x) \mapsto gx$ and $(x, g) \mapsto xg$)

such that $(g, x)g' = g(x, g')$ for all $g, g' \in \mathfrak{g}$ and $x \in D$, and let $f: (D, e_D) \rightarrow (G, e)$ be a smooth map of pointed manifolds such that for all $g, g' \in G$ and $x \in D$

$$f(gxg) = gf(x)g'. \quad (80)$$

Then the pointed manifold (D, e_D, \vdash, \dashv) will be a (bar-unital) dimonoid by setting

$$x \dashv y := f(x)y \text{ and } x \vdash y := xf(y) \quad (81)$$

A general Lie digroup is defined (according to Liu) to be a (bar-unital) dimonoid (D, e, \vdash, \dashv) such that the left unital Lie semigroup (D, e, \vdash) is a right group and the right unital Lie semigroup (D, e, \dashv) is a left group (see Appendix 5 for definitions): Here the right inverse of x with respect to \vdash does in general not coincide with the left inverse of x with respect to \dashv . For an example, take any Lie group G , set $(D, e_D) = ((G \times G, (e, e)))$, define the two canonical G -actions $g(g_1, g_2) := (gg_1, g_2)$ and $(g_1, g_2)g := (g_1, gg_2)$ (for all $g, g_1, g_2 \in G$), and let $f: G \times G \rightarrow G$ be the group multiplication. Then (D, e_D, \vdash, \dashv) will be a general digroup with $(g_1, g_2) \dashv = (g_2^{-1}g_1^{-1}, e)$ and $(g_1, g_2) \vdash = (e, g_2^{-1}g_1^{-1})$.

In [13, Definition 4.1] Kinyon defines a Lie digroup as a general Lie digroup such that in addition for each x its right inverse (w.r.t. to x) is equal to its left inverse (w.r.t. \dashv). This can be shown to be equivalent to demanding that the general Lie digroup (D, e, \vdash, \dashv) be balanced.

Again using the Suschkewitsch decomposition Theorem (which applies in case the underlying manifold is connected), it is not hard to see that the category of all connected Lie digroups (in the sense of Kinyon) is equivalent to the category of all left \mathcal{G} -spaces, i.e. whose objects are pairs (G, X) where G is a connected Lie group and X is a pointed connected left G -space (i.e. the distinguished point of X is a fixed point of the G -action) with obvious morphisms. Recall that the Lie digroup is given by $X \times G$ equipped with the point (e_x, e) and the two multiplications $(x_1, g_1) \dashv (x_2, g_2) = (g_1x_2g_1, g_2)$ and $(x_1, g_1) \vdash (x_2, g_2) = (x_1, g_1g_2)$ for all $x_1, x_2 \in X$ and $g_1, g_2 \in G$.

The following theorem is a direct consequence of the functorial properties of the functor F :

Theorem 3.5: Let (D, e, \vdash, \dashv) be a bar-unital Lie dimonoid (D, e, \vdash, \dashv) .

1. The underlying vector space of the associated coalgebra $F(D)$ to the bar-unital Lie dimonoid (D, e, \vdash, \dashv) equipped with 1 and the multiplications $\mu \dashv, \mu \vdash$ is an associative bar-unital dialgebra.

In case D is balanced, $F(D)$ is a cocommutative Hopf dialgebra.

2. In case (D, e, \vdash, \dashv) is a Lie digroup (in the sense of Kinyon), $F(D)$ is a cocommutative Hopf dialgebra.

(Lie) racks: Recall that a Lie rack is a pointed manifold with multiplication (M, e, m) satisfying the following identities for all $x, y, z \in M$ where the standard notation is $m(x, y) = x \triangleright y$

$$e \triangleright x = x, \quad (82)$$

$$x \triangleright e = e, \quad (83)$$

$$x \triangleright (y \triangleright z) = (x \triangleright y) \triangleright (x \triangleright z) \quad (84)$$

In addition, one demands that (M, e, m) be left-regular, i.e. for all $x \in M$ the left multiplication maps $L_x: y \rightarrow x \triangleright y$ should be a diffeomorphism.

Note the following version of the self-distributivity identity (84) in terms of maps:

$$m \circ (\text{id}_M \times m)$$

$$= m \circ (m \times m) \circ (\text{id}_M \times \tau_{M, M} \times \text{id}_M) \circ (\text{diag}_M \times \text{id}_M \times \text{id}_M) \quad (85)$$

Example 3.1: Note that every pointed differentiable manifold (M, e) carries a trivial Lie rack structure defined for all $x, y \in M$ by

$$x \triangleright y := y, \quad (86)$$

and this assignment is functorial.

Example 3.2: Any Lie group G becomes a Lie rack upon setting for all $g, g' \in G$

$$g \triangleright g' := gg'g^{-1}, \quad (87)$$

again defining a functor from the category of Lie groups to the category of all Lie racks.

Example 3.3: Let G be a Lie group and V be a (smooth) G -module (supposed to be a real or complex vector space). On $X := V \times G$, we define a binary operation \triangleright by

$$(v, g) \triangleright (v', g') = (g(v'), gg'g^{-1})$$

for all $v, v' \in V$ and all $g, g' \in G$. X is a Lie rack with unit $1 := (0, 1)$ which is called a linear Lie rack.

Example 3.4: Let (D, e, \vdash, \dashv) be a (balanced) digroup. Then formula (13) of [5],

$$x \triangleright y := x \dashv y \dashv x^{-1}$$

equips the pointed manifold (D, e, \triangleright) with the structure of a Lie rack.

Any Lie rack (M, e, \triangleright) can be gauged by any smooth map $f: (M, e) \rightarrow (M, e)$ of pointed manifolds satisfying for all $x, y \in M$

$$f(x \triangleright y) = x \triangleright f(y).$$

A straight-forward computation shows that the pointed manifold (M, e) equipped with the gauged multiplication \triangleright_f defined by

$$x \triangleright_f y := f(x) \triangleright y$$

is a Lie rack (M, e, \triangleright_f) .

Furthermore, recall that an augmented Lie rack (M, ϕ, G, ℓ) consists of a pointed differentiable manifold (M, e_M) , of a Lie group G , of a smooth map $\phi: M \rightarrow G$ (of pointed manifolds), and of a smooth left G -action $\ell: G \times M \rightarrow M$ (written $(g, x) \mapsto \ell(g, x) = \ell_g(x) = gx$) such that for all $g \in G, x \in M$

$$g^e_M = e_M \quad (88)$$

$$\phi(gx) = g\phi(x)g^{-1}. \quad (89)$$

It is a routine check that the multiplication \triangleright on M defined for all $x, y \in M$ by

$$x \triangleright y := \ell_{\phi(x)}(y) \quad (90)$$

satisfies all the axioms (82), (83), and (84) of a Lie rack, thus making (M, e_M, \triangleright) into a Lie rack such that the map ϕ is a morphism of Lie racks, i.e. for all $x, y \in M$

$$\phi(x \triangleright y) = \phi(x)\phi(y)\phi(x)^{-1}. \quad (91)$$

A morphism $(\Psi, \psi): (M, \phi, G, \ell) \rightarrow (M', \phi', G', \ell')$ of augmented Lie racks is a pair of maps of pointed differentiable manifolds $\Psi: M \rightarrow M'$ and $\psi: G \rightarrow G'$ such that ψ is homomorphism of Lie groups and such that all reasonable diagrams commute, viz: for all $g \in G$

$$\phi' \circ \Psi = \psi \circ \phi \quad (92)$$

$$\Psi \circ \ell_{g=\ell'} \circ \Psi \tag{93}$$

Note that the trivial Lie rack structure of a pointed manifold (M, e) comes from an augmented Lie rack over the trivial Lie group $G=\{e\}$.

Let (M, e, \triangleright) be a Lie rack. Applying the functor F we get the following

Theorem 3.6: The associated coalgebra $F(M)$ with multiplication μ of the Lie rack (M, e, m) is a rack bialgebra, i.e. satisfying for all $a, b, c \in C$, using the same notation $a \triangleright b$ for $\mu(a \otimes b)$:

$$1 \triangleright a = a, \tag{94}$$

$$a \triangleright 1 = \varepsilon(a)1, \tag{95}$$

$$a \triangleright (b \triangleright c) = \sum_{(a)} (a^{(1)} \triangleright b) \triangleright (a^{(2)} \triangleright c). \tag{96}$$

Proof: 1. By definition, $S \otimes T \in \mathcal{E}'_e(M) \otimes \mathcal{E}'_e(M)$ are sent by $\mu = \triangleright$, to the distribution $f \mapsto F_{2M, M}(S \otimes T)(f^\triangleright)$. We evaluate this formula for $S=1$. This gives the distribution $f \mapsto 1(T^{(2)}(f^\triangleright))$. But $T^{(2)}$ means that the function is seen as function of its second variable, i.e. $T^{(2)}(f^\triangleright)(y) = T(f(y \triangleright -))$. On the other hand, the delta distribution 1 evaluates a function in e , thus

$$1(T^{(2)}(f^\triangleright)) = T(f(e \triangleright -)) = T(f),$$

Because $e \triangleright y = y$ for all $y \in M$. This shows $1T = T$.

2. Exchanging the roles of the two variables in the above computation, we obtain for $T \triangleright 1$ the distribution

$T(1^{(2)}(f^\triangleright))$ or in other words $1(T^{(1)}(f^\triangleright))$, i.e. the above element y is now in the second place. We obtain

$$1(T^{(1)}(f^\triangleright)) = T(f(- \triangleright e)) = T(f(e)) = T(1) = \varepsilon(T).$$

This shows $T \triangleright 1 = \varepsilon(T)1$.

3. As remarked before, the definition of Δ_e , namely $\Delta_e = F_{2M, M}^{-1} \circ \text{diag}_{M^*}$, induces thanks to the naturality relation (73) relations like

$$(id_M \times \text{diag}_M)_* = F_{2M, (M \times M)} \circ (id_{\mathcal{E}'_e(M)} \times F_{2M, M}) \circ (id_{\mathcal{E}'_e(M)} \times \Delta_e) \circ F_{2M, M}^{-1},$$

and

$$\begin{aligned} &\Delta_e \times id_{\mathcal{E}'_e(M)} \times id_{\mathcal{E}'_e(M)} \\ &= (F_{2M, M}^{-1} \times id_{\mathcal{E}'_e(M)} \times id_{\mathcal{E}'_e(M)}) \circ (F_{2(M \times M), M}^{-1} \times id_{\mathcal{E}'_e(M)}) \\ &\circ F_{2(M \times M) \times M, M}^{-1} \circ (\text{diag}_M \times id_M \times id_M)_* \\ &\circ F_{2(M \times M), M} \circ (F_{2M, M} \times id_{\mathcal{E}'_e(M)}). \end{aligned}$$

Therefore, starting from the relation induced on $\mathcal{E}'_e(M)$ by relation (85), one replaces $(\text{diag}_M \times id_M \times id_M)_*$ by the above and obtains finally an equation equivalent to equation (8).

Remark 3.4: This theorem should be compared to Proposition 3.1 in [3]. In [3], the authors work with the vector space $K[M]$ generated by the rack M , while we work with point-distributions on a Lie rack M . Once again, in some sense, we extend their Proposition 3.1 “to all orders”. Observe however that their structure is slightly different (motivated in their Remark 7.2).

We get a similar theorem for an augmented Lie rack: Let \mathfrak{g} denote the Lie algebra of the Lie group G , then we have the:

Theorem 3.7: The associated coalgebra C with multiplication μ of an augmented Lie rack (M, ϕ, G, ℓ) is a cocommutative augmented rack bialgebra $(C, \phi, U(\mathfrak{g}), \ell)$

We shall close the subsection with a geometric explanation of some of the structures appearing in Subsection 2.1: Let $(\mathfrak{h}, [\cdot, \cdot])$ be a real finite-dimensional Leibniz algebra. There is the following Lie rack structure on the manifold \mathfrak{h} defined by

$$x \blacktriangleright y := e^{\text{ad}_x}(y) \tag{97}$$

Moreover, pick a two-sided ideal $\mathfrak{z} \subset \mathfrak{h}$ with $Q(\mathfrak{h}) \subset \mathfrak{z} \subset \mathfrak{z}(\mathfrak{h})$ so that the quotient algebra $\mathfrak{g} := \mathfrak{h}/\mathfrak{z}$ is a Lie algebra. Let $p: \mathfrak{h} \rightarrow \mathfrak{g}$ be the canonical projection. Let G be the connected simply connected Lie group having Lie algebra \mathfrak{g} . Since \mathfrak{g} acts on \mathfrak{h} as derivations, there is a unique Lie group action ℓ of G on \mathfrak{h} by automorphisms of Leibniz algebras. Consider the smooth map

$$\phi: \mathfrak{h} \rightarrow G: x \mapsto \exp(p(x)). \tag{98}$$

Clearly $\phi(g \cdot x) = g\phi(x)g^{-1}$ for all $x \in \mathfrak{h}$ and $g \in G$ whence $(\mathfrak{h}, \phi, G, \ell)$ is an augmented Lie rack, and it is not hard to see that the Lie rack structure coincides with (97).

Theorem 3.8: The C^∞ -rack bialgebra associated to the augmented Lie rack $(\mathfrak{h}, \phi, G, \ell)$ by means of the Serre functor is isomorphic to the universal envelopping algebra of infinite order, $\text{UAR}^\infty(\mathfrak{h})$, see Definition 2.3 and Theorem 2.1.

Proof: First we compute $\phi_* \exp_* \circ p_*$. Since $p: \mathfrak{h} \rightarrow \mathfrak{g}$ is linear, it is easy to see using formula (??) that for all $k \in \mathbb{N}$ and $x_1, \dots, x_k \in \mathfrak{h}$

$$p^*(\Phi_S(x_1 \bullet \dots \bullet x_k)) = \Phi_S(p(x_1) \bullet \dots \bullet p(x_k)) = \Phi_S(p)(x_1 \bullet \dots \bullet x_k),$$

see (??) for a definition of Φ_S . Next, for all $k \in \mathbb{N}$ and $\xi_1, \dots, \xi_k \in \mathfrak{g}$, we shall show the formula (for all $f \in C^\infty(G, \mathbb{K})$)

$$\begin{aligned} (\exp_*(\Phi_S(\xi_1 \bullet \dots \bullet \xi_k)))(f') &= \sum_{\sigma \in S_k} \frac{1}{k!} (\Phi_U(\xi_{\sigma(1)} \cdots \xi_{\sigma(k)}))(f') \\ &= (\Phi_U(\omega(\xi_1 \bullet \dots \bullet \xi_k)))(f') \end{aligned}$$

(see eqn (78) for a definition of Φ_U). Both sides of this equation are symmetric k -linear maps in the arguments ξ_1, \dots, ξ_k , hence by the Polarization Lemma, it suffices to check equality in case $\xi_1 = \dots = \xi_k = \xi$ [19]. Since for each real number t the map $g \mapsto F_t^\xi(g) := g \exp(t\xi)$ is the flow of the left invariant vector field ξ^+ , we get

$$\begin{aligned} (\exp_*(\Phi_S(\xi \bullet \dots \bullet \xi)))(f') &= \frac{\partial^k}{\partial t^k} (f'(\exp(t\xi))) \Big|_{t=0} = \frac{\partial^k}{\partial t^k} (f'(F_t^\xi(e))) \Big|_{t=0} \\ &= \frac{\partial^k}{\partial t^k} (((F_t^\xi)^* f')(e)) \Big|_{t=0} \\ &= ((F_t^\xi)^* ((\mathbb{L}_{\xi^+} \circ \dots \circ \mathbb{L}_{\xi^+})(f')))(e) \Big|_{t=0} \\ &= ((\mathbb{L}_{\xi^+} \circ \dots \circ \mathbb{L}_{\xi^+})(f'))(e) \\ &= (\Phi_U(\omega(\xi_1 \bullet \dots \bullet \xi_k)))(f') \end{aligned}$$

proving the above formula. It follows that

$$\phi_* = \exp_* \circ p_* = \Phi_U \circ \omega \circ S(p) \circ \Phi_S^{-1}. \tag{99}$$

Next, we compute ℓ_* . We get for positive integers k, l , $\xi_1, \dots, \xi_k \in \mathfrak{g}$, $x \in \mathfrak{h}$, and $f \in C(\mathfrak{h}, \mathbb{K})$:

$$(\ell^*(\Phi_U(\xi_1 \dots \xi_k) \otimes \Phi_S(x^l)))(f)$$

$$\begin{aligned}
 &= \frac{\partial^{k+l}}{\partial s_1 \dots \partial s_k \partial t^l} \left(f \left((\ell_{\exp(s_1 \xi_1)} \circ \dots \circ \ell_{\exp(s_k \xi_k)})(tx) \right) \right) \Bigg|_{s_1 = \dots = s_k = 0 = t} \\
 &= \frac{\partial^k}{\partial s_1 \dots \partial s_k} \sum_{j_1 \dots j_l = 1}^{\dim(\mathfrak{h})} \left(\frac{\partial^l f}{\partial x_{j_1} \dots \partial x_{j_l}}(0) \ell_{s_{j_1}}(x) \dots \ell_{s_{j_l}}(x) \right) \Bigg|_{s_1 = \dots = s_k = 0}
 \end{aligned}$$

where in the last line we have used a basis of \mathfrak{h} , have written, y_1, \dots, y_n ($n = \dim(\mathfrak{h})$) for the components of each vector $y \in \mathfrak{h}$, and used the notation ℓ_s for the linear map $\ell_{\exp(s_1 \xi_1)} \circ \dots \circ \ell_{\exp(s_k \xi_k)}$. By induction on k it is easy to prove that

$$\frac{\partial^k}{\partial s_1 \dots \partial s_k} (\ell_{s_1}(x) \bullet \dots \bullet \ell_{s_k}(x)) \Bigg|_{s_1 = \dots = s_k = 0} = \text{ad}_{\xi_1 \dots \xi_k}^s(x^{\bullet}),$$

and using again the Polarisation Lemma, we finally get for all $u \in U(\mathfrak{g})$ and $\alpha \in S(\mathfrak{h})$

$$\ell_*(\Phi_U(u) \otimes \Phi_S(\alpha)) = \Phi_S(\text{ad}_u^s(\alpha)) = \Phi_S(u, \alpha), \tag{100}$$

and the isomorphism with the augmented rack bialgebra $\text{UAR}^\sim(\mathfrak{h}) = S(\mathfrak{h})$ is established.

Remark 3.5: Observe that the Serre functor can be rendered completely algebraic, i.e. for example for an algebraic Lie rack R (meaning that the underlying pointed manifold is a smooth algebraic variety and the rack product is algebraic), one can take as its Serre functor image $F(R)$ the space of derivations along the evaluation map in the distinguished point. The composition of F with the functor of primitives gives then the tangent functor (see text before Remark 3.1). This gives a new and completely algebraic functorial way to associate to a Lie rack its tangent Leibniz algebra.

Some Definitions Around Coalgebras (Appendix)

Let C be a module over a commutative associative unital ring K (which we shall assume to contain \mathbb{Q}). Recall that a linear map $\Delta: C \rightarrow C \otimes_K C = C \otimes C$ is called a coassociative comultiplication iff $(\Delta \otimes \text{id}_C) \circ \Delta = (\text{id}_C \otimes \Delta) \circ \Delta$, and the pair (C, Δ) is called a (coassociative) coalgebra over K . Let (C', Δ') be another coalgebra. Recall that a K -linear map $\Phi: C \rightarrow C'$ is called a homomorphism of coalgebras iff $\Delta' \circ \Phi = (\Phi \otimes \Phi) \circ \Delta$. The coalgebra (C, Δ) is called cocommutative iff $\tau \circ \Delta = \Delta$ where $\tau: C \otimes C \rightarrow C \otimes C$ denotes the canonical flip map. Recall furthermore that a linear map $\varepsilon: C \rightarrow K$ is called a counit for the coalgebra (C, Δ) iff $(\varepsilon \otimes \text{id}_C) \circ \Delta = (\text{id}_C \otimes \varepsilon) \circ \Delta$. The triple (C, Δ, ε) is called a counital coalgebra. Moreover, a counital coalgebra (C, Δ, ε) equipped with an element 1 is called coaugmented iff $\Delta(1) = 1 \otimes 1$ and $\varepsilon(1) = 1 \in K$. Let $C^{\circ} \subset C$ denote the kernel of ε . Recall that a morphism $\phi: (C, \Delta, \varepsilon, 1) \rightarrow (C', \Delta', \varepsilon', 1')$ of counital coaugmented coalgebras over K is a K -linear map satisfying $(\phi \otimes \phi) \circ \Delta = \Delta' \circ \phi$, $\varepsilon' \circ \phi = \varepsilon$, and $\phi(1) = 1'$. Moreover, for any counital coaugmented coalgebra the K -submodule of all primitive elements is defined by

$$\text{Prime}(C) := \{x \in C \mid \Delta(x) = x \otimes 1 + 1 \otimes x\}. \tag{101}$$

Every morphism of counital coaugmented coalgebra clearly maps primitive elements to primitive elements, thus defining a functor Prim from the category of counital coaugmented coalgebras to the category of K -modules. Finally, following Quillen, we shall call a counital coaugmented coalgebra connected iff the following holds: The sequence of submodules $(C_{(r)})_{r \in \mathbb{N}}$ defined by $C_{(0)} = K1$ and recursively by

$$C_{(k+1)} := \{x \in C \mid \Delta(x) - x \otimes 1 - 1 \otimes x \in C_{(k)}\} \tag{102}$$

is easily seen to be an ascending sequence of coaugmented counital

subcoalgebras of $(C, \Delta, \varepsilon, 1)$, and if the union of all the $C_{(k)}$ is equal to C , then $(C, \Delta, \varepsilon, 1)$ is called connected. We refer to each $C_{(k)}$ as the subcoalgebra of order k . Clearly, each $C_{(k)}$ is connected, and $C_{(1)} = K1 \otimes \text{Prim}(C)$. Moreover, each morphism of counital coaugmented coalgebras maps each subcoalgebra of order k to the subcoalgebra of order k of the target coalgebra thus defining a functor $C \rightarrow C_{(k)}$ from the category of coaugmented counital coalgebras to itself. We shall use the following acronyms:

Definition 4.1: We call a coassociative, counital, coaugmented coalgebra a C^3 -coalgebra. In case the C^3 -coalgebra is in addition cocommutative, we shall speak of a C^4 -coalgebra. Finally, a connected C^4 -coalgebra will be coined a C^5 -coalgebra.

Recall also that the tensor product of two counital coaugmented coalgebras $(C, \Delta, \varepsilon, 1)$ and $(C', \Delta', \varepsilon', 1')$ is given by $(C \otimes C', (\text{id}_C \otimes \tau \otimes \text{id}_{C'}) \circ (\Delta \otimes \Delta'), \varepsilon \otimes \varepsilon', 1 \otimes 1')$. Tensor products of connected coalgebras are connected. Recall the standard example: Let V be a K -module and $S(V) = \bigoplus_{r=0}^{\infty} S^r(V)$ be the symmetric algebra generated by V , i.e. the free algebra $T(V)$ (for which we denote the tensor multiplication by suppressing the symbol) modulo the two-sided ideal I generated by $xy - yx$ for all $x, y \in V$. Denoting the commutative associative multiplication in $S(V)$ (which is induced by the free multiplication) by \cdot , i.e.

$$x_1 \bullet \dots \bullet x_k := x_1 \dots x_k \text{ mod } I,$$

we have $\Delta(x) = x \otimes 1 + 1 \otimes x$ for all $x \in V$ and $\Delta(x_1 \bullet \dots \bullet x_k) = (x_1 \otimes 1 + 1 \otimes x_1) \bullet \dots \bullet (x_k \otimes 1 + 1 \otimes x_k)$ for all positive integers k and $x_1, \dots, x_k \in V$. Recall that $S^0(V)$ is the free K -module $K1$ and the counit is defined by $(\lambda 1) = \lambda$ for all $\lambda \in K$ and by declaring that ε vanishes on $\bigoplus_{r=1}^{\infty} S^r(V)$. Moreover, the submodules $(S(V))_{(n)}$ are given by $\sum_{r=0}^{\infty} \lambda_r \varphi^{*r}$, whence $S(V)$ is clearly connected, so it is a C^5 -coalgebra whose submodule of primitive elements equals V .

Moreover, for a given coalgebra (C, Δ) and a given nonassociative algebra (A, μ) where $\mu: A \otimes A \rightarrow A$ is a given K -linear map, recall the convolution multiplication in the K -module $\text{Hom}_K(C, A)$ defined in the usual way for any two K -linear maps $\phi, \psi: C \rightarrow A$ by

$$\phi * \psi := \mu \circ (\phi \otimes \psi) \circ \Delta. \tag{103}$$

In case Δ is coassociative and associative, $*$ will be associative. The following fact is rather important: If C is connected and if the K -linear map $\phi: C \rightarrow A$ vanishes on 1_C , then any convolution power series of ϕ converges, i.e. the evaluation of some formal series $\sum_{r=0}^{\infty} \lambda_r \phi^{*r}$ (with $\lambda_r \in K$ and $\phi^{*0} := 1_A \varepsilon_C$) on $c \in C$ always reduces to a finite number of terms. In particular, let $\psi: C \rightarrow A$ be a K -linear map such that $\psi(1_C) = 1_A$. Then – as has been observed by Takeuchi and Sweedler – ψ has always a convolution inverse, i.e. there is a unique K -linear map $\psi': C \rightarrow A$ such that $\psi * \psi' = 1_A \varepsilon_C = \psi'^* \psi$, where ψ' is defined by the geometric series $\psi' = \sum_{r=0}^{\infty} (1_A \varepsilon_C - \psi)^{*r}$ [20,21].

Semigroups

We collect some properties of semigroups which are very old, but a bit less well-known than properties of groups. The standard reference to these topics is the book [22] by A. H. Clifford and G. B. Preston.

Recall that a semigroup Γ is a set equipped with an associative multiplication $\Gamma \times \Gamma \rightarrow \Gamma$, written $(x, y) \mapsto xy$. An element e of Γ is called a left unit element (resp. a right unit element resp. a unit element) iff for all $x \in \Gamma$ we have $ex = x$ (resp. $xe = x$ resp. iff e is both left and right unit element). A pair (Γ, e) of a semigroup Γ and an element e is called left

unital (resp. right unital resp. unital) iff e is a left unit element (resp. a right unit element resp. a unit element). A unital semigroup is also called a monoid. It is well-known that the unit element of a monoid is the unique unit element (unlike left or right unit elements in general). Let (Γ, e) be a right unital or a left unital semigroup. Recall that for a given element $x \in \Gamma$ an element $y \in \Gamma$ is called a left inverse of x (resp. a right inverse of x resp. an inverse of x) iff $yx=e$ (resp. $xy=e$ resp. iff y is both a left and a right inverse of x). Clearly, a unital semigroup (Γ, e) such that every element has an inverse is a group. In that case it is well-known that for each x there is exactly one inverse element, called x^{-1} .

Note that by a Lemma by L. E. Dickson every left unital semigroup such that each element has at least one left inverse is already a group which can be shown by just using the definitions. Dually, every right unital semigroup such that each element has at least one right inverse is also a group.

More interesting is the case of a left (resp. right) unital semigroup (Γ, e) such that every element x has at least one right (resp. left) inverse element. In that case (which is an equivalent formulation of a so-called right group (resp. left group), the conclusion of Dickson's Lemma does no longer hold. In order to see what is going on, there is first the following useful

Lemma 5.1: Let (Γ, e) be a left-unital semigroup, let a, b, c three elements of Γ such that

$$ab=e \text{ and } bc=e.$$

Then

$$c=ae, be=b,$$

and the left multiplications $L_a: x \rightarrow ax$ and $L_b: x \rightarrow bx$ are invertible. In particular, given the element a , its right inverse b is unique under the above hypotheses.

The proof is straight-forward.

The structure of right (resp. left) groups is completely settled in the Suschkewitsch Decomposition Theorem, 1928: Given a right group (Γ, e) , it can be shown –using the above Lemma and elementary manipulations, see also– that all the left multiplications $L_x: y \rightarrow xy$ (resp. right multiplications $R_x: y \rightarrow yx$) are invertible, that for each element there is exactly one right (resp. left) inverse (whence there is a map $\Gamma \rightarrow \Gamma$ assigning to each element x its right (resp. left) inverse x^{-1}), that the image of this right (resp. left) inverse map is equal to Γe (resp. $e\Gamma$) (which turns out to be a subgroup of (Γ, e)), and that (Γ, e) is isomorphic to the cartesian product $(\Gamma e \times E, (e, e))$ (resp. $(E \times e\Gamma, (e, e))$) where E is the set of all left (resp. right) unit elements in (Γ, e) (coinciding with the set of all idempotent elements). For right groups, the aforementioned isomorphism is given as follows:

$$\phi: \Gamma \times E \rightarrow \Gamma: (a, f) \mapsto af, \tag{104}$$

$$\phi^{-1}: \Gamma \rightarrow \Gamma \times E: x \mapsto (xe, x^{-1}x). \tag{105}$$

Note that both components of ϕ^{-1} are idempotent maps. There is a completely analogous statement for left groups.

Recall that a Lie semigroup is a differentiable manifold Γ equipped with a smooth associative multiplication $m: \Gamma \times \Gamma \rightarrow \Gamma$. All the other definitions of semigroups mentioned above (such as left unital, right unital semigroups, monoids, groups, right groups, left groups etc.) carry over to the Lie, i.e. differentiable, case.

Moreover for right Lie groups, it is easy to see that all the left

multiplications are diffeomorphisms (since their inverse maps are left multiplications with the inverse elements and therefore smooth). This fact and the regular value theorem applied to the equation $xy=e$ imply that the right inverse map is smooth since its graph is a closed submanifold of $\Gamma \times \Gamma$ and the restriction of the projection on the first factor of the graph is a diffeomorphism. As the maps $x \rightarrow xe = (x^{-1})^{-1}$ and $x \rightarrow x^{-1}x$ are smooth and idempotent, it follows that their images, the subgroup Γe , and the semigroup of all left unit elements, E , are both smooth submanifolds of Γ and closed sets provided Γ is connected for a proof [23]. Hence Γe is a connected Lie group, and the Suschkewitsch decomposition $\Gamma \cong \Gamma e \times E$, see Appendix 5, is a diffeomorphism. Conversely, any cartesian product of a Lie group G and a differentiable manifold E equipped with the multiplication $(g, x)(h, y) := (gh, y)$ is easily seen to be a right Lie group. An analogous statement holds for left Lie groups.

It is not hard to see that the category of all connected right Lie groups is equivalent to the product of category of all connected Lie groups and the category of all pointed connected manifolds.

Acknowledgements

F.W. thanks Université de Haute Alsace for an invitation during which the shape of this research project was defined. At some point, the subject of this paper was joint work of S.R. and F.W. with Simon Covez, and we express our gratitude to him for his contributions. Starting from this, S.R. came independently to similar results which we incorporated in this paper. M.B. thanks Nacer Makhlouf for his question about the relations of rack-bialgebras to dialgebras, and Gwénael Massuyeau for his question about universals.

References

- Fenn R, Rourke C (1992) Racks and links in codimension 2. J. Knot Theory Ramifications 1: 343-406.
- Krähmer U, Wagemann F (2014) Racks, Leibniz algebras and Yetter-Drinfel'd modules. Georgian J Math 22: 529-542.
- Carter J, Crans A, Elhamedi M, Saito M (2008) Cohomology of the Categorical Self-Distributivity. J Homotopy Relat Struct 3: 13-63.
- Lebed V (2012) Objets tressés: une étude unificatrice de structures algébriques et une catégorification des tresses virtuelles. doctoral thesis Paris 7.
- Kinyon MK (2007) Leibniz algebras, Lie racks, and digroups. J Lie Theory 17: 99-114.
- Goichot F (2001) Un théorème de Milnor-Moore pour les algèbres de Leibniz. In: Loday JL, Frabetti A, Chapoton F, Goichot F (eds.) Dialgebras and related Operads. Lecture Notes in Mathematics 1763, Springer, Berlin pp: 111-133.
- Loday JL (2001) Dialgebras. In: Loday JL, Frabetti A, Chapoton F, Goichot F (eds.) Dialgebras and related Operads. Lecture Notes in Mathematics 1763, Springer, Berlin, pp: 6-66.
- Covez S (2006) Digroupes de Lie: Une réponse partielle au problème des "coquecigrues", Mémoire de Master 2, Université de Nantes.
- Alexandre C, Bordemann M, Rivière S, Wagemann F, Algebraic deformation quantization of Leibniz algebras. in preparation.
- Blokh AM (1965) On a generalization of the concept of a Lie algebra. Dokl. Akad. Nauk. USSR 165: 471-473.
- Loday JL (1992) Cyclic Homology. Springer, Berlin Vol: 301.
- Daniel Q (1969) Rational Homotopy Theory. Ann Math 90: 205-295.
- Dixmier J (1974) Algèbres enveloppantes. Gauthier-Villars, Paris.
- Mac Lane S (1998) Categories for the Working Mathematician. (2nd edn) Springer, New York.
- Rudin W (1973) Functional Analysis. Tata - McGraw-Hill Publ, New Delhi.
- Serre JP (1964) Lie algebras and Lie groups. Springer Lecture Notes in Mathematics 1500, Springer, Berlin.
- Trèves F (1967) Topological Vector Spaces, Distributions And Kernels. Academic Press, New York.

18. Kolar I, Michor P, Slovák J (1993) Natural Operations in Differential Geometry. Springer, Berlin.
19. Thomas Erik GF (2013) A polarization identity for multilinear maps.
20. Takeuchi M (1971) Free Hopf algebras generated by coalgebras. J Math Soc Japan 23: 561-582.
21. Sweedler M (1969) Hopf Algebras. WA Benjamin, New York.
22. Clifford AH, Preston GB (1961) The Algebraic Theory of Semigroups I. Mathematical Surveys 7, AMS, Providence, Rhode Island.
23. Bröcker T, Jänich K (1973) Einführung in die Differentialtopologie. Springer, Berlin.

Citation: Alexandre C, Bordemann M, Rivière S, Wagemann F (2016) Structure Theory of Rack-Bialgebras. J Generalized Lie Theory Appl 10: 244.
doi:[10.4172/1736-4337.1000244](https://doi.org/10.4172/1736-4337.1000244)

OMICS International: Open Access Publication Benefits & Features

Unique features:

- Increased global visibility of articles through worldwide distribution and indexing
- Showcasing recent research output in a timely and updated manner
- Special issues on the current trends of scientific research

Special features:

- 700+ Open Access Journals
- 50,000+ editorial team
- Rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at major indexing services
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.omicsonline.org/submission>