

## TOTALLY UMBILICAL LIGHTLIKE SUBMANIFOLDS

K. L. DUGGAL AND D. H. JIN

### Abstract

This paper provides new results on a class of totally umbilical lightlike submanifolds in semi-Riemannian manifolds of constant curvature. We prove that the induced Ricci tensor of any such submanifold is symmetric if and only if its screen distribution is integrable.

### 1. Introduction

The theory of submanifolds of a Riemannian or semi-Riemannian manifold is well known (see for example, Chen [4] and O'Neill [12]). However, its counterpart of lightlike (null) submanifolds (for which the local and global geometry is completely different than the non-degenerate case) is relatively new and in a developing stage ([1, 3, 5–9, 11]). In 1996, the first author and Bejancu published their work (see Chapters 4 and 5 of [8]) on lightlike submanifolds  $M$  of semi-Riemannian manifolds. They constructed structure equations for four possible cases of  $M$ , proved the fundamental existence theorem for lightlike submanifolds and found some geometric conditions for the induced connection on  $M$  to be a metric connection. Much of their study was restricted to totally geodesic lightlike submanifolds of semi-Riemannian manifolds. In this paper we study further the geometry of totally umbilical lightlike submanifolds  $M$ .

In Sections 2 and 3, we recall some results for lightlike submanifolds and their structure equations. In Section 4, we prove several new theorems on  $M$  in semi-Riemannian manifolds of constant curvature. Finally, in Section 5, we find conditions for the induced Ricci curvature tensor of  $M$  to be symmetric. The paper contains several simple examples.

### 2. Lightlike submanifolds

Let  $(\bar{M}, \bar{g})$  be a real  $(m+n)$ -dimensional semi-Riemannian manifold of constant index  $q$  such that  $m, n \geq 1$ ,  $1 \leq q \leq m+n-1$  and  $(M, g)$  an  $m$ -dimensional

---

2000 *Mathematics Subject Classification*: 53B25, 53C40, 53C50.

*Keywords*: Degenerate metric, totally umbilical submanifolds.

Received April 12, 2002; revised October 17, 2002.

submanifold of  $\overline{M}$ . In case  $\overline{g}$  is degenerate on the tangent bundle  $TM$  of  $M$  we say that  $M$  is a lightlike submanifold of  $\overline{M}$  [8]. Denote by  $F(M)$  the algebra of smooth functions on  $M$  and by  $\Gamma(E)$  the  $F(M)$  module of smooth sections of a vector bundle  $E$  (same notation for any other vector bundle) over  $M$ . The following range of indices is used:

$$\begin{aligned} i, j, k, \dots &\in \{1, \dots, r\}; & a, b, c, \dots &\in \{r+1, \dots, m\}; \\ A, B, C, \dots &\in \{1, \dots, m\}; & \alpha, \beta, \gamma, \dots &\in \{r+1, \dots, n\}. \end{aligned}$$

For a degenerate tensor field  $g$  on  $M$ , there exists locally a vector field  $\xi \in \Gamma(TM)$ ,  $\xi \neq 0$ , such that  $g(\xi, X) = 0$ , for any  $X \in \Gamma(TM)$ . Then, for each tangent space  $T_x M$  we have

$$T_x M^\perp = \{u \in T_x \overline{M} : \overline{g}(u, v) = 0, \forall v \in T_x M\},$$

which is a degenerate  $n$ -dimension subspace of  $T_x \overline{M}$ . The radical (null) subspace of  $T_x M$ , denoted by  $\text{Rad } T_x M$ , is defined by

$$\text{Rad } T_x M = \{\xi_x \in T_x M; g(\xi_x, X) = 0, X \in T_x M\}.$$

The dimension of  $\text{Rad } T_x M = T_x M \cap T_x M^\perp$  depends on  $x \in M$ . The submanifold  $M$  of  $\overline{M}$  is said to be  $r$ -lightlike submanifold if the mapping

$$\text{Rad } TM : x \in M \rightarrow \text{Rad } T_x M$$

defines a smooth distribution on  $M$  of rank  $r > 0$ , where  $\text{Rad } TM$  is called the radical (null) distribution on  $M$ . Following are four possible cases:

CASE 1.  $r$ -lightlike submanifold.  $1 \leq r < \min\{m, n\}$ .

CASE 2. Co-isotropic submanifold.  $1 \leq r = n < m$ .

CASE 3. Isotropic submanifold.  $1 \leq r = m < n$ .

CASE 4. Totally lightlike submanifold.  $1 \leq r = m = n$ .

We refer [8] for notations and details not mentioned in this paper. For Case 1, there exists a non-degenerate screen distribution  $S(TM)$  which is a complementary vector subbundle to  $\text{Rad } TM$  in  $TM$ . Therefore,

$$(2.1) \quad TM = \text{Rad } TM \oplus S(TM).$$

Although  $S(TM)$  is not unique, it is canonically isomorphic to the factor vector bundle  $TM/\text{Rad } TM$ . Denote an  $r$ -lightlike submanifold by  $(M, g, S(TM), S(TM^\perp))$ , where  $S(TM^\perp)$  is a complementary vector subbundle to  $\text{Rad } TM$  in  $TM^\perp$ . For the dependence of all the induced geometric objects, of  $M$ , on  $\{S(TM), S(TM^\perp)\}$  we refer [8]. Let  $\text{tr}(TM)$  and  $\text{ltr}(TM)$  be complementary (but not orthogonal) vector bundles to  $TM$  in  $T\overline{M}|M$  and to  $\text{Rad } TM$  in  $S(TM^\perp)$  respectively. Then, we obtain

$$(2.2) \quad \text{tr}(TM) = \text{ltr}(TM) \oplus S(TM^\perp),$$

$$(2.3) \quad \begin{aligned} T\bar{M}|_M &= TM \oplus \text{tr}(TM) \\ &= (\text{Rad } TM \oplus \text{ltr}(TM)) \oplus S(TM) \oplus S(TM^\perp). \end{aligned}$$

Consider the following local quasi-orthonormal field of frames of  $\bar{M}$  along  $M$ :

$$(2.4) \quad \{\xi_1, \dots, \xi_r, N_1, \dots, N_r, X_{r+1}, \dots, X_m, W_{r+1}, \dots, W_n\}$$

where  $\{\xi_1, \dots, \xi_r\}$  is a lightlike basis of  $\Gamma(\text{Rad } TM)$ ,  $\{N_1, \dots, N_r\}$  a lightlike basis of  $\Gamma(\text{ltr}(TM))$ ,  $\{X_{r+1}, \dots, X_m\}$  and  $\{W_{r+1}, \dots, W_n\}$  orthonormal basis of  $\Gamma(S(TM)|\mathcal{U})$  and  $\Gamma(S(TM^\perp)|\mathcal{U})$  respectively.

*Example 1.* Consider a surface  $(M, g)$  in  $R_2^4$  given by the equations

$$x^3 = \frac{1}{\sqrt{2}}(x^1 + x^2); \quad x^4 = \frac{1}{2} \log(1 + (x^1 - x^2)^2),$$

where  $(x^1, \dots, x^4)$  is a local coordinate system for  $R_2^4$ . Using a simple procedure of linear algebra, we choose a set of vectors  $\{U, V, \xi, W\}$  given by

$$U = \sqrt{2}(1 + (x^1 - x^2)^2)\partial_1 + (1 + (x^1 - x^2)^2)\partial_3 + \sqrt{2}(x^1 - x^2)\partial_4,$$

$$V = \sqrt{2}(1 + (x^1 - x^2)^2)\partial_2 + (1 + (x^1 - x^2)^2)\partial_3 - \sqrt{2}(x^1 - x^2)\partial_4,$$

$$\xi = \partial_1 + \partial_2 + \sqrt{2}\partial_3,$$

$$W = 2(x^2 - x^1)\partial_2 + \sqrt{2}(x^2 - x^1)\partial_3 + (1 + (x^1 - x^2)^2)\partial_4,$$

so that  $TM$  and  $TM^\perp$  are spanned by  $\{U, V\}$  and  $\{\xi, W\}$  respectively. By direct calculations it follows that  $\text{Rad } TM$  is a distribution on  $M$  of rank 1 and spanned by the lightlike vector  $\xi$ . Choose  $S(TM)$  and  $S(TM^\perp)$  spanned by the timelike vector  $V$  and the spacelike vector  $W$  respectively. Then,

$$\text{ltr}(TM) = \text{Span}\left\{N = -\frac{1}{2}\partial_1 + \frac{1}{2}\partial_2 + \frac{1}{\sqrt{2}}\partial_3\right\},$$

$$\text{tr}(TM) = \text{Span}\{N, W\},$$

where  $N$  is a lightlike vector such that  $g(N, \xi) = 1$ . Thus,  $M$  is a 1-lightlike submanifold of Case 1, with basis  $\{\xi, N, V, W\}$  of  $R_2^4$  along  $M$ .

For Case 2, we have  $\text{Rad } TM = TM^\perp$ . Therefore,  $S(TM^\perp) = \{0\}$  and from (2.2)  $\text{tr}(TM) = \text{ltr}(TM)$ . Thus, (2.3) and (2.4) reduce to

$$(2.5) \quad T\bar{M}|_M = TM \oplus \text{tr}(TM) = (TM^\perp \oplus \text{ltr}(TM)) \oplus S(TM)$$

$$(2.6) \quad \{\xi_1, \dots, \xi_r, N_1, \dots, N_r, X_{r+1}, \dots, X_m\}.$$

*Example 2.* Consider the unit pseudo sphere  $S_1^3$  of Minkowski space  $R_1^4$  given by the equation  $-t^2 + x^2 + y^2 + z^2 = 1$ . Cut  $S_1^3$  by the hypersurface  $t - x = 0$  and obtain a lightlike surface  $(M, g)$  of  $S_1^3$  with  $\text{Rad } TM$  spanned by a lightlike vector  $\xi = \partial_t + \partial_x$ . Clearly,  $\text{Rad } TM = TM^\perp$  and, therefore, this example belongs to Case 2. Consider a screen distribution  $S(TM)$  spanned by a spacelike vector  $X = z\partial_y - y\partial_z$ . Then, we obtain a lightlike transversal vector bundle  $\text{tr}(TM) = \text{ltr}(TM)$  spanned by  $N = (-1/2)\{(1 + t^2)\partial_t + (t^2 - 1)\partial_x + 2ty\partial_y + 2tz\partial_z\}$  such that  $g(N, \xi) = 1$ , with a basis  $\{\xi, N, X\}$  for  $S_1^3$  along  $M$ .

For Case 3, we have  $\text{Rad } TM = TM$ . Therefore,  $S(TM) = \{0\}$ . Therefore, (2.3) and (2.4) reduce to

$$(2.7) \quad T\bar{M}|_M = TM \oplus \text{tr}(TM) = (TM \oplus \text{ltr}(TM)) \oplus S(TM^\perp)$$

$$(2.8) \quad \{\xi_1, \dots, \xi_r, N_1, \dots, N_r, W_{r+1}, \dots, W_n\}.$$

*Example 3.* Suppose  $(M, g)$  is a surface of  $R_2^5$  given by equations

$$x^3 = \cos x^1, \quad x^4 = \sin x^1, \quad x^5 = x^2.$$

We choose a set of vectors  $\{\xi_1, \xi_2, U_1, U_2\}$  given by

$$\xi_1 = \partial_2 + \partial_5, \quad \xi_2 = \partial_1 - \sin x^1 \partial_3 + \cos x^1 \partial_4,$$

$$U_1 = -\sin x^1 \partial_1 + \partial_3, \quad U_2 = \cos x^1 \partial_1 + \partial_4,$$

so that  $\text{Rad } TM = TM = \text{Span}\{\xi_1, \xi_2\}$ ,  $TM^\perp = \text{Span}\{\xi_1, U_1, U_2\}$ . Therefore,  $M$  belongs to Case 3. Construct two null vectors

$$N_1 = \frac{1}{2}\{-\partial_2 + \partial_5\},$$

$$N_2 = \frac{1}{2}\{-\partial_1 - \sin x^1 \partial_3 + \cos x^1 \partial_4\},$$

such that  $g(N_i, \xi_j) = \delta_{ij}$  for  $i, j \in \{1, 2\}$  and  $\text{ltr}(TM) = \text{Span}\{N_1, N_2\}$ . Let  $W = \cos x^1 \partial_3 + \sin x^1 \partial_4$  be a spacelike vector such that  $S(TM^\perp) = \text{Span}\{W\}$ . Thus,  $\{\xi_1, \xi_2, N_1, N_2, W\}$  is a basis of  $R_2^5$  along  $M$ .

For Case 4,  $\text{Rad } TM = TM = TM^\perp$ ,  $S(TM) = S(TM^\perp) = \{0\}$ . Therefore, (2.3) and (2.4) reduce to

$$(2.9) \quad T\bar{M}|_M = TM \oplus \text{ltr}(TM)$$

$$(2.10) \quad \{\xi_1, \dots, \xi_r, N_1, \dots, N_r\}.$$

*Example 4.* Suppose  $(M, g)$  is a surface of  $R_2^4$  given by the equations

$$x^3 = \frac{1}{\sqrt{2}}(x^1 + x^2), \quad x^4 = \frac{1}{\sqrt{2}}(x^1 - x^2).$$

We choose a set of vectors  $\{\xi_1, \xi_2, U, V\}$  given by

$$\begin{aligned}\xi_1 &= \partial_1 + \frac{1}{\sqrt{2}}\partial_3 + \frac{1}{\sqrt{2}}\partial_4, & \xi_2 &= \partial_2 + \frac{1}{\sqrt{2}}\partial_3 - \frac{1}{\sqrt{2}}\partial_4, \\ U &= \partial_1 + \partial_2 + \sqrt{2}\partial_3, & V &= \partial_1 - \partial_2 + \sqrt{2}\partial_4,\end{aligned}$$

so that  $TM$  and  $TM^\perp$  are spanned by  $\{\xi_1, \xi_2\}$  and  $\{U, V\}$  respectively. By direct calculations we check that  $\text{Span}\{\xi_1, \xi_2\} = \text{Span}\{U, V\}$ , that is,  $TM = TM^\perp$ . Finally, the two lightlike transversal vector bundles are:

$$N_1 = \partial_1 + \sqrt{2}\partial_3 + \sqrt{2}\partial_4, \quad N_2 = \partial_2 + \sqrt{2}\partial_3 - \sqrt{2}\partial_4,$$

such that  $g(N_i, \xi_j) = \delta_{ij}$ ,  $i, j = 1, 2$ . Thus,  $M$  is of Case 4, with a basis  $\{\xi_1, \xi_2, N_1, N_2\}$  of  $R^4$  along  $M$ .

On the existence of a local quasi-orthonormal field of frames of  $\bar{M}$  along  $M$  we state (see Chapter 5 of [8] for its proof) the following main result:

**THEOREM 2.1** [8]. *Let  $(M, g, S(TM), S(TM^\perp))$  be an  $r$ -lightlike submanifold of a semi-Riemannian manifold  $(\bar{M}, \bar{g})$ . Then there exists a complementary vector bundle  $\text{ltr}(TM)$  of  $\text{Rad } TM$  in  $S(TM^\perp)^\perp$  and a basis of  $\Gamma(\text{ltr}(TM)|_{\mathcal{U}})$  consisting of smooth sections  $\{N_i\}$  of  $S(TM^\perp)^\perp|_{\mathcal{U}}$ , where  $\mathcal{U}$  is a coordinate neighborhood of  $M$ , such that*

$$(2.11) \quad \bar{g}(N_i, \xi_j) = \delta_{ij}, \quad \bar{g}(N_i, N_j) = 0,$$

where  $\{\xi_1, \dots, \xi_r\}$  is a lightlike basis of  $\Gamma(\text{Rad } TM)$ .

Define locally  $r$  differential 1-forms  $\{\eta_i\}$  on  $\Gamma(TM)$  by

$$(2.12) \quad \eta_i(X) = \bar{g}(X, N_i), \quad \forall X \in \Gamma(TM).$$

Let  $P$  the projection of  $TM$  on  $S(TM)$  with respect to (2.1). Then,

$$(2.13) \quad X = PX + \sum_{i=1}^r \eta_i(X)\xi_i,$$

for every  $X \in \Gamma(TM)$ . According to (2.3) we put

$$(2.14) \quad \bar{\nabla}_X Y = \nabla_X Y + h(X, Y),$$

$$(2.15) \quad \bar{\nabla}_X V = -A(V, X) + \nabla_X^\perp V, \quad \forall X, Y \in \Gamma(TM),$$

$V \in \Gamma(\text{tr}(TM))$ ,  $\{\nabla_X Y, A(V, X)\}$  and  $\{h(X, Y), \nabla_X^\perp V\}$  belong to  $\Gamma(TM)$  and  $\Gamma(\text{tr}(TM))$  respectively. Here  $\bar{\nabla}$  is the metric connection on  $\bar{M}$  but  $\nabla$  (torsion-free) and  $\nabla^\perp$  are linear connections on  $M$  and  $\text{tr}(TM)$  respectively.

Suppose  $S(TM^\perp) \neq \{0\}$ , that is,  $M$  is either an  $r$ -lightlike or a isotropic submanifold of  $\bar{M}$ . According to (2.3) we consider the projection morphisms  $L$  and  $S$  of  $\text{tr}(TM)$  on  $\text{ltr}(TM)$  and  $S(TM^\perp)$  respectively. Then (2.14) and (2.15) become

$$(2.16) \quad \bar{\nabla}_X Y = \nabla_X Y + h^\ell(X, Y) + h^s(X, Y),$$

$$(2.17) \quad \bar{\nabla}_X V = -A_V X + D_X^\ell V + D_X^s V,$$

where we put

$$h^\ell(X, Y) = L(h(X, Y)); \quad h^s(X, Y) = S(h(X, Y)); \quad A_V X = A(V, X), \\ D_X^\ell V = L(\nabla_X^\perp V) = D^\ell(X, V); \quad D_X^s V = S(\nabla_X^\perp V) = D^s(X, V).$$

As  $h^\ell$  and  $h^s$  are  $\Gamma(\text{ltr}(TM))$ -valued and  $\Gamma(S(TM^\perp))$ -valued respectively, we call them the lightlike second fundamental form and the screen second fundamental form of  $M$ . In particular, we derive

$$(2.18) \quad \bar{\nabla}_X N = -A_N X + \nabla_X^\ell N + D^s(X, N),$$

$$(2.19) \quad \bar{\nabla}_X W = -A_W X + \nabla_X^s W + D^\ell(X, W),$$

for any  $X \in \Gamma(TM)$ ,  $N \in \Gamma(\text{ltr}(TM))$  and  $W \in \Gamma(S(TM^\perp))$ .

Next, suppose  $S(TM^\perp) = \{0\}$ , that is,  $M$  is either co-isotropic or totally lightlike. Then, (2.16) and (2.17) become

$$(2.20) \quad \bar{\nabla}_X Y = \nabla_X Y + h^\ell(X, Y),$$

$$(2.21) \quad \bar{\nabla}_X N = -A_N X + \nabla_X^\ell N,$$

for any  $X, Y \in \Gamma(TM)$ . We call (2.14), (2.16), (2.20) the Gauss formulae and (2.15), (2.17)–(2.21) the Weingarten formulae for all cases of a lightlike submanifold  $M$ . Using (2.16)–(2.21), (2.3), (2.5), (2.7) and (2.9), we obtain

$$(2.22) \quad \bar{g}(h^s(X, Y), W) + \bar{g}(Y, D^\ell(X, W)) = g(A_W X, Y),$$

$$(2.23) \quad \bar{g}(h^\ell(X, Y), \xi) + \bar{g}(Y, h^\ell(X, \xi)) + g(Y, \nabla_X \xi) = 0,$$

$$(2.24) \quad \bar{g}(A_N X, N') + \bar{g}(A_{N'} X, N) = 0,$$

for any  $\xi \in \Gamma(\text{Rad } TM)$ ,  $W \in \Gamma(S(TM^\perp))$  and  $N, N' \in \Gamma(\text{ltr}(TM))$ .

Next, suppose  $S(TM) \neq \{0\}$ , that is,  $M$  is either  $r$ -lightlike or co-isotropic. Then according to (2.1) we set

$$(2.25) \quad \nabla_X P Y = \nabla_X^* P Y + h^*(X, P Y),$$

$$(2.26) \quad \nabla_X \xi = -A^*(\xi, X) + \nabla_X^{*t} \xi,$$

for any  $X, Y \in \Gamma(TM)$  and  $\xi \in \Gamma(\text{Rad } TM)$ , where  $\{\nabla_X^* P Y, A^*(\xi, X)\}$  and  $\{h^*(X, P Y), \nabla_X^{*t} \xi\}$  belong to  $\Gamma(S(TM))$  and  $\Gamma(\text{Rad } TM)$  respectively. It follows that  $\nabla^*$  and  $\nabla^{*t}$  are linear connections on  $S(TM)$  and  $\text{Rad } TM$  respectively. By using (2.16), (2.21), (2.25) and (2.26) we obtain

$$(2.27) \quad \bar{g}(h^\ell(X, P Y), \xi) = \bar{g}(A_\xi^* X, P Y)$$

$$(2.28) \quad \bar{g}(h^*(X, P Y), N) = \bar{g}(A_N X, P Y), \quad \forall X, Y \in \Gamma(TM)$$

**THEOREM 2.2** [8]. *Let  $(M, g, S(TM), S(TM^\perp))$  be an  $r$ -lightlike submanifold*

or a co-isotropic submanifold of a semi-Riemannian manifold  $(\bar{M}, \bar{g})$ . Then the following assertions are equivalent:

- (1)  $S(TM)$  is integrable.
- (2)  $h^*$  is symmetric on  $\Gamma(S(TM))$ .
- (3)  $A_N$  is self-adjoint on  $\Gamma(S(TM))$  with respect to  $g$ .
- (4)  $\nabla^*$  is torsion-free linear connection.

*Example 5.* Let  $(R_1^{d+1}, \bar{g})$  be a Minkowski spacetime, where

$$\bar{g}(x, y) = -x^0 y^0 + \sum_{i=1}^d x^i y^i, \quad \forall x, y \in R^{d+1}.$$

Consider a smooth function  $f : D \rightarrow R$ , where  $D$  is an open set of  $R^d$ . Then

$$M = \{(x^0, \dots, x^n) \in R_1^{d+1}; x^0 = f(x^1, \dots, x^d)\},$$

is a hypersurface of  $R_1^{d+1}$  which is called a *Monge hypersurface*. Let natural parameterization on  $M$  be given by

$$x^0 = f(v^0, \dots, v^{d-1}); \quad x^{\alpha+1} = v^\alpha, \quad \alpha \in \{0, \dots, n-1\}.$$

Hence, the natural frames field on  $M$  is globally defined by

$$\partial_{v^\alpha} = f'_{x^{\alpha+1}} \partial_{x^0} + \partial_{x^{\alpha+1}}, \quad \alpha \in \{0, \dots, d-1\}.$$

Then, it follows that  $TM^\perp$  is spanned by a global vector

$$(2.29) \quad \xi = \partial_{x^0} + \sum_{i=1}^d f'_{x^i} \partial_{x^i}.$$

It is known [8] that  $M$  is a lightlike hypersurface if  $TM^\perp = \text{Rad } TM$ . This means that  $\xi$ , given by (2.29), must be a null vector field. Hence, there exists a lightlike Monge hypersurface  $M$ , if the function  $f$  is a solution of the differential equation  $\sum_{i=1}^d (f'_{x^i})^2 = 1$ . The null transversal vector is given by  $N = (1/2)\{-\partial_{x^0} + \sum_{i=1}^d f'_{x^i} \partial_{x^i}\}$ ,  $\bar{g}(N, \xi) = 1$ . Let  $\bar{\nabla}$  be the Levi-Civita connection, with respect to the metric  $\bar{g}$ , on  $R_1^{d+1}$ . Then, for any two vectors  $X, Y \in \Gamma(S(TM))$ , the Lie bracket  $[X, Y] \in \Gamma(S(TM))$ . Indeed,

$$\begin{aligned} \bar{g}([X, Y], N) &= \bar{g}(\bar{\nabla}_X Y - \bar{\nabla}_Y X, \partial_{x^0}) \\ &= -\{\bar{g}(X, \bar{\nabla}_Y \partial_{x^0}) - \bar{g}(Y, \bar{\nabla}_X \partial_{x^0})\} = 0. \end{aligned}$$

Hence,  $S(TM)$  is integrable. Other equivalent assertions follow easily.

### 3. Structure equations

Let  $(M, g, S(TM), S(TM^\perp))$  be an  $m$ -dimensional  $r$ -lightlike submanifold of  $(m+n)$ -dimensional semi-Riemannian manifold  $(\bar{M}, \bar{g})$ . Denote by  $\bar{R}, R$  and  $R^\ell$  the curvature tensors of  $\bar{\nabla}, \nabla$  and  $\nabla^\ell$  respectively. We need following structure equations (see [8] for details on a complete set of equations):

$$\begin{aligned}
(3.1) \quad \bar{R}(X, Y)Z &= R(X, Y)Z \\
&+ A_{h^\ell(X, Z)}Y - A_{h^\ell(Y, Z)}X \\
&+ A_{h^s(X, Z)}Y - A_{h^s(Y, Z)}X \\
&+ (\nabla_X h^\ell)(Y, Z) - (\nabla_Y h^\ell)(X, Z) \\
&+ D^\ell(X, h^s(Y, Z)) - D^\ell(Y, h^s(X, Z)) \\
&+ (\nabla_X h^s)(Y, Z) - (\nabla_Y h^s)(X, Z) \\
&+ D^s(X, h^\ell(Y, Z)) - D^s(Y, h^\ell(X, Z)),
\end{aligned}$$

for any  $X, Y, Z \in \Gamma(TM)$ . Consider the curvature tensor  $\bar{R}$  of type  $(0, 4)$ .

$$\begin{aligned}
(3.2) \quad \bar{R}(X, Y, PZ, PU) &= g(R(X, Y)PZ, PU) \\
&+ \bar{g}(h^*(Y, PU), h^\ell(X, PZ)) - \bar{g}(h^*(X, PU), h^\ell(Y, PZ)) \\
&+ \bar{g}(h^s(Y, PU), h^s(X, PZ)) - \bar{g}(h^s(X, PU), h^s(Y, PZ)),
\end{aligned}$$

$$\begin{aligned}
(3.3) \quad \bar{R}(X, Y, \xi, PU) &= g(R(X, Y)\xi, PU) \\
&+ \bar{g}(h^*(Y, PU), h^\ell(X, \xi)) - \bar{g}(h^*(X, PU), h^\ell(Y, \xi)) \\
&+ \bar{g}(h^s(Y, PU), h^s(X, \xi)) - \bar{g}(h^s(X, PU), h^s(Y, \xi)) \\
&= \bar{g}((\nabla_Y h^\ell)(X, PU) - (\nabla_X h^\ell)(Y, PU), \xi) \\
&+ \bar{g}(h^s(Y, PU), h^s(X, \xi)) - \bar{g}(h^s(X, PU), h^s(Y, \xi)),
\end{aligned}$$

$$\begin{aligned}
(3.4) \quad \bar{R}(X, Y, N, PU) &= -\bar{g}(R(X, Y)PU, N) \\
&+ \bar{g}(A_N Y, h^\ell(X, PU)) - \bar{g}(A_N X, h^\ell(Y, PU)) \\
&+ \bar{g}(h^s(Y, PU), D^s(X, N)) - \bar{g}(h^s(X, PU), D^s(Y, N)) \\
&= \bar{g}((\nabla_Y A)(N, X) - (\nabla_X A)(N, Y), PU) \\
&+ \bar{g}(h^s(Y, PU), D^s(X, N)) - \bar{g}(h^s(X, PU), D^s(Y, N)),
\end{aligned}$$

$$\begin{aligned}
(3.5) \quad \bar{R}(X, Y, W, PU) &= \bar{g}((\nabla_Y A)(W, X) - (\nabla_X A)(W, Y), PU) \\
&+ \bar{g}(h^*(Y, PU), D^\ell(X, W)) - \bar{g}(h^*(X, PU), D^\ell(Y, W)) \\
&= \bar{g}((\nabla_Y h^s)(X, PU) - (\nabla_X h^s)(Y, PU), W) \\
&+ \bar{g}(h^\ell(X, PU), A_W Y) - \bar{g}(h^\ell(X, PU), A_W X),
\end{aligned}$$

$$\begin{aligned}
(3.6) \quad \bar{R}(X, Y, N, \xi) &= \bar{g}(R^\ell(X, Y)N, \xi) \\
&+ \bar{g}(h^\ell(Y, A_N X), \xi) - \bar{g}(h^\ell(X, A_N Y), \xi) \\
&+ \bar{g}(D^s(X, N), h^s(Y, \xi)) - \bar{g}(D^s(Y, N), h^s(X, \xi))
\end{aligned}$$

$$\begin{aligned}
 &= -\bar{g}(R(X, Y)\xi, N) \\
 &\quad + \bar{g}(A_N Y, h^\ell(X, \xi)) - (A_N Y, h^\ell(Y, \xi)) \\
 &\quad + \bar{g}(D^s(X, N), h^s(Y, \xi)) - \bar{g}(D^s(Y, N), h^s(X, \xi)),
 \end{aligned}$$

$X, Y, U \in \Gamma(TM)$ . Let  $R^{*t}$  be the curvature tensor of  $\nabla^{*t}$ . Then,

$$(3.7) \quad g(R(X, Y)\xi, PU) = g((\nabla_Y A^*)(\xi, X) - (\nabla_X A^*)(\xi, Y), PU),$$

$$(3.8) \quad g(R(X, Y)\xi, N = \bar{g}(R^{*t}(X, Y)\xi, N) \\ + g(A_N Y, A_\xi^* X) - g(A_N X, A_\xi^* Y).$$

$$(3.9) \quad g(R(X, Y)PU, N) = \bar{g}((\nabla_X A)(N, Y) - (\nabla_Y A)(N, X), PU) \\ + \bar{g}(h^\ell(X, PU), A_N Y) - \bar{g}(h^\ell(Y, PU), A_N X) \\ = \bar{g}((\nabla_X h^*)(Y, PU) - (\nabla_Y h^*)(X, PU), N).$$

Finally, from (3.6), by using (2.23) and (2.25) we deduce

$$(3.10) \quad \bar{g}(R(X, Y)\xi, N) + \bar{g}(R^\ell(X, Y)N, \xi) = g(A_\xi^* X, A_N Y) \\ - g(A_\xi^* Y, A_N X).$$

*Remark 1.* For structure equations of Case 2, delete all the components involving  $S(TM^\perp)$ . Similarly, one can find the structure equations of the other two cases.

*Remark 2.* In the sequel we denote by  $(M, g)$  a lightlike submanifold for which the results hold for all its four cases. Any result which does not hold for all the cases will be so specified.

#### 4. Totally umbilical lightlike submanifold

Let  $\{N_i, W_\alpha\}$  be a basis of  $\Gamma(\text{tr}(TM)|_{\mathcal{U}})$  on a coordinate neighborhood  $\mathcal{U}$  of  $M$ , where  $N_i \in \Gamma(\text{ltr}(TM)|_{\mathcal{U}})$  and  $W_\alpha \in \Gamma(S(TM^\perp)|_{\mathcal{U}})$ . Then (2.16) becomes

$$(4.1) \quad \bar{\nabla}_X Y = \nabla_X Y + \sum_{i=1}^r h_i^\ell(X, Y)N_i + \sum_{\alpha=r+1}^n h_\alpha^s(X, Y)W_\alpha,$$

$$(4.2) \quad \bar{\nabla}_X Y = \nabla_X Y + \sum_{i=1}^{m<n} h_i^\ell(X, Y)N_i + \sum_{\alpha=m+1}^n h_\alpha^s(X, Y)W_\alpha,$$

for an  $r$ -lightlike or an isotropic submanifold respectively. (2.20) becomes

$$(4.3) \quad \bar{\nabla}_X Y = \nabla_X Y + \sum_{i=1}^{n<m} h_i^\ell(X, Y)N_i,$$

$$(4.4) \quad \bar{\nabla}_X Y = \nabla_X Y + \sum_{i=1}^{n=m} h_i^\ell(X, Y)N_i,$$

for a co-isotropic and a totally lightlike submanifold respectively. We call  $\{h'_i\}$  and  $\{h''_i\}$  the local lightlike second fundamental forms and the local screen second fundamental forms of  $M$  on  $\mathcal{U}$ . Also (2.18) and (2.19) become

$$\begin{aligned}
\bar{\nabla}_X N_i &= -A_{N_i} X + \sum_{j=1}^r \rho_{ij}(X) N_j + \sum_{\alpha=r+1}^n \tau_{i\alpha}(X) W_\alpha, \\
\bar{\nabla}_X W_\alpha &= -A_{W_\alpha} X + \sum_{i=1}^r v_{\alpha i}(X) N_i + \sum_{\beta=r+1}^n \theta_{\alpha\beta}(X) W_\beta, \\
\bar{\nabla}_X N_i &= -A_{N_i} X + \sum_{j=1}^{m<n} \rho_{ij}(X) N_j + \sum_{\alpha=m+1}^n \tau_{i\alpha}(X) W_\alpha, \\
\bar{\nabla}_X W_\alpha &= -A_{W_\alpha} X + \sum_{i=1}^{m<n} v_{\alpha i}(X) N_i + \sum_{\beta=m+1}^n \theta_{\alpha\beta}(X) W_\beta,
\end{aligned}
\tag{4.5}$$

for an  $r$ -lightlike and an isotropic submanifold respectively, where

$$\begin{aligned}
\rho_{ij}(X) &= \bar{g}(\nabla_X^\ell N_i, \zeta_j), \quad \varepsilon_\alpha \tau_{i\alpha}(X) = \bar{g}(D^s(X, N_i), W_\alpha), \\
v_{\alpha i}(X) &= \bar{g}(D^\ell(X, W_\alpha), \zeta_i), \quad \varepsilon_\beta \theta_{\alpha\beta}(X) = \bar{g}(\nabla_X^s W_\alpha, W_\beta),
\end{aligned}
\tag{4.6}$$

and  $\varepsilon_\alpha$  is the signature of  $W_\alpha$ . Similarly, (2.21) becomes

$$\begin{aligned}
\bar{\nabla}_X N_i &= -A_{N_i} X + \sum_{j=1}^r \rho_{ij}(X) N_j, \\
\bar{\nabla}_X N_i &= -A_{N_i} X + \sum_{j=1}^{m<n} \rho_{ij}(X) N_j,
\end{aligned}
\tag{4.7}$$

for a co-isotropic and a totally lightlike submanifold respectively. Then, (2.25) and (2.26) become

$$\begin{aligned}
\nabla_X P Y &= \nabla_X^* P Y + \sum_{i=1}^r h_i^*(X, P Y) \zeta_i, \\
\nabla_X \zeta_i &= -A_{\zeta_i}^* X + \sum_{j=1}^r \mu_{ij}(X) \zeta_j,
\end{aligned}
\tag{4.8}$$

where  $h_i^*(X, P Y) = \bar{g}(h^*(X, P Y), N_i)$  and  $\mu_{ij}(X) = \bar{g}(\nabla_X^{*l} \zeta_i, N_j)$ . Using the equations (2.11) and (4.5)–(4.8) we obtain  $\mu_{ij}(X) = -\rho_{ji}(X)$ . Thus,

$$\nabla_X \zeta_i = -A_{\zeta_i}^* X - \sum_{j=1}^r \rho_{ji}(X) \zeta_j.
\tag{4.9}$$

**DEFINITION 1.** A lightlike submanifold  $(M, g)$  of a semi-Riemannian manifold  $(\bar{M}, \bar{g})$  is said to be totally umbilical in  $\bar{M}$  if there is a smooth transversal

vector field  $\mathcal{H} \in \Gamma(\text{tr}(TM))$  on  $M$ , called the transversal curvature vector field of  $M$ , such that, for all  $X, Y \in \Gamma(TM)$ ,

$$(4.10) \quad h(X, Y) = \mathcal{H}\bar{g}(X, Y)$$

Using (2.16) and (4.1) it is easy to see that  $M$  is totally umbilical, if and only if on each coordinate neighborhood  $\mathcal{U}$  there exist smooth vector fields  $H^\ell \in \Gamma(\text{ltr}(TM))$  and  $H^s \in \Gamma(S(TM^\perp))$ , and smooth functions  $H_i^\ell \in F(\text{ltr}(TM))$  and  $H_i^s \in F(S(TM^\perp))$  such that

$$(4.11) \quad \begin{aligned} h^\ell(X, Y) &= H^\ell \bar{g}(X, Y), & h^s(X, Y) &= H^s \bar{g}(X, Y) \\ h_i^\ell(X, Y) &= H_i^\ell \bar{g}(X, Y), & h_i^s(X, Y) &= H_i^s \bar{g}(X, Y) \end{aligned}$$

for any  $X, Y \in \Gamma(TM)$ . Above definition does not depend on the screen distribution and the screen transversal vector bundle of  $M$ . On the other hand, from the equation (2.22) we obtain the following equation

$$(4.12) \quad g(A_{W_\alpha} X, Y) = \varepsilon_\alpha h_\alpha^s(X, Y) + \sum_{i=1}^r D_i^\ell(X, W_\alpha) \eta_i(Y).$$

Now replace  $Y$  by  $\xi_j$  and obtain

$$(4.13) \quad D_i^\ell(X, W_\alpha) = -\varepsilon_\alpha h_\alpha^s(\xi_i, X).$$

Using (2.22), (2.27), (4.11) and (4.13), we conclude (the relations (4.11) trivially hold in case  $S(TM)$  and or  $S(TM^\perp)$  vanish)

**THEOREM 4.1.** *Let  $(M, g)$  be a lightlike submanifold of  $(\bar{M}, \bar{g})$ . Then  $M$  is totally umbilical, if and only if, on each coordinate neighborhood  $\mathcal{U}$  there exist smooth vector fields  $H^\ell$  and  $H^s$  such that*

$$(4.14) \quad \begin{aligned} D^\ell(X, W) &= 0, & A_{\xi}^* X &= H^\ell P X, & P(A_W X) &= \varepsilon H^s P X, \\ D_i^\ell(X, W_\alpha) &= 0, & A_{\xi_i}^* X &= H_i^\ell P X, & P(A_{W_\alpha} X) &= \varepsilon_\alpha H_\alpha^s P X, \end{aligned}$$

for any  $X \in \Gamma(TM)$ , where  $\varepsilon$  is the signature of  $W \in \Gamma(S(TM^\perp))$ .

*Example 6.* Let  $M$  be a surface of  $R_2^4$ , of Example 1, given by

$$x^3 = \frac{1}{\sqrt{2}}(x^1 + x^2); \quad x^4 = \frac{1}{2} \log(1 + (x^1 - x^2)^2),$$

where  $(x^1, \dots, x^4)$  is a local coordinate system for  $R_2^4$ . As explained in Example 1,  $M$  is a 1-lightlike surface of Case 1 having a local quasi-orthonormal field of frames  $\{\xi, N, V, W\}$  along  $M$ . Denote by  $\bar{\nabla}$  the Levi-Civita connection on  $R_2^4$ . Then, by straightforward calculations, we obtain

$$\begin{aligned} \bar{\nabla}_V V &= 2(1 + (x^1 - x^2)^2)\{2(x^2 - x^1)\partial_2 + \sqrt{2}(x^2 - x^1)\partial_3 + \partial_4\}, \\ \bar{\nabla}_{\xi_1} V &= 0, \quad \bar{\nabla}_X \xi_1 = \bar{\nabla}_X N = 0, \quad \forall X \in \Gamma(TM). \end{aligned}$$

For this example, the equations (4.11) reduce to

$$h^1(X, Y) = H^1 \bar{g}(X, Y); \quad h^2(X, Y) = H^2 \bar{g}(X, Y)$$

where  $h^1$  and  $h^2$  are  $\Gamma(\text{ltr}(TM))$ -valued and  $\Gamma(S(TM^\perp))$ -valued bilinear forms (see equation (2.16)). Using the Gauss and Weingarten formulae we infer

$$h^1 = 0; \quad A_{\xi_1} = 0; \quad A_N = 0; \quad \nabla_X \xi_1 = 0; \quad \rho_i(X) = 0;$$

where for the symbol  $\rho_i$  see the equation (4.7).  $h^2(X, \xi) = 0$ ;

$$H^2(V, V) = 2; \quad \nabla_X V = \frac{2\sqrt{2}(x^2 - x^1)^3}{1 + (x^1 - x^2)^2} X^2 V,$$

$\forall X = X^1 \xi_1 + X^2 V \in \Gamma(TM)$ . Since  $\bar{g}(V, V) = -(1 + (x^1 - x^2)^4)$  we get

$$h^2(V, V) = H^2 \bar{g}(V, V), \quad H^2 = -\frac{2}{(1 + (x^1 - x^2)^4)}.$$

Therefore,  $M$  is totally umbilical 1-lightlike submanifold of  $R_2^4$ .

Note that in case  $M$  is totally umbilical, then due to (2.27)

$$(4.15) \quad h^\ell(X, \xi) = 0, \quad h^s(X, \xi) = 0, \quad A_\xi^* \xi' = 0, \quad A_W \xi = 0.$$

**THEOREM 4.2.** *Let  $(M, g)$  be an  $m$ -dimensional totally umbilical lightlike submanifold of an  $(m+n)$ -dimensional semi-Riemannian manifold of constant curvature  $(\bar{M}(\bar{c}), \bar{g})$ . Then, the functions  $H_i^\ell, H_\alpha^s$  from (4.11) satisfy the following partial differential equations*

$$(4.16) \quad \begin{aligned} \xi_j(H_i^\ell) - H_i^\ell H_j^\ell + \sum_{k=1}^r H_k^\ell \rho_{ki}(\xi_j) &= 0, \\ \xi_j(H_\alpha^s) - H_\alpha^s H_j^\ell + \sum_{i=1}^r H_i^\ell \tau_{i\alpha}(\xi_j) + \sum_{\beta=r+1}^n H_\beta^s \theta_{\beta\alpha}(\xi_j) &= 0, \\ R(X, Y)Z &= \left\{ \bar{c}X + \sum_{i=1}^r H_i^\ell A_{N_i} X + \sum_{\alpha=r+1}^n H_\alpha^s A_{W_\alpha} X \right\} g(Y, Z) \\ &\quad - \left\{ \bar{c}Y + \sum_{i=1}^r H_i^\ell A_{N_i} Y + \sum_{\alpha=r+1}^n H_\alpha^s A_{W_\alpha} Y \right\} g(X, Z), \end{aligned}$$

for any  $X, Y \in \Gamma(TM)$ . Moreover,

$$(4.17) \quad \begin{aligned} PX(H_k^\ell) + \sum_{i=1}^r H_i^\ell \rho_{ik}(PX) &= 0, \\ PX(H_\alpha^s) + \sum_{i=1}^r H_i^\ell \tau_{i\alpha}(PX) + \sum_{\alpha=r+1}^n H_\beta^s \theta_{\beta\alpha}(PX) &= 0. \end{aligned}$$

*Proof.* Taking account of (4.9) in (3.3) and (3.5), and using the fact that  $\bar{M}$  is a space of constant curvature we obtain

$$\begin{aligned}
 & \left\{ X(H_k^\ell) - H_k^\ell \sum_{i=1}^r H_i^\ell \eta_i(X) + \sum_{i=1}^r H_i^\ell \rho_{ik}(X) \right\} g(Y, PU) \\
 & - \left\{ Y(H_k^\ell) - H_k^\ell \sum_{i=1}^r H_i^\ell \eta_i(Y) + \sum_{i=1}^r H_i^\ell \rho_{ik}(Y) \right\} g(X, PU) = 0, \\
 (4.18) \quad & \left\{ X(H_\alpha^s) - H_\alpha^s \sum_{i=1}^r H_i^s \eta_i(X) + \sum_{i=1}^r H_i^s \tau_{ix}(X) + \sum_{\beta=r+1}^n H_\beta^s \theta_{\beta\alpha}(X) \right\} g(Y, PU) \\
 & - \left\{ Y(H_\alpha^s) - H_\alpha^s \sum_{i=1}^r H_i^s \eta_i(Y) + \sum_{i=1}^r H_i^s \tau_{ix}(Y) \right. \\
 & \quad \left. + \sum_{\beta=r+1}^n H_\beta^s \theta_{\beta\alpha}(Y) \right\} g(X, PU) = 0,
 \end{aligned}$$

for any  $X, Y, U \in \Gamma(TM)$ . Take  $X = \zeta_j$  and  $U = Y \in \Gamma(S(TM))$  such that  $g(Y, Y) \neq 0$  on  $\mathcal{U}$  and using (2.12) we obtain (4.16). Then, (4.17) follows from (3.1), (4.18),  $\bar{M}$  a space of constant curvature and (4.16). Setting  $X = PX$  and  $Y = PY$  in (4.18) and using (2.12) we obtain

$$\begin{aligned}
 & \left\{ PX(H_k^\ell) + \sum_{i=1}^r H_i^\ell \rho_{ik}(PX) \right\} PY \\
 & = \left\{ PY(H_k^\ell) + \sum_{i=1}^r H_i^\ell \rho_{ik}(PY) \right\} PX, \\
 & \left\{ PX(H_\alpha^s) + \sum_{i=1}^r H_i^s \tau_{ix}(PX) + \sum_{\alpha=r+1}^n H_\beta^s \theta_{\beta\alpha}(PX) \right\} PY \\
 & = \left\{ PY(H_\alpha^s) + \sum_{i=1}^r H_i^s \tau_{ix}(PY) + \sum_{\alpha=r+1}^n H_\beta^s \theta_{\beta\alpha}(PY) \right\} PX,
 \end{aligned}$$

Now suppose there exists a vector field  $X_o \in \Gamma(TM)$  such that

$$\begin{aligned}
 & PX_o(H_k^\ell) + \sum_{i=1}^r H_i^\ell \rho_{ik}(PX_o) \neq 0, \\
 & PX_o(H_\alpha^s) + \sum_{i=1}^r H_i^s \tau_{ix}(PX_o) + \sum_{\alpha=r+1}^n H_\beta^s \theta_{\beta\alpha}(PX_o) \neq 0
 \end{aligned}$$

at each point  $u \in M$ . Then from the last equations it follows that all vectors

from the fiber  $(S(TM))_u$  are collinear with  $(PX_o)_u$ . This is a contradiction as  $\dim((S(TM))_u) = n - r$ . In particular, if  $r = n$ , that is, if  $S(TM)$  vanishes, then also we have a trivial contradiction. Hence the equations (4.18) in theorem are true at any point of  $\mathcal{U}$ , which completes the proof.

From (3.6), (3.8), (3.10) and  $\bar{M}$  of constant curvature we get

$$2d(\text{Tr}(\rho_{ij}))(X, Y) + \sum_{i=1}^r H_i^\ell \{g(Y, A_{N_i}X) - g(X, A_{N_i}Y)\} = 0$$

where  $\text{Tr}(\rho_{ij})$  is the trace of the matrix  $(\rho_{ij})$ . If  $(M, g)$  is an isotropic or a totally light submanifold, then, we have  $g(Y, A_{N_i}X) = g(X, A_{N_i}Y) = 0$  for every  $X, Y \in \Gamma(TM)$ . Thus, the following holds:

**LEMMA 1.** *Let  $(M, g)$  be an isotropic or a totally lightlike submanifold of a semi-Riemannian manifold  $(\bar{M}(\bar{c}), \bar{g})$  of constant curvature. Then, the trace of each  $\rho_{ij}$ , defined by (4.6), is closed, i.e.,  $d(\text{Tr}(\rho_{ij})) = 0$ .*

In case  $H_i^\ell \neq 0$  and  $H_\alpha^s \neq 0$  on  $\mathcal{U}$  we say that  $M$  is proper totally umbilical. From Theorem 2.2 and the last equation we obtain

**THEOREM 4.3.** *Let  $(M, g, S(TM))$  be a proper totally umbilical  $r$ -lightlike or a co-isotropic submanifold of a semi-Riemannian manifold  $(\bar{M}(\bar{c}), \bar{g})$  of constant curvature  $\bar{c}$ . Then  $S(TM)$  is integrable, if and only if, each 1-form  $\text{Tr}(\rho_{ij})$  induced by  $S(TM)$  is closed, i.e.,  $d(\text{Tr}(\rho_{ij})) = 0$ .*

*Remark 3.* In view of Lemma 1,  $d(\text{Tr}(\rho_{ij})) = 0$  trivially holds for a proper totally umbilical isotropic or a totally lightlike submanifold  $(M, g)$ .

**DEFINITION 2.** Let  $(M, g, S(TM))$  be either an  $r$ -lightlike or a co-isotropic submanifold of a semi-Riemannian manifold  $(\bar{M}, \bar{g})$ . Then, the screen distribution  $S(TM)$  is said to be totally umbilical in  $M$  if there is a smooth vector field  $\mathcal{K} \in \Gamma(\text{Rad } TM)$  on  $M$ , such that

$$h^*(X, PY) = \mathcal{K}g(X, PY) \quad \forall X, Y \in \Gamma(TM).$$

$S(TM)$  is totally umbilical, if and only if, on any coordinate neighborhood  $\mathcal{U} \subset M$ , there exist smooth functions  $K_i$  such that

$$(4.19) \quad h_i^*(X, PY) = K_i g(X, PY) \quad \forall X, Y \in \Gamma(TM).$$

It follows that  $h^*$  is symmetric on  $\Gamma(S(TM))$  and hence from Theorem 2.2,  $S(TM)$  is integrable. In case  $\mathcal{K} = 0$  ( $\mathcal{K} \neq 0$ ) on  $\mathcal{U}$  we say that  $S(TM)$  is totally geodesic (proper totally umbilical). (2.13) and (4.11) imply

$$(4.20) \quad P(A_{N_i}X) = K_i PX, \quad h^*(\xi, PX) = 0, \quad \forall X \in \Gamma(TM).$$

In case  $S(TM)$  is totally umbilical, we have from (2.1), (2.24) and (4.20)

$$(4.21) \quad A_{N_i}X = K_iPX + \sum_{j=1}^r \eta_j(A_{N_i}X)\zeta_j, \quad \eta_i(A_{N_j}X) = -\eta_j(A_{N_i}X).$$

**THEOREM 4.4.** *Let  $(M, g, S(TM))$  be either an  $r$ -lightlike or a co-isotropic submanifold of a semi-Riemannian manifold  $(\bar{M}(\bar{c}), \bar{g})$  of constant curvature  $\bar{c}$ , with a totally umbilical screen distribution  $S(TM)$ . If  $M$  is also totally umbilical, then, the mean curvature vectors  $K_i$  of  $S(TM)$  are a solution of the following partial differential equations*

$$\begin{aligned} X(K_i) - \sum_{j=1}^r K_j \rho_{ji}(X) - K_i \sum_{j=1}^r H_j^\ell \eta_j(X) \\ + \sum_{j=1}^r H_j^\ell \eta_i(A_{N_j}X) + \sum_{\alpha=r+1}^n H_\alpha^s \eta_i(A_{W_\alpha X}) - \bar{c} \eta_i(X) = 0. \end{aligned}$$

*Proof.* Taking account of (4.19) and (4.21) into (3.4) and using (2.12), (2.22), (2.24) and  $\bar{M}$  a space of constant curvature we obtain

$$\begin{aligned} & \left\{ X(K_i) - \sum_{j=1}^r K_j \rho_{ji}(X) - K_i \sum_{j=1}^r H_j^\ell \eta_j(X) + \sum_{j=1}^r H_j^\ell \eta_i(A_{N_j}X) \right. \\ & \left. + \sum_{\alpha=r+1}^n H_\alpha^s \eta_i(A_{W_\alpha X}) - \bar{c} \eta_i(X) \right\} g(Y, PU) \\ & = \left\{ Y(K_i) - \sum_{j=1}^r K_j \rho_{ji}(Y) - K_i \sum_{j=1}^r H_j^\ell \eta_j(Y) + \sum_{j=1}^r H_j^\ell \eta_i(A_{N_j}Y) \right. \\ & \left. + \sum_{\alpha=r+1}^n H_\alpha^s \eta_i(A_{W_\alpha Y}) - \bar{c} \eta_i(Y) \right\} g(X, PU). \end{aligned}$$

Thus by the method of Theorem 4.2 we have the equation in theorem.

**COROLLARY 1.** *Let  $(M, g, S(TM))$  be a totally umbilical co-isotropic lightlike submanifold of a semi-Riemannian manifold  $(\bar{M}(\bar{c}), \bar{g})$ . If  $S(TM)$  is totally geodesic, then  $\bar{c} = 0$ , i.e.,  $\bar{M}$  is semi-Euclidean.*

**COROLLARY 2.** *Under the hypothesis of Corollary 1,  $\nabla$  is a metric connection on  $M$ , if and only if, the mean curvature vectors  $K_i$  of  $S(TM)$  are a solution of the following partial differential equations*

$$X(K_i) - \sum_{j=1}^r K_j \rho_{ji}(X) - \bar{c} \eta_i(X) = 0.$$

By the method of Theorem 4.4, using (4.19) and (4.21) into (3.9), and then (2.12), (2.22), (2.24) and  $M$  a space of constant curvature, we obtain

**THEOREM 4.5.** *Let  $(M(c), g, S(TM))$  be a totally umbilical  $r$ -lightlike or a co-isotropic submanifold of constant curvature  $c$  of a semi-Riemannian manifold  $(\bar{M}, \bar{g})$ . If  $S(TM)$  is totally umbilical, then the mean curvature vectors  $K_i$  of  $S(TM)$  are a solution of the partial differential equations*

$$X(K_i) - \sum_{j=1}^r K_j \rho_{ji}(X) - K_i \sum_{j=1}^r H_j^\ell \eta_j(X) - c\eta_i(X) = 0.$$

**COROLLARY 3.** *Let  $(M(c), g, S(TM))$  be an  $r$ -lightlike or a co-isotropic submanifold of constant curvature  $c$  of a semi-Riemannian manifold  $(\bar{M}, \bar{g})$ . If  $S(TM)$  is totally geodesic, then  $c = 0$ , i.e.,  $M$  is semi-Euclidean.*

**COROLLARY 4.** *Under the hypothesis of Theorem 4.5,  $\nabla$  on  $M$  is metric connection, if and only if, the mean curvature vectors  $K_i$  of  $S(TM)$  are a solution of the following partial differential equations*

$$X(K_i) - \sum_{j=1}^r K_j \rho_{ji}(X) - c\eta_i(X) = 0.$$

Using (3.6), the symmetries of the operators  $A_{\xi_i}^*$  and Lemma 1, we have

**THEOREM 4.6.** *Let  $(M(c), g)$  be a lightlike submanifold of constant curvature  $c$  of a semi-Riemannian manifold  $(\bar{M}, \bar{g})$ , such that  $S(TM)$  is proper totally umbilical or  $S(TM)$  vanishes. Then,  $d(\text{Tr}(\rho_{ij})) = 0$ .*

Let  $x \in M$  and  $\xi$  be a null vector of  $T_x \bar{M}$ . A plane  $\Pi$  of  $T_x \bar{M}$  is called a null plane directed by  $\xi$  if it contains  $\xi$ ,  $\bar{g}_x(\xi, W) = 0$  for any  $W \in \Pi$  and there exists  $W_o \in \Pi$  such that  $\bar{g}(W_o, W_o) \neq 0$ . Following [2], define the null sectional curvature of  $\Pi$  with respect to  $\xi$  and  $\bar{\nabla}$ , as a real number

$$(4.22) \quad \bar{K}_\xi(\Pi) = \frac{\bar{R}(W, \xi, \xi, W)}{g(W, W)}$$

where  $W$  is an arbitrary non-null vector in  $\Pi$ . Similarly, define the null sectional curvature  $K_\xi(\Pi)$  of the null plane  $\Pi$  of the tangent space  $T_x M$  with respect to  $\xi$  and  $\nabla$ , as a real number

$$(4.23) \quad K_\xi(\Pi) = \frac{R(W, \xi, \xi, W)}{g(W, W)}.$$

Taking into account that both null sectional curvatures do not depend on the vector  $W$  and by using (3.3) and (3.7) we obtain

$$(4.24) \quad \bar{K}_\xi(\Pi_i) = \xi_i(H_i^\ell) - (H_i^\ell)^2 + \sum_{k=1}^r H_k^\ell \rho_{ki}(\xi_i) = K_\xi(\Pi_i),$$

where  $\Pi_i$  is a null plane directed by  $\xi_i$ . Thus we have

**THEOREM 4.7.** *Let  $(M, g)$  be a totally umbilical lightlike submanifold of a semi-Riemannian manifold  $(\bar{M}, \bar{g})$ . Then, both the null sectional curvature functions  $\bar{K}_\xi(\Pi_i)$  and  $K_\xi(\Pi_i)$  vanish, if and only if,  $H_i^\ell$  is a solution of the partial differential equation*

$$\xi_i(H_i^\ell) - (H_i^\ell)^2 + \sum_{k=1}^r H_k^\ell \rho_{ki}(\xi_i) = 0.$$

From the equation (4.16) in Theorem 4.2 and Theorem 4.7, we obtain

**THEOREM 4.8.** *Let  $(M, g)$  be a totally umbilical lightlike submanifold of a semi-Riemannian manifold of constant curvature  $(\bar{M}, \bar{g})$ . Then, both the null sectional curvature functions  $\bar{K}_\xi(\Pi_i)$  and  $K_\xi(\Pi_i)$  vanish.*

## 5. Induced Ricci tensor

Consider an  $m$ -dimensional lightlike submanifold  $(M, g)$  of an  $(m+n)$ -dimensional semi-Riemannian manifold  $(\bar{M}, \bar{g})$ . Note that  $h_i^\ell, \rho_{ij}$  and  $\tau_{ix}$  depend on the section  $\xi \in \Gamma(\text{Rad } TM)$ . Indeed, take  $\xi_i^* = \sum_{j=1}^r \alpha_{ij} \xi_j$ , where  $\alpha_{ij}$  are smooth functions with  $\Delta = \det(\alpha_{ij}) \neq 0$  and  $A_{ij}$  be the co-factors of  $\alpha_{ij}$  in the determinant of  $\Delta$ . It follows that  $N_i^* = (1/\Delta) \sum_{j=1}^r A_{ij} N_j$ . Hence by straightforward calculation and using (4.1)–(4.4) and (4.6) we obtain  $h_i^{\ell*} = \sum_{j=1}^r \alpha_{ij} h_j^\ell$ . Denote  $\rho_{ij}^*$  and  $\tau_{ix}^*$  by affinely combinations of  $\rho_{ij}$  and  $\tau_{ix}$  with coefficients  $\alpha_{ij}, A_{ij}$  and  $X(A_{ij})$ . Moreover,

$$\text{Tr}(\rho_{ij})(X) = \text{Tr}(\rho_{ij}^*)(X) + X(\log \Delta), \quad \forall X \in \Gamma(TM).$$

Thus, using the formula  $d\rho(X, Y) = (1/2)\{X(\rho(Y)) - Y(\rho(X)) - \rho([X, Y])\}$  of a differential 2-form, we obtain

**THEOREM 5.1.** *Let  $(M, g)$  be a lightlike submanifold of a semi-Riemannian manifold  $(\bar{M}, \bar{g})$ . Suppose  $\text{Tr}(\rho_{ij})$  and  $\text{Tr}(\rho_{ij}^*)$  are 1-forms on  $\mathcal{U}$  with respect to  $\xi_i$  and  $\xi_i^*$ . Then  $d(\text{Tr}(\rho_{ij}^*)) = d(\text{Tr}(\rho_{ij}))$  on  $\mathcal{U}$ .*

To find local expression of Ricci tensor of  $M$ , consider the frames field

$$\{\xi_1, \dots, \xi_r; N_1, \dots, N_r; X_{r+1}, \dots, X_m; W_{r+1}, \dots, W_n\}$$

on  $\bar{M}$ . Denote by  $\{F_A\} = \{\xi_1, \dots, \xi_r, X_{r+1}, \dots, X_m\}$  the induced frames field on  $M$ . Then,

$$\begin{aligned}
\bar{R}_{ABCD} &= \bar{g}(\bar{R}(F_D, F_C)F_B, F_A), & R_{ABCD} &= g(R(F_D, F_C)F_B, F_A), \\
\bar{R}_{iBCD} &= \bar{g}(\bar{R}(F_D, F_C)F_B, N_i), & R_{iBCD} &= \bar{g}(R(F_D, F_C)F_B, N_i), \\
\bar{R}_{\alpha BCD} &= \bar{g}(\bar{R}(F_D, F_C)F_B, W_\alpha), & R_{\alpha BCD} &= \bar{g}(R(F_D, F_C)F_B, W_\alpha), \\
\bar{R}_{i\alpha CD} &= \bar{g}(\bar{R}(F_D, F_C)W_\alpha, N_i), & R_{i\alpha CD} &= \bar{g}(R(F_D, F_C)W_\alpha, N_i).
\end{aligned}$$

Using above we obtain the following local expression for the Ricci tensor:

$$\text{Ric}(X, Y) = \sum_{a,b=r+1}^m g^{ab} g(R(X, X_a)Y, X_b) + \sum_{i=1}^r \bar{g}(R(X, \xi_i)Y, N_i).$$

By using the symmetries of curvature tensor and the first Bianchi identity and taking into account (3.2) and (3.9) we obtain

$$\begin{aligned}
&\text{Ric}(X, Y) - \text{Ric}(Y, X) \\
&= \sum_{a,b=r+1}^m g^{ab} \{ \bar{g}(h^*(X, X_b), h^\ell(Y, X_a)) - \bar{g}(h^*(Y, X_b), h^\ell(X, X_a)) \} \\
&\quad + \sum_{i=1}^r \{ g(A_{\xi_i}^* X, A_{N_i} Y) - g(A_{\xi_i}^* Y, A_{N_i} X) + \bar{g}(R^{*t}(X, Y)\xi_i, N_i) \}.
\end{aligned}$$

Replacing  $X, Y$  by  $X_A, X_B$  respectively, using (2.27), (2.28), (4.9) and

$$\begin{aligned}
\sum_{i=1}^r \bar{g}(R^{*t}(X, Y)\xi_i, N_i) &= -2 \sum_{i,j=1}^r \bar{g}(d(\rho_{ji})(X, Y)\xi_j, N_i) \\
&= -2d(\text{Tr}(\rho_{ij}))(X, Y),
\end{aligned}$$

we have

$$R_{AB} - R_{BA} = 2d(\text{Tr}(\rho_{ij}))(X_A, X_B)$$

where  $R_{AB} = \text{Ric}(X_B, X_A)$ . Thus, using Theorem 5.1, we conclude

**THEOREM 5.2.** *Let  $(M, g, S(TM))$  be an  $r$ -lightlike or a co-isotropic submanifold of a semi-Riemannian manifold  $(\bar{M}, \bar{g})$ . Then the Ricci tensor of the induced connection  $\nabla$  on  $M$  is symmetric, if and only if, each 1-forms  $\text{Tr}(\rho_{ij})$  induced by  $S(TM)$  is closed, i.e., on any  $\mathcal{U} \subset M$ ,*

$$d(\text{Tr}(\rho_{ij})) = 0.$$

Using Theorems 4.3, 4.6 and 5.2, we obtain the following theorem:

**THEOREM 5.3.** *Let  $(M, g, S(TM))$  be a proper totally umbilical  $r$ -lightlike or a co-isotropic submanifold of a semi-Riemannian manifold  $(\bar{M}(\bar{c}), \bar{g})$  of a constant curvature  $\bar{c}$ . Then, the induced Ricci tensor on  $M$  is symmetric, if and only if its screen distribution  $S(TM)$  is integrable.*

**COROLLARY 5.** *Let  $(M(c), g, S(TM))$  be an  $r$ -lightlike or a co-isotropic submanifold of constant curvature  $c$  of a semi-Riemannian manifold  $(\bar{M}, \bar{g})$ , such that  $S(TM)$  is proper totally umbilical. Then the Ricci tensor of the induced connection  $\nabla$  on  $M$  is symmetric.*

Suppose the Ricci tensor of  $\nabla$  is symmetric. Theorem 5.3 and Poincare lemma implies  $\text{Tr}(\rho_{ij}(X)) = X(f)$ , where  $f$  is a smooth function. Let  $\Delta = \exp f$  and obtain  $\text{Tr}(\rho_{ij}^*(X)) = 0 \forall X \in \Gamma(TM|_{\mathcal{U}})$ . Thus we have

**THEOREM 5.4.** *Let  $(M, g, S(TM))$  be an  $r$ -lightlike or a co-isotropic submanifold of a semi-Riemannian manifold  $(\bar{M}, \bar{g})$ . Then Ricci tensor of  $M$  is symmetric and there exists a pair of frames field  $\{\xi_i^*, N_i^*\}$  on  $\mathcal{U}$  such that the corresponding 1-forms  $\text{Tr}(\rho_{ij}^*)$  induced by  $S(TM)$  vanishes.*

*Remark 4.* Lemma 1 and Theorem 5.2 imply that the induced Ricci tensor of either an isotropic or a totally lightlike  $M$  of  $\bar{M}(\bar{c})$  is always symmetric. This clarifies the fact that Theorems 5.1–5.4 will trivially hold for an isotropic or a totally lightlike  $M$ , since for these two cases  $d(\text{Tr}(\rho_{ij})) = 0$ .

*Examples.* Minkowski [3], de Sitter [1], Schwarzschild and Robertson-Walker spacetimes (see [10] and pages 225–230 of [8]) all have lightlike hypersurfaces with an integrable 2-dimensional screen distribution.

## REFERENCES

- [1] M. A. AKIVIS AND V. V. GOLDBERG, The geometry of lightlike hypersurfaces of the de Sitter space, *Acta Appl. Math.*, **53** (1998), 297–328.
- [2] J. K. BEEM, P. E. EHRLICH AND K. L. EASLEY, *Global Lorentzian Geometry*, Second Edition, Monogr. Textbooks Pure Appl. Math. 202, Marcel Dekker, New York, 1996.
- [3] W. B. BONNOR, Null hypersurfaces in Minkowski space-time, *Tensor (N.S.)*, **24** (1972), 329–345.
- [4] B.-Y. CHEN, *Geometry of Submanifolds*, Pure and Applied Mathematics 22, Marcel Dekker, New York, 1973.
- [5] K. L. DUGGAL, Warped product of lightlike manifolds, *Nonlinear Anal.*, **47** (2001), 3061–3072.
- [6] K. L. DUGGAL, Constant scalar curvature and warped product globally null manifolds, *J. Geom. Phys.*, **43** (2002), 327–340.
- [7] K. L. DUGGAL AND A. BEJANCU, *Lightlike Submanifolds of Semi-Riemannian Manifolds and Applications*, Math. Appl. 364, Kluwer Academic Publishers, Dordrecht, 1996.
- [8] K. L. DUGGAL AND D. H. JIN, Geometry of null curves, *Math. J. Toyama Univ.*, **22** (1999), 95–120.
- [9] K. L. DUGGAL AND D. H. JIN, Half lightlike submanifolds of codimension 2, *Math. J. Toyama Univ.*, **22** (1999), 121–161.
- [10] S. W. HAWKING AND G. F. R. ELLIS, *The Large Scale Structure of Space-Time*, Cambridge Monographs on Mathematical Physics 1, Cambridge University Press, London, 1973.

- [11] T. IKAWA, On curves and submanifolds in an indefinite-Riemannian manifold, *Tsukuba J. Math.*, **9** (1985), 353–371.
- [12] B. O'NEILL, *Semi-Riemannian Geometry, With Applications to Relativity*, Pure Appl. Math. 103, Academic Press, New York, 1983.

DEPARTMENT OF MATHEMATICS AND STATISTICS  
UNIVERSITY OF WINDSOR  
WINDSOR, ONTARIO, CANADA N9B3P4  
e-mail: yq8@uwindsor.ca

DEPARTMENT OF MATHEMATICS  
DONGGUK UNIVERSITY  
KYONGJU, KOREA, 780-714  
e-mail: jindh@dongguk.ac.kr