

## THE MAXIMAL QUOTIENT RING OF A LEFT H-RING

Dedicated to Professor Hiroyuki Tachikawa on his sixtieth birthday

JIRO KADO

(Received March 20, 1989)

In [2], M. Harada has introduced two new artinian rings which are closely related to  $QF$ -rings; one is a left artinian ring whose non-small left module contains a non-zero injective submodule and the other is a left artinian ring whose non-cosmall left module contains a non-zero projective summand. K. Oshiro called the first ring a *left H-ring* and the second one a *left co-H-ring* ([3]). However, later in [5], he showed that a ring  $R$  is a left  $H$ -ring if and only if it is a right co- $H$ -ring.  $QF$ -rings and Nakayama (artinian serial) rings are left and right  $H$ -rings ([3]). As the maximal quotient rings of Nakayama rings are Nakayama, it is natural to ask whether the maximal quotient rings of left  $H$ -rings are left  $H$ -rings. In this note, we show that this problem is affirmative, by determining the structure of the maximal quotient rings of left  $H$ -rings.

### 1. Preliminaries

Throughout this paper, we assume that all rings  $R$  considered are associative rings with identity and all  $R$ -modules are unital. Let  $M$  be a  $R$ -module. We use  $J(M)$  and  $S(M)$  to denote its Jacobson radical and its socle, respectively.

Definition [3]. A module is *non-small* if it is not a small submodule of its injective hull. We say that a ring  $R$  is a *left H-ring* if  $R$  is a left artinian ring satisfying the condition that every non-small left  $R$ -module contains a non-zero injective submodule.

We note that a left  $H$ -ring is also right artinian by [7, Th. 3]. In [5], for a left  $H$ -ring  $R$ , K. Oshiro gave the following theorem, by using M. Harada's results of [2, Th. 3.6.]: a ring  $R$  is a left  $H$ -ring if and only if it is left artinian and its complete set  $E$  of orthogonal primitive idempotents is arranged as  $E = \{e_{11}, \dots, e_{1n(1)}, \dots, e_{m1}, \dots, e_{mn(m)}\}$  for which

- (1) each  $e_{ii}R$  is injective,
- (2) for each  $i$ ,  $e_{ik-1}R \cong e_{ik}R$  or  $J(e_{ik-1}R) \cong e_{ik}R$  for  $k=2, \dots, n(i)$ , and
- (3)  $e_{ik}R \not\cong e_{ji}R$  if  $i \neq j$ .

As a left  $H$ -ring is a  $QF$ -3 ring by [4], the maximal left quotient ring and the maximal right quotient ring of a left  $H$ -ring coincide by [9, Th. 1.4]. From now on, let  $Q$  be the maximal quotient ring of a left  $H$ -ring  $R$ . We shall study the structure of  $Q$ . Since maximal quotient rings and left  $H$ -rings are Morita-invariant [7], in order to investigate the problem whether  $Q$  is a left  $H$ -ring or not, we may restrict our attention to basic left  $H$ -rings. Therefore, hereafter, we assume that  $R$  is a basic left  $H$ -ring and  $E$  is a complete set of orthogonal primitive idempotents of  $R$ . Then  $E$  is arranged as  $E = \{e_{11}, \dots, e_{1n(1)}, \dots, e_{m1}, \dots, e_{mn(m)}\}$  for which

- (1) each  $e_{i1}R$  is injective,
- (2) for each  $i, J(e_{ik-1}R) \cong e_{ik}R$  for  $k=2, \dots, n(i)$ .

Definition [10, p. 153]. A primitive idempotent  $e$  is called *S-primitive* if the simple module  $eR/eJ(R)$  is isomorphic to a minimal right ideal.

We shall use the H.H. Storrer's characterization of the maximal quotient ring of a perfect ring [10].

Since each  $e_{i1}R$  ( $i=1, \dots, m$ ) is injective, there exists a unique  $g_i$  in  $E$  such that  $(e_{i1}R; Rg_i)$  is an *injective pair*, that is,  $S(e_{i1}R) \cong g_iR/J(g_iR)$  and  $S(Rg_i) \cong Re_{i1}/J(Re_{i1})$  (cf. K.R. Fuller [1, Th. 3.1]). Each pair  $\{e_{i1}, g_i\}$  ( $i=1, \dots, m$ ) is very important for studying left  $H$ -rings.

Now we shall determine all  $S$ -primitive idempotents in  $E$ . Let  $e$  be an idempotent in  $E$ . It is known that  $e$  is  $S$ -primitive if and only if  $S(R_R)e \neq 0$  [10, Lemma 2.3]. Since  $S(R_R) = \bigoplus_{i=1}^m \bigoplus_{k=1}^{n(i)} S(e_{ik}R)$ ,  $S(e_{ik}R) \cong S(e_{j1}R)$  for  $i \neq j$  and  $S(e_{ik}R) \cong S(e_{i1}R)$ , we have  $S(R_R)e \neq 0$  if and only if  $S(e_{i1}R)e \neq 0$  for a unique  $i$ . Therefore  $e$  is an  $S$ -primitive idempotent if and only if  $e = g_i$  for some  $i$ . Then  $E' = \{g_1, \dots, g_m\}$  is the set of all  $S$ -primitive idempotents in  $E$ . Put  $g = g_1 + \dots + g_m$  and  $D = RgR$ . Storrer has shown that  $D = RgR$  is the minimal dense ideal of  $R$  and  $Q$  is isomorphic to  $\text{Hom}_R(D_R, D_R) = \text{Hom}_R(D_R, R_R)$  by [10, Prop. 1.2 and Th. 2.5]. Since  $R$  is a two-sided artinian ring,  $Q$  is a left artinian ring by [10, Prop. 3.1].

**Lemma 1.** *For each  $e$  in  $E$ ,  $e$  is also a primitive idempotent in  $Q$ . Therefore  $S(eQ)$  is a simple  $Q$ -module.*

Proof. Since  $eR$  is a uniform right ideal,  $eQ$  is also a uniform right ideal of  $Q$  by [30, Prop. 4.4]. Thus  $eQ$  is indecomposable.

By the above lemma, we know that  $E = \{e_{11}, \dots, e_{1n(1)}, \dots, e_{m1}, \dots, e_{mn(m)}\}$  is also a complete set of orthogonal primitive idempotents of  $Q$ . We shall prove that  $Q$  is left  $H$ -ring by showing that  $E$  satisfies the conditions (1), (2) and (3) of left  $H$ -rings. We again note that left  $H$ -rings are also right artinian by

[7, Th. 3] and the maximal quotient ring  $Q$  of  $R$  is a left artinian ring.

**Proposition 2.** *In the maximal quotient ring  $Q$ ,  $(e_{i1}Q; Qg_i)$  is an injective pair for  $i=1, \dots, m$ . Consequently  $e_{i1}Q$  and  $Qg_i$  are injective  $Q$ -modules.*

Proof. By assumption, let  $\phi: g_iR \rightarrow S(e_{i1}R)$  be an epimorphism.  $\phi$  extends uniquely to a  $Q$ -homomorphism  $\phi^*: g_iQ \rightarrow S(e_{i1}R)Q$  by [10, Prop. 4.3]. Since  $S(e_{i1}R)Q = S(e_{i1}Q)$ ,  $\phi^*$  is also an epimorphism and hence  $g_iQ/J(g_iQ) \cong S(e_{i1}Q)$ . Since  $Q$  is the maximal left quotient ring of  $R$ , we have symmetrically that  $Qe_{i1}/J(Qe_{i1}) \cong S(Qg_i)$ . By [1, Th. 3.1],  $(e_{i1}Q; Qg_i)$  is an injective pair for  $i=1, \dots, m$ .

Next we shall study isomorphisms among the indecomposable right ideals  $e_{ik}Q$ . Let  $f_1, f_2$  be idempotents in  $E$  and we assume that there exists a monomorphism  $\theta: f_1R \rightarrow f_2R$  such that  $\text{Im } \theta = J(f_2R)$ . Then by [10, Prop. 4],  $\theta$  can be uniquely extended to a  $Q$ -homomorphism  $\theta^*: f_1Q \rightarrow f_2Q$ . We shall prove the following result.

**Proposition 3.** (1) *If  $f_2$  is not  $S$ -primitive, then the extension  $\theta^*: f_1Q \rightarrow f_2Q$  is an isomorphism.*

(2) *If  $f_2$  is  $S$ -primitive, then  $\theta^*: f_1Q \rightarrow f_2Q$  is a monomorphism such that  $\text{Im } \theta^* = J(f_2Q)$ .*

Proof. From  $0 \rightarrow f_1R \xrightarrow{\theta} f_2R \rightarrow M \rightarrow 0$ , where  $M = f_2R/J(f_2R)$ , we have the following exact sequence

$$0 \rightarrow f_1Q = \text{Hom}(D, f_1R) \xrightarrow{\theta^*} f_2Q = \text{Hom}(D, f_2R) \rightarrow \text{Hom}(D, M).$$

(1) It is sufficient to prove that  $\text{Hom}(D, M) = 0$ . We assume that there exists a non-zero homomorphism  $\phi: D \rightarrow M$ . Since  $D = R(g_1 + \dots + g_m)R$  by [10, Th. 2.5], there exist some  $i$  and some  $x \in R$  such that  $xg_iR \not\subseteq \text{Ker } \phi$ . Then  $g_iR/J(g_iR) \cong M$ . Therefore  $g_iR \cong f_2R$ . This contradicts that  $f_2$  is not  $S$ -primitive. Consequently we have that  $\text{Hom}(D, M) = 0$ , and so  $\theta^*$  is an isomorphism.

(2) First we shall show that  $\text{Im } \theta^* \neq f_2Q$ . Since  $f_2$  is  $S$ -primitive, we have that  $f_2R \subset D$  and so  $D = f_2R \oplus (D \cap (1-f_2)R)$ . Therefore the projection  $\alpha: D \rightarrow f_2R$  is not contained in  $\text{Im } \theta^* \subseteq \text{Hom}(D, J(f_2R))$ . For any  $\phi \in J(f_2Q)$ ,  $\phi$  is not an epimorphism as  $R$ -homomorphism. In fact, we shall show that any epimorphism  $\alpha: D \rightarrow f_2R$  generates  $f_2Q$ . Let  $\beta$  be any homomorphism  $D \rightarrow f_2R$  and  $\alpha': f_2R \rightarrow D$  the split homomorphism of  $\alpha$ . Then we have  $\beta = \alpha\alpha'\beta$ . Therefore any  $\phi \in J(f_2Q)$  is contained in  $\text{Im } \theta^*$  and so  $\text{Im } \theta^* = J(f_2Q)$ , because  $J(f_2Q)$  is the unique maximal submodule of  $f_2Q$ .

Now we shall prove our main theorem.

**Theorem 4.** *Let  $R$  be a left  $H$ -ring. Then the maximal quotient ring  $Q$  of  $R$  is also an  $H$ -ring.*

Proof. Let  $E = \{e_{11}, \dots, e_{in(i)}, \dots, e_{m1}, \dots, e_{mn(m)}\}$  be a complete set of orthogonal primitive idempotents of  $R$  such that

- (1) each  $e_{ii}R$  is injective,
- (2) for each  $i, J(e_{ik-1}R) \cong e_{ik}R$  for  $k=2, \dots, n(i)$ .

We have already known that  $Q$  is a left artinian ring and  $E$  is also a complete set of orthogonal primitive idempotents of  $Q$ . By Proposition 2, each  $e_{ii}Q$  is an injective  $Q$ -module and by Proposition 3,  $e_{ik}Q \cong e_{ik-1}Q$  or  $e_{ik}Q \cong J(e_{ik-1}Q)$   $k=2, \dots, n(i)$  for each  $i$ . We shall show that  $e_{ik}Q \not\cong e_{jt}Q$  if  $i \neq j$ . If  $e_{ik}Q \cong e_{jt}Q$  for some  $i \neq j, k, t$ , then  $S(e_{ik}Q) \cong S(e_{jt}Q)$ . Since  $S(e_{ik}Q) = S(e_{ik}R)Q$  and  $S(e_{jt}Q) = S(e_{jt}R)Q$ , we have  $S(e_{ik}R) \cong S(e_{jt}R)$  as  $R$ -modules by [10, Th. 4.5]. This contradicts the assumption of  $E$ .

We recall that  $g_i$  is the element of  $E$  such that  $(e_{ii}R; Rg_i)$  is an injective pair for  $i=1, \dots, m$ . Here we define two mappings

$$\begin{aligned} \sigma: \{1, \dots, m\} &\rightarrow \{1, \dots, m\} \\ \rho: \{1, \dots, m\} &\rightarrow \{1, \dots, n(1)\} \cup \dots \cup \{1, \dots, n(m)\} \end{aligned}$$

by the rule  $\sigma(i) = k$  and  $\rho(i) = t$  if  $g_i = e_{kt}$ . We note that  $\{\sigma(1), \dots, \sigma(m)\} \subseteq \{1, \dots, m\}$  and  $1 \leq \rho(i) \leq n(\sigma(i))$ .

Here we shall define a special left  $H$ -ring.

Definition [7, p. 94]. A left  $H$ -ring is *Type (\*)* if  $\{\sigma(1), \dots, \sigma(m)\}$  is a permutation of  $\{1, \dots, m\}$  and  $\rho(i) = n(\sigma(i))$  for all  $i=1, \dots, m$ .

**Cororally.** *Let  $R$  be a left  $H$ -ring. Then the maximal quotient ring  $Q$  of  $R$  is a QF-ring if and only if  $R$  is Type (\*).*

Proof. It is easy by Proposition 3.

**Example.** Let  $T$  be a local QF-ring,  $J=J(T)$  and  $S=S(T)$ .

Put  $V = \begin{pmatrix} T & T & T \\ J & T & T \\ J & J & T \end{pmatrix}$  and  $W = \begin{pmatrix} 0 & 0 & S \\ 0 & 0 & S \\ 0 & 0 & S \end{pmatrix}$ . The factor ring  $R = V/W$  is a left

$H$ -ring such that  $e_1R$  is injective,  $J(e_1R) \cong e_2R$  and  $J(e_2R) \cong e_3R$ , where  $e_i$  is the matrix such that its  $(i, i)$ -position is 1 and all other entries are zero.  $R$  is repre-

sented as follows:  $\begin{pmatrix} T & T & \tilde{T} \\ J & T & \tilde{T} \\ J & J & \tilde{T} \end{pmatrix}$  where  $\tilde{T} = T/S$ . Since  $(e_1R; Re_2)$  is injective pair by

[8, § 2], the minimal dense ideal is  $Re_2R$ . Therefore the maximal quotient ring  $Q$  of  $R$  is a left  $H$ -ring such that  $e_1Q$  is an injective module,  $e_1Q \cong e_2Q$  and  $J(e_2Q) \cong e_3Q$ . Since  $e_1Q/J(e_1Q) \cong S(e_1Q)$ , we have that  $\text{Hom}_Q(e_1Q, J(e_1Q)) \cong$

$J(e_1 Q e_1), \text{Hom}_Q(J(e_1 Q), e_1 Q) \cong e_1 Q e_1 / S(e_1 Q e_1), \text{Hom}_Q(J(e_1 Q), J(e_1 Q)) \cong e_1 Q e_1 / S(e_1 Q e_1)$ .  
 Moreover, since  $e_1 Q e_1 = e_1 R e_1 = T$  by [10, Lemma 4.2],  $Q$  is

represented as a matrix ring 
$$\begin{pmatrix} T & T & \tilde{T} \\ T & T & \tilde{T} \\ J & J & \tilde{T} \end{pmatrix}.$$

---

### References

- [1] K.R. Fuller: *On indecomposable injectives over artinian rings*, Pacific. J. Math. **29** (1969), 115–135.
- [2] M. Harada: *Non-small modules and non-cosmall modules*, Ring Theory, Proceedings of 1978 Antwerp Conference, Marcel Dekker Inc., 1979, 669–689.
- [3] K. Oshiro: *Lifting modules, extending modules and their applications to QF-rings*, Hokkaido Math. J. **13** (1984), 310–338.
- [4] K. Oshiro: *Lifting modules, extending modules and their applications to generalized uniserial rings*, Hokkaido Math. J. **13** (1984), 339–346.
- [5] K. Oshiro: *On Harada-rings I*, to appear in Math. J. of Okayama Univ.
- [6] K. Oshiro: *On Harada-rings II*, to appear in Math. J. of Okayama Univ.
- [7] K. Oshiro and S. Masumoto: *The self-duality of H-rings and Nakayama automorphisms of QF-rings*, Proceedings of the 18th Symposium of Ring Theory, 1985, 84–107.
- [8] K. Oshiro and K. Shigenaga: *On H-rings with homogeneous socles*, to appear in Math. J. of Okayama Univ.
- [9] C.M. Ringel and H. Tachikawa: *QF-3 rings*, J. Reine Angew. Math. **272** (1975), 49–72.
- [10] H.H. Storrer: *Rings of quotients of perfect rings*, Math. Z. **122** (1971), 151–165.

Department of Mathematics  
 Osaka City University  
 Sugimoto, Sumiyosi-ku  
 Osaka, Japan

