

RSK in last passage percolation: a unified approach*

Duncan Dauvergne

*Department of Mathematics, University of Toronto,
Room 6290, 40 St. George Street, Toronto, ON, Canada M5S 2E4
e-mail: duncan.dauvergne@utoronto.ca*

Mihai Nica

*Department of Mathematics & Statistics, University of Guelph,
50 Stone Road East Room 437, MacNaughton Building, Guelph, ON Canada, N1G 2W1
e-mail: nicam@uoguelph.ca*

Bálint Virág

*Department of Mathematics, University of Toronto,
Room 6290, 40 St. George Street, Toronto, ON, Canada M5S 2E4
e-mail: balint@math.toronto.edu*

Abstract: We present a version of the RSK correspondence based on the Pitman transform and geometric considerations. This version unifies ordinary RSK, dual RSK and continuous RSK. We show that this version is both a bijection and an isometry, two crucial properties for taking limits of last passage percolation models.

We use the bijective property to give a non-computational proof that dual RSK maps Bernoulli walks to nonintersecting Bernoulli walks.

MSC2020 subject classifications: Primary 05A05; secondary 60K35.

Keywords and phrases: RSK bijection, Pitman transform, Robinson-Schensted correspondence, Greene’s theorem, last passage percolation, KPZ universality class, Young tableaux.


Received June 2021.

Contents

1	Introduction	66
2	Percolation across cadlag functions	71
2.1	Basic definitions	71
2.2	Multiple paths	72
2.3	Basic geometric properties	72
3	The melon	75
3.1	The Pitman transform	75
3.2	Repeated Pitman transforms, cars, and the melon	78

*D.D. and M.N. were supported by NSERC postdoctoral fellowships. B.V. was supported by the Canada Research Chair program and an NSERC Discovery Accelerator Grant.

4	Minimal and maximal elements	81
5	The infinite bijection	83
6	The finite bijection	88
6.1	Bijectivity for lattice specializations and other restrictions	96
7	Preservation of uniform measure	97
8	Embedding classical versions of RSK	101
8.1	Young tableaux	101
8.2	Classical RSK via Greene’s theorem	102
8.3	Classical RSK and the melon map	103
8.4	Dual RSK	106
A	Appendix: technical proofs	108
	References	110



1. Introduction

The Robinson-Schensted-Knuth (RSK) correspondence originates in the study of representations of the symmetric group, Robinson (1938). In probability, it has been mainly used for understanding different models of two-dimensional directed last passage percolation.

Across these models, several versions of RSK have been used in the literature. RSK has also been used to construct the directed landscape, an object that is expected to be the universal limit of such models, see Dauvergne, Ortmann and Virág (2018). The goal of this paper is to introduce a version of RSK that unifies three commonly used versions: ordinary RSK, dual RSK, and continuous RSK. We use this unified setting to prove some basic important properties of RSK: bijectivity and isometry. The results in this paper are central to establishing the scaling limit of the longest increasing subsequence and other KPZ models in Dauvergne and Virág (2021). Our approach keeps probability applications in mind and avoids representation theory concepts entirely. We work with a global perspective, using last passage values as opposed to local bumping algorithms.

The paper is centred around first fully understanding an infinite-time version of the RSK bijection for cadlag paths. Infinite time versions of RSK have previously been studied by O’Connell (2003b); Biane, Bougerol and O’Connell (2005), see Remark 5.10 for more discussion. Surprisingly, this version turns out to be much simpler than ordinary RSK! We also obtain parallel descriptions for




FIG 1. The blue path is to the left of the red path and the two paths are essentially disjoint

RSK and its inverse, allowing us to give a purely global geometric description for the RSK inverse map.

Let \mathcal{D}^n be the space of n -tuples of cadlag functions from $[0, \infty) \rightarrow \mathbb{R}$ with no negative jumps. Each $f = (f_1, \dots, f_n) \in \mathcal{D}^n$ defines a finitely additive signed measure df on $[0, \infty) \times \{1, \dots, n\}$ through

$$df([x, y] \times \{i\}) = f_i(y) - f_i(x^-).$$

When visualizing f and this measure, we will think in matrix coordinates, so that line 1 (i.e. the function f_1) is on top and line n is on the bottom, see Figure 1. For $(p, q) = (x, n; y, m) \in (\mathbb{R} \times \mathbb{Z})^2$, we write $p \nearrow q$ if $x \leq y$ and $n \geq m$. For $p \nearrow q$, a path π from p to q is a union of closed intervals

$$[t_i, t_{i-1}] \times \{i\}, \quad i = m, m+1, \dots, n, \quad x = t_n \leq t_{n-1} \leq \dots \leq t_m \leq t_{m-1} = y.$$

Two paths are called **essentially disjoint** if the corresponding intervals have disjoint interiors, see Figure 1. Define the **length** of a path π by

$$|\pi|_f = df(\pi).$$

For $u = (p; q) = (x, n; y, m) \in (\mathbb{R} \times \mathbb{Z})^2$ define the **distance** in f from p to q by

$$f[u] = f[p \rightarrow q] = f[(x, n) \rightarrow (y, m)] = \sup_{\pi} |\pi|_f, \quad (1.1)$$

where the supremum is taken over all paths π from p to q . If $p \not\nearrow q$ then this set of paths is empty. It will still be useful to define $f[u]$ in this case, so here we simply set $f[u] = -\infty$.

We also define $f[p^k \rightarrow q^k] = \sup df(\pi_1 \cup \dots \cup \pi_k)$, where the supremum is over tuples of k essentially disjoint paths from p to q . We call a tuple that achieves $f[p^k \rightarrow q^k]$ a **k -disjoint optimizer** from p to q .

Define the **melon map** $W : \mathcal{D}^n \rightarrow \mathcal{D}^n$ by

$$(Wf)_k(y) = f[(0, n)^k \rightarrow (y, 1)^k] - f[(0, n)^{k-1} \rightarrow (y, 1)^{k-1}] \quad (1.2)$$

with the convention that $f[p^0 \rightarrow q^0] = 0$, see the end of the introduction for discussion on how this is related to the standard presentation of RSK. Throughout the paper we will write $Wf_k = (Wf)_k$ to avoid cluttered notation. We use this same convention for components of all other operators.

We summarize some of the remarkable properties of the map W in the next theorem.

Theorem 1.1. *Consider the melon map W on \mathcal{D}^n .*

- (i) **Isometry.** For $p = (x, n)$ and $q = (y, 1)$, we have $f[p \rightarrow q] = Wf[p \rightarrow q]$.
- (ii) **Idempotent property.** $WW = W$.
- (iii) **Image.** $\text{Im } W = \mathcal{D}_{\uparrow}^n$, the set of all $f \in \mathcal{D}^n$ such that $f_{i-1}(y^-) \geq f_i(y)$ for all i, y .
- (iv) **Bijection.** W is a bijection between $\mathcal{D}_{\downarrow}^n$ and \mathcal{D}_{\uparrow}^n , see below.

The ordering in Theorem 1.1 (iii) gives the sequence Wf_1, \dots, Wf_n the appearance of stripes on a watermelon. For this reason, we call Wf the **melon** of f . The isometry property extends to multi-point last passage values as in (1.2), see Proposition 3.12.

The set of functions $\mathcal{D}_{\downarrow}^n \subset \mathcal{D}^n$ on which W is a bijection can be explicitly described. Define $\mathcal{D}_{\downarrow-}^n$ as the set of functions on which the following holds: the k lowest constant paths $[0, \infty) \times \{i\}, i = k - n + 1, \dots, n$ are a local limit as $t \rightarrow \infty$ of a sequence of k -disjoint optimizers from $(0, n)$ to $(t, 1)$, see Section 5. The set $\mathcal{D}_{\downarrow}^n$ is the closure of $\mathcal{D}_{\downarrow-}^n$ in the uniform topology.

For the inverse of W , let R transform the signed measure df up to time t by rotating the base space $[0, t] \times \{1, \dots, n\}$ by 180 degrees. We write $Rf = g$ if $R(df) = R(dg)$, and let

$$Mf = \lim_{t \rightarrow \infty} RWRf. \quad (1.3)$$

Theorem 1.2 (Explicit inverse). *M is well-defined on \mathcal{D}^n . Moreover we have*

- (i) **Isometry.** For $p = (x, n)$ and $q = (y, 1)$, we have $f[p \rightarrow q] = Mf[p \rightarrow q]$.
- (ii) **Idempotent property.** $MM = M$.
- (iii) **Image.** $\text{Im } M = \mathcal{D}_{\uparrow}^n$.
- (iv) **Bijection.** M is a bijection between \mathcal{D}_{\uparrow}^n and $\mathcal{D}_{\downarrow}^n$ with inverse W .

The following proposition provides a simple sufficient condition for f to be in $\mathcal{D}_{\downarrow}^n$. It implies that classical examples, such as i.i.d. random walks or Brownian motion paths are a.s. in $\mathcal{D}_{\downarrow}^n$. In other words, if $F \in \mathcal{D}^n$ is a sequence of i.i.d. random walks or Brownian motion paths, then a.s. we can reconstruct F from WF .

Proposition 1.3. *If $f \in \mathcal{D}^n$ and for all j , the function $(f_{j+1} - f_j)^+$ is unbounded, then $f \in \mathcal{D}_{\downarrow}^n$.*

Remark 1.4. One useful perspective on our description of RSK is to focus purely on the isometry. Put an equivalence relation on \mathcal{D}^n by letting $f \sim g$ if $f[(x, n) \rightarrow (y, 1)] = g[(x, n) \rightarrow (y, 1)]$ for all x, y . From this point of view, W maps f to the element of its equivalence class with the leftmost disjoint optimizers, and M maps f to the element of its equivalence class with the

FIG 2. Bernoulli walks f and Wf

rightmost disjoint optimizers, see Section 4 for a more precise setup. When thinking in these terms, idempotence and bijectivity fall out naturally.

We can embed a finite time RSK correspondence into the melon map W , see Section 6 for details. Bijectivity and other properties in the finite setting can be deduced from the simpler infinite case. Restrictions of this finite time bijection recover the usual RSK and dual RSK correspondences, see Section 8.

Bijectivity is the reason that certain measures on \mathcal{D}^n have tractable push-forwards under W , and this is why RSK is useful in probability. For example, if $B \in \mathcal{D}^n$ consists of n independent standard Brownian motions, then WB is simply n independent standard Brownian motions conditioned so that $B_1(t) \geq \dots \geq B_n(t)$ at all times t .

These results are traditionally proven using the Burke property and dynamical symmetry, e.g. see O’Connell and Yor (2002), König, O’Connell and Roch (2002), Draief, Mairesse and O’Connell (2005), Biane, Bougerol and O’Connell (2005). We give a proof for the case of Bernoulli walks that instead directly uses the bijectivity of RSK in Theorems 7.1 and 7.2. Indeed, the only other ingredient we require is a soft monotonicity property for Bernoulli Gibbs measures from Corwin and Hammond (2014). This approach can be adapted to work for all the different integrable models of RSK. In the following theorem, we use the piecewise linear embedding of Bernoulli walks in the space of continuous functions, see Figure 2.

Theorem 1.5. *Let $Y \in \mathcal{D}^n$ consist of independent Bernoulli random walks of drift $d \in [0, 1]^n$. Then the law of $WY \in \mathcal{D}^n$ is nonintersecting Bernoulli walks with drift given by $d^\circ = (d_1^\circ \geq \dots \geq d_n^\circ)$, the components of d sorted in decreasing order.*

While Theorem 1.5 is not new (i.e. it follows from results of O’Connell (2003a) or a combination of the ideas of O’Connell (2003b) and König, O’Connell and Roch (2002)) we believe the proof we present is.

There are several subsets $A \subset \mathcal{D}^n$ of paths so that RSK is a bijection between $\mathcal{D}_\downarrow^n \cap A$ and $\mathcal{D}_\uparrow^n \cap A$. The following table informally summarizes these sets and natural measures on them. The measures then correspond to classical integrable last passage percolation models, see Sections 5 and 7. “Unit jumps” means piecewise constant functions that have jumps of size 1 only; Bernoulli paths

are continuous and linear of slope 0 or 1 on $[j, j + 1]$, $j \in \{0, 1, 2, \dots\}$ and the S-J model is the Seppäläinen-Johansson model, see Seppäläinen (1998). In all of these cases, the image under W of the natural measure on paths can be interpreted as the same measure conditioned to fall in \mathcal{D}_\uparrow^n .

A	independent walk measure on \mathcal{D}^n	LPP model
continuous functions	Brownian motions	Brownian LPP
unit jumps	Poisson counting processes	Poisson lines LPP
\mathbb{N} jumps at \mathbb{N} times	discrete-time geometric random walks	geometric LPP
\mathbb{R}^+ jumps at \mathbb{N} times	discrete-time exponential random walks	exponential LPP
Bernoulli paths	piecewise linear Bernoulli walks	S-J model

These examples also show how versions of classical RSK embed into the present framework of RSK. More precisely, usual RSK corresponds to piecewise constant nonnegative integer jumps at integer times, dual RSK corresponds to Bernoulli paths, and continuous RSK along the lines of Biane, Bougerol and O’Connell (2005) embeds as continuous paths. See Section 8 for proofs and more details.

Background

A version of RSK first appeared in Robinson (1938). The bijection was later extended in Schensted (1961), and in Knuth (1970). Classical treatments of RSK can be found in Stanley (1999), Fulton (1997), Romik (2015) and Sagan (2013). Schensted (1961) and Greene (1974) tied RSK to longest increasing subsequences, and therefore last passage percolation, see Vershik and Kerov (1977) and Logan and Shepp (1977). We use Greene’s description as the *definition* of RSK. An independent line of research started with the discovery of Pitman’s $2M - X$ theorem, Pitman (1975). The two ideas were first unified in depth and in great generality in Biane, Bougerol and O’Connell (2005), building on ideas from Bougerol and Jeulin (2002); O’Connell (2003a,b); O’Connell and Yor (2002) and other works. That work has versions of many of the results presented here, and is rooted in representation theory – part of our goal is to give a treatment where concepts of representation theory are not prerequisite.

Versions of the isometry property and its strengthening (Proposition 3.12.(i)) were shown in Noumi and Yamada (2002), and also in Biane, Bougerol and O’Connell (2005), and Dauvergne, Ortmann and Virág (2018).

There are other generalizations for RSK. Geometric RSK is a finite temperature version of ordinary RSK initiated by Kirillov (2001), see also Noumi and Yamada (2002), Corwin, O’Connell, Seppäläinen and Zygouras (2014). Noumi and Yamada (2002) have finite and zero temperature versions of many of the results presented here, obtained using matrix methods. In particular, isometry in the geometric setting was shown in Noumi and Yamada (2002), see also Corwin (2020) and Dauvergne (2020).

Further extensions, including randomized versions are studied in O’Connell and Pei (2013), Bufetov and Matveev (2018), Garver, Patrias and Thomas (2018), Aigner and Frieden (2020), and Dauvergne (2020).

2. Percolation across cadlag functions

2.1. Basic definitions

Recall that a function f from an interval I to \mathbb{R} is **cadlag** if for all $x \in I$, we have

$$\lim_{y \rightarrow x^+} f(y) = f(x), \quad \text{and} \quad \lim_{y \rightarrow x^-} f(y) \text{ exists.} \quad (2.1)$$

Note that either one of these limits may not be defined if x is an endpoint of I . We write $f(x^-)$ for the second limit in (2.1). When $f(x^-) \neq f(x)$, we say that $f(x) - f(x^-)$ is a **jump** of f and that x is a **jump location**. Cadlag functions can only have countably many jumps. Let \mathcal{D}^n be the space of all functions

$$f : [0, \infty) \times \{1, \dots, n\} \rightarrow \mathbb{R}, \quad (x, i) \mapsto f_i(x).$$

so that each f_i is a cadlag function whose jumps are all positive and satisfies $f_i(0) \geq 0$. We impose that $f_i(0^-) = 0$ for all i . If $f_i(0) > 0$, we interpret this as f having a jump at 0. The boundary condition $f_i(0^-) = 0$ is simply a convention for us since we will only care about the increments of f . We will often think of f as a sequence of functions f_1, \dots, f_n . When visualizing f we will think in matrix coordinates, so that line 1 (i.e. the function f_1) is on top and line n is on the bottom, see Figure 1.

We associate to any $f \in \mathcal{D}^n$ a finitely additive signed measure df on $[0, \infty) \times \{1, \dots, n\}$ given by

$$df([x, y] \times \{i\}) = f_i(y) - f_i(x^-)$$

for $x \leq y$ and $i \in \{1, \dots, n\}$. Our boundary convention $f_i(0^-) = 0$ means that we can always reconstruct $f \in \mathcal{D}^n$ from its measure df .

Now, for $p = (x, n), q = (y, m) \in [0, \infty) \times \mathbb{Z}$ with $x \leq y, n \geq m$, a **path** π from p to q is a union of closed intervals

$$[t_i, t_{i-1}] \times \{i\} \subset \mathbb{R} \times I, \quad i = m, m+1, \dots, n, \quad (2.2)$$

where

$$x = t_n \leq t_{n-1} \leq \dots \leq t_m \leq t_{m-1} = y, \quad (2.3)$$

see Figure 1. The points $t_i, i = m, \dots, n-1$ are called the **jump times** of π . For $f \in \mathcal{D}^n$ and a path π contained in $\mathbb{R} \times \{1, \dots, n\}$, we can define the **length** of π with respect to f by

$$|\pi|_f = df(\pi) = \sum_{i=m}^n f_i(t_{i-1}) - f_i(t_i^-).$$

This definition is chosen so that all the jumps of f that lie along the path π are accounted for. For $f \in \mathcal{D}^n$ and $u = (p; q) = (x, n; y, m) \in (\mathbb{R} \times \{1, \dots, n\})^2$ define the **last passage value** across f from p to q by

$$f[u] = f[p \rightarrow q] = f[(x, n) \rightarrow (y, m)] = \sup_{\pi} |\pi|_f, \quad (2.4)$$

where the supremum is taken over all paths π from p to q . If no path from p to q exists (i.e. if $p \not\curvearrowright q$), we set $f[u] = -\infty$. We call a path π from p to q a **geodesic** if $|\pi|_f = f[p \rightarrow q]$.

2.2. Multiple paths

Next, we generalize last passage values to multiple paths. Recall that we think in matrix coordinates, so that line 1 is on top and line n is on the bottom. First, for two paths π and ρ , we say that π is **to the left of** ρ and write $\pi \leq \rho$ if for every $(x, \ell) \in \pi$ there exists $(y, m) \in \rho$ such that $\ell \leq m$ and $x \leq y$. This defines a partial order on paths. Equivalently, we say that ρ is **to the right of** π . Pictorially, if ρ is a path from (x, n) to (y, m) , then π is to the left of ρ if it lies in the region of the plane bounded on the right by ρ and $\{y\} \times \mathbb{R}$, and bounded below (in matrix coordinates) by ρ and $\mathbb{R} \times \{n\}$. We say π, ρ are **essentially disjoint** if the set $\pi \cap \rho$ is finite. See Figure 1 for examples of all these definitions.

Let $\mathbf{p} = (p_1, \dots, p_k), \mathbf{q} = (q_1, \dots, q_k) \in ([0, \infty) \times \mathbb{Z})^k$. A **disjoint k -tuple (of paths)** $\pi = (\pi_1, \dots, \pi_k)$ from \mathbf{p} to \mathbf{q} is defined by the following properties:

- π_i is a path from p_i to q_i ,
- π_i is to the left of π_j for $i < j$,
- π_i and π_j are essentially disjoint for all $i \neq j$.

For $f \in \mathcal{D}^n$ and a disjoint k -tuple π let $\cup \pi := \pi_1 \cup \dots \cup \pi_k \subset [0, \infty) \times \{1, \dots, n\}$, and define the **length** of π by

$$|\pi|_f = df(\cup \pi).$$

For $\mathbf{u} = (\mathbf{p}, \mathbf{q})$, we can then define the **multi-point last passage value**

$$f[\mathbf{u}] = f[\mathbf{p} \rightarrow \mathbf{q}] = \sup_{\pi} |\pi|_f,$$

where the supremum is over disjoint k -tuples from \mathbf{p} to \mathbf{q} . Again, if no such k -tuples exist, we set $f[\mathbf{u}] = -\infty$. We call a k -tuple π satisfying $f[\mathbf{p} \rightarrow \mathbf{q}] = |\pi|_f$ a **disjoint optimizer**.

2.3. Basic geometric properties

Next, we collect some basic geometric facts about last passage paths. We start by showing that disjoint optimizers always exist. As in the previous section, $f \in \mathcal{D}^n$ and let $\mathbf{p} = (p_1, \dots, p_k), \mathbf{q} = (q_1, \dots, q_k) \in ([0, \infty) \times \mathbb{Z})^k$.

Lemma 2.1. *If there is at least one disjoint k -tuple from \mathbf{p} to \mathbf{q} , then there exists a disjoint optimizer for f from \mathbf{p} to \mathbf{q} .*

Lemma 2.1 is an immediate consequence of the following two observations. Each of these next lemmas is also useful in its own right.

Lemma 2.2 (Compactness). *The space of disjoint k -tuples from \mathbf{p} to \mathbf{q} is compact in the Hausdorff topology on $([0, \infty) \times \{1, \dots, n\})^k$.*

Proof. Let $[a, b]$ be any interval such that $p_i, q_i \in [a, b] \times \{1, \dots, n\}$ for all i . The space of disjoint k -tuples from \mathbf{p} to \mathbf{q} is a subset of the space of closed sets in the compact metric space $([a, b] \times \{1, \dots, n\})^k$ endowed with the Hausdorff topology. The Hausdorff topology on closed sets in any compact metric space is compact, e.g. see Henrikson (1999). Moreover, it is easy to check that the property of being a disjoint k -tuple from \mathbf{p} to \mathbf{q} is closed under Hausdorff limits. \square

Lemma 2.3 (Upper semicontinuity). *For any $f \in \mathcal{D}^n$, the function $\pi \mapsto |\pi|_f$ mapping disjoint k -tuples to their f -length is upper semicontinuous in the Hausdorff topology.*

Lemma 2.3 is a consequence of the fact that if $a_m \rightarrow a, b_m \rightarrow b$, then since $f \in \mathcal{D}^n$ has only positive jumps,

$$\limsup_{m \rightarrow \infty} df([a_m, b_m] \times \{i\}) \leq df([a, b] \times \{i\}).$$

Next, we give a metric composition law. The proof is again immediate from the definitions. For this lemma and in the sequel, we use the shorthand notation

$$(t, \mathbf{m}) = \left((t, m_1), \dots, (t, m_k) \right) \quad \text{and similarly} \quad (\mathbf{x}, \ell) = \left((x_1, \ell), \dots, (x_k, \ell) \right).$$

Lemma 2.4 (Metric composition law). *Let $f \in \mathcal{D}^n$, let $(\mathbf{p}, \mathbf{q}) = (\mathbf{x}, \ell; \mathbf{y}, m) \in ([0, \infty) \times \{1, \dots, n\})^{2k}$ and let $i \in \{m+1, \dots, \ell\}$. Then*

$$f[\mathbf{p} \rightarrow \mathbf{q}] = \max_{\mathbf{z} \in [0, \infty)^k} f[\mathbf{p} \rightarrow (\mathbf{z}, i)] + f[(\mathbf{z}, i-1) \rightarrow \mathbf{q}].$$

More general versions of the metric composition law exist, though we do not require them here. Lemma 2.4 implies certain triangle inequalities for last passage values, reinforcing the idea that the last passage structure is best thought of as a metric.

We end this section with an extremely useful **quadrangle inequality** for multi-point last passage values. This inequality generalizes a well-known quadrangle inequality for single-path last passage values, e.g. see Proposition 3.8 in Dauvergne, Ortman and Virág (2018).

Lemma 2.5. *Let $(\mathbf{p}, \mathbf{q}) = (\mathbf{x}, m; \mathbf{y}, \ell), (\mathbf{p}', \mathbf{q}') = (\mathbf{x}', m; \mathbf{y}', \ell) \in ([0, \infty) \times \{1, \dots, n\})^{2k}$ be such that $x_i \leq x'_i, y_i \leq y'_i$ for all i . Then*

$$f[\mathbf{p} \rightarrow \mathbf{q}'] + f[\mathbf{p}' \rightarrow \mathbf{q}] \leq f[\mathbf{p} \rightarrow \mathbf{q}] + f[\mathbf{p}' \rightarrow \mathbf{q}'].$$

This is a special case of Lemma 2.4 in Dauvergne and Zhang (2021) generalized to the cadlag setting.

Proof. Let π be a disjoint optimizer from \mathbf{p} to \mathbf{q}' , and let π' be a disjoint optimizer from \mathbf{p}' to \mathbf{q} . Let $t_{i,m}, \dots, t_{i,\ell-1}, t'_{i,m}, \dots, t'_{i,\ell-1}$ be the jump times specified in (2.3) for the paths π_i, π'_i . For each i , let τ_i^L be the path from p_i to q_i with jump times $t_{i,m} \wedge t'_{i,m}, \dots, t_{i,\ell-1} \wedge t'_{i,\ell-1}$, and let τ_i^R be the path from p'_i to q'_i with jump times $t_{i,m} \vee t'_{i,m}, \dots, t_{i,\ell-1} \vee t'_{i,\ell-1}$. We can think of τ_i^L, τ_i^R

as taking π_i, π'_i and sorting them into a leftmost path and a rightmost path. With this construction, it is straightforward to check that τ_i^L and τ_i^R are paths from p_i to q_i and p'_i to q'_i , respectively. Moreover, for every $j \in \{\ell - 1, \dots, m\}$ the points $t_{i,j} \wedge t'_{i,j}$ and $t_{i,j} \vee t'_{i,j}$ are increasing in i since both $t_{i,j}$ and $t'_{i,j}$ are increasing in i . Therefore the paths $\tau_1^L, \dots, \tau_k^L$ and $\tau_1^R, \dots, \tau_k^R$ are ordered from left to right.

To conclude that τ^L is a disjoint k -tuple from \mathbf{p} to \mathbf{q} and τ^R is a disjoint k -tuple from \mathbf{p}' to \mathbf{q}' we just need to check essential disjointness. We do this for τ^L ; the proof for τ^R is identical. Fix $i < i^*$. To check that τ_i^L and $\tau_{i^*}^L$ are essentially disjoint, it is enough to show that for all $j \in \{\ell, \dots, m\}$ we have $t_{i,j-1} \wedge t'_{i,j-1} \leq t_{i^*,j} \wedge t'_{i^*,j}$. This follows from the inequalities $t_{i,j-1} \leq t_{i^*,j}$ and $t'_{i,j-1} \leq t'_{i^*,j}$, which are guaranteed by the essential disjointness and ordering of π_i, π_{i^*} .

To complete the proof, we just need to show that

$$|\pi|_f + |\pi'|_f \leq |\tau^L|_f + |\tau^R|_f. \quad (2.5)$$

Indeed, the left side of (2.5) equals $f[\mathbf{p} \rightarrow \mathbf{q}'] + f[\mathbf{p}' \rightarrow \mathbf{q}]$ and the right side is less than or equal to $f[\mathbf{p} \rightarrow \mathbf{q}] + f[\mathbf{p}' \rightarrow \mathbf{q}']$. We can write

$$|\pi|_f + |\pi'|_f = df((\cup\pi) \cup (\cup\pi')) + df((\cup\pi) \cap (\cup\pi')),$$

and similarly for $|\tau^L|_f + |\tau^R|_f$. By construction, it is easy to see that $(\cup\pi) \cup (\cup\pi') = (\cup\tau^L) \cup (\cup\tau^R)$. Therefore to prove (2.5) it is enough to show that $(\cup\pi) \cap (\cup\pi') \subset (\cup\tau^L) \cap (\cup\tau^R)$ and that the difference $S := [(\cup\tau^L) \cap (\cup\tau^R)] \setminus [(\cup\pi) \cap (\cup\pi')]$ is a finite set, since df assigns nonnegative weight to all finite sets.

Suppose that a point $p = (x, j)$ is contained in both $\cup\pi$ and $\cup\pi'$. Then for some i, i^* we have $p \in \pi_i$ and $p \in \pi'_{i^*}$. Without loss of generality, assume $i \leq i^*$. Then by the ordering of the paths in π, π' , we have

$$t_{i,j} \leq z \leq t_{i,j-1} \leq t_{i^*,j-1}, \quad t'_{i,j} \leq t'_{i^*,j} \leq z \leq t'_{i^*,j-1}.$$

These inequalities show that $p \in \tau_i^R, p \in \tau_{i^*}^L$. Next, observe that $S \cap (\cup\pi)$ is always a union of sets of the form $I \times \{j\}$, where I is an interval in \mathbb{R} and $I \times \{j\}$ is a subset of some particular π_i or π'_i . Therefore if S is infinite, without loss of generality there exists $i \in \{1, \dots, k\}, j \in \{m, \dots, \ell\}$ and $a < b$ such that $[a, b] \times \{j\} \subset \pi_i \cap S$. By essential disjointness of π , we have $(a, b) \times \{j\} \notin \pi_{i^*}$ for all $i^* \neq i$. Therefore $(a, b) \times \{j\} \notin \tau_{i^*}^L \cup \tau_{i^*}^R$ for all $i^* \neq i$, so $(a, b) \times \{j\} \subset \tau_i^L \cap \tau_i^R$. This implies that $(a, b) \times \{j\} \subset \pi_i \cap \pi'_i$, contradicting that $(a, b) \times \{j\} \subset S$. \square

A similar proof idea to Lemma 2.5 shows that rightmost and leftmost optimizers always exist. Again, this is a generalization of Lemma 2.2 in Dauvergne and Zhang (2021) to the cadlag setting. To state the lemma, for disjoint k -tuples λ, π , we write $\lambda \leq \pi$ and say that λ is **to the left** of π if λ_i is to the left of π_i for every i .

Lemma 2.6. *Let $(\mathbf{p}, \mathbf{q}) = (\mathbf{x}, n; \mathbf{y}, m)$ be such that there is at least one disjoint k -tuple from \mathbf{p} to \mathbf{q} . Then for any $f \in \mathcal{D}^n$, there are unique optimizers ρ, λ*

across f from \mathbf{p} to \mathbf{q} such that for any optimizer π from \mathbf{p} to \mathbf{q} , we have $\rho \leq \pi \leq \lambda$. We call ρ, λ the **rightmost** and **leftmost** optimizers from \mathbf{p} to \mathbf{q} .

Proof. Consider the set S of all optimizers from \mathbf{p} to \mathbf{q} with the partial order \leq . Let T be a totally ordered subset of S in the order \leq . We claim that T has a lower bound in S . Any k -tuple π in T is characterized by the jump times $\pi_{i,j}, i \in \{m-1, \dots, n\}$ of each of the constituent paths π_i , see (2.3). For all i, j define $\pi_{i,j}^* = \inf\{\pi_{i,j} : \pi \in T\}$. Since T is totally ordered, it is straightforward to check that these points are jump times for a k -tuple of paths π^* , and that π^* is a Hausdorff limit of k -tuples in T . Lemma 2.2 then implies that π^* is a disjoint k -tuple from \mathbf{p} to \mathbf{q} , and Lemma 2.3 implies that π^* is an optimizer. Hence π^* is a lower bound for T contained in S .

Therefore by Zorn's lemma, S contains at least one minimal element. Suppose that π, π' are both minimal elements. Construct disjoint k -tuples τ^L, τ^R from \mathbf{p} to \mathbf{q} as in the proof of Lemma 2.5 with $(\mathbf{p}, \mathbf{q}) = (\mathbf{p}', \mathbf{q}')$ so that $\tau^L \leq \pi, \pi'$ and

$$2f[\mathbf{p} \rightarrow \mathbf{q}] = |\pi|_f + |\pi'|_f \leq |\tau^L|_f + |\tau^R|_f. \quad (2.6)$$

Since $|\tau^L|_f, |\tau^R|_f \leq f[\mathbf{p} \rightarrow \mathbf{q}]$ by definition of $f[\mathbf{p} \rightarrow \mathbf{q}]$, (2.6) can only hold if both τ^R, τ^L are optimizers from \mathbf{p} to \mathbf{q} . Therefore $\tau^L \in S$ so by the minimality of π, π' we have $\pi = \tau^L = \pi'$. Therefore S contains a unique minimal element, the leftmost optimizer. The existence of the rightmost optimizer follows by a symmetric argument. \square

3. The melon

Recall that the melon $Wf \in \mathcal{D}^n$ of a function $f \in \mathcal{D}^n$ is given by

$$Wf_k(y) = f[(0, n)^k \rightarrow (y, 1)^k] - f[(0, n)^{k-1} \rightarrow (y, 1)^{k-1}]. \quad (3.1)$$

Here recall that for $p \in \mathbb{R} \times \{1, \dots, n\}$ we write $p^k = (p, \dots, p) \in (\mathbb{R} \times \{1, \dots, n\})^k$. We use the convention that $f[p^0 \rightarrow q^0] = 0$. We say that $f, g \in \mathcal{D}^n$ are **isometric** if

$$f[(\mathbf{x}, n) \rightarrow (\mathbf{y}, 1)] = g[(\mathbf{x}, n) \rightarrow (\mathbf{y}, 1)]$$

for all k -tuples \mathbf{x}, \mathbf{y} . In other words, last passage values between the top and bottom boundaries in the environments defined by f and g are equal. Isometry is an equivalence relation on \mathcal{D}^n , which we denote by $f \sim g$.

The main goal of the section is to show that f is isometric to Wf . To do this, we will show that Wf agrees with iterated applications of 2-line melon maps to f , alternately known as **Pitman transforms**.

3.1. The Pitman transform

From the definition of the melon map for $n = 2$ we immediately get the following result.

Lemma 3.1 (The Pitman transform). *For $f \in \mathcal{D}^2$, we have*

$$Wf_1(x) = f[(0, 2) \rightarrow (x, 1)] \quad (3.2)$$

$$Wf_2(x) = f_1(x) + f_2(x) - Wf_1(x). \quad (3.3)$$

The terminology Pitman transform arises from considering the special case when $f = B$ is a standard Brownian motion on \mathbb{R}^2 . In this case, the 2M-X theorem of Pitman (1975) characterizes the law of WB : it implies that $WB_1 - WB_2$ is a Bessel-3 process, independent of the Brownian motion $WB_1 + WB_2 = B_1 + B_2$. We will return to this setting when we consider cadlag RSK with random input in Section 7.

Our main goal in Section 3.1 is to prove that in the 2-line case, $Wf \sim f$. The first step is Lemma 3.2, which shows that W preserves single-path last passage values. This has the most technical proof in the paper, and consists of careful manipulations of the definitions. The proof is complicated by the fact that the function f is only cadlag, rather than continuous, see Lemma 4.2 in Dauvergne, Ortmann and Virág (2018) for the easier continuous case. We defer the proof to Appendix A.

Lemma 3.2. *The map W maps \mathcal{D}^2 to itself. Moreover, for $f \in \mathcal{D}^2$ and $0 \leq x \leq y$, we have that*

$$f[(x, 2) \rightarrow (y, 1)] = Wf[(x, 2) \rightarrow (y, 1)] \quad (3.4)$$

Lemma 3.2 and the definition of W allow us to identify both the image of the two-line map W , and the fact that it is idempotent.

Definition 3.3. We say that two functions $f_1, f_2 \in \mathcal{D}$ are **Pitman ordered**, if

$$f_2(t) \leq f_1(t^-) \quad \text{for all } t.$$

When this is the case, we will write $f_2 \preceq f_1$.

Remark 3.4. With this definition the set \mathcal{D}_\uparrow^n introduced in Theorem 1.1 (iii) can be written simply as $\mathcal{D}_\uparrow^n = \{f \in \mathcal{D}^n : f_n \preceq \dots \preceq f_1\} \subset \mathcal{D}^n$. Since the lines are ordered in this way, it is clear that for $f \in \mathcal{D}_\uparrow^n$ there is a disjoint optimizer from $(0, n)^k \rightarrow (y, 1)^k$ following the leftmost possible paths, i.e. $f_1(x) + \dots + f_k(x) = f[(0, n)^k \rightarrow (x, 1)^k]$. In short,

$$Wf = f \quad (3.5)$$

for $f \in \mathcal{D}_\uparrow^n$.

Lemma 3.5. *Let $f = (f_1, f_2) \in \mathcal{D}^2$. The following are equivalent:*

- (i) $(Wf)_1 = f_1$,
- (ii) $(Wf)_2 = f_2$,
- (iii) $Wf = f$,
- (iv) $f_2 \preceq f_1$, i.e. $f \in \mathcal{D}_\uparrow^2$.

Moreover, the Pitman transform is idempotent, $W^2 = W$ and the image of the Pitman transform is precisely the set of all Pitman ordered functions:

$$W(\mathcal{D}^2) = \mathcal{D}_\uparrow^2 = \{f \in \mathcal{D}^2 : f_2 \preceq f_1\}.$$

Proof. By Lemma 3.1, $Wf_1 + Wf_2 = f_1 + f_2$. This gives the equivalence of (i), (ii) and (iii). For (iv), observe that $f_2 \preceq f_1$ if and only if

$$f[(0, 2) \rightarrow (x, 1)] = f_1(x) + \sup_{0 \leq y \leq x} f_2(y) - f_1(y^-) = f_1(x)$$

for all x , or in other words $Wf_1 = f_1$, giving the equivalence of (i) and (iv). Lemma 3.2 then implies the claims about idempotency and image. \square

Next, we extend Lemma 3.2 to deal with disjoint k -tuples. For the proof, we need the notion of a path from p^+ to q . This is defined the same way as the path π from $p = (x, n)$ to $q = (y, m)$ as in (2.2), with the vertical line $\{x\} \times \{m, \dots, n\}$ removed. The last passage value $f[p^+ \rightarrow q]$ is the supremum of $|\pi|_f$ over all paths π from p^+ to q . Analogously, we define paths from p to q^- and p^+ to q^- paths, and last passage values $f[p \rightarrow q^-]$ and $f[p^+ \rightarrow q^-]$. We can think of the last passage value $f[p \rightarrow q]$ as $f[p^- \rightarrow q^+]$.

Corollary 3.6. *Let $p = (x, 2)$, $q = (y, 1)$ with $0 \leq x < y$. Then*

$$\begin{aligned} Wf[p^+ \rightarrow q] &= f[p^+ \rightarrow q], & Wf[p \rightarrow q^-] &= f[p \rightarrow q^-], \\ Wf[p^+ \rightarrow q^-] &= f[p^+ \rightarrow q^-]. \end{aligned}$$

Proof. Since $\lim_{z \rightarrow x^+} Wf$ and $\lim_{z \rightarrow x^+} f$ both exist, we have

$$\begin{aligned} f[p^+ \rightarrow q] &= \lim_{z \rightarrow x^+} f[(z, 2) \rightarrow q], & \text{and} \\ Wf[p^+ \rightarrow q] &= \lim_{z \rightarrow x^+} Wf[(z, 2) \rightarrow q]. \end{aligned}$$

Lemma 3.2 then implies the first claim. The others are proven analogously, using that the functions Wf, f also have left limits. \square

Lemma 3.7 (Isometry of the Pitman transform). *For $f \in \mathcal{D}^2$, $Wf \sim f$. That is, for every pair of k -tuples $(\mathbf{p}, \mathbf{q}) = (\mathbf{x}, 2; \mathbf{y}, 1)$, we have*

$$f[\mathbf{p} \rightarrow \mathbf{q}] = Wf[\mathbf{p} \rightarrow \mathbf{q}].$$

Proof. Define $N : \mathbb{R} \rightarrow \{0, 1, \dots\}$ by $N(z) = \#\{i : z \in [x_i, y_i]\}$. Disjointness of a k -tuple π from \mathbf{p} to \mathbf{q} implies the following.

- Let $Z = N^{-1}(\{2, \dots\})$, that is the set of points z that are contained in at least two intervals $[x_i, y_i]$. Then $Z \times \{1, 2\} \subset \cup \pi$.
- Let I_1, \dots, I_ℓ be the connected components of $N^{-1}(1)$, let $\hat{a}_j \leq \hat{b}_j$ be the endpoints of I_j and set $a_j = (\hat{a}_j, 2), b_j = (\hat{b}_j, 1)$. Then $I_j \times \{1, 2\} \cap (\cup \pi)$ is a path from a_j^* to b_j^* for all j . Here $*$ $\in \{+, -\}$, depending on whether the corresponding end of I_j is open or closed.

- $(\cup\pi) \cap (N^{-1}(\{0\}) \times \{1, 2\}) = \emptyset$.

Therefore letting ρ_i be the path $I_j \times \{1, 2\} \cap (\cup\pi)$ from a_j^* to b_j^* , we can write

$$\cup\pi = Z \times \{1, 2\} \cup \bigcup_{i=1}^{\ell} \rho_i. \quad (3.6)$$

Moreover, given arbitrary $a_i^* \rightarrow b_i^*$ paths ρ_i , there is a $\mathbf{p} \rightarrow \mathbf{q}$ path π so that (3.6) holds. Therefore

$$f[\mathbf{p} \rightarrow \mathbf{q}] = df(Z \times \{1, 2\}) + \sum_{i=1}^{\ell} f[a_i^* \rightarrow b_i^*].$$

Since $f_1 + f_2 = Wf_1 + Wf_2$, the first term does not change when we apply W . By Corollary 3.6, nor does the remaining sum. \square

Before moving beyond the two-line case, we note that following simple monotonicity for later use:

$$Wf_2 \leq f_2 \wedge f_1 \leq f_1 \vee f_2 \leq Wf_1. \quad (3.7)$$

3.2. Repeated Pitman transforms, cars, and the melon

Next, we build the n -line melon map. We first extend the Pitman transform to functions $f \in \mathcal{D}^n$ by applying it to two lines at a time. For $i \in \{1, \dots, n-1\}$, define $\sigma_i : \mathcal{D}^n \rightarrow \mathcal{D}^n$ by

$$\sigma_i f = (f_1, \dots, f_{i-1}, W(f_i, f_{i,i+1})_1, W(f_i, f_{i,i+1})_2, f_{i+2}, \dots, f_n).$$

Proposition 3.8 (Isometry property of σ_i). *Let $f \in \mathcal{D}^n$, $n, k \in \mathbb{N}$, and let $(\mathbf{p}, \mathbf{q}) = (\mathbf{x}, n; \mathbf{y}, 1)$. Then for $m \in \{1, \dots, n-1\}$, we have*

$$f[\mathbf{p} \rightarrow \mathbf{q}] = \sigma_m f[\mathbf{p} \rightarrow \mathbf{q}]. \quad (3.8)$$

Proof. We first assume $m \in \{2, \dots, n-2\}$. By the metric composition law, Lemma 2.4, applied twice, we can write

$$\begin{aligned} f[\mathbf{p} \rightarrow \mathbf{q}] &= \sup_{\mathbf{z}, \mathbf{w} \in \mathbb{R}^k} \left(f[\mathbf{p} \rightarrow (\mathbf{z}, m+2)] \right. \\ &\quad \left. + f[(\mathbf{z}, m+1) \rightarrow (\mathbf{w}, m)] + f[(\mathbf{w}, m-1) \rightarrow \mathbf{q}] \right). \end{aligned} \quad (3.9)$$

For a fixed pair (\mathbf{z}, \mathbf{w}) , when we apply σ_m to f , the first and third terms under the supremum in (3.9) do not change since the relevant components f_i do not change. The middle term is preserved under the transformation σ_m by Lemma 3.7. Hence the right hand side of (3.9) is also preserved under the map $f \mapsto \sigma_m f$. The cases $m = 1, n-1$ are similar with one of the terms in (3.9) removed. \square

For $m \leq n$, define $\tau_m : \mathcal{D}^n \rightarrow \mathcal{D}^n$ by

$$\tau_m = \sigma_m \sigma_{m+1} \cdots \sigma_{n-1}.$$

We will build the n -line Pitman transform from composing the maps τ_i .

Remark 3.9 (Cars). The maps τ_m can be used to give a connection between particle systems and last passage percolation as follows. The functions $t \mapsto \tau_k f_k(t)$, with $k = 1, \dots, n$ can be thought of as deterministic versions of the totally asymmetric simple exclusion process, tasep, or equivalently as a simple model of traffic flow.

Informally, think of a line of n cars on a single-lane highway moving in the same, typically negative direction. The cars cannot pass each other. Letting $c_1(t), \dots, c_n(t)$ denote the positions of these cars, we will define dynamics where $c_n \leq c_{n-1} \leq \cdots \leq c_1$ so that car n is at the front of the line and car 1 is at the back.

The derivative $f'_k(t)$ is the desired velocity of car k at time t (for the heuristic picture it may be helpful to think of all the f_k as differentiable with $f'_k \leq 0$). However, cars cannot always move at their desired velocity. These are conscientious and focused drivers, so they will always slow down to avoid hitting the car immediately in front (i.e. creating a situation where $c_i(t) < c_{i+1}(t)$). On the other hand, cars ignore cars behind them.

Since car n is at the front of the line, it always moves at its desired velocity and so $c_n(t) = f_n(t) = \tau_n f_n(t)$. The condition that the remaining cars slow down exactly according to the movements of the car immediately in front gives us the recursive equation

$$c_k(t) = \sup_{s \in [0, t]} f_k(t) - f_k(s^-) + c_{k+1}(s),$$

which amounts to saying that $c_k = (\sigma_k \tau_{k-1} f)_k = \tau_k f_k$. If we introduce the possibility that the functions f_k may have jumps or positive derivatives, then while the traffic flow interpretation may not be as concrete, the mathematical model still makes sense. The key feature of these models is that the *interaction*, rather than the direction of movement, is totally asymmetric.

A different model for the movement of the cars is that they ignore the cars ahead of them and are forced to speed up to avoid collisions from cars behind them. For this model, the location of car k is then given by $\sigma_{k-1} \cdots \sigma_1 f_k(t)$. This version has slight differences in the setting where the f_i s have jumps; it corresponds to push-asep and a type of first passage percolation.

Stochastic models that fit in this setting include tasep, discrete-time tasep, push-asep (which also has totally asymmetric interactions), Brownian tasep and the Hammersley process.

In light of the remark, the following lemma states a deterministic equivalence between exclusion processes and last passage percolation.

Lemma 3.10. *Let $f \in \mathcal{D}^n$ and $1 \leq m \leq n \in \mathbb{N}$. Then for $y \geq 0$, we have*

$$f[(0, n) \rightarrow (y, m)] = \tau_m f_m(y). \quad (3.10)$$

Proof. We show this by induction on $n - m$. The $n = m$ case is true by definition of $\tau_n = \text{id}$. For $m < n$, we have by the metric composition law, Lemma 2.4,

$$\begin{aligned} f[(0, n) \rightarrow (y, m)] &= \sup_{z \leq y} f[(0, n) \rightarrow (z, m+1)] - f_m(z^-) + f_m(y) \\ &= \sup_{z \leq y} \tau_{m+1} f_{m+1}(z) - f_m(z^-) + f_m(y) \\ &= \sigma_m \tau_{m+1} f_m(y), \end{aligned}$$

where the second equality follows from the inductive hypothesis, and the third equality is simply the definition of σ_m . By the definition of τ_m this equals the right hand side of (3.10). \square

Definition 3.11. We will build the n -line map $\omega : \mathcal{D}^n \rightarrow \mathcal{D}^n$ from the functions τ_m as follows. Define

$$\omega = \tau_{n-1} \tau_{n-2} \cdots \tau_1 = \sigma_{n-1} \sigma_{n-2} \sigma_{n-1} \cdots \sigma_1 \sigma_2 \cdots \sigma_{n-1}. \quad (3.11)$$

Informally speaking, if we think of σ_i as “sorting” the i -th and $i+1$ -st functions f_i and f_{i+1} , then ω is precisely the “bubble sort” algorithm to sort the entire ensemble f .

In the following proposition we show that $\omega = W$, the melon map from (3.1).

Proposition 3.12. *Let ωf be the melon of a function $f \in \mathcal{D}^n$. Then:*

- (i) (Isometric equivalence) $\omega f \sim f$.
- (ii) (Idempotence) $\omega^2 = \omega$.
- (iii) (Image) We have $\omega(\mathcal{D}^n) = \mathcal{D}_\uparrow^n = \{f \in \mathcal{D}^n : f_n \leq \cdots \leq f_1\}$.
- (iv) $\omega = W$ from (3.1).

To prove Proposition 3.12 we need two short lemmas.

Lemma 3.13. *Let $m < n \in \mathbb{N}$. Then $\tau_m^2 = \tau_{m+1} \tau_m$.*

Proof. Let $g = \tau_{m+1} \tau_m f$. We want to show that $g = \sigma_m g$. The functions g and $\sigma_m g$ can only differ in the coordinates $m, m+1$. By Lemma 3.5 it suffices to show that $g_m = \sigma_m g_m$.

By Lemma 3.10, we have

$$\sigma_m g_m(y) = \tau_m \tau_m f_m(y) = \tau_m f[(0, n) \rightarrow (y, m)] = f[(0, n) \rightarrow (y, m)],$$

where the last equality is by Proposition 3.8 repeatedly applied to (f_m, \dots, f_n) . Since τ_{m+1} does not change coordinate m , Lemma 3.10 gives

$$g_m(y) = \tau_{m+1} \tau_m f_m(y) = \tau_m f_m(y) = f[(0, n) \rightarrow (y, m)]. \quad \square$$

Lemma 3.14. *For $m = 1, \dots, n-1$ we have $\sigma_m \omega = \omega$.*

Proof. The statement for $m = n-1$ follows since $\sigma_{n-1}^2 = \sigma_{n-1}$ by Lemma 3.5. Now assume $m < n-1$. When $i - j \geq 2$, the functions σ_i and σ_j commute because they act on disjoint coordinates. Thus, by Lemma 3.13 we have

$$\sigma_m \omega = \tau_{n-1} \cdots \tau_{m+2} \sigma_m \tau_{m+1} \tau_m \cdots \tau_1$$

$$\begin{aligned}
&= \tau_{n-1} \cdots \tau_{m+2} \tau_m^2 \cdots \tau_1 \\
&= \tau_{n-1} \cdots \tau_{m+2} \tau_{m+1} \tau_m \cdots \tau_1 = \omega. \quad \square
\end{aligned}$$

Proof of Proposition 3.12. Part (i) follows immediately from Proposition 3.8. Part (ii) follows from Lemma 3.14.

For (iii), Lemma 3.5 (iii) implies that ω is the identity on \mathcal{D}_\uparrow^n , so $\omega(\mathcal{D}^n) \supset \mathcal{D}_\uparrow^n$. For the other direction, note that σ_i does not change ωf by Lemma 3.14. By Lemma 3.5 (iv), this implies that $f_{i+1} \preceq f_i$ for all i , so $\omega(\mathcal{D}^n) \subset \mathcal{D}_\uparrow^n$.

Finally, for (iv),

$$Wf = W\omega f = \omega f,$$

where the first equality is a consequence of part (i), and the second is a consequence of (3.5), since $\omega f \in \mathcal{D}_\uparrow$ by (iii). \square

Remark 3.15. We mention without proof that by applying a similar framework in the $n = 3$ case, ω can alternately be defined as $\sigma_1\sigma_2\sigma_1$, instead of $\sigma_2\sigma_1\sigma_2$. This yields a braid relation for the operators σ_i , which implies that for larger n , $W = \omega = \sigma_{i_1} \cdots \sigma_{i_{\binom{n}{2}}}$ whenever the product of transpositions $(i_1 \ i_1+1) \cdots (i_{\binom{n}{2}} \ i_{\binom{n}{2}}+1)$ is a reduced word for the reverse permutation.

Remark 3.16 (Historical notes). The map ω coincides with the generalized Pitman transform Γ_n first introduced in O’Connell and Yor (2002) and studied there in the context of continuous paths and paths with unit jumps. It was subsequently studied in many further papers. O’Connell (2003b) defined the map ω for n -tuples of functions from $\mathbb{Z} \rightarrow \mathbb{Z}^+$ with positive jumps, and showed that $\omega = W$ in that setting, thereby related ω to the usual RSK correspondence. The relation between a version of ω and dual RSK is discussed in O’Connell (2003a) after Theorem 4.11. The map ω was given a representation-theoretic generalization in Biane, Bougerol and O’Connell (2005). The braid relation discussed in Remark 3.15 and a special case of the isometry property were shown in that paper in the setting of continuous functions f .

4. Minimal and maximal elements

In this section, we show that the melon map picks out the minimal element in the equivalence class of isometric environments under a certain natural preorder. Recall that a preorder is partial order without the antisymmetry requirement, i.e. $f \leq g$ and $g \leq f$ are both allowed for $f \neq g$. While this perspective is not necessary for the proofs in later sections, it is helpful for developing intuition about the melon map and its inverse.

For $f \in \mathcal{D}^n$, let $\lambda_f(\mathbf{x}, \mathbf{y}), \rho_f(\mathbf{x}, \mathbf{y})$ be the leftmost and rightmost optimizers in f from (\mathbf{x}, n) to $(\mathbf{y}, 1)$, see Lemma 2.6. We say that $f \preceq g$ if:

$$\lambda_f(\mathbf{x}, \mathbf{y}) \leq \lambda_g(\mathbf{x}, \mathbf{y}) \quad \text{and} \quad \rho_f(\mathbf{x}, \mathbf{y}) \leq \rho_g(\mathbf{x}, \mathbf{y}) \quad \text{for all } \mathbf{x}, \mathbf{y}.$$

Our goal is to prove the following.

Proposition 4.1. *For every $f \in \mathcal{D}^n$ with $f(0) = 0$, the melon Wf is the unique minimal element with respect to \trianglelefteq in the isometry class of f .*

Remark 4.2. The proof of Proposition 4.1 shows that $Wf \trianglelefteq g$ for any $f, g \in \mathcal{D}^n$ with $f \sim g$. The condition that $f(0) = 0$ is only necessary for the uniqueness statement. Indeed, consider the example where $f_i(x) = 1$ for all $x \geq 0$ so that df consists of a line of atoms at $\{0\} \times \{1, \dots, n\}$. Then $Wf_1 = n$ and $Wf_i = 0$ for all $i \geq 2$ so $Wf \neq f$. However, it is not difficult to check that $\lambda_f(\mathbf{x}, \mathbf{y}) = \lambda_{Wf}(\mathbf{x}, \mathbf{y})$, $\rho_f(\mathbf{x}, \mathbf{y}) = \rho_{Wf}(\mathbf{x}, \mathbf{y})$ for all \mathbf{x}, \mathbf{y} so we have both $f \trianglelefteq Wf$ and $Wf \trianglelefteq f$.

The main lemma needed for Proposition 4.1 concerns the Pitman transform $W : \mathcal{D}^2 \rightarrow \mathcal{D}^2$. Its proof is in a similar vein to the proof of Lemma 3.2. As a result, we defer it to the appendix where it is proven with Lemma 3.2.

Lemma 4.3. *Let $f \in \mathcal{D}^2$ and let $x \leq y$. Then*

$$\rho_{Wf}(x, y) \leq \rho_f(x, y) \quad \text{and} \quad \lambda_{Wf}(x, y) \leq \lambda_f(x, y).$$

We can extend this to general optimizers using the same ideas as in Lemma 3.7. As the proof is essentially identical, we omit it.

Lemma 4.4. *For all $f \in \mathcal{D}^2$, we have $Wf \trianglelefteq f$.*

We can extend this to σ_i by a simple application of the metric composition law. The proof is essentially identical to the proof of Proposition 3.8 so we omit it.

Lemma 4.5. *For all $f \in \mathcal{D}^n$ and $i \in \{1, \dots, n-1\}$, we have $\sigma_i f \trianglelefteq f$.*

Proof of Proposition 4.1. By Lemma 4.5 and the definition, we have $Wf \trianglelefteq f$ for all f . Now, for any g in the same isometry class as f , we have $Wf = Wg \trianglelefteq g$. Since \trianglelefteq is only a preorder, to prove that Wf is the *unique* minimal element we still need to check that if $f \trianglelefteq Wf$ then $f = Wf$.

Indeed, since Wf is Pitman ordered the leftmost optimizer $\lambda_{Wf}(0^k, x^k)$ simply follows the top k paths Wf_1, \dots, Wf_k on the interval $(0, x)$. Since no disjoint k -tuple is to the left of this optimizer, the same must be true for $\lambda_f(0^k, x^k)$ if $f \trianglelefteq Wf$. Therefore for all x, k we have

$$\sum_{i=1}^k Wf_i(x) = f[(0^k, n) \rightarrow (x^k, 1)] = \sum_{i=1}^k f_i(x) - f_i(0^-) + \sum_{i=k+1}^n f_i(0) - f_i(0^-).$$

Since $f(0^-) = 0$ by definition and $f(0) = 0$ by assumptions, this implies that $Wf = f$. \square

Just as Wf identifies a minimal element of the isometry class of f , we would also like to identify a maximal element. We can do this with a straightforward symmetric definition if we first restrict to functions defined on $[0, t]$ rather than $[0, \infty)$.

Let \mathcal{D}_t^n be the set of n -tuples of cadlag functions $f_1, \dots, f_n : [0, t] \rightarrow \mathbb{R}$ with only positive jumps. For $f \in \mathcal{D}_t^n$, we can define the truncated melon map $W_t : \mathcal{D}_t^n \rightarrow \mathcal{D}_t^n$ by (3.1). Again, we can think of $W_t f$ as picking out a distinguished minimal element in the isometry class of f in \mathcal{D}_t^n . In the finite setting, a conjugation of the map W_t produces a maximal element. For $f \in \mathcal{D}_t^n$, let $Rf \in \mathcal{D}_t^n$ be the rotation of f by 180 degrees. More precisely, we set

$$Rf_i(x) = f_{n+1-i}(t) - f_{n+1-i}((t-x)^-).$$

The function Rf is the unique element of \mathcal{D}_t^n for which the measure $d(Rf)$ is the pushforward of the measure df under the 180-degree rotation R of $[0, t] \times \{1, \dots, n\}$ given by $R(x, i) = (t-x, n+1-i)$. Define $M_t = RW_t R$. Then we have the following.

Proposition 4.6. *For every $f \in \mathcal{D}_t^n$ with $f(t^-) = f(t)$, $M_t f \in \mathcal{D}_t^n$ is the unique maximal element in the isometry class of f with respect to \leq .*

Proof. The main idea is that the rotation operator R sends leftmost geodesics to right-most geodesics and vice versa. For any f and \mathbf{x}, \mathbf{y} , we have

$$\lambda_{Rf}(R\mathbf{x}, R\mathbf{y}) = R[\rho_f(\mathbf{x}, \mathbf{y})] \quad \text{and} \quad \rho_{Rf}(R\mathbf{x}, R\mathbf{y}) = R[\lambda_f(\mathbf{x}, \mathbf{y})].$$

Therefore

$$\rho_{RW_t Rf}(\mathbf{x}, \mathbf{y}) = R[\lambda_{W_t Rf}(R\mathbf{x}, R\mathbf{y})] \geq R[\lambda_{Rf}(R\mathbf{x}, R\mathbf{y})] = \rho_f(\mathbf{x}, \mathbf{y}),$$

where the inequality uses that $W_t Rf \leq Rf$. A similar calculation for $\lambda_{RW_t Rf}$ shows that $f \leq M_t f$. Using that $M_t f = M_t g$ for all g isometric to f yields the that $g \leq M_t f$ for all $g \sim f$. Finally, uniqueness follows from the same reasoning as in the proof of Proposition 4.1, but working with rightmost instead of leftmost optimizers. \square

5. The infinite bijection

In this section we find the inverse of the melon map W . Let \mathcal{D}_t^n be the set of n -tuples of cadlag functions $f_1, \dots, f_n : [0, t] \rightarrow \mathbb{R}$ with only positive jumps. For $f \in \mathcal{D}_t^n$, we can define the truncated melon map $W_t : \mathcal{D}_t^n \rightarrow \mathcal{D}_t^n$ by (3.1). Recall from the introduction (1.3) that M_t is the conjugate of W_t by the 180 degree rotation R , namely $M_t = RW_t R$. By the definition of W , this can be written as

$$M_t f_{n-k+1}(x^-) + \dots + M_t f_n(x^-) = f[(0, n)^k \rightarrow (t, 1)^k] - f[(x, n)^k \rightarrow (t, 1)^k]. \quad (5.1)$$

For $f \in \mathcal{D}^n$, we define the (infinite) **lemon map** by the limiting formula

$$Mf = \lim_{t \rightarrow \infty} M_t f. \quad (5.2)$$

We note in passing that M records a collection of multi-point Busemann functions for the metric environment defined by f . Busemann functions are an important object of study in first and last passage percolation, e.g. see Georgiou et al. (2017). Before studying M , we must justify why the limit exists.

Lemma 5.1. *For any $f \in \mathcal{D}^n$, the limit (5.2) exists in the topology of uniform convergence on compact sets. In particular, the map M is well-defined from \mathcal{D}^n to \mathcal{D}^n .*

Proof. For $g \in \mathcal{D}^n$, we let $\Sigma_k g = g_{n-k+1} + \dots + g_n$. For every $k \in \{1, \dots, n\}$, $x_1 \leq x_2 \leq t_1 \leq t_2 \in [0, \infty)$, we have

$$\Sigma_k M_{t_2} f(x_1^-) - \Sigma_k M_{t_1} f(x_1^-) \geq \Sigma_k M_{t_2} f(x_2^-) - \Sigma_k M_{t_1} f(x_2^-). \quad (5.3)$$

This is a consequence of the definition (5.1) and the quadrangle inequality (Lemma 2.5), with $\mathbf{p} = (x_1, n)^k$, $\mathbf{p}' = (x_2, n)^k$, $\mathbf{q} = (t_1, 1)^k$, $\mathbf{q}' = (t_2, 1)^k$.

Plugging in $x_1 = 0$, the left hand side of (5.3) equals 0. Therefore $\Sigma_k M_t f(x_2^-)$ is nonincreasing in t for every fixed x_2 . Moreover, from (5.1) we can conclude that

$$\Sigma_k M_t f(x^-) \geq \Sigma_k f(x^-) \quad (5.4)$$

since any disjoint k -tuple from $(x, n)^k$ to $(t, 1)^k$ can be extended to a disjoint k -tuple from $(0, n)^k$ to $(t, 1)^k$ by appending on the segments $[0, x) \times \{i\}$, $i = n, \dots, n - k + 1$. Combining these facts implies that there is some collection of real-valued functions $Mf = (Mf_1, \dots, Mf_n)$ such that $\Sigma_k M_t f \downarrow \Sigma_k Mf$ pointwise for all k .

Next, we establish uniform convergence. Returning to (5.3), we can replace M_{t_2} with M , to get that $g_t(x) := \Sigma_k Mf(x) - \Sigma_k M_t f(x)$ is monotone in x for every t . Moreover, we have already shown $g_t(x) \rightarrow 0$ pointwise in x . Any sequence of functions g_t with these two properties must converge uniformly on compact sets to 0. This implies that $M_t f \rightarrow Mf$ uniformly on compact sets.

Finally, the space \mathcal{D}^n is closed in the topology of uniform-on-compact convergence so $Mf \in \mathcal{D}^n$, yielding the final part of the lemma. \square

The definition of $M_t = RW_t R$ and Lemma 5.1 suggest that many properties of the melon map W in Proposition 3.12 should have parallels for the lemon map M . This is indeed the case.

Lemma 5.2. *The lemon map M has the following properties.*

- (i) (Isometry) $Mg \sim g$.
- (ii) (Idempotence) $M^2 = M$.

Proof. Part (i) for M_t is immediate from the definition of $M_t = RW_t R$ and the corresponding property of W_t , Proposition 3.12(i). Now, for any path π from (x, n) to $(y, 1)$, from the definition of path length we have

$$\left| |\pi|_{M_t f} - |\pi|_{Mf} \right| \leq 2n \sup_{(z,i) \in [x^-, y] \times \{1, \dots, n\}} |M_t f_i(x) - Mf_i(x)|.$$

This implies the following bound on any k -path last passage value from $\mathbf{p} = (\mathbf{x}, n)$ to $\mathbf{q} = (\mathbf{y}, 1)$:

$$\left| M_t f[\mathbf{p} \rightarrow \mathbf{q}] - Mf[\mathbf{p} \rightarrow \mathbf{q}] \right| \leq 2nk \sup_{(z,i) \in [x_1^-, y_k] \times \{1, \dots, n\}} |M_t f_i(x) - Mf_i(x)|.$$

The right side above converges to 0 by the uniform-on-compact convergence in Lemma 5.1, giving part (i) for M . Now by part (i), replacing f with Mf in (5.1) does not change the value, and hence $M_t M = M_t$. Letting $t \rightarrow \infty$ gives that $M^2 = M$. \square

To define the image of M , we need the following notion. First, we say that a k -tuple $\pi = (\pi_1, \dots, \pi_k)$ is a disjoint k -tuple from $(0, n)^k$ to $(\infty, 1)^k$ if π is the jump-time limit of disjoint k -tuples π^ℓ from $(0, n)^k$ to $(\ell, 1)^k$ for some sequence $\ell \rightarrow \infty$. Here a jump-time limit means that the jump times (2.3) of each individual path of the k -tuple π^ℓ converge to the jump times of each individual path in π . The point $+\infty$ is considered a valid limit; in this case the limiting path will not intersect all lines. For example, consider the paths π^ℓ from $(0, n)$ to $(\ell, 1)$ with jump times given by $t_i^\ell = \ell(1 - i/n)$. For each $i < n$, the sequence $t_i^\ell \rightarrow \infty$ so the limiting jump times of this path are $(t_n, \dots, t_1, t_0) = (0, \infty, \dots, \infty)$. The limit of the sequence π^ℓ is the path $\pi = [0, \infty) \times \{n\}$; the remaining path segments are empty sets, formally given by $(\infty, \infty) \times \{i\}$ for $i < n$. This path π is the **rightmost path** from $(0, n)$ to $(\infty, 1)$.

Notions of left and right naturally extend to these k -tuples of semi-infinite paths. We say that π is an **(infinite) quasi-optimizer** for g if the π^ℓ can be chosen so that

$$\lim_{\ell \rightarrow \infty} |\pi^\ell|_g - g[(0, n)^k \rightarrow (\ell, n)^k] = 0, \quad (5.5)$$

and π is an **(infinite) optimizer** if the π^ℓ can be chosen so that $|\pi^\ell|_g = g[(0, n)^k \rightarrow (\ell, n)^k]$ for all large enough ℓ . For every $k \in \{1, \dots, n\}$, by taking a subsequential limit of a sequence of optimizers from $(0, n)^k \rightarrow (\ell, n)^k$, there is at least one optimizer from $(0, n)^k$ to $(\infty, 1)^k$.

Let $\mathcal{D}_\downarrow^n \subset \mathcal{D}^n$ be the space where the **rightmost** k -tuple from $(0, n)^k$ to $(\infty, 1)^k$ is a **quasi-optimizer** for all k . This is the k -tuple $\pi = (\pi_1, \dots, \pi_k)$, where $\pi_i = \{0\} \times \{n, \dots, n - k + i + 1\} \cup [0, \infty) \times \{n - k + i\}$ (here the first part of this union is empty for $i = k$). Similarly, let $\mathcal{D}_{\downarrow-}^n$ be the space where the rightmost k -tuple from $(0, n)^k$ to $(\infty, 1)^k$ is an **optimizer** for all k .

The image \mathcal{D}_\uparrow^n of W can be thought of as the space where the **leftmost** k -tuple from $(0, n)^k$ to $(\infty, 1)^k$ is an **optimizer** for all k . This leftmost k -tuple is given by $\pi = (\pi_1, \dots, \pi_k)$ where $\pi_i = \{0\} \times \{n, \dots, i + 1\} \cup [0, \infty) \times \{i\}$. The analogue for M is given by the following two lemmas.

Lemma 5.3. $M(\mathcal{D}^n) = \mathcal{D}_\downarrow^n$.

Proof. Since M idempotent by Lemma 5.2(ii), g is in its image if and only if $Mg = g$. We use the notation $\Sigma_k g = g_{n-k+1} + \dots + g_n$. By (5.1), $Mg = g$ if and only if

$$\lim_{t \rightarrow \infty} g[(0, n)^k \rightarrow (t, 1)^k] - g[(x, n)^k \rightarrow (t, 1)^k] - \Sigma_k g(x^-) = 0 \quad (5.6)$$

for every x, k . We now prove that these g are exactly \mathcal{D}_\downarrow^n by proving both inclusions.

Only if: $Mg = g \Rightarrow g \in \mathcal{D}_\downarrow^n$. If (5.6) holds for every x, k , then by a diagonalization argument we can find $x_\ell \rightarrow \infty$, $x_\ell \leq \ell$ with $\ell \in \mathbb{N}$ such that

$$\lim_{\ell \rightarrow \infty} g[(0, n)^k \rightarrow (\ell, 1)^k] - g[(x_\ell, n)^k \rightarrow (\ell, 1)^k] - \Sigma_k g(x_\ell^-) = 0 \quad (5.7)$$

for every k . For $\ell \in \mathbb{N}$, let π^ℓ be the concatenation of the bottom k paths on the interval $[0, x_\ell]$ with an optimizer from $(x_\ell, n)^k \rightarrow (\ell, 1)^k$. We have $|\pi^\ell|_g = g[(x_\ell, n)^k \rightarrow (\ell, 1)^k] + \Sigma_k g(x_\ell^-)$, so by (5.7), the equation (5.5) is satisfied for this sequence π^ℓ . Moreover, since $x_\ell \rightarrow \infty$ with ℓ and π^ℓ simply follows the bottom k paths up to time x_ℓ , it converges to the rightmost k -tuple π_k from $(0, n)^k$ to $(\infty, 1)^k$. Therefore π_k is a quasi-optimizer, so $g \in \mathcal{D}_\downarrow^n$.

If: $g \in \mathcal{D}_\downarrow^n \Rightarrow Mg = g$. If the rightmost k -tuple π_k from $(0, n)^k$ to $(\infty, 1)^k$ is a quasi-optimizer, then, by the definition of \mathcal{D}_\downarrow^n , there is a sequence $\ell \rightarrow \infty$ and a sequence of disjoint k -tuples π^ℓ from $(0, n)^k$ to $(\ell, n)^k$ that converge to π_k and satisfy (5.5). Since $\pi^\ell \rightarrow \pi_k$, the optimizer π^ℓ uses the bottom k paths up to some time $x_\ell \rightarrow \infty$. This implies (5.6) for any fixed x as long as we take the limit only over the sequence ℓ , rather than over all $t \in [0, \infty)$. We can pass to the full limit in t since we know this limit exists by Lemma 5.1. \square

Lemma 5.4. $\mathcal{D}_\downarrow^n = \overline{\mathcal{D}_{\downarrow-}^n}$, where the closure is taken in the uniform norm.

Proof. Inclusion $\mathcal{D}_\downarrow^n \subset \overline{\mathcal{D}_{\downarrow-}^n}$. Let $g \in \mathcal{D}_\downarrow^n$, and for $\epsilon > 0$, define $g_i^\epsilon(t) = g_i(t) + \epsilon i \arctan(t)$, so that $g^\epsilon \rightarrow g$ uniformly as $\epsilon \rightarrow 0$. This approximation g^ϵ adds more weight onto lower lines $n, n-1, \dots$, encouraging optimizers to use these lines. Specifically, if a path τ from p to q is to the left of a path π from p to q and $\tau \neq \pi$ then

$$|\tau|_{g^\epsilon} - |\tau|_g < |\pi|_{g^\epsilon} - |\pi|_g. \quad (5.8)$$

We claim that $g^\epsilon \in \mathcal{D}_{\downarrow-}^n$ for all $\epsilon > 0$.

Fix k and let π^ℓ be the sequence in (5.5) for g converging to the rightmost k -tuple π_k from $(0, n)^k$ to $(\infty, 1)^k$. Let τ^ℓ be any sequence of optimizers in g^ϵ from $(0, n)^k$ to $(\ell, n)^k$. Suppose, for the sake of contradiction, that $\tau^\ell \not\rightarrow \pi_k$, and by choosing a subsequence if necessary, suppose $\tau^\ell \rightarrow \tau \neq \pi_k$. Since paths in τ are to the left of paths in π_k , (5.8) guarantees that there exists $\delta > 0$ such that for all large enough ℓ , we have

$$\delta + |\tau^\ell|_{g^\epsilon} - |\tau^\ell|_g < |\pi^\ell|_{g^\epsilon} - |\pi^\ell|_g.$$

On the other hand, the liminf of $|\pi^\ell|_g - |\tau^\ell|_g$ is at least 0 by the quasi-optimality of π^ℓ , implying that $|\pi^\ell|_{g^\epsilon} > |\tau^\ell|_{g^\epsilon} + \delta$ for large enough ℓ , contradicting the optimality of τ^ℓ . Hence $\tau^\ell \rightarrow \pi_k$, and so $g^\epsilon \in \mathcal{D}_{\downarrow-}^n$.

Inclusion $\mathcal{D}_\downarrow^n \supset \overline{\mathcal{D}_{\downarrow-}^n}$. As in the previous argument, let π_k be the rightmost k -tuple from $(0, n)^k$ to $(\infty, 1)^k$. Suppose $g \in \overline{\mathcal{D}_{\downarrow-}^n}$, and suppose $g_m \in \mathcal{D}_{\downarrow-}^n$ converges uniformly to g . Fix $k \in \{1, \dots, n\}$, and for each m , let π_m^ℓ be a sequence of disjoint optimizers in g_m from $(0, n)^k$ to $(\ell, 1)^k$, $\ell \in \mathbb{N}$ that converges along a subsequence to π_k . By a diagonalization argument, we can find a sequence $\ell_m \rightarrow \infty$ such that $\pi_m^{\ell_m} \rightarrow \pi_k$ as $m \rightarrow \infty$ and $|\pi_m^{\ell_m}|_{g_m} - |\pi_m^{\ell_m}|_g \rightarrow 0$. The sequence $\pi_m^{\ell_m}$ is quasi-optimal for g , so $g \in \mathcal{D}_\downarrow^n$. \square

Later, we will also use the following closely related observation.

Corollary 5.5. *Let \mathcal{D}_{*t}^n be the set of $f \in \mathcal{D}^n$ such that for every k there is a quasi-optimizer τ_k from $(0, n)^k \rightarrow (\infty, 1)^k$ that agrees with the rightmost k -tuple from $(0, n)^k \rightarrow (\infty, 1)^k$ up to time t . For $f \in \mathcal{D}_{*t}^n$, we have $Mf_{[0,t]} = f|_{[0,t]}$.*

Proof. Just as in the ‘if’ part of the proof of Lemma 5.3, the existence of such a quasi-optimizer τ_k implies (5.6) for $x \leq t$, giving the result. \square

The fact that M and W are inverses follows from an abstract lemma.

Lemma 5.6. *Let $A, B : X \rightarrow X$ be two idempotent maps satisfying $AB = A$ and $BA = B$. Then $A|_{B(X)}$ is a bijection between $B(X)$ and $A(X)$ with inverse $B|_{A(X)}$.*

Proof. Since B is idempotent, B is the identity on $B(X)$. Therefore $BA = B$ is also the identity on $B(X)$. Similarly, AB is the identity on $A(X)$. Finally, $A(X) = AB(X)$ and $B(X) = BA(X)$, yielding the result. \square

Proposition 5.7. *We have $MW = M$ and $WM = W$. Moreover, the restriction $W|_{\mathcal{D}_{\downarrow}^n}$ is a bijection between $\mathcal{D}_{\downarrow}^n$ and \mathcal{D}_{\uparrow}^n with inverse $M|_{\mathcal{D}_{\uparrow}^n}$.*

Proof. The first sentence is immediate from the isometric properties of W and M and the fact that both maps are defined in terms of last passage from line n to line 1. The second sentence follows from Lemma 5.6. \square

It may seem that elements of $\mathcal{D}_{\downarrow}^n$ in \mathcal{D}^n should be somewhat rare and special. Surprisingly, this is not the case! The following proposition gives a natural condition for an element of \mathcal{D}^n to be in $\mathcal{D}_{\downarrow}^n$ (in fact, in $\mathcal{D}_{\downarrow-}^n$).

Proposition 5.8. *Let $f \in \mathcal{D}^n$, and suppose that*

$$\limsup_{t \rightarrow \infty} f_{j+1}(t) - f_j(t) = \infty \quad (5.9)$$

for $j = 1, \dots, n-1$. Then $f \in \mathcal{D}_{\downarrow-}^n$.

Proof. Suppose for the sake of contradiction that $f \notin \mathcal{D}_{\downarrow-}^n$. Then for some $k \in \{1, \dots, n\}$, there is an optimizer $\pi = (\pi_1, \dots, \pi_k)$ from $(0, n)^k$ to $(\infty, 1)^k$ such that $\pi_i \neq \{0\} \times \{n, \dots, n-k+i+1\} \cup [0, \infty) \times \{n-k+i\}$ for some i . Let i be the maximal index for which this holds, let $j < n-k+i$ be the index of the line on which π_i contains a semi-infinite interval, and let $t \in [0, \infty)$ be the time when π_i jumps to line j , so that $(t, j+1) \in \pi_i$ and $[t, \infty) \times \{j\} \subset \pi_i$. Since $\pi_\ell = [0, \infty) \times \{n+k-\ell\}$ for $\ell > i$, we have that

$$(t, \infty) \times \{j+1\} \cap (\cup \pi) = \emptyset. \quad (5.10)$$

Since π is an optimizer, for any $s > t$, restricting each of the π_i to a compact set $[t, s] \times \{1, \dots, n\}$ must also yield an optimizer. In particular, (5.10) implies

that the path $\pi_i \cap [t, s] \times \{j, j+1\}$ must always be a geodesic from $(t, j+1)$ to (s, j) . Therefore

$$\sup_{r \in [t, s]} f_{j+1}(r) - f_j(r^-) = \sup_{r \in [t, s]} (f_{j+1} - f_j)(r) + [f_j(r) - f_j(r^-)] = f_{j+1}(t) - f_j(t^-) \quad (5.11)$$

for all $s > t$. Taking $s \rightarrow \infty$ in (5.11) shows that the left hand side of (5.9) equals $f_{j+1}(t) - f_j(t^-) < \infty$, a contradiction of (5.9). \square

Remark 5.9. Natural measures on \mathcal{D}^n , such as i.i.d. random walks or i.i.d. Lévy processes, satisfy the conditions of Proposition 5.8 almost surely. Many of these measures, such as Brownian motion, piecewise constant geometric random walks, or piecewise linear versions of Bernoulli walks, appear in integrable models. See Section 7.

Remark 5.10 (Historical notes). In O’Connell (2003b), an infinite time RSK correspondence is built for nondecreasing functions with domain and range in \mathbb{Z} . See Lemma 2.2 in that paper for the description of the inverse map and Corollary 3.2 in that same paper for the relation between that map and the usual RSK correspondence. In Section 4 of Biane, Bougerol and O’Connell (2005), an infinite time version of RSK is developed in for continuous functions in greater representation-theoretic generality. In the language of that paper, our melon map W is a generalized Pitman operator and the lemon map M is a co-Pitman operator.

6. The finite bijection

In Section 5, we studied the bijectivity of the melon map in the infinite-time setting. The goal of this section is to construct a version in the finite-time setting. This version is more similar to classical RSK: an additional Gelfand-Tsetlin pattern will play the role of the second Young tableaux.

Like the infinite case, it is straightforward to check that the maps W_t and M_t are inverses of each other on appropriate domains. However, unlike the infinite case, in the finite case, the image of M_t is a rare subset of \mathcal{D}_t^n and so the bijection loses usefulness. Instead, our strategy is to take an element of \mathcal{D}_t^n and map it to an element of \mathcal{D}_{*t}^n , see Corollary 5.5. This allows us to define the RSK correspondence on the finite domain $[0, t]$ in terms of the maps W, M defined on the infinite domain $[0, \infty)$.

For $f \in \mathcal{D}_t^n$, and a k -tuple of functions $g = (g_1, \dots, g_n)$ where each $g_i : [t, \infty) \rightarrow \mathbb{R}$ and $g(t) = f(t)$, define the **concatenation** $f \oplus g \in \mathcal{D}^n$ to be equal to f on $[0, t]$ and g on $[t, \infty)$. Note that our notation implicitly depends on t . For $\alpha > 0$, we write $f \oplus \alpha$ to mean $f \oplus g_\alpha$, where dg_α is purely atomic on $(t, \infty) \times \{1, \dots, n\}$ with atoms of size αi at locations $(t+i, i)$. The concatenations $f \oplus \alpha$ are set up so that for large enough α , $f \oplus \alpha \in \mathcal{D}_{*t}^n$. We prove this formally in Lemma 6.2, though it may be more useful to simply study the cartoon example in Figure 3 to see why this holds.

Given this, one direction of the finite RSK correspondence will essentially be the map $f \mapsto W(f \oplus \alpha)$ for large α . We can then invert this correspondence using M :

$$MW(f \oplus \alpha)|_{[0,t]} = M(f \oplus \alpha)|_{[0,t]} = (f \oplus \alpha)|_{[0,t]} = f. \quad (6.1)$$

Here the first equality follows from Proposition 5.7, the second follows from Corollary 5.5, and the third is by definition.

While (6.1) clearly hints at the existence of a bijection, there are a few more things to check. We need to figure out what information about f is actually used in the map $f \mapsto W(f \oplus \alpha)$, identify the image of this map, and then show that if we start with data in that space, then WM is also the identity on that space. The rest of this section is devoted to doing this. We will also present our version of the finite RSK correspondence in a way that more closely resembles the classical RSK bijection between matrices and Young tableaux. Our correspondence will be related to classical RSK and dual RSK in Section 8.

First, let $\mathcal{D}_{t\uparrow}^n = W_t(\mathcal{D}_t^n)$ be the space of Pitman ordered sequences $w_n \succ \dots \preceq w_1$ in \mathcal{D}_t^n . For $n \in \mathbb{N}$, let GT_n be the space of triangular arrays of real numbers $g = \{g_i(j) : i \leq j \in \{1, \dots, n\}\}$ satisfying the inequalities

$$g_i(j) \geq g_i(j-1), \quad g_i(j-1) \geq g_{i+1}(j), \quad (6.2)$$

for all i, j where these quantities are defined. Such an array is called a **Gelfand-Tsetlin pattern** of depth n . These inequalities amount to saying that the j th row $g(j)$ is an ordered i -tuple, and that consecutive rows interlace.

Next, define the space

$$\mathcal{G}_t^n = \{h = (w, g) \in \mathcal{D}_{t\uparrow}^n \times \text{GT}_n : w(t) = g(n)\}.$$

The space \mathcal{G}_t^n is the analogue of the set of pairs of Young tableaux with the same shape, see Section 8. The finite- t RSK correspondence, RSK_t , maps \mathcal{D}_t^n into \mathcal{G}_t^n , where $\text{RSK}_t(f) = (W_t f, G_t f)$, and for $k \leq s \in \{1, \dots, n\}$ we have

$$\sum_{i=1}^k G_t f_i(s) = f[(0, n)^k \rightarrow (t, n-s+1)^k]. \quad (6.3)$$

Lemma 6.1. *The map $\text{RSK}_t : \mathcal{D}_t^n \rightarrow \mathcal{G}_t^n$ is well-defined. That is, $\text{RSK}_t f \in \mathcal{G}_t^n$ for any $f \in \mathcal{D}_t^n$.*

Proof. By Proposition 3.12 (iii), $W_t f \in \mathcal{D}_{t\uparrow}^n$, and $W_t f(t) = G_t f(n)$ by definition. It remains to verify the Gelfand-Tsetlin inequalities (6.2) for $G_t f$. Both of these types of inequalities have similar proofs and follow from quadrangle inequalities that are similar in spirit to Lemma 2.5. We prove only the first inequality; the second one is similar. By (6.3), the inequality $G_t f_i(s) \geq G_t f_i(s-1)$ is equivalent to the quadrangle inequality

$$\begin{aligned} & f[(0, n)^i \rightarrow (t, n-s+1)^i] + f[(0, n)^{i-1} \rightarrow (t, n-s+2)^{i-1}] \\ & \geq f[(0, n)^i \rightarrow (t, n-s+2)^i] + f[(0, n)^{i-1} \rightarrow (t, n-s+1)^{i-1}]. \end{aligned} \quad (6.4)$$

To prove (6.4), let $\pi = (\pi_1, \dots, \pi_i), \pi' = (\pi'_1, \dots, \pi'_{i-1})$ be disjoint optimizers from $(0, n)^i \rightarrow (t, n - s + 2)^i$ and $(0, n)^{i-1} \rightarrow (t, n - s + 1)^{i-1}$, respectively. Similarly to the proof of Lemma 2.5, we will use π, π' to construct a disjoint i -tuple τ^L from $(0, n)^i \rightarrow (t, n - s + 1)^i$ and a disjoint $(i - 1)$ -tuple τ^R from $(0, n)^{i-1} \rightarrow (t, n - s + 2)^{i-1}$.

First, define a disjoint $(i - 1)$ -tuple $\tilde{\pi}$ from $(0, n)^{i-1}$ to $(t, n - s + 1)^{i-1}$ by letting $\tilde{\pi}_j = \pi_j \cup \{(t, n - s + 1)\}$. Form disjoint $(i - 1)$ -tuples $\hat{\tau}^L$ and $\hat{\tau}^R$ from $(0, n)^{i-1} \rightarrow (t, n - s + 1)^{i-1}$ from $\tilde{\pi}$ and π' exactly as in the proof of Lemma 2.5. We form τ^L from $\hat{\tau}^L$ by letting $\tau_j^L = \hat{\tau}_j^L$ for $j \leq i - 1$ and letting $\tau_i^L = \pi_i \cup \{(t, n - s + 1)\}$. Since $\hat{\tau}_j^L \leq \pi_j$ for all $j \leq i - 1$ and π is a disjoint i -tuple, τ^L is a disjoint i -tuple from $(0, n)^i \rightarrow (t, n - s + 1)^i$. We form τ^R from $\hat{\tau}^R$ by restricting to the lines $\{n - s + 2, \dots, n\}$. Our construction guarantees that this only removes the single point $(t, n - s + 1)$ from all paths $\hat{\tau}_j^R$ and so we are left with a disjoint $(i - 1)$ -tuple from $(0, n)^{i-1} \rightarrow (t, n - s + 2)^{i-1}$. Exactly as in the proof of Lemma 2.5, we have

$$|\pi|_f + |\pi'|_f \leq |\tau^L|_f + |\tau^R|_f.$$

Now, $|\tau^L|_f + |\tau^R|_f$ is bounded above by the left side of (6.4) and $|\pi|_f + |\pi'|_f$ equals the right side of (6.4), yielding the result. \square

We now move to understanding the map $f \mapsto W(f \oplus \alpha)$ for large α .

Lemma 6.2. *Let $f \in \mathcal{D}_t^n$. For $\alpha \geq (n - 1)(Wf_1(t) - Wf_n(t))$, we have $f \oplus \alpha \in \mathcal{D}_{*t}^n$ and $W(f \oplus \alpha) = W_t f \oplus \Delta_{t,\alpha} G_t f$, for some function $\Delta_{t,\alpha} : \text{GT}_n \rightarrow \mathcal{E}^{n,t}$. Here $\mathcal{E}^{n,t}$ is the space of k -tuples of cadlag paths from $[t, \infty) \rightarrow \mathbb{R}$.*

Note that the cadlag paths in $\mathcal{E}^{n,t}$ are, a priori, allowed to have negative jumps. In Lemma 6.3(iii) we will show that this does not happen in the cases we care about.

Proof. On $[0, t]$, the formula (3.1) implies that $W(f \oplus \alpha) = Wf$. Next, we show that $W(f \oplus \alpha)$ is an explicit function of $G_t f$ on $[t, \infty)$. The action of W applied to $f \oplus \alpha$ is illustrated in Figure 3. Indeed, we can see that for $k, \ell \in \{1, \dots, n\}$, there will be a disjoint k -tuple π across $f \oplus \alpha$ from $(0, n)^\ell$ to $(t + k, 1)^\ell$ that picks up large α -weights at locations $(1 + (k - \ell)^+, t + (k - \ell)^+ + 1), \dots, (k, t + k)$. By ensuring that these weights are chosen and that π is also optimal on $[0, t]$, we can ensure that π has weight

$$\begin{aligned} & \sum_{i=1+(k-\ell)^+}^k \alpha i + f[(0, n)^\ell \rightarrow (t, (k - \ell)^+ + 1)^\ell] \\ &= \sum_{i=1+(k-\ell)^+}^k \alpha i + \sum_{i=1}^{\ell} G_t f_i(n - (k - \ell)^+). \end{aligned}$$

No other disjoint k -tuple τ from $(0, n)^\ell$ to $(t + k, 1)^\ell$ can improve the α -part of the sum above, and any disjoint k -tuple τ can only improve the f -part of the




FIG 3. An example of $f \oplus \alpha$ (on the left) and $W(f \oplus \alpha)$ (on the right) for a function $f \in \mathcal{D}_t^4$ to give the reader a visual aid to the proofs of Lemmas 6.2 and 6.3. The white circles represent atoms of the measures df and $dW(f \oplus \alpha)$, or equivalently, jumps on the lines. In $W(f \oplus \alpha)$, the ten weights all have a contribution of a single α plus a contribution related to $G_t f$, not contained in the figure.

sum by at most

$$f[(0, n)^\ell \rightarrow (t, 1)^\ell] - f[(0, n)^\ell \rightarrow (t, (k - \ell)^+ + 1)^\ell] \quad (6.5)$$

at the expense of at least one α . Now, the quantity (6.5) equals 0 for $\ell \geq k$. For $\ell \leq k - 1$, the first term equals $Wf_1(t) + \dots + Wf_\ell(t) \leq \ell Wf_1(t)$ by (3.1). The second term is always bounded below by the weight of the rightmost disjoint k -tuple from $(0, n)^\ell$ to $(t, k - \ell + 1)^\ell$, giving

$$f[(0, n)^\ell \rightarrow (t, k - \ell + 1)^\ell] \geq \sum_{i=n-\ell+1}^n f_i(\ell) + \sum_{i=k-\ell+1}^{n-\ell+2} f_i(\ell) - f_i(\ell^-) \geq \sum_{i=n-\ell+1}^n f_i(\ell).$$

Now, by iterated applications of the two-line bound (3.7) we have the inequality $f_i \geq Wf_n$ for all i , giving that the right hand side above is bounded below by $\ell Wf_n(t)$, and hence (6.5) is bounded above by $\ell(Wf_1(t) - Wf_n(t)) \leq (k - 1)(Wf_1(t) - Wf_n(t))$.

Since $\alpha \geq (n - 1)(Wf_1(t) - Wf_n(t))$, we have $|\pi|_{f \oplus \alpha} \geq |\tau|_{f \oplus \alpha}$, so π is an optimizer. The story for optimizers up to time $t + k$ when $k \notin \{1, \dots, n\}$ is similar by rounding down to the nearest integer time. Therefore $W(f \oplus \alpha)$ is an explicit function of $G_t f$ on $[t, \infty)$. Moreover, from the construction of optimizers above from $(0, n)^\ell$ to $(t + k, 1)^\ell$ for $k \geq n$, we can see that $f \oplus \alpha \in \mathcal{D}_{*t}^n$. \square

The proof of Lemma 6.2 allows us to explicitly construct the function $\Delta_{t, \alpha}$. While this will be necessary to fill in some proof details regarding the invertibility of RSK_t , for now it will be easier to think of the map $\Delta_{t, \alpha}$ abstractly, as a linear map from $\mathbb{R}^{\binom{n+1}{2}}$ (which contains GT_n) onto an $\binom{n+1}{2}$ -dimensional linear subspace of $\mathcal{E}^{n, t}$, the space of n -tuples of cadlag functions from $[t, \infty) \rightarrow \mathbb{R}$. It has the following properties.

Lemma 6.3. *For every $t, \alpha > 0$, the map $\Delta_{t, \alpha} : \text{GT}_n \rightarrow \mathcal{E}^{n, t}$ satisfies the following:*

- (i) $\Delta_{t,\alpha}$ is one-to-one.
- (ii) $\Delta_{t,\alpha}g(t) = g(n)$.
- (iii) For $\alpha \geq \alpha_g := (n-1)(g_1(n) - g_n(n))$, paths in $\Delta_{t,\alpha}g$ are cadlag with only positive jumps
- (iv) The paths in $\Delta_{t,\alpha}g$ are Pitman ordered: $\Delta_{t,\alpha}g_{i+1}(s) \leq \Delta_{t,\alpha}g_i(s^-)$ for all $s \in [t, \infty)$, $i \in \{1, \dots, n-1\}$. For this to make sense when $s = t$ we define $\Delta_{t,\alpha}g(t^-) = \Delta_{t,\alpha}g(t)$.
- (v) For $\alpha \geq \alpha_g$ and $0 < r \leq n$ we have

$$\Delta_{t,\alpha}g[(t+r, n)^k \rightarrow (t+n, 1)^k] = \sum_{i=\lceil r \rceil \vee (n-k+1)}^n \alpha_i. \quad (6.6)$$

Note that properties (ii)-(v) above are easy to see when $g = G_t f$ for some f . In this case, properties (ii)-(iv) follow from the fact that $Wf \oplus \Delta_{t,\alpha}g = W(f \oplus \alpha) \in \mathcal{D}_\uparrow^n$ (Lemma 6.2) and property (v) follows immediately from the fact that $Wf \oplus \Delta_{t,\alpha}g$ is isometric to $f \oplus \alpha$.

We postpone the proof of Lemma 6.3 for now, and use it to show the invertibility of RSK_t . Let $O_t : \mathcal{G}_t^n \rightarrow \mathcal{D}^n$ be the map taking a pair $(w, g) \mapsto w \oplus \Delta_{t,\alpha_g}g$, where $\alpha_g = (n-1)(g_1(n) - g_n(n))$. The concatenation here is well-defined by Lemma 6.3(ii) and the fact that $w(t) = g(n)$. Also let $\Gamma_t : \mathcal{D}^n \rightarrow \mathcal{D}_t^n$ be the restriction of a function to $[0, t]$.

Proposition 6.4. *The map $\text{RSK}_t : \mathcal{D}_t^n \rightarrow \mathcal{G}_t^n$ is a bijection with inverse $\text{RSK}_t^{-1} := \Gamma_t M O_t$.*

Proof. First, for $(w, g) \in \mathcal{G}_t^n$ we claim that

$$M(w \oplus \Delta_{t,\alpha_g}g) = f \oplus \alpha_g, \quad (6.7)$$

for some $f \in \mathcal{D}_t^n$. To see this, using that the function $w \oplus \Delta_{t,\alpha_g}g$ is constant on the interval $[t+n, \infty)$, for any $r > 0$ and $s \geq t+n \vee r$ we have

$$\begin{aligned} & \sum_{i=n-k+1}^n M_s(w \oplus \Delta_{t,\alpha_g}g)_i((t+r)^-) \\ &= (w \oplus \Delta_{t,\alpha_g}g)[(0, n)^k \rightarrow (s, 1)^k] - (w \oplus \Delta_{t,\alpha_g}g)[(t+r, n)^k \rightarrow (s, 1)^k] \\ &= c_k - \Delta_{t,\alpha_g}g[(t+r, n)^k \rightarrow (t+n, 1)^k] \mathbf{1}(r \leq n), \end{aligned} \quad (6.8)$$

where $c_k = (w \oplus \Delta_{t,\alpha_g}g)[(0, n)^k \rightarrow (t+n, 1)^k]$. The final expression (6.8) does not depend on s , and so the same equation holds with the limit M in place of M_s . Next, Lemma 6.3(v) gives that for $r \in (0, n]$ we have

$$\Delta_{t,\alpha_g}g[(t+r, n)^k \rightarrow (t+n, 1)^k] = \sum_{i=\lceil r \rceil \vee (n-k+1)}^n \alpha_g i.$$

We can use this formula along with (6.8) to compute the function $r \mapsto M(t+r) - M(t)$, $r \geq 0$. This computation gives that $dM(w \oplus \Delta_{t,\alpha_g}g)$ is purely atomic on

$(t, \infty) \times \{1, \dots, n\}$ with atoms of size α_i at locations $(t+i, i)$ for all $i \in \{1, \dots, n\}$. In other words, we have the representation (6.7) for some $f \in \mathcal{D}_t^n$. Next, by Proposition 5.7,

$$w \oplus \Delta_{t, \alpha_g} g = WM(w \oplus \Delta_{t, \alpha_g} g) = W(f \oplus \alpha_g).$$

In particular, $W_t f = w$. Moreover, $W_t f(t) = \Delta_{t, \alpha_g} g(t) = g(n)$ by Lemma 6.3(ii), so $\alpha_g = (n-1)(W f_1(t) - W f_n(t))$. Therefore by Lemma 6.2, $W(f \oplus \alpha_g) = W_t f \oplus \Delta_{t, \alpha_g} G_t f$, so $\Delta_{t, \alpha_g} G_t f = \Delta_{t, \alpha_g} g$. Since Δ_{t, α_g} is one-to-one (Lemma 6.3(i)), we get that $G_t f = g$. Putting all this together gives that $\text{RSK}_t \Gamma_t MO_t(w, g) = (w, g)$.

On the other hand, $\Gamma_t MO_t \text{RSK}_t f = f$ via the computation (6.1). \square

Proof of Lemma 6.3. We first find an explicit formula for $\Delta_{t, \alpha}$. Following from the proof of Lemma 6.2, we can see that $\Delta_{t, \alpha}$ is given by the following two rules:

- $\Delta_{t, \alpha} g(t) = g(n)$, and on $(t, \infty) \times \{1, \dots, n\}$, the finitely additive measure $d\Delta_{t, \alpha} g$ is purely atomic with support contained in the set of points $(t+k, \ell)$, $\ell \leq k \in \{1, \dots, n\}$.
- For $k, \ell \in \{1, \dots, n\}$,

$$\sum_{i=1}^{\ell} \Delta_{t, \alpha} g_i(t+k) = \sum_{i=1+(k-\ell)^+}^k \alpha_i + \sum_{i=1}^{\ell} g_i(n - (k-\ell)^+). \quad (6.9)$$

Proof of (i, ii). By rearranging equation (6.9) it is verified from these formulas that $\Delta_{t, \alpha} g$ determines g , giving (i). Part (ii) follows from the first bullet point.

Proof of (iii). Now, for $k \in \{1, \dots, n\}$ and $\ell < k$, from equation (6.9), $d\Delta_{t, \alpha} g$ has an atom of size

$$\alpha - \sum_{i=1}^{\ell} [g_i(n-k+\ell+1) - g_i(n-k+\ell)] + \sum_{i=1}^{\ell-1} [g_i(n-k+\ell) - g_i(n-k+\ell-1)] \quad (6.10)$$

at $(t+k, \ell)$, and has an atom of size $\alpha + \sum_{i=1}^{k-1} g_i(n) - g_i(n-1)$ at the points $(t+k, k)$ for $k \in \{1, \dots, n\}$. Note that for $\ell = 1$ the second sum in (6.10) is empty. To prove (iii), we first show that all these atoms have positive weight. For the atoms at $(t+k, k)$, $k = 2, \dots, n$, this is true by the Gelfand-Tsetlin inequalities (6.2). For the remaining atoms, by (6.2), all terms in both sums in (6.10) are nonnegative and bounded above by $g_1(n) - g_n(n)$. Since at most $n-1$ terms in (6.10) come with a negative sign, since $\alpha \geq \alpha_g = (n-1)(g_1(n) - g_n(n))$, these remaining atoms are positive. This shows that the paths in $\Delta_{t, \alpha} g$ have positive jumps.

Proof of (iv). Next, we check that $\Delta_{t, \alpha} g_{i+1}(s) \leq \Delta_{t, \alpha} g_i(s^-)$ for all $i \leq k-1$, $s \in [t, \infty)$. This inequality holds at time t since $g(n)$ is ordered by (6.2). For $s > t$, it suffices to show that for $k, \ell \in \{1, \dots, n\}$,

$$\Delta_{t, \alpha} g_{\ell+1}(t+k) \leq \Delta_{t, \alpha} g_{\ell}(t+k-1).$$

Writing out both sides of the inequality in terms of g and cancelling common terms, this inequality is equivalent to

$$g_{\ell+1}(n - (k - \ell - 1)^+) \leq g_\ell(n - (k - \ell)^+).$$

For $\ell < k$, this is the second Gelfand-Tsetlin inequality in (6.2). For $\ell \geq k$, this follows since the vector $g(n)$ is ordered.

Proof of (v). Let $0 < r \leq n$ and fix $k \in \{1, \dots, n\}$. First, for $i = 1, \dots, k$, let π_i be the unique path from $(t + r, n)$ to $(t + n, 1)$ containing the sets

$$[t + r, t + (i + n - k) \vee r] \times \{i + n - k\}, \quad [t + (i + n - k) \vee r, t + n] \times \{i\}.$$

Then $\pi = (\pi_1, \dots, \pi_k)$ is a disjoint k -tuple from $(t + r, n)^k$ to $(t + n, 1)^k$, see Figure 4. Moreover, from the formula (6.10) we can see that $\alpha(\sum_{i=\lceil r \rceil \vee (n-k+1)}^n i) = |\pi|_{\Delta_{t,\alpha}g}$ (it may also help to examine Figure 4 here). For $k > n - \lceil r \rceil$, this k -tuple collects all the atoms of $d\Delta_{t,\alpha}g$ in the box $[t + r, t + n] \times \{1, \dots, n\}$, and so must be an optimizer.

To complete the proof in the case when $k \leq n - \lceil r \rceil$, it is enough to show that for every $\ell \in \{1, \dots, k\}$ we have

$$\begin{aligned} \Delta_{t,\alpha}g[(t + r, n)^{k-\ell+1} \rightarrow (t + n - \ell + 1, 1)^{k-\ell+1}] \\ \leq \Delta_{t,\alpha}g[(t + r, n)^{k-\ell} \rightarrow (t + n - \ell, 1)^{k-\ell}] + \alpha(n - \ell + 1), \end{aligned} \quad (6.11)$$

where we use the convention that $\Delta_{t,\alpha}g[p^0 \rightarrow q^0] = 0$. Indeed, by induction (6.11) implies that $\Delta_{t,\alpha}g[(t + r, n)^k \rightarrow (t + n, 1)^k] \leq \alpha(\sum_{i=n-k+1}^n i)$, as desired. Fix ℓ , and let $\tau = (\tau_1, \dots, \tau_{k-\ell+1})$ be a disjoint optimizer from $(t + r, n)^{k-\ell+1}$ to $(t + n - \ell + 1, 1)^{k-\ell+1}$. The collection τ remains a disjoint k -tuple if we push $\tau_{k-\ell+1}$ further right. In particular, we may replace $\tau_{k-\ell+1}$ with the rightmost path $\tau'_{k-\ell+1}$ from $(t + r, n)$ to $(t + n - \ell + 1, 1)$ given by

$$\tau'_{k-\ell+1} = [t + r, t + n - \ell + 1] \times \{n\} \cup \{t + n - \ell + 1\} \cup \{1, \dots, n - 1\}.$$

Letting $\tau' = (\tau_1, \dots, \tau_{k-\ell}, \tau'_{k-\ell+1})$, we claim that

$$|\tau|_{\Delta_{t,\alpha}g} \leq |\tau'|_{\Delta_{t,\alpha}g}. \quad (6.12)$$

Before proving (6.12), let us explain how it gives (6.11). For this, we write $\tau'|_{[a,b]}$ for the restriction of all paths in a k -tuple to the set $[a, b] \times \{1, \dots, n\}$. We have that

$$\begin{aligned} |\tau'|_{\Delta_{t,\alpha}g} &= |\tau'|_{[t+r, t+n-\ell]}|_{\Delta_{t,\alpha}g} + |\tau'|_{(t+n-\ell, t+n-\ell+1]}|_{\Delta_{t,\alpha}g} \\ &= |(\tau_1, \dots, \tau_{k-\ell})|_{[t+r, t+n-\ell]}|_{\Delta_{t,\alpha}g} + \alpha(n - \ell + 1). \end{aligned}$$

Here the second equality uses that $\tau'_{k-\ell+1}$ picks up no weight in the interval $[t + r, t + n - \ell]$, and pick ups all atoms of $d\Delta_{t,\alpha}g$ at time $t + n - \ell + 1$; these atoms have total mass $\alpha(n - \ell + 1)$. To conclude (6.11) we can observe that

$$|(\tau_1, \dots, \tau_{k-\ell})|_{[t+r, t+n-\ell]}|_{\Delta_{t,\alpha}g} \leq \Delta_{t,\alpha}g[(t + r, n)^{k-\ell} \rightarrow (t + n - \ell, 1)^{k-\ell}],$$




FIG 4. An aid for the proof of Lemma 6.3(v). The left side gives an example of the k -tuple π in the first part of the proof when $n = 6, r = 2.5, k = 3$. The right side gives an example of the $\sigma \mapsto \sigma'$ path change in the final step in the proof.

since, after possibly adding a few points at time $t + n - \ell$, the collection $(\tau_1, \dots, \tau_{k-\ell})|_{[t+r, t+n-\ell]}$ is a disjoint $(k - \ell)$ -tuple from $(t + r, n)^{k-\ell} \rightarrow (t + n - \ell, 1)^{k-\ell}$. It remains to check (6.12). For this, first observe that by the ordering on the paths in τ , the set difference $\tau'_{k-\ell+1} \setminus \tau_{k-\ell+1}$ is disjoint from $\cup \tau$. Therefore

$$\begin{aligned} |\tau'|_{\Delta_{t,\alpha g}} - |\tau|_{\Delta_{t,\alpha g}} &= d\Delta_{t,\alpha g}(\cup \tau' \setminus \cup \tau) - d\Delta_{t,\alpha g}(\cup \tau \setminus \cup \tau') \\ &\geq d\Delta_{t,\alpha g}(\tau'_{k-\ell+1} \setminus \tau_{k-\ell+1}) - d\Delta_{t,\alpha g}(\tau_{k-\ell+1} \setminus \tau'_{k-\ell+1}) \\ &= |\tau'_{k-\ell+1}|_{\Delta_{t,\alpha g}} - |\tau_{k-\ell+1}|_{\Delta_{t,\alpha g}}, \end{aligned}$$

so to prove (6.12) it suffices to show that $\tau'_{k-\ell+1}$ is a geodesic from $(t + r, n)$ to $(t + n - \ell + 1, 1)$. Since $d\Delta_{t,\alpha g}$ is purely atomic with all atoms located at times in $t + \mathbb{Z}$, it is enough to show that $|\tau'_{k-\ell+1}|_{\Delta_{t,\alpha g}}$ is maximal among paths from $(t + r, n)$ to $(t + n - \ell + 1, 1)$ whose jumps are all in $t + \mathbb{Z}$. Let σ be any such path. Then there exists a unique $j \in \{[r], \dots, n - \ell + 1\}$ such that $(t + j, j) \in \sigma$. If $j = n - \ell + 1$, then σ picks up the same set of atoms as $\tau'_{k-\ell+1}$ and hence $|\tau'_{k-\ell+1}|_{\Delta_{t,\alpha g}} = |\sigma|_{\Delta_{t,\alpha g}}$. If not, then there exists $i \in \{1, \dots, j\}$ such that the set

$$J = \{t + j\} \times \{i, \dots, j\} \cup (t + j, t + j + 1) \times \{i\}$$

is contained in σ . Let $\tilde{J} = \{(z, p) \in \sigma : z \leq t + j + 1, p \geq i\}$. Define a new path σ' by

$$\sigma' = (\sigma \setminus \tilde{J}) \cup K, \quad \text{where} \quad K = [t + r, t + j + 1] \times \{n\} \cup \{t + j + 1\} \times \{i, \dots, n - 1\}.$$

See Figure 4 for an illustration of σ, σ' . Using (6.10), we can calculate that

$$|\sigma'|_{\Delta_{t,\alpha g}} - |\sigma|_{\Delta_{t,\alpha g}} = g_i(n - j + i) - g_i(n - j + i - 1),$$

which is nonnegative by the first inequality in (6.2). Moreover, σ' picks up the atom at $(t + j + 1, j + 1)$. Therefore by induction, (i.e continuing to push the path further to the right), we get that $|\tau'_{k-\ell+1}|_{\Delta_{t,\alpha g}} \geq |\sigma|_{\Delta_{t,\alpha g}}$ as desired. \square

Remark 6.5. Just as we can build the melon map W as a composition of 2-line Pitman transforms, see Definition 3.11, we can also build the n -line RSK correspondence by composing $\binom{n}{2}$ 2-line correspondences. More precisely, we

build up the melon Wf using the maps $\sigma_i f$ as in (3.11), but every time we apply one of the maps σ_i to an intermediate function g , we also record the additional value $g_{i+1}(t)$.

The map $(g_i, g_{i+1}) \mapsto (\sigma_i g_i, \sigma_i g_{i+1}; g_{i+1}(t))$ is a 2-line RSK_t correspondence and hence is invertible, therefore so is the whole correspondence. Moreover, the $\binom{n}{2}$ additional values that we record with this procedure correspond to the $\binom{n}{2}$ entries in the Gelfand-Tsetlin pattern $G_t g_i(j)$, $1 \leq i \leq j \leq n-1$ that cannot be read off of $W_t f$.

Though this basic idea is fairly simple, we found that the method we chose to present is more straightforward and geometrically intuitive.

Remark 6.6. Our RSK_t map is based on one method of embedding \mathcal{D}_t^n into \mathcal{D}_{*t}^n by adding heavy weights after time t . There are clearly many ways to do this, and different methods will result in different bijections. One common feature of these bijections is that the key data that they see about f beyond its melon Wf will be a collection of left-to-right last passage values from time 0 to time t . Though we will not prove it here, all left-to-right last passage values are contained in $G_t f$, just as all bottom-to-top last passage values are contained in $W_t f$ by Proposition 3.12 (i).

Another option for constructing an RSK-like bijection would be to add heavy weights *before* time 0, essentially embedding f as an element of \mathcal{D}_\uparrow^n . We could also add weights to both sides of $[0, t]$ to embed f as a different distinguished element of an isometry class.

Bijections related to RSK exploring the use of different left-to-right or bottom-to-top last passage values have been constructed in Dauvergne (2020) and Garver, Patrias and Thomas (2018).

6.1. Bijectivity for lattice specializations and other restrictions

Bijectivity of the cadlag RSK correspondence $\text{RSK}_t : \mathcal{D}_t^n \rightarrow \mathcal{G}_t^n$ naturally implies that for any subset $A \subset \mathcal{D}_t^n$, that RSK_t is also a bijection from A to $\text{RSK}_t(A)$. For certain subsets A , we can explicitly identify $\text{RSK}_t(A)$, allowing us to recover previously known bijections and identify some new ones. In the next set of examples, we gather together the restricted bijections that correspond to classical integrable models of last passage percolation. In the $t = \infty$ setting where the Gelfand-Tsetlin pattern is dropped, these examples correspond exactly to those introduced immediately after Theorem 1.5.

For these examples, we say that a cadlag function f with positive jumps is **pure-jump** if df is an atomic measure.

Example 6.7. Let $t > 0$.

1. **Continuous functions.** If A is the set of continuous functions $f \in \mathcal{D}_t^n$ then $\text{RSK}_t(A)$ is the set of pairs $(w, g) \in \mathcal{G}_t^n$ such that w is also in A (e.g. w is continuous as well). This setting of continuous functions is studied in detail in Biane, Bougerol and O’Connell (2005).

2. **Unit jumps.** Let A be the set of pure-jump functions $f \in \mathcal{D}_t^n$, such that every jump of each f_i has size 1, and such that all jumps of f_i, f_j are at distinct locations for $i \neq j$. The $\text{RSK}_t(A)$ is the set of pairs $(w, g) \in \mathcal{G}_t^n$, where w is also in A , and all entries of g are nonnegative integers. In this setting, the space $\text{RSK}_t(A)$ is equivalent to the decorated Young Tableau defined in Nica (2017).
3. **Integer jumps at integer times.** Let A be the set of pure-jump functions $f \in \mathcal{D}_t^n$, such that every f_i only jumps at integer times and all jumps have integer values. Then $\text{RSK}_t(A)$ is the set of pairs $(w, g) \in \mathcal{G}_t^n$, where w is also in A , and all coordinates of g are nonnegative integers.
4. **Real jumps at integer times.** Let A be the set of pure-jump functions $f \in \mathcal{D}_t^n$, such that every f_i only jumps at integer times. Then $\text{RSK}_t(A)$ is the set of pairs $(w, g) \in \mathcal{G}_t^n$, where w is also in A .
5. **Bernoulli paths.** Suppose that additionally, $t \in \mathbb{N}$. Let A be the set of all functions $f \in \mathcal{D}_t^n$ that are linear with slope in $\{0, 1\}$ on every integer interval $[i, i + 1]$. Then $\text{RSK}_t(A)$ is the set of pairs $(w, g) \in \mathcal{G}_t^n$, where w is also in A , and all coordinates of g are nonnegative integers.

It is easy to verify each of the five examples above from the explicit formulas for RSK_t and RSK_t^{-1} in Section 6. Example 4 above corresponds to the usual RSK correspondence via Corollary 8.4 and Example 5 corresponds to the dual RSK correspondence via Corollary 8.7.


7. Preservation of uniform measure

In each of the five examples in Example 6.7, there are natural measures on A that push forward tractable measures on $\text{RSK}_t(A)$. By taking a limit as $t \rightarrow \infty$, we can also get tractable pushforward measures under the original melon map $W : \mathcal{D}^n \rightarrow \mathcal{D}^n$. Each of these measures corresponds to a classical integrable model of last passage percolation. This is summarized in the following table, essentially repeated from the introduction.

Example	Measure on \mathcal{D}^n	LPP model
6.7.1	Independent Brownian motions	Brownian LPP
6.7.2	Independent Poisson counting processes	Poisson lines LPP
6.7.3	Independent discrete-time geometric random walks	Geometric LPP
6.7.4	Independent discrete-time exponential random walks	Exponential LPP
6.7.5	Piecewise linear walks with independent Bernoulli slopes	S-J model

In all five examples above, the pushforward of these measures under W is the nonintersecting version of these objects. This is known in all cases, e.g. see O’Connell (2003a) and references therein, or Section 6 of Dauvergne, Nica and Virág (2019). The standard proofs of these facts require explicit computations involving determinants and Doob transforms. Here we give an alternate approach that is computation-free. We demonstrate this in the case of Bernoulli walks, Example 6.7.5.

We start with a more precise setup. For $k \in \mathbb{N} \cup \{\infty\}$, a function $f : [0, k] \rightarrow \mathbb{R}$ is a **Bernoulli path** if on every integer interval $[n, n + 1]$, f is linear with slope

FIG 5. 10 Bernoulli walks f and their melon Wf , 20 steps

in $\{0, 1\}$. A **Bernoulli walk** of drift $d \in [0, 1]$ is a random Bernoulli path whose slopes are independent Bernoulli random variables of mean d , and an **n -dimensional Bernoulli walk** with drift vector $d = (d_1, \dots, d_n)$ is an element of \mathcal{D}^n whose components are independent Bernoulli walks of drift d_i . See Figure 5 for an illustration.

Now, for $t \in \mathbb{N}$ and an ordered vector $x = (x_1 \geq \dots \geq x_n) \in \{0, 1, \dots\}^n$, let $\nu_t(x)$ denote the uniform measure on n -tuples of ordered Bernoulli paths $f : [0, t] \rightarrow \mathbb{R}$, $f = (f_1 \geq \dots \geq f_n)$ that satisfy $f(0) = 0$, $f(t) = x$. There are only finitely many such k -tuples, so uniform measure is well-defined. A measure μ on the space of n -tuples of ordered Bernoulli paths

$$X = (X_1 \geq X_2 \geq \dots \geq X_n), \quad X_i : [0, \infty) \rightarrow \mathbb{R}$$

is a **Bernoulli Gibbs measure** if for any integer $t > 0$, the conditional distribution under μ of $X|_{[0, t]}$ given $X|_{[t, \infty)}$ is $\nu_t(X(t))$. We start by showing that W maps Bernoulli walks to Bernoulli Gibbs measures.

Theorem 7.1. *Let $Y \in \mathcal{D}^n$ be a Bernoulli walk of drift d . Then the law of $WY \in \mathcal{D}_\uparrow^n$ is a Bernoulli Gibbs measure satisfying*

$$\lim_{t \rightarrow \infty} WY(t)/t = d^\circ \tag{7.1}$$

almost surely, where $d^\circ = (d_1^\circ \geq \dots \geq d_n^\circ)$ has the same components as d listed in decreasing order.

Proof. Fix $t > 0$ and let $A, \text{RSK}_t(A)$ be as in Example 6.7.5. The map RSK_t applied to Y up to time $t \in \mathbb{N}$ gives an n -tuple of ordered paths $W_t Y$ and a Gelfand-Tsetlin pattern $G_t Y$. We first show that the law of $W_t Y$ given $G_t Y$ is $\nu_t(WY(t))$, and then use this to deduce the Bernoulli Gibbs property. We first consider the case $d_i = 1/2$ for all i , so that the law of $Y|_{[0, t]}$ is the uniform measure on A .

By the bijectivity of RSK_t in Example 6.7.5, the law of $\text{RSK}_t Y$ is uniform on $\text{RSK}_t(A)$. Therefore, conditionally on $G_t Y$, which determines $WY(t)$, the law of $W_t Y$ is $\nu_t(WY(t))$. As an aside, the conditional law of $G_t Y$ given $W_t Y$ is also independent and uniform on Gelfand-Tsetlin patterns with n th row $WY(t)$.

Now for general $d \in [0, 1]^n$, the law of Y up to time $t \in \mathbb{N}$ is the uniform measure on A biased by the Radon-Nikodym derivative

$$2^{nt} \prod_{i=1}^n d_i^{Y_i(t)} (1 - d_i)^{t - Y_i(t)}.$$

Since this derivative only depends on $Y(t)$, the conditional law of Y given $Y(t)$ does not depend on the original drift vector d . Moreover, for any $i \in \{1, \dots, n\}$ we have

$$Y_1(t) + \dots + Y_i(t) = \sum_{j=1}^i G_t Y_j(i),$$

and so $Y(t)$ can be expressed from the the Gelfand-Tsetlin pattern $G_t Y$. Therefore conditionally on $G_t Y$, the law of $W_t Y = WY|_{[0,t]}$ does not depend on the drift d . Therefore as in the $d_i = 1/2$ case, the conditional law of $WY|_{[0,t]}$ given $G_t Y$ is still $\nu_t(WY(t))$.

We now use this conditional law to prove the Bernoulli Gibbs property. First, this conditional law implies the stronger claim that for any integers $s \leq t$, the conditional law of $WY|_{[0,s]}$ given $WY|_{[s,t]}$ and $G_t Y$ is still $\nu_t(WY(s))$. Therefore it suffices to show that as $t \rightarrow \infty$, that $WY|_{[0,s]}$ and $G_t Y$ are asymptotically independent. For this, it is enough to show that for any $k, \ell \in \{1, \dots, n\}$ with $\ell \geq n - k + 1$, for all large enough t we have

$$Y[(0, n)^k \rightarrow (\ell, t)^k] = Y[(s, n)^k \rightarrow (\ell, t)^k] + \sum_{i=n-k+1}^n Y_i(s). \quad (7.2)$$

Indeed, the right side of (7.2) only depends on $Y|_{[s,\infty)} - Y(s)$ which is independent of $WY|_{[0,s]}$, and $G_t Y$ can be expressed from the left hand side by varying ℓ, k . Equation (7.2) is equivalent to the claim that for large enough t , the right-most disjoint optimizer from $(0, n)^k$ to $(\ell, t)^k$ follows the bottom k paths up to time s . This follows from Remark 5.9.

We now show that WY satisfies (7.1). Define operators $\Lambda_k : \mathcal{D}^n \rightarrow \mathcal{D}^n$ by $\Lambda_k f(t) = f(kt)/k$. By the law of large numbers, as $k \rightarrow \infty$, $\Lambda_k Y(t) \rightarrow y(t) := dt$ uniformly on compact sets. Since W is continuous with respect to the uniform-on-compact topology and commutes with Λ_k by definition of last passage, we have

$$\lim_{k \rightarrow \infty} \Lambda_k WY(1) = \lim_{k \rightarrow \infty} W \Lambda_k Y(1) = W y(1).$$

Finally, W applied to linear functions just sorts them, so $W y(t) = d^\circ t$. \square

Next, we show that there is a unique Bernoulli Gibbs measure satisfying (7.1) for every possible d° .

Theorem 7.2. *For any $d \in [0, 1]^n$ with $d_1 \geq \dots \geq d_n$ there is a unique Gibbs measure μ_d on ordered n -tuples of Bernoulli paths in \mathcal{D}^n so that for $Y \sim \mu_d$,*

$$\lim_{t \rightarrow \infty} WY(t)/t = d \quad a.s. \quad (7.3)$$

Proof. Let μ_d denote the law of WY^d , where Y^d is a Bernoulli walk of drift $d = (d_1 \leq \dots \leq d_n)$.

Now let X be a sample from an arbitrary Bernoulli Gibbs measure satisfying (7.3). To show that $X \sim \mu_d$, it suffices check that $X|_{[0,s]} \stackrel{d}{=} WY^d|_{[0,s]}$ for all $s \in \mathbb{N}$. Let $\epsilon > 0$ and $v = (n, n-1, \dots, 1)$, set

$$\bar{d}_i = (d_i + \epsilon v_i) \wedge 1, \quad \underline{d}_i = (d_i - \epsilon v_i) \vee 0$$

and let $\underline{X} \sim \mu_{\underline{d}}$ and $\bar{X} \sim \mu_{\bar{d}}$. Consider any coupling of $X, \underline{X}, \bar{X}$. When $\underline{d}_i < d_i$, by (7.1) and (7.3), for all i we have $\mathbb{P}(\underline{X}_i(t) \leq X_i(t)) \rightarrow 1$ as $t \rightarrow \infty$. Otherwise $d_i = \underline{d}_i = 0$, but in this case $\underline{X}_i = 0$ as well, so $\underline{X}_i \leq X_i$ a.s. Thus, after a symmetric upper bound, we get that

$$\mathbb{P}A_t \rightarrow 1 \quad \text{as } t \rightarrow \infty, \quad \text{where } A_t = \{\underline{X}(t) \leq X(t) \leq \bar{X}(t)\}. \quad (7.4)$$

Now, the proof of Lemmas 2.6/2.7 in Corwin and Hammond (2014) shows that if $x \leq x'$ coordinatewise, $t \in \mathbb{N}$, and $Y \sim \nu_t(x), Y' \sim \nu_t(x')$ then we can couple Y and Y' so that $Y_i(s) \leq Y'_i(s)$ for all $i \in \{1, \dots, n\}, s \in [0, t]$. Therefore for every $t \in \mathbb{N}$, there is a new, t -dependent, coupling of $X, \underline{X}, \bar{X}$ such that for all $s \leq t$ and $i \in \{1, \dots, n\}$ we have

$$\underline{X}_i|_{[0,s]} \leq X|_{[0,s]} \leq \bar{X}_i|_{[0,s]} \quad (7.5)$$

with probability $\mathbb{P}A_t$. Now fixing s and taking $t \rightarrow \infty$, by compactness there exists a coupling of $X, \underline{X}, \bar{X}$ where (7.5) holds with probability $\lim_{t \rightarrow \infty} \mathbb{P}A_t = 1$. Now let $\epsilon \rightarrow 0$. The laws μ_d restricted to $[0, s]$ are continuous in d in the total variation norm, since the laws of $Y^d|_{[0,s]}$ are themselves continuous in d in total variation. Therefore there exists a coupling where (7.5) holds even when $\epsilon = 0$. In this case \underline{X} and \bar{X} both have distribution μ_d , and so in any coupling where (7.5) holds we must have $\underline{X} = X = \bar{X}$ on $[0, s]$, and hence $X|_{[0,s]} \stackrel{d}{=} WY^d|_{[0,s]}$, as required. \square

Theorems 7.1 and 7.2 and Proposition 3.12(i) yield the following immediate corollary.

Corollary 7.3 (Metric Burke property). *Last passage percolation across an n -dimensional Bernoulli walk ignores the order of the drift vector. More precisely, if $Y, Z \in \mathcal{D}^n$ are Bernoulli walks with drifts d, e satisfying $d^\circ = e^\circ$, then $WY \stackrel{d}{=} WZ$, and as functions of $x \leq y$ we have*

$$Y[(x, n) \rightarrow (y, 1)] \stackrel{d}{=} Z[(x, n) \rightarrow (y, 1)].$$

Burke's theorem normally refers to a certain invariance between arrivals and departures in a queuing processes; Corollary 7.3 is a kind of Burke property because it shows an invariance in the last passage value under exchanging the rows of the underlying environment. See O'Connell and Yor (2002).

The proof framework in this section goes through essentially verbatim if we start with a vector of independent geometric random walks, as in Example 6.7.4.

In this setting, the random walks Y are embedded in \mathcal{D}^n as pure-jump paths with jumps at integer times. The Pitman ordering condition on WY means that the output is a vector of geometric walks conditioned so that $WY_i(t) \geq WY_{i+1}(t+1)$ for all i, t . Measure-preservation for the remaining three examples in Example 6.7 can be deduced by standard limiting procedures. We leave the details of this to the interested reader.

8. Embedding classical versions of RSK

In this section, we relate our RSK map RSK_t to the usual RSK and dual RSK correspondences for nonnegative matrices. These correspondences are connected to last passage percolation in the lattice \mathbb{Z}^2 . We start with the connection to the standard RSK correspondence.

8.1. Young tableaux

We recall some basic combinatorial objects, see e.g. Stanley (1999) for a detailed reference. A **partition** is a weakly decreasing sequence $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k)$ of positive integers. The **size** of the partition is $|\lambda| = \sum_{i=1}^k \lambda_i$. To any partition λ , the **Young diagram** associated to λ is the set of squares $Y(\lambda) = \{(i, j) \in \mathbb{Z}^2 : 1 \leq i \leq \lambda_j\}$. A **semistandard Young tableau** of shape λ is a filling of the corresponding Young diagram with positive integers such that the entries are strictly increasing along columns and weakly increasing along rows.

There is a natural correspondence between Young tableaux and Pitman ordered cadlag paths with only integer-valued positive jumps at positive integer times as in Example 6.7.4. Consider a Young tableau T of shape $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k)$. Define $w \in \mathcal{D}_\uparrow^k$ by setting

$$w_i(t) = \# \text{ of entries in row } i \text{ of } T \text{ that are } \leq t.$$

In other words, the path w_i has jumps precisely at the times t which are equal to the entries of the i -th row of T . It is straightforward to check that with this definition, each w_i is a cadlag path with positive integer jumps at integer times. The fact that the entries of T are strictly increasing along columns implies that $w_i(t^-) \geq w_{i+1}(t)$ for all i, t , and so the $w \in \mathcal{D}_\uparrow^k$. This map from Young tableaux to Pitman ordered paths on this space is invertible. Moreover, for $n > k$ we can extend the collection (w_1, \dots, w_k) to a collection (w_1, \dots, w_n) of Pitman ordered paths by setting $w_i = 0$ for $i > k$.

There is also a well-known correspondence between Young tableaux and Gelfand-Tsetlin patterns with nonnegative integer entries. Namely, for a Young tableau T of shape λ whose largest entry is less than or equal to m , define a Gelfand-Tsetlin pattern $g = \{g_i(j) : i \leq j \leq m\}$ by setting $g_i(j)$ to be equal to the number of entries in row i of T that are less than or equal to j .

8.2. Classical RSK via Greene's theorem

The RSK correspondence is a map between the space of nonnegative matrices with integer entries and pairs of semistandard Young tableaux (P, Q) of equal shape. Typically it is described using a local *row insertion* algorithm. However, the RSK bijection can alternately be described using last passage percolation. For the restriction of RSK to permutation matrices (the Robinson-Schensted correspondence) this is due to Greene (1974). A version of Greene's theorem for RSK is also well-known, but appears to be folklore and we do not know of an original reference. See, for example, Theorem 24 in Hopkins (2014) or Krattenthaler (2006), Theorem 8.

In the following, we describe RSK based on this connection with Greene's theorem in the language of last passage values. For two points $p = (x, n), q = (y, m) \in \mathbb{Z}^2$ with $x \leq y$ and $n \geq m$, we say that a sequence of vertices $\pi = (\pi_1 = p, \dots, \pi_k = q)$ is a **directed path from p to q** if $\pi_i - \pi_{i-1} \in \{(1, 0), (0, -1)\}$ for all i . For an array $A = \{A_u : u \in \mathbb{Z}^2\}$ of nonnegative numbers, we can define the weight of any path π from p to q by

$$|\pi|_A = \sum_{v \in \pi} A_v. \quad (8.1)$$

We also define the last passage value

$$A[p \rightarrow q] = \max_{\pi} |\pi|_A, \quad (8.2)$$

where the maximum is taken over all possible paths π from p to q . More generally, for vectors $\mathbf{p} = (p_1, \dots, p_k), \mathbf{q} = (q_1, \dots, q_k)$, define the **multi-point last passage value**

$$A[\mathbf{p} \rightarrow \mathbf{q}] = \max_{\pi_1, \dots, \pi_k} |\pi_1|_A + \dots + |\pi_k|_A \quad (8.3)$$

where the maximum now is taken over all possible k -tuples of *disjoint* paths, where each π_i is a path from p_i to q_i . This is defined so long as a disjoint k -tuple exists. We also introduce the shorthand $A[p^{*k} \rightarrow q^{*k}]$ for the k -point last passage value from

$$(p - (0, k - 1), \dots, p - (0, 1), p) \rightarrow (q, q + (0, 1), \dots, q + (0, k - 1)).$$

The value $A[p^{*k} \rightarrow q^{*k}]$ is best thought of as a last passage value with k disjoint paths from p to q , hence the similar notation to the corresponding object in the cadlag setting. We are forced to stagger the start and end points of the paths to allow for disjointness.

Now for an $n \times m$ matrix of nonnegative integers A (equivalently, a restriction of a nonnegative array to the set $\{1, \dots, m\} \times \{1, \dots, n\}$), we can define a semistandard Young tableau, called the **recording tableau** Q with at most $n \wedge m$ rows and entries in $\{1, \dots, m\}$ by letting

$$A[(1, n)^{*k \wedge i} \rightarrow (i, 1)^{*k \wedge i}] = \# \text{ of entries in rows } 1, \dots, k \text{ of } Q \text{ that are } \leq i.$$

Similarly, define a semistandard Young tableau, called the **insertion tableau** P with at most $n \wedge m$ rows and entries in $\{1, \dots, n\}$ by letting

$$A[(1, n)^{*k \wedge i} \rightarrow (m, n - i + 1)^{*k \wedge i}] = \# \text{ of entries in rows } 1, \dots, k \text{ of } P \text{ that are } \leq i.$$

The RSK correspondence is the map $A \mapsto (Q, P)$. Observe that with these definitions P and Q have the same shape determined by the last passage values $A[(1, n)^{*k} \rightarrow (m, 1)^{*k}]$, $k = 1, \dots, m \wedge n$. By the correspondences between semistandard Young tableaux and Pitman ordered collections of cadlag paths and Gelfand-Tsetlin patterns we can associate to (Q, P) a pair $(WA, GA) \in \mathcal{G}_m^n$. Unravelling the bijections in Section 8.1, we get that for all $t \in [1, m]$ and $k \in \{1, \dots, n\}$, we have

$$\sum_{j=1}^k WA_j(t) = A[(1, n)^{*k \wedge [t]} \rightarrow ([t], 1)^{*k \wedge [t]}], \quad (8.4)$$

and for $t < 1$, we have $WA_j(t) = 0$. Also, for $1 \leq k \leq i \leq n$ we have

$$\sum_{j=1}^k GA_j(i) = A[(1, n)^{*k} \rightarrow (m, n - i + 1)^{*k}]. \quad (8.5)$$

8.3. Classical RSK and the melon map

For a nonnegative $n \times m$ matrix A , define $f^A \in \mathcal{D}_m^n$ by

$$f_k^A(0^-) = 0, \quad \text{and} \quad f_k^A(t) - f_k^A(s) = \sum_{r \in (s, t]} A_{r, k}. \quad (8.6)$$

We will show that discrete last passage values across A equal last passage values across f^A .

Proposition 8.1. *For all tuples of points \mathbf{p}, \mathbf{q} such that $A[\mathbf{p} \rightarrow \mathbf{q}]$ is defined, we have*

$$A[\mathbf{p} \rightarrow \mathbf{q}] = f^A[\mathbf{p} \rightarrow \mathbf{q}], \quad (8.7)$$

To prove Proposition 8.1, we will show that lattice last passage values can be equivalently defined using unions of possibly overlapping paths.

We first prove this for endpoints that lie in a packed staircase configuration.

Lemma 8.2. *Let \mathbf{p}, \mathbf{q} be such that $p_i = p_{i-1} + (1, 1)$ and $q_i = q_{i-1} + (1, 1)$ for all i . Then*

$$A[\mathbf{p} \rightarrow \mathbf{q}] = \max_{\pi_1, \dots, \pi_k} \sum_{v \in \bigcup_i \pi_i} A_v, \quad (8.8)$$

where the maximum is over all k -tuples of paths π_i from p_i to q_i , without any disjointness condition enforced. In the union in (8.8), weights on multiple paths are only counted once.




FIG 6. Example of how the overlapping paths in the proof of Lemma 8.2 are moved down to find a better configuration of non-intersecting paths.

For the proof, it will be easier to imagine the coordinate system as rotated clockwise by 45 degrees, and scaled up by $\sqrt{2}$ as in Figure 6. After this rotation, all the points p_i lie on a common vertical line $\mathbb{Z} \times \{p^*\}$. Similarly, all the points q_i lie on a common vertical line $\mathbb{Z} \times \{q^*\}$. Moreover, with this rotation any path π from p_i to q_i for some i gets transformed to the graph of a function $\hat{\pi} : \{p^*, \dots, q^*\} \rightarrow \mathbb{Z}$ with steps of ± 1 . That is, π gets transformed to a simple random walk path $\hat{\pi}$.

Proof. The fact that $\text{LHS} \leq \text{RHS}$ in (8.8) follows since we are maximizing over a smaller set on the left. To achieve the opposite inequality, we just need to show that there is a set of disjoint paths π_i that achieves the maximum on the right side of (8.8). Without loss of generality, by passing to order statistics, we may assume that the maximum is achieved on a k -tuple of paths τ satisfying

$$\hat{\tau}_1(x) \leq \hat{\tau}_2(x) \leq \dots \leq \hat{\tau}_k(x), \quad (8.9)$$

for all $x = p^*, p^* + 1, \dots, q^*$.

Now consider the set S of all k -tuples which achieve the maximum in (8.8) and satisfy (8.9). We put a partial order on this set by saying that $\pi \leq \tau$ if $\hat{\pi}_i(x) \leq \hat{\tau}_i(x)$ for all $i \in \{1, \dots, k\}, x = p^*, p^* + 1, \dots, q^*$. Let π be a minimal element of the finite set S . We show that π consists of disjoint paths.

Suppose not. Then there exists an $i < j$ and a value x such that $\hat{\pi}_i(x) = \hat{\pi}_j(x)$. We may also assume that i is the minimal such index where there is such a conflict, and hence that

$$\hat{\pi}_i(x) - 2 \geq \hat{\pi}_\ell(x) \quad \text{for all } i > \ell. \quad (8.10)$$

Let $I = \{a, \dots, b\} \subset \{p^*, \dots, q^*\}$ be the largest interval containing x such that $\hat{\pi}_i = \hat{\pi}_j$ on I . Since the start and endpoints of $\hat{\pi}_i, \hat{\pi}_j$ are distinct, we have $p^* < a, b < q^*$. Therefore $\hat{\pi}_i, \hat{\pi}_j$ are well-defined at $a - 1$ and $b + 1$ and satisfy

$$\hat{\pi}_i(a - 1) = \hat{\pi}_j(a - 1) - 2, \quad \hat{\pi}_i(b + 1) = \hat{\pi}_j(b + 1) - 2.$$

Therefore the function $\hat{\pi}'_i$ which is equal to $\hat{\pi}_i$ on I^c , and shifted down by 2 units, $\hat{\pi}'_j = \hat{\pi}_j - 2$, on I is also a simple random walk path, see Figure 6. Thus the k -tuple $\pi' = (\pi_1, \dots, \pi'_i, \pi_{i+1}, \dots, \pi_k)$ also consists of paths from p_i to q_i . Moreover, the vertices covered by π' contain all the vertices covered by π , so because the weights are all non-negative, π' must also achieve the maximum in (8.8). Finally, by (8.10), the k -tuple π' still satisfies inequalities in (8.9), so $\pi' \in S$. On the other hand, $\pi' \leq \pi$ by construction, contradicting the minimality of π . \square

We can now extend this to general endpoints.

Lemma 8.3. *For any \mathbf{p}, \mathbf{q} such that $A[\mathbf{p} \rightarrow \mathbf{q}]$ is defined, we have*

$$A[\mathbf{p} \rightarrow \mathbf{q}] = \max_{\pi_1, \dots, \pi_k} \sum_{v \in \bigcup_i \pi_i} A_v, \quad (8.11)$$

where the maximum is over all k -tuples of paths π_i from p_i to q_i , without any disjointness condition enforced.

Proof. We can find a pair of vectors $(\mathbf{p}', \mathbf{q}')$ that are of the form in Lemma 8.2 such that there are sets of disjoint paths π from \mathbf{p}' to \mathbf{p} and τ from \mathbf{q} to \mathbf{q}' . Let H be a nonnegative array which is equal to 1 for $x \in \cup \pi \cup \tau$, and zero otherwise, and let $A' = A + sH$. Then for large enough s , letting $m = |\cup \pi \cup \tau|$ we have

$$A'[\mathbf{p}' \rightarrow \mathbf{q}'] = ms + A[\mathbf{p} \rightarrow \mathbf{q}], \quad (8.12)$$

since any optimal disjoint paths from \mathbf{p}' to \mathbf{q}' will necessarily follow π and τ . By Lemma 8.2, we similarly have that

$$A'[\mathbf{p}' \rightarrow \mathbf{q}'] = ms + R, \quad (8.13)$$

where R denotes the right hand side of (8.11). Equating (8.12) and (8.13) completes the proof. \square

Proof of Proposition 8.1. Any disjoint lattice paths from \mathbf{p} to \mathbf{q} can be mapped to disjoint cadlag paths, so we have $A[\mathbf{p} \rightarrow \mathbf{q}] \leq f^A[\mathbf{p} \rightarrow \mathbf{q}]$. Now let

$$f^A\{\mathbf{p} \rightarrow \mathbf{q}\} = \max_{\pi} |\pi|_f,$$

where the maximum is now over k -tuples from \mathbf{p} to \mathbf{q} with the disjointness condition removed. In $|\pi|_f$ we only count weights once even if they are covered by multiple paths. Let π be a k -tuple that achieves this maximum, and define a new k -tuple $\lfloor \pi \rfloor$ by setting $\lfloor \pi \rfloor_i(t) = \pi_i(\lfloor t \rfloor)$ for all i, t . Since f^A has only positive jumps and is constant between integer times, $\lfloor \pi \rfloor$ also achieves this maximum. Each $\lfloor \pi \rfloor_i$ corresponds to a discrete lattice path π'_i from p_i to q_i , and we have the equality

$$|\lfloor \pi \rfloor|_{f^A} = \sum_{v \in \bigcup_i \pi'_i} A_v.$$

Therefore by Lemma 8.3, $f^A\{\mathbf{p} \rightarrow \mathbf{q}\} \leq A[\mathbf{p} \rightarrow \mathbf{q}]$. Since $f^A[\mathbf{p} \rightarrow \mathbf{q}] \leq f^A\{\mathbf{p} \rightarrow \mathbf{q}\}$, we have that $f^A[\mathbf{p} \rightarrow \mathbf{q}] \leq A[\mathbf{p} \rightarrow \mathbf{q}]$ as well. \square

Finally, we can show that the usual RSK bijection is a special case of the cadlag RSK bijection.

Corollary 8.4. *Let A be an $n \times m$ matrix. Define $f^A \in \mathcal{D}_m^n$ via the formula in (8.6). Then with (WA, GA) as in (8.4) and (8.5) we have that*

$$(WA, GA) = (W_m f^A, G_m f^A) = \text{RSK}_m(f^A).$$

Proof. By Proposition 8.1 and tracing through the definitions, it suffices to show that

$$f^A[(0, n)^k \rightarrow (t, j)^k] = f^A[(1, n)^{*k \wedge [t]} \rightarrow ([t], j)^{*k \wedge [t]}] \quad (8.14)$$

for all t, j, k with $k \leq n + 1 - j$. For $k \geq [t]$, both sides pick up all weights of A in the box $\{1, \dots, [t]\} \times \{1, \dots, n\}$. For $k < [t]$, notice that since $f^A(t) = 0$ for all $t < 1$ and f^A is unchanging between integer times, that $f^A[(0, n)^k \rightarrow (t, j)^k] = f^A[(1, n)^k \rightarrow ([t], j)^k]$. Moreover, essential disjointness at times 1 and $[t]$ implies that any disjoint k -tuple from $(1, n)^k$ to $([t], j)^k$ has the same length as some disjoint k -tuple from $(1, n)^{*k} \rightarrow ([t], j)^{*k}$. \square

Remark 8.5. While the RSK correspondence is defined only for matrices with nonnegative integer entries, the maps (8.5) and (8.4) are still defined for matrices A with nonnegative real entries; there is just no longer a connection with Young tableaux. Proposition 8.1, Lemma 8.2, and Corollary 8.4 still hold in this generality and the proofs go through verbatim.

We can also use the framework of cadlag RSK to construct an infinite RSK map for arrays A indexed by $\{1, \dots, n\} \times \mathbb{N}$. Such a map was first constructed in O’Connell (2003b). Again, formula (8.6) takes such an array to a function $f^A \in \mathcal{D}^n$, and the infinite melon Wf^A can be expressed in terms of lattice passage values as in Corollary 8.4. The map $A \mapsto Wf^A$ is invertible if $f^A \in \mathcal{D}_\downarrow^n$, which is a fairly mild constraint, see Proposition 5.8. The fact that this map is typically invertible can be thought of as saying that in the $n \times m$ matrix RSK correspondence, the insertion tableau P contains less and less information about the first $O(1)$ -many columns in the matrix as $m \rightarrow \infty$. This is natural, since P is expressed in terms of last passage values across all m columns, and so we should not be able to glean much information about any set of $O(1)$ -many columns from P alone.

8.4. Dual RSK

When restricting to permutation matrices, the dual RSK correspondence is given by replacing the row insertion RSK algorithm with a *column insertion* algorithm. Dual RSK does not extend from permutation matrices to all matrices with nonnegative entries, but it does extend to arbitrary $0 - 1$ matrices. As in the case of the usual RSK bijection, there is a version of Greene’s theorem for dual RSK (see Theorem 10 in Krattenthaler (2006)) that represents dual RSK in terms of last passage percolation, and therefore allows us to show that dual RSK is a special case of cadlag RSK. As the details connecting cadlag RSK and

dual RSK are similar to the case of the usual RSK correspondence, we only include theorem statements here.

Let A be an $n \times m$ matrix of 0s and 1s. For two points $p = (x, k), q = (y, \ell)$ with $x \leq y$ and $k \geq \ell$, we say that $\pi = (\pi_1 = p, \dots, \pi_k = q)$ is a **dual path from p to q** if $\pi_i - \pi_{i-1} \in \{(1, s) : s \in \mathbb{Z}_{\leq 0}\}$ for all i . That is π is a path that moves strictly to the right and weakly up at every step. Definitions (8.1), (8.2), and (8.3) still make sense for dual paths and we write $A\{\mathbf{p} \rightarrow \mathbf{q}\}$ for a last passage value with dual paths.

Now, for a filling Q of a Young diagram Y , we write Q^T for the transposed filling of the transposed Young diagram Y^T , i.e. a cell $(a, b) \in Y$ if and only if $(b, a) \in Y^T$ and $Q^T(b, a) = Q(a, b)$. For an $n \times m$ matrix A of 0s and 1s, we define a semistandard Young tableau P with at most $n \wedge m$ rows and entries in $\{1, \dots, m\}$ by letting

$$A\{(1, n)^{*k \wedge i} \rightarrow (i, 1)^{*k \wedge i}\} = \# \text{ of entries in rows } 1, \dots, k \text{ of } Q^T \text{ that are } \leq i.$$

Also define a semistandard Young tableau P with at most $n \wedge m$ rows and entries in $\{1, \dots, n\}$ by letting

$$A\{(1, n)^{*k \wedge i} \rightarrow (m, n-i+1)^{*k \wedge i}\} = \# \text{ of entries in rows } 1, \dots, k \text{ of } P \text{ that are } \leq i.$$

The dual RSK correspondence is the map $A \mapsto (Q, P)$ which maps 0–1 matrices to pairs of semistandard Young tableaux such that the shapes of Q and P are **conjugate**, i.e. Q^T has the same shape as P . Observe that with the above definitions Q^T and P have the same shape.

The fact that Q , rather than Q^T , is a semistandard Young tableau is a consequence of the differences in the definition of paths and dual paths. Nonetheless, to connect this definition to cadlag RSK it is still Q^T that we want to write as a collection of Pitman ordered paths (WA_1, \dots, WA_n) . To do this, we embed Q^T not as a collection of cadlag paths with jumps, but rather as a collection of paths with piecewise linear increments. For all $t \in \{1, \dots, m\}$ and $k \in \{1, \dots, n\}$, we write

$$\sum_{j=1}^k WA_j(t) = A\{(1, n)^{*k \wedge t} \rightarrow (t, 1)^{*k \wedge t}\}. \quad (8.15)$$

We also set $WA(0) = 0$, and let each line WA_i be linear on every interval $[t, t+1]$ with $t \in \mathbb{Z}$. Since Q is a semistandard Young tableau, with this definition each line WA_i either has slope 0 or slope 1 on every interval. We also turn P into a Gelfand-Tsetlin pattern G in the usual way. For $1 \leq k \leq i \leq n$ we have

$$\sum_{j=1}^k GA_j(i) = A\{(1, n)^{*k} \rightarrow (m, n-i+1)^{*k}\}. \quad (8.16)$$

We now connect this description to cadlag RSK. For an $n \times m$ $\{0, 1\}$ -matrix A , define $\ell^A \in \mathcal{D}_m^n$ by letting

$$\ell_k^A(0) = 0, \quad \text{and} \quad \ell_k^A(t) - \ell_k^A(s) = \sum_{r \in [s+1, t]} A_{r, k} \quad \text{for } s, t \in \mathbb{Z}, \quad (8.17)$$

and by letting each ℓ_k^A be linear between integers. We then have the following analogue of Proposition 8.1.

Proposition 8.6. *Let \mathbf{p}, \mathbf{q} be such that $A\{\mathbf{p} \rightarrow \mathbf{q}\}$ is defined. Then*

$$A\{\mathbf{p} \rightarrow \mathbf{q}\} = \ell^A[\mathbf{p} \rightarrow \mathbf{q}], \quad (8.18)$$

The proof of Proposition 8.6 is similar to the proof of Proposition 8.1. Proposition 8.6 leads to an analogue of Corollary 8.4.

Corollary 8.7. *Let A be an $n \times m$ matrix of 0s and 1s. Then with (WA, GA) as in (8.15) and (8.16) we have that*

$$(WA, GA) = (W_m \ell^A, G_m \ell^A) = \text{RSK}_m(\ell^A).$$

Remark 8.8. While the dual RSK correspondence is defined only for matrices with $\{0, 1\}$ entries, the maps (8.16) and (8.15) are still defined for matrices A with arbitrary real entries; there is just no longer a connection with Young tableaux. Proposition 8.6 and Corollary 8.7 still hold in this generality.

Appendix A: Appendix: technical proofs

Proof of Lemma 3.2. Set

$$s(x, y) = \sup_{z \in [x, y]} f_2(z) - f_1(z^-) = \sup_{z \in [x, y]} s(z, z),$$

so that we have

$$s(x, y^-) = \sup_{z \in [x, y)} f_2(z) - f_1(z^-).$$

For each x , the function $s(x, \cdot)$ is increasing. Also, since the functions f_i are cadlag with positive jumps, we have that $s(x, \cdot)$ is cadlag. (Note that this would not hold if we allowed negative jumps in f_1 .) We also have

$$s(x, y) - s(x, y^-) = [s(y, y) - s(x, y^-)]^+ \leq f_2(y) - f_2(y^-). \quad (\text{A.1})$$

We can explicitly write last passage values across f as

$$f[(x, 2) \rightarrow (y, 1)] = f_1(y) - f_2(x^-) + s(x, y). \quad (\text{A.2})$$

Specializing to the case $x = 0$, and using that $Wf_1 + Wf_2 = f_1 + f_2$, we have

$$Wf_1(t) = f_1(t) + s(0, t) \quad \text{and} \quad Wf_2(t) = f_2(t) - s(0, t). \quad (\text{A.3})$$

From the fact that $s(0, \cdot)$ is cadlag and increasing, the function Wf_1 is cadlag with only positive jumps. Also, by (A.1), the function Wf_2 is cadlag with only positive jumps, so W maps \mathcal{D}^2 to itself. The last passage value across Wf is

$$Wf[(x, 2) \rightarrow (y, 1)] = Wf_1(y) - Wf_2(x^-) + \sup_{z \in [x, y]} Wf_2(z) - Wf_1(z^-).$$

Substituting the formulas (A.3) we get that this equals

$$f_1(y) + s(0, y) - f_2(x^-) + s(0, x^-) + \sup_{z \in [x, y]} [s(z, z) - s(0, z) - s(0, z^-)]. \quad (\text{A.4})$$

By comparing with (A.2), we can see that the lemma will follow from the equality

$$s(x, y) - s(0, y) - s(0, x^-) = \sup_{z \in [x, y]} [s(z, z) - s(0, z) - s(0, z^-)]. \quad (\text{A.5})$$

To prove (A.5), we divide into cases. First suppose that $s(0, x^-) = s(0, y)$. In this case, since $s(0, \cdot)$ is nondecreasing, we have that $s(0, z) = s(0, z^-) = s(0, x^-) = s(0, y)$ for all $z \in [x, y]$. Therefore

$$\begin{aligned} \sup_{z \in [x, y]} [s(z, z) - s(0, z) - s(0, z^-)] &= \sup_{z \in [x, y]} s(z, z) - s(0, y) - s(0, x^-) \\ &= s(x, y) - s(0, y) - s(0, x^-). \end{aligned}$$

We turn to the case when $s(0, x^-) < s(0, y)$. By definition,

$$s(0, y) = s(0, x^-) \vee s(x, y), \quad \text{so} \quad s(0, y) = s(x, y). \quad (\text{A.6})$$

Set

$$z_0 = \sup\{z \in [x, y] : s(0, z^-) = s(0, x^-)\}.$$

The function $s(0, \cdot^-)$ is left continuous, so this is in fact a maximum. In particular, since s is nondecreasing, for each $z_1 > z_0$

$$s(0, z_1) \geq s(0, z_1^-) > s(0, z_0^-).$$

So we have, by definition of s

$$s(0, z_1) = s(0, z_0^-) \vee s(z_0, z_1) = s(z_0, z_1).$$

By the right continuity of $s(0, \cdot)$ and $s(z_0, \cdot)$, as $z_1 \downarrow z_0$ we get $s(0, z_0) = s(z_0, z_0)$. By choosing $z = z_0$ in the supremum on the right hand side of (A.5) we get

$$-s(0, x^-) = -s(0, z_0^-) \leq \sup_{z \in [x, y]} [s(z, z) - s(0, z) - s(0, z^-)] \quad (\text{A.7})$$

Since $s(z, z) \leq s(0, z)$, and the fact that s is nondecreasing, the right hand side can be upper bounded by

$$\sup_{z \in [x, y]} [-s(0, z^-)] = -s(0, x^-).$$

so (A.7) is in fact an equality. Since $s(0, y) = s(x, y)$ by (A.6), this proves the preservation of last passage values in (A.5). \square

Proof of Lemma 4.3. We use the notation from the proof of Lemma 3.2 in this appendix. Recall that

$$s(x, y) = \sup_{z \in [x, y]} f_2(z) - f_1(z^-) = \sup_{z \in [x, y]} s(z, z).$$

Let

$$I_f(x, y) = \{z \in [x, y] : s(z, z) = s(x, y)\}.$$

This is the set of all possible jump times from line 2 to 1 for geodesics from $(x, 2)$ to $(y, 1)$ in f . Also set $r(x, y) = \sup_{z \in [x, y]} [s(z, z) - s(0, z) - s(0, z^-)]$ and let

$$I_{Wf}(x, y) = \{z \in [x, y] : r(z, z) = r(x, y)\}.$$

By (A.4) this is the set of all possible jump times from line 2 to 1 for geodesics from $(x, 2)$ to $(y, 1)$ in Wf . Then the desired ordering on geodesics holds if and only if

$$\inf I_{Wf}(x, y) \leq \inf I_f(x, y), \quad \text{and} \quad \sup I_{Wf}(x, y) \leq \sup I_f(x, y).$$

Again, we first deal with the case when $s(0, x^-) = s(0, y)$. In this case, for all $w \in [x, y]$ we have

$$r(w, w) = s(w, w) - 2s(0, x^-),$$

so $I_f(x, y) = I_{Wf}(x, y)$. Now suppose $s(0, x^-) < s(0, y)$. Define

$$A = \sup\{z \in [x, y] : s(0, z^-) = s(0, x^-)\}, \quad B = \inf\{z \in [x, y] : s(0, z) = s(0, y)\}.$$

We clearly have $A \leq B$. Moreover, $s(0, y) = s(x, y)$ in this case, and so for $z < I$, we must have $s(z, z) < s(x, y)$. Hence $I_f(x, y) \subset [B, y]$. To complete the proof, we show $I_{Wf}(x, y) \subset [x, A]$. Since (A.7) is an equality in this case, see the discussion following that inequality, at every point $w \in I_{Wf}(x, y)$, we have $r(w, w) = -s(0, x^-)$. Moreover, for $w > A$, we have

$$r(w, w) = s(w, w) - s(0, w) - s(0, w^-) \leq -s(0, w^-) < -s(0, x^-).$$

Therefore $I_{Wf}(x, y) \subset [x, A]$. □

References

- Aigner, F. and Frieden, G. (2020). qRSt: A probabilistic Robinson–Schensted correspondence for Macdonald polynomials, *arXiv:2009.03526*. [MR4311960](#)
- Biane, P., Bougerol, P. and O’Connell, N. (2005). Littelmann paths and Brownian paths, *Duke Mathematical Journal* **130**(1): 127–167. [MR2176549](#)
- Bougerol, P. and Jeulin, T. (2002). Paths in Weyl chambers and random matrices, *Probability theory and related fields* **124**(4): 517–543. [MR1942321](#)
- Bufetov, A. and Matveev, K. (2018). Hall–Littlewood RSK field, *Selecta Mathematica* **24**(5): 4839–4884. [MR3874706](#)

- Corwin, I. (2020). Invariance of polymer partition functions under the geometric RSK correspondence, *arXiv:2001.01867*.
- Corwin, I. and Hammond, A. (2014). Brownian Gibbs property for Airy line ensembles, *Inventiones mathematicae* **195**(2): 441–508. [MR3152753](#)
- Corwin, I., O’Connell, N., Seppäläinen, T. and Zygouras, N. (2014). Tropical combinatorics and whittaker functions, *Duke Mathematical Journal* **163**: 513–563. [MR3165422](#)
- Dauvergne, D. (2022). Hidden invariance of last passage percolation and directed polymers, *Annals of Probability* **50**(1): 18–60.
- Dauvergne, D., Nica, M. and Virág, B. (2019). Uniform convergence to the Airy line ensemble, *arXiv preprint arXiv:1907.10160*. [MR4260467](#)
- Dauvergne, D., Ortmann, J. and Virág, B. (2018). The directed landscape, *arXiv:1812.00309*. To appear in *Acta Mathematica*. [MR4035169](#)
- Dauvergne, D. and Virág, B. (2021). The scaling limit of the longest increasing subsequence, *arXiv:2104.08210*. [MR4083455](#)
- Dauvergne, D. and Zhang, L. (2021). Disjoint optimizers and the directed landscape, *arXiv:2102.00954*. [MR4035169](#)
- Draief, M., Mairesse, J. and O’Connell, N. (2005). Queues, stores, and tableaux, *Journal of Applied Probability* **42**(4): 1145–1167. [MR2203829](#)
- Fulton, W. (1997). *Young tableaux: with applications to representation theory and geometry*, Vol. 35, Cambridge University Press. [MR1464693](#)
- Garver, A., Patrias, R. and Thomas, H. (2018). Minuscule reverse plane partitions via quiver representations, *arXiv:1812.08345*. [MR4098265](#)
- Georgiou, N., Rassoul-Agha, F. and Seppäläinen, T. (2017). Stationary cocycles and Busemann functions for the corner growth model, *Probability Theory and Related Fields* **169**(1): 177–222. [MR3704768](#)
- Greene, C. (1974). An extension of Schensted’s theorem, *Advances in Mathematics* **14**(2): 254–265. [MR0354395](#)
- Henrikson, J. (1999). Completeness and total boundedness of the Hausdorff metric, *MIT Undergraduate Journal of Mathematics* **1**: 69–80.
- Hopkins, S. (2014). RSK via local transformations, *Unpublished notes*, <http://www-users.math.umn.edu/~shopkins/docs/rsk.pdf>.
- Kirillov, A. (2001). Introduction to tropical combinatorics, *Physics and combinatorics*, World Scientific, pp. 82–150. [MR1872253](#)
- Knuth, D. (1970). Permutations, matrices, and generalized Young tableaux, *Pacific journal of mathematics* **34**(3): 709–727. [MR0272654](#)
- König, W., O’Connell, N. and Roch, S. (2002). Non-colliding random walks, tandem queues, and discrete orthogonal polynomial ensembles, *Electronic Journal of Probability* **7**. [MR1887625](#)
- Krattenthaler, C. (2006). Growth diagrams, and increasing and decreasing chains in fillings of Ferrers shapes, *Advances in Applied Mathematics* **37**(3): 404–431. [MR2261181](#)
- Logan, B. F. and Shepp, L. A. (1977). A variational problem for random Young tableaux, *Advances in mathematics* **26**(2): 206–222. [MR1417317](#)
- Nica, M. (2017). Decorated Young tableaux and the Poissonized Robinson–Schensted process, *Stochastic Processes and their Applications* **127**(2): 449–

474. [MR3583759](#)
- Noumi, M. and Yamada, Y. (2002). Tropical Robinson-Schensted-Knuth correspondence and birational Weyl group actions, *math-ph/0203030*. [MR2074600](#)
- O’Connell, N. (2003a). Conditioned random walks and the RSK correspondence, *Journal of Physics A: Mathematical and General* **36**(12): 3049. [MR1986407](#)
- O’Connell, N. (2003b). A path-transformation for random walks and the Robinson-Schensted correspondence, *Transactions of the American Mathematical Society* **355**(9): 3669–3697. [MR1990168](#)
- O’Connell, N. and Pei, Y. (2013). A q -weighted version of the Robinson-Schensted algorithm, *Electronic Journal of Probability* **18**. [MR3126578](#)
- O’Connell, N. and Yor, M. (2002). A representation for non-colliding random walks, *Electronic Communications in Probability* **7**: 1–12. [MR1887169](#)
- Pitman, J. W. (1975). One-dimensional Brownian motion and the three-dimensional Bessel process, *Advances in Applied Probability* **7**(3): 511–526. [MR0375485](#)
- Robinson, G. d. B. (1938). On the representations of the symmetric group, *American Journal of Mathematics* **60**(3): 745–760. [MR1507943](#)
- Romik, D. (2015). *The surprising mathematics of longest increasing subsequences*, Vol. 4, Cambridge University Press. [MR3468738](#)
- Sagan, B. E. (2013). *The symmetric group: representations, combinatorial algorithms, and symmetric functions*, Vol. 203, Springer Science & Business Media. [MR1824028](#)
- Schensted, C. (1961). Longest increasing and decreasing subsequences, *Canadian Journal of mathematics* **13**: 179–191. [MR0121305](#)
- Seppäläinen, T. (1998). Exact limiting shape for a simplified model of first-passage percolation on the plane, *The Annals of Probability* **26**(3): 1232–1250. [MR1640344](#)
- Stanley, R. P. (1999). Enumerative combinatorics, volume 2, *Cambridge University Press*. [MR1676282](#)
- Vershik and Kerov (1977). Asymptotics of the Plancherel measure of the symmetric group and the limiting form of Young tableaux, *Soviet. Math. Dokl.*, Vol. 18, pp. 527–531. [MR0480398](#)