ALMOST PERIODIC DINI THEOREMS

GRAY H. MEISTERS†

Introduction. Dini's classical theorem [4] states that if a monotone sequence of real valued continuous functions converges pointwise to a continuous limit on a compact space, then the convergence is uniform. It is, of course, well known that the corresponding statement fails on a non-compact space. However, in light of the Bohr compactification of the real line $\mathbf{R}$, one might expect that the corresponding statement for Bohr almost periodic functions would be true. (See e.g., [7].) Alas, this too is false as can be seen by examples such as $f_n(t) = [(2 + \sin\sqrt{2}t + \cos t)/4]^n$, which does not converge uniformly on $\mathbf{R}$ even though $f_n \downarrow 0$ pointwise on $\mathbf{R}$ (and even though the mean values $M(f_n)$ tend to zero). Nevertheless, by adding some appropriate additional condition one does obtain true versions of a Dini-type theorem for almost periodic functions. One such version has been given by Luigi Amerio in [1] and [2]. Moreover, Salomon Bochner has given in [3] a rather general Dini-type theorem for almost automorphic functions which includes both Amerio's theorem and the classical Dini theorem (although in the latter case, as Bochner points out, his functions must be defined on a sequentially compact space rather than on a compact space). Finally Mario Dolcher [5] has given a proof of Amerio's theorem based on the Bohr compactification of the real line.

The purpose of this note is to show that Amerio's theorem can be deduced briefly from an abstract Dini theorem (Proposition 3.4 on page 91 in [10]), which was evidently first stated by Krein in [8] for partially ordered normed linear spaces, and which is already known to contain the classical Dini theorem (cf. [10] pp. 92-93). We also give another criterion, different from Amerio's, for the uniform convergence of monotone sequences of a. p. functions on $\mathbf{R}$.

The Abstract Dini Theorem. In order to make our argument complete and self-contained (and for the convenience of the reader) we first state and prove a simplified version of the abstract Dini theorem which on the one hand is entirely adequate for the deduction of Amerio's result and on the other hand can be given a rather short

---

†Supported in part by NSF Grant GP-38029.

Received by the Editors December 7, 1973 and in revised form April 3, 1974.

proof (different from others we have seen, but closely related to arguments appearing on page 65 of [11]).

By an *ordered topological vector space* (OTVS) X we shall mean a real locally convex topological vector space X equipped with a partial order relation, $f \leq g$, which satisfies the following three properties:

- (i) $f \leq g \Rightarrow f + h \leq g + h$ for all $h \in X$.
- (ii) $f \leq g \Rightarrow \alpha f \leq \alpha g$ for all nonnegative $\alpha \in \mathbf{R}$.
- (iii) To each neighborhood U of 0 in X , there corresponds a neighborhood V of 0 such that if $0 \leq f \leq g$, and $g \in V$, then $f \in U$.

Namioka [9] calls a partially ordered topological vector space satisfying (iii) "locally full", while Peressini [10] calls it "an OTVS with a normal cone". See Theorems 4.8 and 4.9 of [9] and Propositions 1.3 and 1.5 on pages 62 and 63 of [10]. Such spaces were evidently first introduced by Krein in [8]. Clearly the Banach space A of all real valued Bohr almost periodic functions $f(t)$ on the real line $\mathbf{R}$ is an OTVS when equipped with the supremum norm and the natural partial order, $f \leq g$ whenever $f(t) \leq g(t)$ for all real t .

AN ABSTRACT DINI THEOREM. *If a decreasing sequence f_n in an OTVS X converges weakly to an element f in X , with all $f_n \geq f$, then f_n converges to f in the original ("strong") topology of X .*

PROOF. A separation form of the Hahn-Banach theorem (e.g., Theorem 3.4 (b) in [11]) implies the well-known important fact that the weak closure of a convex subset C of a LCTVS X is equal to the original closure of C . (See e.g., Theorem 3.12 of [11].) Therefore, for each neighborhood U of 0 in X , there exists an integer N and non-negative scalars $\alpha_1, \dots, \alpha_N$ such that $\sum_{k=1}^N \alpha_k = 1$ and $\sum_{k=1}^N \alpha_k f_k$ belongs to $f + V$, where V is the neighborhood of 0 which corresponds to U by property (iii). But for all $m > N$ we also have $f \leq f_m \leq f_N \leq \sum_{k=1}^N \alpha_k f_k$ which yields $0 \leq f_m - f \leq (\sum_{k=1}^N \alpha_k f_k) - f \in V$. Property (iii) now gives $f_m \in f + U$, which proves the theorem.

We remark that the hypothesis "with all $f_n \geq f$ " can be omitted, but then the proof is slightly longer (and more technical) and we do not need this stronger version in this note.

Amerio's Condition. A sequence $s_m \in \mathbf{R}$ is called *regular* for a function $f: \mathbf{R} \rightarrow \mathbf{C}$ if the sequence of functions $f(t + s_m)$ converges uniformly for $t \in \mathbf{R}$. A continuous function $f: \mathbf{R} \rightarrow \mathbf{C}$ is *Bohr almost periodic* (a.p.) if every sequence $s_m \in \mathbf{R}$ contains a subsequence s_m' which is regular for f . Let

$$(1) \quad f_n \geq f_{n+1} \geq f$$

be a decreasing sequence of real valued a.p. functions converging pointwise on R to an a.p. limit f . By the definition of almost periodicity and by the diagonal process, every sequence $s_m \in R$ contains a sequence s_m' which is regular for all the functions $f, f_1, f_2, \dots$. Define

$$(2) \quad f_{s'}(t) = \lim_{m \rightarrow \infty} f(t + s_m'),$$

and

$$(3) \quad f_{n,s'}(t) = \lim_{m \rightarrow \infty} f_n(t + s_m').$$

Then also, since the convergence in (3) is uniform,

$$(4) \quad f_n(t) = \lim_{m \rightarrow \infty} f_{n,s'}(t - s_m').$$

Equations (2) and (3) applied to (1) yield

$$f_{n,s'} \geq f_{n+1,s'} \geq f_{s'},$$

and hence

$$(5) \quad f_{n,s'} \geq \Phi_{s'} \geq f_{s'},$$

where $\Phi_{s'}$ is the pointwise limit of $f_{n,s'}$ as n tends to infinity. Amerio's Condition is to assume that $\Phi_{s'}$ is a. p. for every sequence s' which is regular for all the functions $f, f_1, f_2, \dots$.

AMERIO'S THEOREM. *If f_n is a decreasing sequence of a. p. functions which converges pointwise on R to an a. p. limit f , and if $\Phi_{s'}$ is a. p. for every sequence s' in R which is regular for all the functions $f, f_1, f_2, \dots$, then the convergence is uniform.*

PROOF. As in the discussion above, let s_m be an arbitrary sequence or real numbers and let s_m' be a subsequence which is regular for all the functions $f, f_1, f_2, \dots$. Since $\Phi_{s'}$ is a.p. there exists a subsequence $-s_m''$ (of $-s_m'$) which is regular for $\Phi_{s'}$. Of course s_m'' remains regular for all the functions $f, f_1, f_2, \dots$. Define

$$(6) \quad F(t) \equiv \lim_{m \rightarrow \infty} \Phi_{s'}(t - s_m'').$$

Then, since this convergence is uniform, we also have

$$(7) \quad \Phi_{s'}(t) = \lim_{m \rightarrow \infty} F(t + s_m'').$$

Now (4) and (6) applied to the first part of (5) yields $f_n \geq F$, which

entails $f \geq F$. This last inequality along with (2) and (7) gives $f_{s'} \geq \Phi_{s'}$. But this inequality with the last part of (5) yields $f_{s'} = \Phi_{s'}$, or

$$\lim_{m \rightarrow \infty} f(t + s_m') = \lim_{n \rightarrow \infty} \lim_{m \rightarrow \infty} f_n(t + s_m').$$

In particular (for $t = 0$) we may state that: Every sequence s_m in $\mathbf{R}$ contains a subsequence s_m' such that

$$\lim_{m \rightarrow \infty} f(s_m') = \lim_{n \rightarrow \infty} \lim_{m \rightarrow \infty} f_n(s_m').$$

But this last statement is equivalent to the weak convergence of f_n to f in the Banach space A of a.p. functions on $\mathbf{R}$. (cf. [6], Chapter IV, 13.41, pp. 345-346.) The Abstract Dini Theorem now applies to conclude the proof of Amerio's theorem.

This method of proof of Amerio's Theorem suggests that there may be other Dini-type theorems for a. p. functions. All that is needed is a condition which, together with pointwise convergence and monotonicity, entails weak convergence.

Asymptotic a. p. sequences. Some years ago, while a Research-Instructor at Duke University, we found that the following condition is both necessary and sufficient for the uniform convergence on $\mathbf{R}$ of a pointwise-convergent monotone sequence of a. p. functions f_n . Call a sequence of functions $f_n: \mathbf{R} \rightarrow C$ *asymptotically almost periodic* (a.a.p.) if for each positive real number ϵ there exists an integer $N \geq 1$ such that all the functions f_n with $n \geq N$ have a relatively dense set of ϵ -translation numbers in common.

THEOREM 1. *If a sequence of functions f_n converges uniformly on $\mathbf{R}$ to a continuous function f , then a necessary and sufficient condition that f be a. p. is that f_n be a.a.p.*

PROOF OF NECESSITY. Let $\epsilon > 0$ be given. Since the convergence is uniform, there exists an integer N such that $\|f_n - f\|_\infty \leq \epsilon/3$ provided only that $n \geq N$. Since f is a. p. it possesses a relatively dense set of $\epsilon/3$ -translation numbers τ . Hence for these τ it follows from the inequality $|f_n(t + \tau) - f_n(t)| \leq |f_n(t + \tau) - f(t + \tau)| + |f(t + \tau) - f(t)| + |f(t) - f_n(t)|$ that $|f_n(t + \tau) - f_n(t)| \leq \epsilon$ for all t in $\mathbf{R}$, provided only that $n \geq N$.

PROOF OF SUFFICIENCY. Let $\epsilon > 0$ be given. Then there exists an integer $N \geq 1$ and a relatively dense set T of $\epsilon/3$ -translation numbers τ common to all f_n with $n \geq N$. Also there exists an integer $M \geq 1$ such that $\|f - f_n\|_\infty \leq \epsilon/3$ whenever $n > M$. Therefore for $\tau \in T$ and

for $n > \max\{M, N\}$ it follows from the inequality

$$|f(t + \tau) - f(t)| \leq |f(t + \tau) - f_n(t + \tau)| + |f_n(t + \tau) - f_n(t)| \\ + |f_n(t) - f(t)|,$$

that $|f(t + \tau) - f(t)| \leq \epsilon$ for all t in $\mathbf{R}$. Thus f is Bohr a. p. on $\mathbf{R}$.

COROLLARY. *In order that a sequence of a. p. functions f_n converge uniformly on $\mathbf{R}$ it is necessary that f_n be a. a. p.*

PROOF. The uniform limit of a sequence of a. p. functions is itself a. p., so the corollary follows from the necessity part of Theorem 1.

THEOREM 2. *If f_n is a decreasing sequence of continuous functions which converges pointwise on $\mathbf{R}$ to a continuous limit f , and if the sequence f_n is a. a. p., then the convergence is uniform.* (It then follows from Theorem 1 that the limit f is a. p.)

PROOF. Let $\epsilon > 0$ be given. Then there exists an integer N and a number $L > 0$ such that for each $t \in \mathbf{R}$ there exists an $\epsilon/3$ -translation number τ_t common to all the functions f_n for $n \geq N$ and satisfying $0 \leq t + \tau_t \leq L$. Furthermore, since the convergence of f_n to f is uniform on $[0, L]$ (by the classical Dini Theorem) there exists an integer M such that for $m, n > M$, $|f_n(x) - f_m(x)| \leq \epsilon/3$ for all x in $[0, L]$. Therefore, if m and n are greater than $\max\{M, N\}$ the inequality

$$|f_n(t) - f_m(t)| \leq |f_n(t) - f_n(t + \tau_t)| + |f_n(t + \tau_t) - f_m(t + \tau_t)| \\ - f_m(t + \tau_t)| + |f_m(t + \tau_t) - f_m(t)|$$

yields $\|f_n - f_m\|_\infty \leq \epsilon$. That is, the convergence is uniform.

A Counterexample. For each real λ , the mapping $f \rightarrow \hat{f}(\lambda) \equiv M_t[f(t)e^{-i\lambda t}]$ is a continuous linear functional on A . However, not all continuous linear functionals on A can be represented in this manner. Nevertheless, one might ask whether f_n converges to zero weakly if both

- (1) $\lim_{n \rightarrow \infty} \hat{f}_n(\lambda) = 0$ for all $\lambda \in \mathbf{R}$, and
- (2) $f_n(t)$ decreases to zero (as $n \rightarrow \infty$) for all $t \in \mathbf{R}$.

The following example shows that it need not. Let $\phi(t)$ denote any nonnegative continuous function on $\mathbf{R}$ whose support is contained in $[0, 1]$, and which is not identically zero. Define

$$f_n(t) = \sum_{k \in \mathbf{Z}} \phi(t - 3^nk - \frac{1}{2}(3^n - 1)),$$

for $n \geq 1$. Then each f_n is continuous and periodic with period 3^n , the sequence f_n decreases to zero pointwise on $\mathbf{R}$, but obviously does

not converge to zero uniformly on $\mathbf{R}$ (although, of course, it does converge to zero uniformly on compact subsets of $\mathbf{R}$). We now calculate $\hat{f}_n(\lambda)$ for arbitrary real λ (and for $n \geq 1$) in order to show that (1) holds. Since f_n has period 3^n , it follows that $\hat{f}_n(\lambda) = 0$ unless $\lambda = 2\pi m 3^{-n}$, for some integer m ; and in this case a calculation yields

$$\hat{f}_n(\lambda) = 3^{-n} \exp(i\lambda/2) \cos m\pi \int_0^1 \phi(s) e^{-i\lambda s} ds.$$

Now fix m and p and let $\lambda = 2\pi m 3^{-p} = 2\pi m 3^{n-p} 3^{-n}$ for $n \geq p$. Then

$$|\hat{f}_n(\lambda)| \leq 3^{-n} \int_0^1 |\phi(s)| ds,$$

for all $n \geq p$ so that (1) follows.

Another Counterexample. Not only does monotone pointwise convergence of a sequence of a. p. functions to an a. p. limit not imply uniform convergence, it does not even imply convergence in the mean. That is, there exists a sequence g_n of nonnegative Bohr a. p. functions such that g_n decreases pointwise to zero on $\mathbf{R}$, but the sequence $M(g_n)$ of their mean-values does not converge to zero. In other words, the mean-value $f \rightarrow M(f)$ is not an abstract Daniell integral on the vector lattice of real valued a. p. functions. Let λ_n be a sequence of real numbers ≥ 1 such that the set $\{1/\lambda_1, 1/\lambda_2, \dots\}$ is linearly independent over the rationals. Define f_n to have period $p_n = 2^{n+1}n\lambda_n$ and define f_n in the interval $[0, p_n]$ by the equations

$$\begin{aligned} f_n(t) &= t/2n\lambda_n, & \text{for } 0 \leq t \leq 2n\lambda_n, \\ f_n(t) &= 1, & \text{for } 2n\lambda_n \leq t \leq (2^n - 1)2n\lambda_n, \text{ and} \\ f_n(t) &= (2^{n+1}n\lambda_n - t)/2n\lambda_n, & \text{for } (2^n - 1)2n\lambda_n \leq t \leq 2^{n+1}n\lambda_n. \end{aligned}$$

Then $0 \leq f_n(t) \leq 1$, f_n has Fourier exponents of the form $2\pi k/2^{n+1}n\lambda_n$, and

$$(8) \quad 0 \leq f_n(t) \leq \frac{1}{2}, \quad \text{for all } |t| \leq n \leq n\lambda_n.$$

Also,

$$M(f_n) = \lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T f_n(s) ds = \frac{1}{p_n} \int_0^{p_n} f_n(s) ds = (1 - 2^{-n}).$$

Now define $g_n(t) = f_1(t) \cdot f_2(t) \cdots f_n(t)$. Then each g_n is a. p., $0 \leq g_{n+1} \leq g_n \leq 1$, and (8) implies that g_n decreases pointwise to zero on $\mathbf{R}$. But it follows from the Multiplication Theorem for almost

periodic Bohr-Fourier series and from the linear independence of the numbers $\{1/\lambda_n: n \geq 1\}$ that

$$\begin{aligned} M(g_n) &= M(f_1) \cdot M(f_2) \cdots M(f_n) \\ &= \prod_{k=1}^n (1 - 2^{-k}) \geq e^{-2} > 0. \end{aligned}$$

Some questions. What are the (sub) vector-lattices of real-valued a.p. functions in which the mean-value $M(f)$ is an abstract Daniell integral; that is, in which $f_n \downarrow 0$ implies $M(F_n) \rightarrow 0$? Each subspace consisting of all periodic functions of a given fixed period is such a vector lattice (by the classical Dini Theorem), but the last example above shows the entire space of all Bohr a. p. functions is not such a vector lattice.

The author believes, but has been unable to prove, that the notion of "asymptotic almost periodic sequence" is distinct from the notion of "uniformly almost periodic sequence". Is every a. a. p. sequence of a. p. functions eventually uniformly almost periodic? That is, given an a. a. p. sequence of a. p. functions f_k , does there exist (in every case) an integer N (independent of ϵ) such that for every $\epsilon > 0$ the functions $f_N, f_{N+1}, \dots$ have a relatively dense set of ϵ -translation numbers in common? We think not.

REFERENCES

1. L. Amerio, *Quasi-periodicità degli integrali ed energia limitata dell' equazione delle onde con termine noto quasi-periodico*, I, II, III, Rend. Acc. Naz. Lincei **28** (1960), February, March, April.
2. L. Amerio and G. Prouse, *Almost-Periodic Functions and Functional Equations*, Van Nostrand Reinhold Company, New York, 1971, page 61.
3. S. Bochner, *Uniform Convergences of Monotone Sequences of Functions*, Proceedings of the National Academy of Sciences **47** (1961), 582-585.
4. U. Dini, *Fondamenti per la teorica delle funzioni di variabili reali*. Pisa, 1878, pages 110-112.
5. M. Dolcher, *Su un criterio di convergenza uniforme per la successioni monotone di funzioni quasi-periodiche*, Rend. Sem. Mat. Padova **34** (1964), MR 29 #3828.
6. N. Dunford and J. T. Schwartz, *Linear Operators, Part I*, Interscience Pub., New York, 1958.
7. W. Greve, *Dini's theorem for almost periodic functions*, J. Australian Mathematical Society, Vol. II — Part 2 (1961), 143-146.
8. M. G. Krein, *Propriétés fondamentales des ensembles coniques normaux dans l'espace de Banach*, Doklady Akad. Nauk. SSSR **28** (1940), 13-17.
9. Isaac Namioka, *Partially ordered linear topological spaces*, Memoirs of the American Mathematical Society **24** (1957).

10. A. L. Peressini, *Ordered topological vector spaces*, Harper & Row Publishers, New York, 1967. MR 37 #3315.
11. W. Rudin, *Functional Analysis*, McGraw-Hill, New York, 1973.

UNIVERSITY OF NEBRASKA, LINCOLN, NEBRASKA 68508